

New Class of Squalene-based releasable nanoassemblies of paclitaxel, podophyllotoxin, camptothecin and epothilone A


Stella Borrelli,^a Michael S. Christodoulou,^a Ilaria Ficarra,^a Alessandra Silvani,^a Graziella Cappelletti,^b Daniele Cartelli,^b Giovanna Damia,^c Francesca Ricci,^c Massimo Zucchetti,^c Franco Dosio,^{d*} Daniele Passarella^{a*}

Supporting Information


Table of Contents

- ¹H-NMR and ¹³C-NMR spectra of 12a;
- ¹H-NMR and ¹³C-NMR spectra of 12b;
- ¹H-NMR and ¹³C-NMR spectra of 13a;
- ¹H-NMR and ¹³C-NMR spectra of 13b;
- ¹H-NMR and ¹³C-NMR spectra of 14a;
- ¹H-NMR of 14b;
- ¹H-NMR and ¹³C-NMR spectra of 15a;
- ¹H-NMR and ¹³C-NMR spectra of 15b.



$^1\text{H-NMR}$ and $^{13}\text{C-NMR}$ spectra of 12a






$^1\text{H-NMR}$ and $^{13}\text{C-NMR}$ spectra of 12b






$^1\text{H-NMR}$ and $^{13}\text{C-NMR}$ spectra of 13a




^1H -NMR and ^{13}C -NMR spectra of 13b





¹H-NMR and ¹³C-NMR spectra of 14a



¹H-NMR of 14b



$^1\text{H-NMR}$ and $^{13}\text{C-NMR}$ spectra of 15a



$^1\text{H-NMR}$ and $^{13}\text{C-NMR}$ spectra of 15b

