



# UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on:

T. S. Rocha, C. Tramuta, C. Catania, A. Matucci, M. G. Giuffrida, C. Baro, M.  
Profiti, L. Bertolotti, S. Rosati  
Cloning, expression, and antigenic characterization of recombinant protein of  
Mycoplasma gallisepticum expressed in Escherichia coli.  
POULTRY SCIENCE (9999)  
DOI: 10.3382/ps/pev012

The definitive version is available at:

<http://ps.oxfordjournals.org/cgi/doi/10.3382/ps/pev012>

## **Cloning, expression, and antigenic characterization of recombinant protein of *Mycoplasma gallisepticum* expressed in *Escherichia coli***

T. S. Rocha,<sup>\*,1</sup> C. Tramuta,<sup>\*</sup> S. Catania,<sup>†</sup> A. Matucci,<sup>†</sup> M. G. Giuffrida,<sup>‡</sup> C. Baro,<sup>‡</sup> M. Profiti,<sup>\*</sup> L. Bertolotti,<sup>\*</sup> and S. Rosati<sup>\*</sup>

<sup>\*</sup>University of Turin (UNITO), Department of Veterinary Science. 10100 Torino, Italy;

<sup>†</sup>Istituto Zooprofilattico Sperimentale delle Venezie, Diagnostic Service Avian Medicine Laboratory–Mycoplasma Unit, 35020 Legnaro, Padova, Italy; and

<sup>‡</sup>Istituto di Scienze delle Produzioni Alimentari–Consiglio Nazionale delle Ricerche (ISPA–CNR), Bioindustry Park S. Fumero, 10100 Giacosa, Torino, Italy

<sup>1</sup>Corresponding author: ticrocha@yahoo.com.br

### **ABSTRACT**

*Mycoplasma gallisepticum* (MG) is a member of the most important avian mycoplasmas, causing chronic respiratory disease in chickens and leading to important economic losses in the poultry industry. Recombinant technology represents a strategic approach used to achieve highly reliable and specific diagnostic tests in veterinary diseases control: in particular this aspect is crucial for confirming mycoplasma infection and for maintaining mycoplasma-free breeder flocks. In this study, we identified a component of the pyruvate dehydrogenase dihydrolipoamide acetyltransferase (i.e., E2) protein by 2-dimensional electrophoresis (2-DE), characterized it in immunoblotting assays, and analyzed its recombinant (r-E2) in a rec-ELISA test. For full-length protein expression in *Escherichia coli* (EC) a point mutation was introduced. A rabbit antiserum produced against r-E2 was tested in a Western Blot using different samples of *Mycoplasma* species. The results showed the applicability of site directed mutagenesis, with a good yield of the r-E2 after purification. Also, anti-E2 serum reacted with all the tested MG strains showing no cross reaction with other mycoplasmas. The developed E2 ELISA test was capable of detecting MG antibodies in the sera examined. Those results demonstrate the antigenic stability of the E2 protein which could represent a recombinant antigen with potential diagnostic applications.

Key words: *Mycoplasma gallisepticum*, recombinant protein, E2

### **INTRODUCTION**

*Mycoplasma gallisepticum* (MG) is an important pathogen of poultry worldwide that represents considerable economic losses to the poultry industry, causing chronic respiratory disease in chickens and turkeys, reduction in egg production and quality, poor hatchability and feed efficiency, and increase in mortality and carcass condemnations besides the medication costs. (Ley et al., 2009; Ghorashi et al., 2010).

The genus *Mycoplasma* (class Mollicutes) has evolved by degenerative or reductive evolution, accompanied by significant loss of genomic sequences and progressive adaptation to their specific hosts. They are distinguishable from other bacteria by their minute size and lack of a cell wall, and because of that, the surface proteins (lipoproteins) anchored or embedded in the cell membrane play a crucial role in the interactions between the mycoplasmas and their hosts (Benčina et al., 1994). In the absence of a cell wall, the surface of the mycoplasma membrane constitutes a critical interface in the infectious process, mediating basic functions such as the transport of nutrients as well as more complex interactions with the host cells and the host immune defenses.

The chronic nature of *Mycoplasma* infections demonstrates a failure of the host immune system to deal effectively with these organisms. The well-documented antigenic variation of surface proteins is a major contributor to the complexity in implementing efficient preventive control measures and accurate diagnostic tests. It also allows MG to evade the host's immune response through the generation of escape variants (Glew et al., 2000; Hudson et al., 2006; Citti et al., 2010) and may contribute to resistance versus antimicrobial therapy by intracellular invasion and survival within eukaryotic cells (Winner et al., 2000).

The knowledge derived from studies of MG proteins undoubtedly has considerable potential for diagnostic application. Since serological monitoring of poultry is used extensively as a first-line diagnostic test and will continue to be performed on a large scale in the future, there will be an increasing need for well-defined and characterized antigens to enable the development of efficacious and cost-effective control

measures (Benčina et al., 1994). The currently used tests, such as the commercial rapid serum agglutination (RSA) and ELISA, as well as hemagglutination inhibition (HI), have greatly contributed to the control of *Mycoplasma* infections, but previous work reported a lack of specificity, mainly due to the intrinsic ability of various *Mycoplasma* species to cross-react (Büyüktanır et al., 2010).

As a consequence, recombinant technology could significantly help to alleviate drawbacks, allowing the production of unlimited amounts of multiple and more specific antigens (Ben Abdelmoumen Mardassi et al., 2008) although identification of immunogenic antigens and genetic manipulations of *Mycoplasma* in general are more laborious than any other prokaryotic genome as far as to successfully express MG proteins in heterologous systems such as *Escherichia coli* (EC), the TGA codons that encode tryptophan in MG genes and, at the same time, are used in EC as a stop codons, need to be replaced by the TGG codon, unless the protein would be produced in a truncated form (Rosati et al., 2000).

Numerous studies have focused on immunogenic proteins of MG, particularly those involved in hemagglutination, haemadsorption, and cytoadhesion (Milosevic Berlic et al., 2000). Among specific antigens characterized so far, researchers have identified a large cell surface-exposed lipoprotein (i.e., pMGA) involved in haemagglutination and also pvpA, a putative variable adhesin. Both proteins present polymorphisms and size variation in different MG strains, as well as a phase variable expression (Markham et al., 1992; Boguslavsky et al., 2000; Liu et al., 2001). Considering the limited size of *Mycoplasma* genomes, the number of genes involved in generating antigenic diversity on their cell surface is surprisingly large (Razin et al., 1998). In previous works, a surface protein called P52 was described and found to be similar to the dihydrolipoamide acetyltransferase (i.e., E2) from several mollicutes (Jan et al., 1996, 2001). This protein is a membrane-associated component of the pyruvate dehydrogenase complex which has been shown to be required for MG *in vivo* growth and survival in the host. Also, antibodies directed against P52 induced growth inhibition of cells and activation of the complement system. Partial characterization of P52 protein was limited to amino terminal peptide sequencing and a full antigenic characterization may be required to evaluate its diagnostic potential. The availability of the full genome sequence of MG allows the application of an immune proteome-based strategy to identify potential antigenic sites, useful to further improve current diagnostic tests.

The aim of the present study was to detect, identify, and characterize novel MG immunogenic proteins. For that, the proteome of MG was resolved on 2-dimensional electrophoresis (2-DE), blotted on membrane, and probed against monospecific sera. Along

with the known haemagglutinin precursor, a second immunostained protein was identified by matrix-assisted laser desorption/ionization time-of-flight/time-of-flight (MALDI-TOF/TOF) mass spectrometry as E2 protein. The corresponding gene was amplified, mutagenized, and cloned in a prokaryotic expression vector. Preliminary antigenic characterization

of recombinant E2 (r-E2) suggests that this protein is well-conserved among and specific within MG isolates, and may be considered for the development of a specific diagnostic assay.

## **MATERIALS AND METHODS**

### ***Two-Dimensional Gel Electrophoresis of Fractionated Mycoplasma Cell Proteins***

*Mycoplasma gallisepticum* American Type Culture Collection (ATCC) Strain 15302, cultured according to Kleven et al. (1998) was centrifuged at 6,000 rpm for 10min at 4°C and the pellet was washed in a 1X phosphate-buffered saline (PBS) solution 3 times as described (Regula et al., 2000). One milliliter preheated sodium dodecyl sulfate (SDS) 1.0% and 10 µL phenyl methylsulfonyl fluoride (PMSF), 0.1M, were added to the bacterial pellet and boiled for 3min. The sample was sonicated 3 times in ice and centrifuged at 10,000 rpm for 10min.

Supernatant was collected and the protein concentration was determined by using a DC protein assay quantification kit (Biorad, Hercules, CA) and purified using the 2D clean up kit (GE Healthcare, Little Chalfont, UK).

The sample was resuspended in a lysis buffer containing 8 M urea, 2 M thiourea, 2.5% 3-[(3-cholamidopropyl)dimethylammonio]-1 propanesulfonate (CHAPS), 1.0% IPG buffer and 20 mM dithiothreitol (DTT, GE Healthcare, Little Chalfont, UK). Strips over the pH range of 4 to 7 (11 cm, Biorad) were subjected to isoelectric focusing (IEF) using the following parameters: rehydration at 50V for 12h; running at 500V for 1h, 1,000V for 1h, 8,000V for 1h, 1,000 to 8,000V for 4h, and 500 V for 2 h. After isoelectric focusing the gel strips were equilibrated 2× in a buffer containing 7 M urea, 500mM Tris-HCl pH 6.8, 2% SDS, and 10% glycerol. In the first equilibration step 2% DTT was added, and in the second 2.5% iodoacetamide, and incubation was performed with 15 min each. Immediately following equilibration, the IEF strips were transferred onto a 10% SDS-polyacrylamide gel. Electrophoresis was carried out at 170 V for 1 h.

All 2D gel electrophoresis experiments were run in triplicate. On 2 gels, 200 µg proteins were loaded and visualized by Coomassie Blue staining; the third gel was used to run 50 µg proteins and visualized by silver staining. All gel stains were made with MALDI-compatible colorants. Proteins from one of the Coomassie gels were electro transferred to nitrocellulose membranes for Western immunoblot analyses using a monospecific MG anti serum (Biovac, Beaucazoué, France) diluted at 1/100. The secondary antibody used was anti-chicken IgY, HRP conjugate (Sigma-Aldrich, Milano, Italy) diluted at 1/1,000 (Bencina et al., 1994). After the immunoblot, the Coomassie gel and the blotted membrane were compared in order to detect immunoreactive proteins.

### ***Protein Identification***

The spots selected by immunochemical analyses were subjected to MALDI-TOF/TOF mass spectrometry for identification using an Ultraflex III MALDI TOF/TOF instrument (Bruker Daltonics, Bremen). The spots were cut out of the 2-DE, destained overnight with a 4.2 g/L solution of ammonium bicarbonate and 400 mL/L ethanol, and then washed 3 times for 10 min with acetonitrile and dried in a Speedvac. Proteins were in gel digested with trypsin (Promega, Madison, WI), and spectra were acquired as previously described (Zava et al., 2009). Mascot and MS-Fit software packages were used to search against the NCBI nr.2011.10.10 database using the peptide-mass fingerprinting (PMF) method (Pappin et al., 1993). The following parameters were set for the searches: S-carbamidomethyl derivative on cysteine as fixed modification, oxidation on methionine as variable modification, and 2 missed cleavage sites for trypsin digestion. The peptide mass tolerance was 20 ppm. For in silico analysis, the Basic Local Alignment Search Tool (BLAST) program (Altschul et al., 1990) was used for nucleotide and protein homology search, as far as for the detection of tryptophan codons.

### ***Site-directed Polymerase Chain Reaction-Mediated Mutagenesis***

*Mycoplasma gallisepticum* American Type Culture Collection (ATCC) Strain 15302 was used for genomic DNA extraction.

A single tryptophan codon UGA was detected in the whole E2 gene and point-mutated to convert into UGG, using two polymerase chain reaction (PCR) runs as described (Villaruel and Regalado, 1997; Rosati et al., 2000). Briefly, in the first run, the following primer sets (mutated positions are boxed) were used to generate 2 overlapping fragments [offset by 30 base pair (bp)]: MGfow 5' TTGGATCCTTTGAATATAAATTTACAGACGTTG and MGmut- rev 5' -TTATCAGCACCATCGAT CA TCTGTGGTCAGCTGC for the amplification of the first fragment and MGmut- fow 5' -TTGCAGC TGACCACAGAT GATCGATGGTGCTGAT and MGrev 5' -TTGAATTCAATTAGTAAACCGTTAAGA TTTTC for the second fragment. The MGfow and MGrev primers contained restriction sites for BamHI and EcoRI, respectively (underlined in the primer sequences), to facilitate the following cloning step.

Forward MGfow primer was designed excluding the protein leader peptide sequence. The first PCR reaction (50  $\mu$ L volume) contained 10 mM Tris-HCl, pH 8.8, 50 mM KCl, 1.5 mM MgCl<sub>2</sub>, 200  $\mu$ M deoxynucleotide triphosphates (dNTPs), 10 pmol each primers, 50 ng of template, 1 U Hotstar Taq DNA polymerase (Qiagen, Venlo, Netherlands). PCR was performed in 7 min at 95°C followed by 30 cycles of 60 s at 95°C, 60 s at 50°C, and 60 s at 72°C, and a final extension of 7 min at 72°C. Amplified products were resolved and visualized after agarose gel electrophoresis and ethidium bromide staining, gel purified, and combined in equimolar concentrations to form the template for the second PCR run, using the most external primers. During the annealing step, overlapping fragments joined each other accomplishing the full length synthesis of the gene encoding the mature protein. The second reaction mixture (50  $\mu$ L volume) contained the same buffer as above, 200  $\mu$ M dNTPs, 50 pmol each primer, and 1 U Taq polymerase. Amplification was performed as described above in 40 cycles.

#### ***Cloning and Sequencing of Mutated Genes***

Amplified product of the expected length was digested with appropriate restriction enzymes (Thermo Scientific, Waltham, MA), purified, and cloned into pSER-6HIS expression vector, which allows expression of the recombinant protein fused to a 6 $\times$  His tag at the N-terminus. The recombinant plasmid was used to transform *Escherichia coli* BL21 (i.e., DE3) competent cells, plated on Luria-Bertani (LB) agar with ampicillin (50  $\mu$ g/mL) and screened by colony PCR, using primers specific for the pSER vector (Robino et al., 2005). Positive colonies were grown in 2 mL LB with ampicillin (50  $\mu$ g/mL). Plasmids were extracted using QIAprep Spin Miniprep Kit (Qiagen, Venlo, Netherlands) and sequenced in order to confirm the induced point mutation (BMR Genomics, Padova, Italy).

#### ***Expression and Purification of Recombinant Protein***

To express the recombinant protein, early log phase culture [optical density (OD)<sub>600</sub> = 0.6] of a positive clone was induced for 2 h with 0.1 mM isopropyl  $\beta$ -thiogalactopyranoside (IPTG) under agitation. Bacterial cells were recovered by centrifugation, washed 1 $\times$  with PBS, and lysed by conventional physico-chemical methods. Soluble fractions were separated by centrifugation and sequentially extracted in a buffer containing 20 mM sodium phosphate pH 7.4, 0.5 M NaCl, 10 mM imidazole, and increasing urea concentrations (urea 0 M : S0 fraction; urea 1 M: S1 fraction; urea 4 M: S2 fraction; urea 8 M: S3 fraction).

Fraction analysis, carried out by SDS-PAGE, revealed that recombinant protein was located in the S3 fraction. Purification was conducted by immobilized metal chelate affinity chromatography using an high performance liquid chromatography (HPLC) system. (GE Healthcare, Little Chalfont, UK) under denaturing condition (Smith and Johnson, 1988). Purity of each fraction of eluted protein was analyzed by SDS-PAGE and protein concentration estimated by a DC protein assay (BioRad, Hercules, CA).

#### ***Western Blot Analysis***

To confirm the presence and the molecular mass of the recombinant proteins, Western blotting was carried out using an anti-6× His monoclonal antibody as probe. Purified recombinant proteins were separated by 10% SDS-PAGE, transferred to a nitrocellulose membrane, and subsequently blocked with 5% non fat powdered milk in PBS for 2h. After washing with PBS-Tween 0.05%, the membrane was incubated with anti-6× His monoclonal antibody (GE Healthcare, Little Chalfont, UK) for 1 h at room temperature. After a second washing step, the membrane was incubated with horseradish peroxidase (HRP)-conjugate anti-mouse IgG (KPL Laboratories, Gaithersburg, UMDSA) for 1 h at room temperature. Immunoreactive protein bands were stained with 10 mg 3,3'-diaminobenzidine tetrahydrochloride (DAB, Sigma-Aldrich, Milano, Italy).

To evaluate the expression, antigenic stability, and specificity of the corresponding native protein, a specific rabbit antiserum was elicited using the purified recombinant antigen E2 as an immunogen (Khiari and Mardassi, 2012). The rabbit serum was probed against whole lysate of 7 different field MG strains as well as different avian mycoplasmas [2 *Mycoplasma synoviae* (MS) and 1 *Mycoplasma iowae* (MI) samples]. All samples are field isolates that were identified and characterized by denaturing gradient gel electrophoresis (DGGE) PCR in our laboratory.

#### ***Serum Samples and ELISA Procedure***

For the evaluation of the developed ELISA protocol 12 serum samples were collected from naturally MG infected chickens flocks (confirmed by PCR and bacterial isolation) and 32 from specific-pathogen-free (SPF) subjects. ELISA was carried out as previously described with minor modifications (Rosati et al., 2000), using a commercially available monospecific MG antiserum (Biovac, Beaucauzé, France) as positive control and a SPF serum as negative control.

Briefly, microplate was coated with 100 ng/well of recombinant antigen. Serum samples, diluted 1/10 were incubated, in duplicate, for 1h at room temperature. After 3 washing steps, 100 of anti-chicken IgY (Sigma Aldrich, Milano, Italy) was added to each well and plate incubated as above. Following the final washing steps, reactions were developed with 2,20-azino-bis-[3 ethylbenzthiazoline-6-sulfonic acid (ABTS, Chemicon, Temecuta, CA)] and read at 405 nm. Absorbance was

expressed as percentage of reactivity of the positive control serum. Sera with reactivity > 40% of the positive control serum were classified as positive. Differences between absorbance in negative and positive sera were tested by nonparametric statistics (Wilcoxon 2-sample test) using R software (R Core Team, 2013).

## **RESULTS**

### ***Two-Dimensional Gel Electrophoresis and Mass Spectrometry Analysis and Protein Identification***

Following fractionation of total cell lysate and 2-DE, a companion gel was electro blotted on membrane and immunostained with reference MG positive sera. Two major spots were recognized (Figure 1) and analyzed by MALDI-TOF/TOF. Peptide mass fingerprinting resulted in the identification of pMGA and E2 protein (a total of 3 peptides recognized in a total of 27 with a coverage of 11%). Since pMGA is already a well-known protein, with a large potential for generation of antigenic variants (Bencina, 2002), we decided to further characterize only the E2 protein.

#### ***Recombinant E2 Expression***

The complete gene of E2, lacking the most 5' terminal region corresponding to the leader peptide, was then retrieved from the database Genbank amplified in 2 different PCR runs to accomplish site-directed mutagenesis to convert a single TGA codon into TGG and allow expression of full product in EC. As expected, after the second PCR run a single band of 1.3 kb was detected. The amplicon was cloned into pSER-6HIS plasmid and sequence analysis

conducted with vector primers confirmed that mutagenesis was truly accomplished and the E2 amplicon was cloned in the correct frame and orientation. When positive clone was grown and induced, a new protein band was clearly detected in SDS-PAGE with an apparent molecular weight of 49 kDa, corresponding to E2 protein. The recombinant E2 fusion protein was found to be soluble in 8M urea and purified under denatured condition.

#### ***Western Blot and ELISA Analyses***

The purified recombinant protein was immunostained in Western Blot using an anti-6× His MAb which probes the C' terminal tag of fusion protein. The purified recombinant protein was used as immunogen to elicit a specific immunoresponse in rabbit and the monospecific serum was used in a second Western Blot experiments using total cell lysates of different avian mycoplasmas. The antiserum was able to recognize the native protein with the expected molecular weight in all MG samples tested; meanwhile no reaction was observed in the MS and MI control samples (Figure 2).

Recombinant E2 was tested in ELISA at the optimal conditions described above. A panel including 12 positive and 32 negative samples was tested. Results showed a significant difference between OD recorded in the 2 groups. In more details, all positive sera tested positive in ELISA; among negative samples, 3 sera were erroneously classified as positive, showing low OD values. Differences between OD values of sera belonging to positive and negative sample sets were statistically significant as expected (Wilcoxon rank sum test  $P < 0.001$ ). A boxplot of ELISA OD values obtained by r E2 ELISA is shown in Figure 3.

#### **DISCUSSION**

The identification and characterization of novel immunogenic proteins are important for the development of both improved diagnostic assays and subunit vaccines. In order to reach new and useful results in this research field, the 2-DE is a technique that can be routinely applied for parallel quantitative expression profiling of large sets of complex protein mixtures. This technique is able to detect and characterize immunodominant proteins, and it was used to improve the knowledge about MG proteins as potential candidates for the diagnosis and control of infections (Görg et al., 2004).

The pMGA gene family is a large group of surface proteins mediating the phase variation of the bacterial surface architecture and they are believed to be involved in evasion of the host immune system (Glew et al., 2000; Bearson et al., 2003; Szczepanek et al., 2010). Metabolic pathways have also been shown to contribute to MG virulence and host-interaction and, in our study, a component of the pyruvate dehydrogenase was identified by 2-DE analysis, the dihydrolipoamide acetyltransferase (i.e., E2). This protein was first described by Jan and colleagues in 1996 and fully sequenced by Szczepanek and colleagues in 2010. E2 was proven to be highly immunogenic and antibodies directed against it induced growth inhibition of cells and activation of the complement system. Moreover, topological experiments using in situ treatment with proteases and growth inhibition in the presence of anti-E2 serum provided evidence of the surface exposition of the polypeptide (Jan et al., 1996), but to our knowledge no serological and diagnostic potential of this protein was not yet evaluated. In this study, with the aim to fully characterize the E2, we directly mutagenized its gene and successfully produced recombinant protein in a prokaryotic system. The results obtained after the Western Blot using the E2 rabbit antiserum with different strains of mycoplasmas showed that E2 protein is stably expressed in all MG tested strains isolated in different parts of Italy, while no cross reaction was found in any MS and MI strains tested. This aspect is particularly important given that MG is equally pathogenic for poultry compared to MS and MI, causing important economic losses. Under the diagnostic point of view it is fundamental to overcome the possibility of cross reactions in serological tests, especially in field samples. These results also agree with the ones found by Jan and coauthors in 2001. Indeed the anti-

E2 serum revealed a polypeptide with an apparent molecular mass close to 52 kDa in all of the MG strains assayed, demonstrating the antigenic stability of the protein and reinforcing the idea that it might be used to develop serologic diagnostic tests based on the detection of anti-E2 antibodies in infected hosts.

In the ELISA test, r-E2 was capable of detecting all the positive serum tested, even though there were some showing low OD values, slightly above the cutoff. The presence of 3 false-positive results among negative samples revealed a nonabsolute specificity of the test. It is known that many parameters are involved in stimulating the avian immune system and the obtained results can be explained by the variation of the immune response of the host, according to individual response to infection and heterogeneity of the antigen presentation by the MG (Levisohn et al., 1995; Grodio et al., 2012). Moreover, Western Blot supports the specificity of the

E2 against sera positive for other than MG mycoplasma strains. Given this, our results seem to suggest that the false positive outcomes could be justified by a non ideal ELISA test optimization. Further investigations about serum dilution and conjugate concentration will be performed in order to obtain higher signal-to-noise ratio and, consequently, better diagnostic performances.


In conclusion, the protein identified by the 2-DE analysis was successfully mutagenized demonstrating that the used method is fast and highly efficient. Over all, E2 can represent a recombinant antigen with potential diagnostic applications, being discriminative, cost effective, and less time-consuming. Moreover, E2 can improve diagnostic tools together with the well-known pMGA and pvpA antigens, increasing the spectrum of MG antibodies detected in serological procedures and representing an important update in the diagnostic of mycoplasma infection in poultry.




## REFERENCES

- Altschul, S. F., W. Gish, W. Miller, E. W. Myers, and D. J. Lipman. 1990. Basic Local Alignment Search Tool. *J Mol Biol.* 5:403–410.
- Bearson, S. M. D., S. D. Collier, B. L. Bearson, S. L. Branton, and A. S. D. Collier. 2003. Induction of a *Mycoplasma gallisepticum* pMGA gene in the chicken tracheal ring organ culture model. *Avian Dis.* 47:745–749.
- Ben Abdelmoumen Mardassi, B., A. A. B'ejoui, L. Oussaeif, B. Mlik, and F. Amouna. 2008. A recombinant antigen-based ELISA for the simultaneous differential serodiagnosis of *Mycoplasma gallisepticum*, *Mycoplasma synoviae*, and *Mycoplasma meleagridis* infections. *Avian Dis.* 52:214–221.
- Bencina, D. 2002. Haemagglutinins of pathogenic avian *Mycoplasma* spp. *Avian Pathol.* 31:535–547.
- Bencina, D., S. H. Kleven, M. G. Elfaki, A. Snoj, P. Dovč, D. Dorrer, and I. Russ. 1994. Variable expression of epitopes on the surface of *Mycoplasma gallisepticum* demonstrated with monoclonal antibodies. *Avian Pathol.* 23:19–36.
- Boguslavsky, S., D. Menaker, I. Lysnyansky, T. Liu, S. Levisohn, R. Rosengarten, and M. Garcia. 2000. Molecular characterization of the *Mycoplasma gallisepticum* pvpA gene which encodes a putative variable cytoadhesin protein. *Infect. Immun.* 68:3956–3964.
- Bu'yu'ktanir, O., O. Genc, and N. Yurdusev. 2010. Bi-antigenic immunoassay models based on the recombinant PvpA proteins for *Mycoplasma gallisepticum* diagnosis in chickens. *J. Vet. Diagn. Invest.* 22:908–913.
- Citti, C., L.-X. Nouvel, and E. Baranowski. 2010. Phase and antigenic variation in mycoplasmas. *Future Microbiol.* 5:1073–1085.
- Ghorashi, S. a, A. H. Noormohammadi, and P. F. Markham. 2010. Differentiation of *Mycoplasma gallisepticum* strains using PCR and high-resolution melting curve analysis. *Microbiology* 156:1019–1029.
- Glew, M. D., G. F. Browning, P. F. Markham, and I. D. Walker. 2000. pMGA phenotypic variation in *Mycoplasma gallisepticum* occurs in vivo and is mediated by trinucleotide repeat length variation. *Infect. Immun.* 68:6027–6033.
- Go'rg, A., W. Weiss, and M. J. Dunn. 2004. Current two-dimensional electrophoresis technology for proteomics. *Proteomics* 4:3665–3685.
- Grodio, J. L., D. M. Hawley, E. E. Osnas, D. H. Ley, K. V. Dhondt, A. A. Dhondt, and K. A. Schat. 2012. Pathogenicity and immunogenicity of three *Mycoplasma gallisepticum* isolates in house finches (*Carpodacus mexicanus*). *Vet. Microbiol.* 155:53–61.
- Hudson, P., T. S. Gorton, L. Papazisi, K. Cecchini, S. Frasca, and S. J. Geary. 2006. Identification of a virulence-associated determinant, dihydrolipoamide dehydrogenase (lpd), in *Mycoplasma gallisepticum* through in vivo screening of transposon mutants. *Infect. Immun.* 74:931–939.
- Jan, G., C. Brenner, and H. Wroblewski. 1996. Purification of *Mycoplasma gallisepticum* membrane proteins p52, p67 (pMGA), and p77 by high-performance liquid chromatography. *Protein Expr. Purif.* 7:160–166.
- Jan, G., M. Le Henaff, C. Fontenelle, and H. Wroblewski. 2001. Biochemical and antigenic characterization of *Mycoplasma gallisepticum* membrane proteins P52 and P67 (pMGA). *Arch. Microbiol.* 177:81–90.
- Khiari, A. B., and B. Ben Abdelmoumen Mardassi. 2012. Characterization of the antigenic and functional domains of a *Mycoplasma synoviae* variant vIhA gene. *Vet. Microbiol.* 156:322–329.
- Kleven, S. H. 1998. Mycoplasmosis. In: D. E. Swayne, J. R. Glisson, M. W. Jackwood, J. E. Pearson, and W. M. Reed (Eds.), *A Laboratory Manual for the Isolation and Identification of Avian Pathogens*. American Association of Avian Pathologists, Kennett Square, PA, pp. 74–80.


- Levisohn, S., R. Rosengarten, and D. Yogev. 1995. In vivo variation of *Mycoplasma gallisepticum* antigen expression in experimentally infected chickens. *Vet. Microbiol.* 45:219–231.
- Ley, R. E., M. Hamady, C. Lozupone, P. Turnbaugh, R. Roy, J. S. Bircher, M. L. Schlegel, T. A. Tucker, D. Mark, R. Knight, and J. I. Gordon. 2009. Evolution of mammals and their gut microbes. *320*:1647–1651.
- Liu, T., M. García, S. Levisohn, D. Yogev, and S. H. Kleven. 2001. Molecular variability of the adhesin-encoding gene *pvpA* among *Mycoplasma gallisepticum* strains and its application in diagnosis. *J. Clin. Microbiol.* 39:1882–1888.
- Markham, P. F., M. D. Glew, M. R. Brandon, I. D. Walker, and K. G. Whithear. 1992. Characterization of a major hemagglutinin protein from *Mycoplasma gallisepticum*. *Infect. Immun.* 60:3885–3891.
- Milosevic Berlic, T., D. Bencina, and P. Dovc. 2000. Sequence polymorphisms within the pMGA genes and pMGA antigenic variants in *Mycoplasma gallisepticum*. *FEMS Microbiol. Lett.* 184:133–139.
- Pappin, D. J. C., P. Hojrup, and A. J. Bleasby. 1993. Rapid identification of proteins by peptide-mass fingerprinting. *Curr. Biol.* 3:327–332.
- R Core Team (2013). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, <http://www.R-project.org/>.
- Razin, S., D. Yogev, and Y. Naot. 1998. Molecular biology and pathogenicity of *Mycoplasmas*. *Microbiol. Mol. Biol. Rev.* 62:1094–1156.
- Regula, J. T., B. Ueberle, G. Boguth, A. Görg, M. Schnörlzer, R. Herrmann, and R. Frank. 2000. Towards a two-dimensional proteome map of *Mycoplasma pneumoniae* proteomics and 2-DE. *Electrophoresis* 21:3765–3780.
- Robino, P., A. Alberti, M. Pittau, B. Chessa, M. Miciletta, P. Nebbia, D. Le Grand, and S. Rosati. 2005. Genetic and antigenic characterization of the surface lipoprotein P48 of *Mycoplasma bovis*. *Vet. Microbiol.* 109:201–209.
- Rosati, S., P. Robino, M. Fadda, S. Pozzi, A. Mannelli, and M. Pittau. 2000. Expression and antigenic characterization of recombinant *Mycoplasma agalactiae* P48 major surface protein. *Vet. Microbiol.* 71:201–210.
- Smith, D. B., and K. S. Johnson. 1988. Single-step purification of polypeptides expressed in *Escherichia coli* as fusions with glutathione S-transferase. *Gene* 67:31–40.
- Szczepanek, S. M., E. R. Tulman, T. S. Gorton, X. Liao, Z. Lu, J. Zinski, F. Aziz, S. Frasca, G. F. Kutish, and S. J. Geary. 2010. Comparative genomic analyses of attenuated strains of *Mycoplasma gallisepticum*. *Infect. Immun.* 78:1760–1771.
- Villarroel, A., and M. P. Regalado. 1997. A fast and simple method to introduce multiple distant point mutations. *Tech. Tips Online* 2:24–26.
- Winner, F., R. Rosengarten, and C. Citti. 2000. In vitro cell invasion of *Mycoplasma gallisepticum*. *Infect. Immun.* 68:4238–4244.
- Zava, S., C. Barello, A. Pessione, L. P. Garoffo, P. Fattori, G. Montorfano, A. Conti, C. Giunta, E. Pessione, B. Berra, and M. G. Giuffrida. 2009. Mare's colostrum globules stimulate fibroblast growth in vitro: A biochemical study. *J. Med. Food* 12:836–845.



**Figure 1.** 2D gel electrophoresis and Western blot of *Mycoplasma gallisepticum* total proteins. MG proteins were separated on pH 4 to 7 linear strips and 10% acrylamide gels followed by (A) Coomassie staining, (B) silver staining (B), and (C) immunoblotting using monospecific serum. Proteins in the molecular mass standard are indicated to the right of the gels. Spot position of the E2 protein is indicated by white arrow.



**Figure 2.** Western Blot using the E2 rabbit antiserum. Lanes M: molecular weight marker; Lanes 1 and 7: r-E2 (positive control); Lanes 2 to 6, 8, and 9: whole lysate of different field *Mycoplasma gallisepticum* strains; Lanes 10 and 11: whole lysate of different *Mycoplasma synoviae* strains; Lane 12: whole lysate of *Mycoplasma iowae*; Lane 13: negative control.



**Figure 3.** Distributions of D-values in ELISA test. OD values are reported as percentage of reactivity of the positive control serum.