



# UNIVERSITÀ DEGLI STUDI DI TORINO

***This is an author version of the contribution published on:***

*Questa è la versione dell'autore dell'opera:*

*[Book of abstracts 11th Conference of the European Foundation for Plant Pathology – Healthy plants – healthy people, 2014]*

***The definitive version is available at:***

*La versione definitiva è disponibile alla URL:*

*[<http://www.kongres-fitopatologiczny.pl/kongres.php?id=book>]*

## **An integrated approach to monitor and control the invasive fungal pathogen *Heterobasidion irregulare* in European forest stands**

Paolo Gonthier<sup>1</sup>, Naldo Anselmi<sup>2</sup>, Paolo Capretti<sup>3</sup>, Filippo Bussotti<sup>3</sup>, Matteo Feducci<sup>3</sup>, Luana Giordano<sup>1</sup>, Tommaso Honorati<sup>2</sup>, Guglielmo Lione<sup>1</sup>, Nicola Luchi<sup>4</sup>, Marco Michelozzi<sup>5</sup>, Bruno Papparatti<sup>2</sup>, Martina Pollastrini<sup>3</sup>, Fabiano Sillo<sup>1</sup>, Anna Maria Vettraino<sup>2</sup>, Matteo Garbelotto<sup>6</sup>

<sup>1</sup>Department of Agricultural, Forest and Food Sciences, University of Torino, Grugliasco, ITALY; <sup>2</sup>Department for Innovation in Biological, Agro-food and Forest Systems, University of Tuscia, Viterbo, ITALY; <sup>3</sup>Department of Agri-Food Production and Environmental Sciences, University of Firenze, Firenze, ITALY; <sup>4</sup>CNR - Institute for Plant Protection, Sesto Fiorentino, ITALY; <sup>5</sup>IBBR-FI/CNR - Institute of Biosciences and Bioresources, Sesto Fiorentino, ITALY; <sup>6</sup>Department of Environmental Science, Policy and Management, University of California at Berkeley, Berkeley, USA  
[paolo.gonthier@unito.it](mailto:paolo.gonthier@unito.it)

The North American fungal pathogen *Heterobasidion irregulare* is currently distributed in pine and oak stands along 103 km of coastline of central Italy. This paper reviews the pathways of introduction and invasion, the factors driving the invasion, and the dispersal abilities of this pathogen in Italy. Furthermore, an integrated disease management program to minimize the risk of spread of the fungus in Europe is suggested, based both on published literature and on new unpublished data. Observational and genetic evidence support a single introduction through infected wood during WWII, and a subsequent invasion through spore dispersal. Experimental evidence suggests transmission potential of the pathogen rather than hyper-susceptibility of native hosts is the major determinant of invasion. The current range of *H. irregulare* is too vast to suggest eradication, however we recommend minimizing the risk of spread of *H. irregulare* outside the zone of infestation while reducing the magnitude of infestations within its current range. We provide evidence suggesting the most cost-effective management approach hinges on preventing the saprobic establishment of the fungus in stumps in a “buffer” area surrounding the current zone of infestation.