



# UNIVERSITÀ DEGLI STUDI DI TORINO

***This is an author version of the contribution published on:***

*Questa è la versione dell'autore dell'opera:*

*Ecology of Food and Nutrition, Volume 53, Issue 5, 2014,*

*doi:10.1080/03670244.2014.883976*

***The definitive version is available at:***

*La versione definitiva è disponibile alla URL:*

*<http://www.tandfonline.com/doi/full/10.1080/03670244.2014.883976>*

Randomized Controlled Trials Evaluating Effect of Television Advertising on Food Intake  
in Children: Why Such a Sensitive Topic is Lacking Top-Level Evidence?

Authors

Dario Gregori<sup>1</sup>, Simonetta Ballali<sup>2</sup>, Maria Gabriella Vecchio<sup>2</sup>, Antonella Silvia Scirè<sup>2</sup>, Francesca  
Foltran<sup>1</sup>, Paola Berchialla<sup>3</sup>

<sup>1</sup>Unit of Biostatistics, Epidemiology and Public Health, Department of Cardiac, Thoracic and  
Vascular Sciences, University of Padova, Padova, Italy

<sup>2</sup>PROCHILD Onlus, Trieste, Italy

<sup>3</sup>Department of Clinical and Biological Sciences, University of Torino, Torino, Italy

Address correspondence to

Dario Gregori,

Unit of Biostatistics, Public Health and Epidemiology,

Department of Cardiac, Thoracic and Vascular Sciences,

University of Padova,

Via Loredan, 18, 35131

Padova, Italy.

E-mail: [dario.gregori@unipd.it](mailto:dario.gregori@unipd.it)

## ABSTRACT

The aim of this study was to perform a systematic review of evidence coming from randomized controlled trials (RCT) aimed at assessing the effect of television advertising on food intake in children from 4 to 12 years old. Randomized controlled trials were searched in PubMed database and included if they assessed the effect of direct exposure to television food advertising over the actual energy intake of children. Seven studies out of 2166 fulfilled the inclusion criteria. The association between television advertising and energy intake is based on a very limited set of randomized researches lacking a solid ground of first-level evidence.

**KEYWORDS** child obesity, food consumption, food preference, snacks, TV advertising

## INTRODUCTION

In recent years, obesity has reached epidemic status, affecting not only developed country but also developing ones (WHO 2000). Obesity levels don't show an homogeneous distribution, ranging from below 5% in China, Japan and certain African nations, to over 75% in urban Samoa, and even within the same countries, a high variance can be seen, as for example in China, where figures reach almost 20% in some cities.

Where national data are available, adiposity has increased in both preschool and school-aged children in nearly every country studied (von Kries et al. 2012). However, large variations in secular trends do appear across countries, and these relate to the degree of economic development and urbanisation (Wang and Lobstein 2006). Childhood obesity and overweight are at the roots of adult diseases development, including type 2 diabetes, fatty liver disease, and endocrine and orthopaedic disorder (Lobstein, Baur, and Uauy 2004).

The complexity underlying obesity has been often simplified to a dichotomy between energy intake and energy expenditure (Crowley, Yeo, and O'Rahilly 2002), considering unregulated food consumption as the main responsible of increased energy intake. Easy access to energy-rich foods

and drinks, food marketing through television and other media, increased use of video entertainment and computer games playing, have all been suggested as promoting the “toxic” or “obesogenic” environment (Ebbeling, Pawlak, and Ludwig 2002; Egger and Swinburn 1997; Kumanyika et al. 2002).

Television is part of worldwide home setting; therefore TV-related sedentary time has been investigated in order to assess its relation with obesity development. Results have not being homogeneous, ranging from proven association (Lobstein et al. 2004) to no influence at all on energy intake (Jackson et al. 2009). Television is suspected to be linked to a reduction in physical activity whilst advertising seems to promote an overconsumption of food high in fat and sugar (Caroli et al. 2004; Carter et al. 2011). Most of the studies focusing on this subject have been conducted in the Anglo-Saxon context (Adams et al. 2009; Adams et al. 2012; Crespo et al. 2001), showing a correlation between time spent watching television and nutritional status of the subjects involved. Dietz and Gortmaker demonstrated in 1985 a significant, positive association between hours of television viewed and obesity in children and adolescents, and since then these results were confirmed in several studies (Boone et al. 2007; Crespo et al. 2001; Jackson et al. 2009; Jago et al. 2005; Mendoza, Zimmerman, and Christakis 2007). There are two main aspects considered when investigating the effects of TV on childhood obesity: (1) reduced energy expenditure linked to screen time (Scaglioni et al. 2011) and (2) augmented energy intake driven by advertising and snacking (Rennie, Johnson, and Jebb 2005). When considering the problem from the latter perspective, the main hypothesis is that the effect of TV viewing time on childhood obesity may be independent of physicalactivity status and may be attributed to the increased total energy intake during TV watching (Manios et al. 2009), statement supported for example from Jackson’s results (Jackson et al. 2009) but contradicted in Schmidt and colleagues’ (2012) systematic review, which highlights that the studies on obesity intervention considered have reduced screen-media use in children, but points out as well that there are several research gaps, including a relative paucity of

studies targeting young children and minorities, limited long-term follow-up data, and few studies targeting removing TVs from children's bedrooms.

From the point of view of advertising, several important reviews reported already on the effect that TV advertising has on influencing children's preferences and choices (Hastings et al. 2003; Kunkel et al. 2004).

Nevertheless, very small attention has been given at today to the effect of TV advertising on the actual intake of children. Indeed, even since the very seminal experimental work (Gorn and Goldberg 1982) on effects of TV advertising on child behaviors with respect to food, the causal link between food-choice and food-consumption has been always hypothesized but seldom investigated as a research outcome. The preference for a certain food cannot be however directly translated in terms of actual consumption of that food, being modulated by a series of demographic, socioeconomic (Drewnowski 1997) and perhaps genetic factors (Cecil et al. 2008).

The aim of this article is to review evidences on experimental research focused on the relationship between children's exposure to TV advertising and their actual food intake.

## METHODS

Randomized controlled trials (RCTs) assessing the direct influence of food advertising on food consumption in children during TV viewing, were evaluated. The medical subject-headings terms that were combined and researched on the PubMed database were: TV, television, advertisements, screen time, weight, overweight, intake, consumption, eating, and child.

Kilocalories (Kcal) and grams (gr) were chosen as selection criterion to quantify children's food intake. No restrictions on time or language were applied.

All included abstracts were independently identified by two reviewers, and paper selection was independently performed, in order to select the articles fulfilling the inclusion criteria. All studies whose outcome was food intake's quantification during or immediately after screen time, were chosen. A summary of the selection criteria and search strategy is provided in table 1.

For each paper included in the review, authors, study design, population (dimension and age group), aim of the study, setting and intervention, outcome of interest, and results, were recorded. The methodological quality of the studies was assessed using the Jadad scale for RCTs (Jadad et al. 1996).

## RESULTS

Up to March 2012, 2,166 items were found. Seven studies, targeted to investigate the food intake during or after exposure to food advertising (either on TV or videogames) compared to neutral (non-food) commercials, were identified as pertinent. The main characteristics of the studies are shown in table 2. Quality of studies, evaluated on the 0- to 5-point Jadad scale, was low, having an average score of 1.57 and never being higher than 2 (table 3).

The results of Halford and colleagues (Halford et al. 2007; Halford et al. 2008) showed that total kcal intake was significantly higher after exposure to food advertising. Also, Harris and colleagues (2009) and Dovey and colleagues (2011) observed a relevant higher consumption of snacks during food advertising.

One study from Anschutz and colleagues (2009) reported on the potential existence of a gender effect, demonstrating that snack intake was different among girls and boys. Indeed, food intake in boys was higher when watching food advertising, whereas snack intake in girls was slightly lower when exposed to food advertising.

In a study by Halford and colleagues (2004), the effect of food advertising on food intake was reported as being potentially modulated by the overweight or obese status of the child. Children were divided into three weight status categories: lean, overweight, obese; and food intake in the three groups was estimated further to two categories: an exposure to food, and non-food advertising. Results showed a major consumption of snacks among obese and overweight children, both in the food advertising group and in non-food one.

The role of maternal pressure to control weight gain has been also investigated as a potential modulator of the children's eating as a reaction to food advertising (Anschutz, Engels, and Van Strien. 2010). The study showed that kids, who were encouraged from their mothers to be thinner, ate more especially when exposed to light food commercials than when exposed to neutral advertising. In contrast, children who perceived no maternal pressure ate more when exposed to neutral commercials than when exposed to either energy-dense or light food commercials.

## DISCUSSION

A first finding from the current review—in spite of the broad coverage on the subject, attesting to intense research activity—was that very few papers presenting RCT as a chosen study design (7 out of 2,166), were targeted to actual food intake as an outcome. This finding confirmed the initial impression at the ground of our investigation that, although there's global interest on the subject and strong recommendation to provide higher-level evidence (Barton 2000), studies evaluating actual food consumption using a higher-level study design, are still lacking.

The fact that this situation is an incontrovertible pitfall, is in line with Fenton and Fenton's (2011) considerations, namely that nutrition science should not and need not be complacent with lower levels of evidence than is required for other branches of medical science. In this perspective, also in the nutrition field, RCT in one of its variant forms is the study design that best permits strong causal inference concerning the relationship between a factor and any specific outcome (Blumberg et al. 2010). In addition, all the RCTs considered in this review showed an association between TV advertising and energy intake; but all the authors stressed the fact that their study was only a proof of concept, taken the small samples considered and the very specific age groups, and lacking the ability to prove causality (Adamson 2004).

Gender seemed to influence eating behavior in several studies (Anschutz et al. 2009; Halford et al. 2008), showing that an increased intake after exposure to food commercials occurred mostly in boys. A possible explanation for this finding was that boys may have a higher tendency to eat in response to visual food stimuli than do girls (Snoek et al. 2007; Van Strien and Bazelier 2007).

Boys are often found to have less self-control than do girls (Turner and Piquero 2002), and low self-control has been found to be related to a higher intake of saturated fat (Wills et al. 2007). Previous research showed that even very young girls experience more sociocultural pressure, eating disturbances, and/or body dissatisfaction than do their male peers (Gardner, Sorter, and Friedman 1997; Wood, Becker, and Thompson 1996). Thus, boys may actually have shown a normal response to food commercials, whereas girls may have suppressed their natural response to the food commercials (Doeschka et al. 2009). Another concept that appeared influencing food intake was food neophobia, considered especially in a study by Dovey and colleagues (2011); the study showed that acceptance or rejection of healthy foods has critical consequences for health during childhood and throughout the lifespan.

Paradigms which examine the role of food neophobia in determining the nature of a child's response to television food advertising, may enable the identification of children potentially at risk of increased weight gain, earlier in their development (Dovey et al. 2011). The nature of children's response to food advertisements for healthy and unhealthy foods, in part determined by food neophobia, may increase caloric intake (Dovey et al. 2011).

Parents' influence was analyzed in depth in a paper by Anschutz and colleagues (2010) and as seen also in other studies (Brown et al. 2008; Scaglioni, Salvioni, and Galimberti 2008), a restraining behavior was peered by an increased intake of food. Children appeared to be more influenced by TV advertising when the mothers were encouraging them to be thinner, while no significant association was found when considering maternal concern for children's weight and TV commercials.

As considered at the very beginning, methodological issues in this field of behavioral nutrition research appear to be exposed to potentially highly bias results, limiting therefore the possibility to extend these results to the population level. Similar considerations have been recently made from Vyth and colleagues (2012), again considering a specific and debated issues pertaining to behavioral nutrition. With their first study providing an overview of the methodological quality of



current front-of-packaging labeling research, Vyth and colleagues concluded that few methodologically sound studies are presently available, moreover suggesting the urgent need of longitudinal, randomized, controlled design in a real-life setting. In line with what was observed by Vyth and colleagues, the quality of the seven studies considered in this review was much lower than commonly accepted standards in other research fields.

In the same stream sets the present review, pointing out that standardized methods to assess the interaction between TV advertising and food intake in children are needed, in order to provide rigorous evidence to rulers and policy makers, in a field yet strongly influenced by familiar behavioral aspects and biology.

In conclusion, considering the insufficient data incident to this topic and the poor methodological quality of the few available evidences, more RCTs addressing the relationship between TV advertising and food intake in children, perhaps using strong outcomes like energy intake instead of food preferences, are required to deepen knowledge on causal relationships and therefore to set the root for incisive prevention.

## FUNDING

The work has been partially covered by an unrestricted grant from the University of Padova and by an unrestricted grant from Prochild ONLUS (Italy). In addition, the research did benefit by an unrestricted grant from the Italian Ministry of Foreign Affairs under the program “Programmi di alta rilevanza scientifica e tecnologica Italia-Messico.”

## AUTHORS' CONTRIBUTION

Dario Gregori conceptualized and drafted the manuscript, Simonetta Ballali, Maria Gabriella Vecchio, Silvia Scirè, and Francesca Foltran reviewed the literature and contributed to drafting the manuscript, Paola Berchiolla critically revised the manuscript and contributed to drafting its last version.

## REFERENCES

- Adams, J., K. Hennessy-Priest, S. Ingimarsdottir, J. Sheeshka, T. Ostbye and M. White. 2009. Food advertising during children's television in Canada and the UK. *Archives of Disease in Childhood* 94 (9): 658–662.
- Adams, J., R. Tyrrell, A. J. Adamson and M. White. 2012. Socio-economic differences in exposure to television food advertisements in the UK: A cross-sectional study of advertisements broadcast in one television region. *Public Health Nutrition* 15 (3): 487–494.
- Adamson, J. 2004. Evaluation of studies of causation (etiology). *Evidence Based Nursing* 7 (2): 36–40.
- Anschutz, D. J., R. C. Engels and T. Van Strien. 2009. Side effects of television food commercials on concurrent nonadvertised sweet snack food intakes in young children. *American Journal of Clinical Nutrition* 89 (5): 1328–1333.
- Anschutz, D. J., R. C. M. E. Engels and T. Van Strien. 2010. Maternal encouragement to be thin moderates the effect of commercials on children's snack food intake. *Appetite* 55 (1): 117–123.
- Barton, S. 2000. Which clinical studies provide the best evidence? The best RCT still trumps the best observational study. *BMJ* 321 (7256): 255–256.
- Boone, J. E., P. Gordon-Larsen, L. S. Adair and B. M. Popkin. 2007. Screen time and physical activity during adolescence: Longitudinal effects on obesity in young adulthood. *International Journal of Behavioral Nutrition and Physical Activity* 4:26.
- Blumberg, J., R. P Heaney, M. Huncharek, T. Scholl, M. Stampfer, R. Vieth, C. M Weaver, and S. H Zeisel. 2010. Evidence-based criteria in the nutritional context. *Nutrition Reviews* 68 (8): 478–484.
- Brown, K. A., J. Ogden, C. Vogele and E. L. Gibson. 2008. The role of parental control practices in explaining children's diet and BMI. *Appetite* 50 (2–3): 252–259.

- Caroli, M., L. Argentieri, M. Cardone and A. Masi. 2004. Role of television in childhood obesity prevention. *International Journal of Obesity and Related Metabolic Disorders* 28 (Suppl 3): S104–108.
- Carter, O. B., L. J. Patterson, R. J. Donovan, M. T. Ewing and C. M. Roberts. 2011. Children's understanding of the selling versus persuasive intent of junk food advertising: Implications for regulation. *Social Science & Medicine* 72 (6): 962–968.
- Cecil, J. E., R. Tavendale, P. Watt, M. M. Hetherington and C. N. A. Palmer. 2008. An obesity-associated FTO gene variant and increased energy intake in children. *New England Journal of Medicine* 359 (24): 2558–2566.
- Crespo, C. J., E. Smit, R. P. Troiano, S. J. Bartlett, C. A. Macera and R. E. Andersen. 2001. Television watching, energy intake, and obesity in US children: Results from the third National Health and Nutrition Examination Survey, 1988–1994. *Archives of Pediatrics & Adolescent Medicine* 155 (3): 360–365.
- Crowley, V. E., G. S. Yeo and S. O'Rahilly. 2002. Obesity therapy: Altering the energy intake-and-expenditure balance sheet. *Nature Reviews Drug Discovery* 1 (4): 276–286.
- Dietz, W. H. and S. L. Gortmaker. 1985. Do we fatten our children at the television set? Obesity and television viewing in children and adolescents. *Pediatrics* 75 (5): 807–812.
- Doeschka J Anschutz, Rutger CME Engels and Tatjana Van Strien. 2009. Side effects of television food commercials on concurrent nonadvertised sweet snack food intakes in young children. *American Journal of Clinical Nutrition* 89:1328–1333.
- Dovey, T. M., L. Taylor, R. Stow, E. J. Boyland and J. C. Halford. 2011. Responsiveness to healthy television (TV) food advertisements/commercials is only evident in children under the age of seven with low food neophobia. *Appetite* 56 (2): 440–446.
- Drewnowski, A. 1997. Taste preferences and food intake. *Annual Review of Nutrition* 17 (1): 237–253.

Ebbeling C. B., D. B. Pawlak, and D. S. Ludwig. 2002. Childhood obesity: Public health crisis, common sense cure. *Lancet* 360 (9331): 473–482.

Egger, G., and B. Swinburn. 1997. An “ecological” approach to the obesity pandemic. *BMJ* 315 (7106): 477–480.

Fenton, T. R., and C. J. Fenton. 2011. Nutrition science mustn’t accept a lower level of evidence. *Nutrition Reviews* 69 (7): 413–414; reply 415–416.

Gardner, R. M., R. G. Sorter, and B. N. Friedman. 1997. Developmental changes in children’s body images. *Journal of Social Behavior & Personality* 12:1019–1036.

Gorn, G. J., and M. E. Goldberg. 1982. Behavioral evidence of the effects of televised food messages on children. *Journal of Consumer Research* 9 (2): 200–205.

Halford, J. C., E. J. Boyland, G. Hughes, L. P. Oliveira, and T. M. Dovey. 2007. Beyond-brand effect of television (TV) food advertisements/commercials on caloric intake and food choice of 5–7-year-old children. *Appetite* 49 (1): 263–267.

Halford, J. C., E. J. Boyland, G. M. Hughes, L. Stacey, S. McKean, and T. M. Dovey. 2008. Beyond-brand effect of television food advertisements on food choice in children: The effects of weight status. *Public Health Nutrition* 11 (9): 897–904.

Halford, J. C., J. Gillespie, V. Brown, E. E. Pontin, and T. M. Dovey. 2004. Effect of television advertisements for foods on food consumption in children. *Appetite* 42 (2): 221–225.

Harris, J. L., J. A. Bargh, and K. D. Brownell. 2009. Priming effects of television food advertising on eating behavior. *Health Psychology* 28 (4): 404–413.

Hastings, G., M. Stead, L. McDermott, A. Forsyth, A. M. MacKintosh, M. Rayner, C. Godfrey, M. Caraher, and K. Angus. 2003. Review of research on the effects of food promotion to children. Glasgow, Scotland: Center for Social Marketing, University of Strathclyde Glasgow.

Jackson, D. M., K. Djafarian, J. Stewart, and J. R. Speakman. 2009. Increased television viewing is associated with elevated body fatness but not with lower total energy expenditure in children. *American Journal of Clinical Nutrition* 89 (4): 1031–1036.

Jadad, A. R., R. A. Moore, D. Carroll, C. Jenkinson, D. J. M. Reynolds, D. J. Gavaghan, and H. J. McQuay. 1996. Assessing the quality of reports of randomized clinical trials: Is blinding necessary? *Controlled Clinical Trials* 17 (1): 1–12.

Jago, R., T. Baranowski, J. C. Baranowski, D. Thompson and K. A. Greaves. 2005. BMI from 3–6 y of age is predicted by TV viewing and physical activity, not diet. *International Journal of Obesity (London)* 29 (6): 557–564.

Kumanyika, S., R. W. Jeffery, A. Morabia, C. Ritenbaugh, and V. J. Antipatis. 2002. Public Health Approaches to the Prevention of Obesity (PHAPO) Working Group of the International Obesity Task Force (IOTF). *International Journal of Obesity & Related Metabolic Disorders* 26 (3): 425–436.

Kunkel, D., B. L. Wilcox, J. Cantor, E. Palmer, S. Linn, and P. Dowrick. 2004. Report of the APA Task Force on Advertising and Children, 40. Washington, DC: American Psychological Association.

Lobstein, T., L. Baur, and R. Uauy. 2004. Obesity in children and young people: A crisis in public health. *Obesity Reviews* 5 (Suppl 1): 4–104.

Manios, Y., G. Kourlaba, K. Kondaki, E. Grammatikaki, A. Anastasiadou, and E. Roma-Giannikou. 2009. Obesity and television watching in preschoolers in Greece: The GENESIS study. *Obesity (Silver Spring)* 17 (11): 2047–2053.

Mendoza, J. A., F. J. Zimmerman, and D. A. Christakis. 2007. Television viewing, computer use, obesity, and adiposity in US preschool children. *International Journal of Behavioral Nutrition & Physical Activity* 4:44.

Rennie, K. L., L. Johnson, and S. A. Jebb. 2005. Behavioral determinants of obesity. *Best Practice & Research Clinical Endocrinology & Metabolism* 19 (3): 343–358.

Scaglioni, S., C. Arrizza, F. Vecchi, and S. Tedeschi. 2011. Determinants of children’s eating behavior. *American Journal of Clinical Nutrition* 94 (6 Suppl): 2006S–2011S.

Scaglioni, S., M. Salvioni, and C. Galimberti. 2008. Influence of parental attitudes in the development of children eating behavior. *British Journal of Nutrition* 99 (Suppl 1): S22–25.

Schmidt, M. E., J. Haines, A. O'Brien, J. McDonald, S. Price, B. Sherry, and E. M. Taveras. 2012. Systematic review of effective strategies for reducing screen time among young children. *Obesity (Silver Spring)* 20 (7): 1338–1354.

Snoek H. M., T. Van Strien, J. M. A. M. Janssens, and R. C. M. E Engels. 2007. Emotional, external, restrained eating and overweight in Dutch adolescents. *Scandinavian Journal of Psychology* 48:23–32.

Turner, M. G., and A. R. Piquero. 2002. The stability of self-control. *Journal of Criminal Justice* 30:457–471.

Van Strien, T., and F. G. Bazelier. 2007. Perceived parental control of food intake is related to external, restrained and emotional eating in 7–12-year-old boys and girls. *Appetite* 49:618–625.

von Kries, R., H. Reulen, O. Bayer, C. Riedel, K. Diethelm, and A. E. Buyken. 2012. Increase in prevalence of adiposity between the ages of 7 and 11 years reflects lower remission rates during this period. *Pediatric Obesity* 8 (1): 13–20.

Vyth, E. L., I. H. Steenhuis, H. E. Brandt, A. J. Roodenburg, J. Brug, and J. C. Seidell. 2012. Methodological quality of front-of-pack labeling studies: A review plus identification of research challenges. *Nutrition Reviews* 70 (12): 709–720.

Wang, Y., and T. Lobstein. 2006. Worldwide trends in childhood overweight and obesity. *International Journal of Pediatric Obesity* 1 (1): 11–25.

WHO. 2000. Obesity: Preventing and managing the global epidemic. Report of a WHO consultation. Technical Report Series 894, i–xii; 1–253. Rome: World Health Organization.

Wills, T.A., C. R. Isasi, D. Mendoza, and M. G. Ainette. 2007. Self-control constructs related to measures of dietary intake and physical activity in adolescents. *Journal of Adolescent Health* 41:551–558.

Wood, K. C., J. A. Becker, and J. K. Thompson. 1996. Body image dissatisfaction in preadolescent children. *Journal of Applied Developmental Psychology* 17:85–100.







Table 1. Studies Selection Strategy

String on PubMed	No. of obtained articles	Selection criterion to quantify children's food intake	Restrictions on time or language	Outcome	No. of adequate articles
TV, television, advertisements, screen time, weight, overweight, intake, consumption, eating and child	2166	Kilocalories (Kcal) and grams(gr)	None	Food intake's quantification during or immediately after screen time	7

Table 2. Studies Included in the Analysis

Reference (authors, year)	Country	Participants (N, age)	Aim of the study	Setting and intervention	Outcome of interest	Results
Dovey 2011	UK	66 children aged 5–7 years old (male: 34, 51.5%)	Assessment of the effects of advertisement type (healthy, unhealthy foods and toy) and food neophobia status on caloric intake by food item and total intake after viewing television.	School. Children watched a 14-min cartoon embedded with a selection of each type of advert. After watching, they were offered a snack consisting of pre-weighed individual servings of six snack foods on a plate.	Caloric intake after viewing a cartoon embedded with each advertisement type: total kcal consumed in each condition and individually for each food item.	For all children, total kcal intake was higher following the unhealthy food ads compared to both the healthy food ads and toy ads.
Anschutz 2010	Netherlands	121 children aged 8–12 years old (male: 8.8%)	Assessment of the effects of adult targeted food commercials (energy-dense and light food products) on actual intake in young children while watching television and the moderating effect of maternal encouragement on the relation between commercial condition and food intake.	A setting that resembled a living room was created at school. Children could freely eat from a pre-weighed bowl with chocolate-coated peanuts, while watching a 20-min movie clip interrupted by commercials of different content across the experimental conditions.	Snack food intake (in g) during watching television.	Children who perceived maternal encouragement to be thin ate more when exposed to light food commercials compared to neutral ads. Children who perceived no maternal encouragement to be thin ate more when exposed to neutral commercials compared to either energy-dense or light food commercials. Neither commercial condition nor maternal encouragement to be

						thin had a significant main effect on food intake.
Harris 2009	US	118 children aged 7–11 years old (male: 62, 52.5%)	Experiment testing whether exposure to food advertising during TV viewing may prime, or directly activate, an automatic increase in snack food consumption in elementary-school-age children.	School or camp. The children watched a 14-min episode of a cartoon, included commercials: one-half were randomized to the “food commercial condition” and one-half to the “nonfood commercial condition”. Children received a pre-weighed bowl of snacks to consume while watching.	Amount of snack foods consumed (g) during TV advertising exposure.	The group exposed to cartoon and ads ate considerably more snacks, compared to the group exposed to non-food advertisement.
Anschutz 2009	Netherlands	120 children aged 8–12 years old (male: 46.7%)	Assessment of the general effect of food commercials on immediate snack food intake in children while watching television.	School. Children were randomly assigned over conditions (food ads and neutral ads). They watched a 20-min movie clip interrupted by commercial breaks and they could freely eat from a pre-weighed bowl containing chocolate-coated peanuts during TV viewing.	Amount of snack food eaten (g) during the experiment.	No effect of commercial condition on food intake, while a significant main effect was found for sex. The interaction between advertisement condition and sex was significant. Snack intake by boys was higher when they watched the food ads than when they watched the neutral ads whereas snack food intake by girls was slightly lower in the food commercial

						condition than in the neutral commercial condition.
Halford 2008	UK	59 children aged 9–11 years old (male: 32, 54.2%)	Assessment of the influence of food advertisement exposure on caloric intake and food preferences after TV viewing.	School. Children were exposed to a collection of 10 non-food or food ads, followed by a 10-min cartoon. After viewing, each child was presented with five plates, containing one of each of the five food categories (low-fat savory, low-fat sweet, high-fat savory, high-fat sweet, low density energy) in either or a standard portion size or pre-weighed. They could eat as little or as much food as they liked. The remaining uneaten food was re-weighed.	Amount of food eaten in kcal after advertisement exposure.	Total energy intake was significantly higher after exposure to food advertisements than after the control (toy) advertisements. A significant interaction with BMI status and advertisement type was observed. The significant difference in intake between the three weight status groups was in the food advertisement condition only as all participants consumed a similar amount of food following the control (toy) advertisements. Specifically, following food ads, OB children consumed more food than the NW and OW children.
Halford 2007	UK	93 children aged 5–7 years old (male: 39, 42%)	Assessment of the influence of food advertisement exposure on caloric intake and food	School. Children were exposed to a collection of 10 non-food or food ads, followed by a	Amount of food eaten in kcal after food advertisement exposure	A main effect of advertisement exposure on total kcal intake was found. Total kcal intake

			preferences after TV viewing	10-min cartoon. After viewing, each child was presented with pre-weighed individual servings of five food categories (low-fat savory, low-fat sweet, high-fat sweet, high-fat savory and fruit). They could eat as little or as much food as they liked. The remaining uneaten food was re-weighed.		was significantly higher after exposure to food advertisements (FA) than after the non-food advertisements (NA). There was no main effect of weight status on intake despite the fact that mean intake in the overweight and the obese children was higher in both experimental conditions (692.1 in the FA condition; 594.9 in the NA condition), nor was there any significant interaction with weight status.
Halford 2004	UK	42 children aged 9-11 years old (male: 18, 43%)	Assessment of the ability to recognize food and non-food related advertisements and assessment of the impact of exposure to TV food advertisements on total food intake and on food choices after TV viewing	School. Children watched a collection of non-food or food ads, followed by a 10-min cartoon. After viewing, children were presented with a list of 16 ads (of which 8 had been shown) and asked to identify which ads they had seen. Then each child was presented with a pre-weighed set of four food categories (low-fat savory, low-fat	Amount of food eaten in g after food advertisement exposure	The obese and overweight groups ate significantly more than the healthy-weight group, both with food ads (FA) and with non-food ads (NA). The difference between FA and NA conditions across all children was highly reliable

				sweet, high-fat sweet, high-fat savory). They could eat as little or as much food as they liked. The remaining uneaten food was reweighed.		
--	--	--	--	--	--	--

Table 3. Quality Assessment of the Included Studies

Study	Study design	Quality score
Anschutz 2009	RCT	2
Halford 2004	Counterbalanced design	1
Halford 2007	Counterbalanced design	1
Anschutz 2010	RCT	2
Dovey 2011	Counterbalanced cross-over study	1
Harris 2009	RCT	2
Halford 2008	Counterbalanced design	2