

## Expression of NOX2 protein in neutrophils of patients with ALS

De Marco G<sup>1</sup>, Lomartire A<sup>1</sup>, Casale F<sup>1</sup>, Marrali G<sup>1</sup>, Cammarosano S<sup>1</sup>, Ilardi A<sup>1</sup>, Canosa A<sup>1</sup>, Moglia C<sup>1</sup>, Calvo A<sup>1</sup>, Rinaudo MT<sup>2</sup>, Chiò A<sup>1</sup>

<sup>1</sup>Rita Levi Montalcini' Department of Neuroscience, ALS Centre, University of Torino, Via Cherasco 15, 10126, Turin, Italy

<sup>2</sup>Department of Oncology, University of Torino, Via Michelangelo 27bis, 10125, Turin, Italy

Oxidative stress, more precisely the damage caused by an abnormal production of reactive oxygen species (ROS), is a key mechanism in ALS pathogenesis. NADPH oxidases are a family of enzymes responsible for the production of ROS. In our previous study, an increased survival of ALS-affected individuals has been related to low activity of NADPH oxidase-2 (NOX2) in their neutrophils (1). Based on this finding, this work aimed to evaluate the protein levels of NOX2 in neutrophils of patients affected by ALS, in order to verify a possible correlation between levels and activity.

The analysis was performed in a cohort of 40 subjects using Western immunoblot technique. The attention was mainly focused on ALS patients carrying mutations of copper-zinc superoxide dismutase-1 (SOD1) gene, since mutations of SOD1 are referred to alter NOX2 activity (2).

(1) Marrali G et al. NADPH oxidase (NOX2) activity is a modifier of survival in ALS. *J. Neurol.* 2014

(2) Harraz et al. SOD1 mutations disrupt redox-sensitive Rac regulation of NADPH oxidase in a familial ALS model. *J Clin Invest.* 2008