



The premier source for peer-reviewed plant pathology research since 1911.

[Journals Home](#)[APS Home](#)[IS-MPMI Home](#)[My Profile](#)[Subscribe](#)[Search](#)[Advanced Search](#)[Help](#)[Share](#) |[Subscribe](#) [Free alerts](#) [RSS](#)

About the cover for November 2015

Impact Factor: 3.02

ISSN: 0191-2917

SEARCH

Enter Keywords

- Phytopathology
- Plant Disease
- MPMI

search

[Advanced Search](#)

Resources

[About Plant Disease](#)[First Look](#)[Most Downloaded Articles](#)[Submit a Manuscript](#)[Customer Care](#)[About My Password](#)[Copyright and Permissions](#)[Plagiarism and Ethics](#)[Advertise](#)[e-Xtra](#)[Open Access](#)

plant disease

Editor-in-Chief: Mark L. Gleason

Published by The American Phytopathological Society

[Home](#) > [Plant Disease](#) > [Table of Contents](#) > [Full Text HTML](#)[Previous Article](#) | [Next Article](#)

November 2015, Volume 99, Number 11

Page 1648

<http://dx.doi.org/10.1094/PDIS-01-15-0026-PDN>

DISEASE NOTES

First Report of *Sclerotinia sclerotiorum* Causing Postharvest Sclerotinia Rot on Highbush Blueberry in Europe

M.P. Bustos Lopez, D. Spadaro, M.L. Gullino, AGROINNOVA—Centre of Competence for the Innovation in the Agro-Environmental Sector; and Department of Agricultural, Forestry and Food Sciences, University of Torino, 10095 Grugliasco, Italy.

[Citation](#)[Open Access](#).

ABSTRACT

Northern highbush blueberry (*Vaccinium corymbosum* L.) is widely cultivated in different European countries. Its distribution in Italy is steadily growing in northern and central regions, but it is mainly limited by the requirement of acidic soil. Blueberries are available on the market from June to September and they have a shelf life of 21 days. During June 2012, a postharvest fruit rot was observed on highbush blueberry 'Duke' cultivated in Saluzzo (44°38'50.64" N; 07°29'14" N), Piedmont region, Italy, and stored 7 days at 2°C after harvesting. Infected fruit showed brown, soft, rotted lesions, covered with grayish white mycelium under high humidity. These symptoms were similar to Sclerotinia rot caused by *Sclerotinia sclerotiorum* (Libert) de Bary in Japan (Umamoto et al. 2007), or mummy berry disease caused by *Monilinia vaccinii-corymbosi* in United States and Europe (Hildebrand et al. 1995). Tissues were excised from the margin between the healthy and diseased tissues, cultured on potato dextrose agar (PDA, Merck) amended with 25 mg streptomycin per liter, and incubated at 24°C (16 h of light and 8 h of darkness). Fungal colonies initially appeared white with a fairly flat sheet of aerial mycelia and after 7 days produced black sclerotia measuring 3 to 8 mm in diameter on the colony surface mainly near the edge of the culture plates. Typical microconidia of the genus *Monilia* (syn. *Monilinia*) were not observed (Hildebrand et al. 1995). The pathogen was identified as *Sclerotinia sclerotiorum* based on the size of sclerotia and the DNA sequences of the ribosomal region ITS1-5.8S-ITS2, obtained with PCR amplification with primers ITS1 and ITS4. A BLAST search of one sequence in GenBank showed 99% sequence coverage and 99% (487/489 bp) similarity to two ribosomal sequences of *S. sclerotiorum* isolated from *V. corymbosum* in Japan (Umamoto et al. 2007). The sequence (490 bp) was deposited in GenBank with Accession No. JX442064. *S. sclerotiorum* was isolated from 6% of the berries showing postharvest rots. Pathogenicity was tested on 20 highbush blueberries cv 'Duke'. Mycelial blocks (9 mm²) were placed on artificial wounds (1-mm diameter, 1-mm depth) generated with a needle on the equatorial region of the berries and incubated at 20°C with 12 h of light. Control berries were wounded and a sterile agar block was placed

Quick Links

[Add to favorites](#)[E-mail to a colleague](#)[Alert me when new articles cite this article](#)[Download to citation manager](#)[Related articles found in APS Journals](#)

Article History

Print: 3 Nov 2015

Ahead of Print: 8 Sep 2015


Accepted: 25 Feb 2015

This Journal is brought to you via a subscription from the Univ Di Torino

CALL FOR PAPERS
Special Focus Issue

Disease Management
in the Genomics Era

Submissions Due:
February 1, 2016



on the wounds. Seven days after inoculation, the rot symptoms occurred and *S. sclerotiorum* was reisolated from inoculated fruit on PDA and identified based on morphology and rDNA sequencing. Control fruits were symptomless. Previously, *S. sclerotiorum* was reported causing shoot blight, flower cluster rots, or berry rots on highbush blueberry in Canada, New Zealand, Japan (Umemoto et al. 2007), and Argentina (Perez et al. 2011). To our knowledge, this is the first report of *S. sclerotiorum* causing a postharvest fruit rot of blueberry in Europe.

References:Section:

-
- Hildebrand, P. D.**, et al. 1995. Page 11 in: Compendium of Blueberry and Cranberry Disease. **Caruso, F. L.**, and **Ramsdell, D. C.**, eds. APS Press, St Paul.
- Perez, B. A.**, et al. 2011. Plant Dis. 95: 774. 10.1094/PDIS-01-11-0074
- Umemoto, S.**, et al. 2007. J. Gen. Plant Pathol. 73:290. 10.1007/s10327-007-0004-1

[Citation](#)