


Contents lists available at [ScienceDirect](http://ScienceDirect.com)

Medical Hypotheses

journal homepage: www.elsevier.com/locate/mehy

Skin phenotypes can offer some insight about the association between telomere length and cancer susceptibility

S. Ribero MD, PhD^{a,b,*}, M. Mangino PhD^a, V. Bataille MD, PhD^{a,c}^a Department of Twin Research and Genetic Epidemiology, King's College London, UK^b Department of Medical Sciences, University of Turin, Turin, Italy^c Department of Dermatology, West Herts NHS Trust, Herts, UK

ARTICLE INFO

Article history:

Received 26 August 2016

Accepted 18 October 2016

ABSTRACT

The role of telomere biology in cancer has been studied for a wide variety of different cancers but the association with telomere length has been controversial. This is because some cancers have been found to be associated with longer telomeres in circulating white cells whilst other cancer types are more common in individuals with shorter telomeres. Hence, there has been some skepticism as to whether telomere length may be helpful in estimating cancer risk. For melanoma, however, results have been fairly consistent showing that longer telomeres are associated with an increased risk. This link was first discovered because of a link between longer telomeres and a high number of naevi. In contrast, for cutaneous squamous cell carcinomas, the relationship is reversed with higher risk in individuals with shorter telomeres. Differences in skin phenotypes with the presence of high number of naevi versus photoageing with solar elastosis and solar keratoses have already been valuable for dermatologists as the former phenotype is associated with melanoma whilst the latter is more common in patients with squamous cell carcinoma of the skin. The hypothesis is that the differences in cutaneous phenotypes already observed by dermatologists for skin cancers may, in fact, be useful as well for cancer prediction in general as it may reflect underlying telomere biology. This manuscript will address the evidence for links between telomere biology, skin phenotypes and cancer risk.

© 2016 Elsevier Ltd. All rights reserved.

Telomeres are repeat TTAGGG sequences at the end of linear chromosomes, which guard against loss of genetic material during cellular replication. Due to an inherent end replication problem, chromosomes are exposed to a potential loss of genetic material so telomeres act as a buffer against loss of chromatin [1]. In normal human somatic cells, telomeres range from 9 to 15 kb initially, with a progressive loss in mean telomere length of 15–66 bp per year [2]. It has been estimated that 17.5% of the inter-individual variation in leukocyte telomere length is due to the ageing process [3]. Repeated cell cycles eventually lead to a critically shortened telomere length which then triggers apoptosis. In cell cultures, the replicative potential of human cells is estimated at an average of 52 mitoses per cell known as the Hayflick limit, by which stage, the critical telomere shortening will lead to the activation of genes pushing the cells in cell cycle arrest [4]. This arrest in proliferation is thought to protect against malignant transformation and a failure to do so results in catastrophic genomic instability and carcino-

genesis. Telomeres are thus important in managing genomic stability. This central role in genome maintenance makes telomeres key players in carcinogenesis and an attractive candidate for tumour profiling at the molecular level. Although telomere length has a strong inheritable component [5], it is also influenced by a wide variety of environmental factors such as oxidative stress, chronic inflammation, smoking and obesity [2,6–8]. Telomeres have been estimated to be 240 bp shorter in obese women compared to women with a BMI below 25 and also 5 bp shorter for every cigarette pack year smoked [2].

The link between cancer susceptibility and telomere length has not been easy to elucidate perhaps because there was an assumption that all cancers would show associations with telomere length in the same direction whatever the cancer type. However, when grouping all cancers together it appears that conflicting results emerge [9,10]. It is likely that cancer types need to be analysed separately to observe trends with telomere length. Most studies looking at associations between telomere length and cancer have used circulating leukocytes as blood sampling is easily accessible and telomere length in white cells correlates to telomere length in other human healthy tissues (but very different from neoplastic

* Corresponding author at: Department of Twin Research and Genetic Epidemiology, King's College London, St Thomas' Campus, Westminster Bridge Road, London SE1 7EH, UK.

E-mail address: simone.ribero@unito.it (S. Ribero).

tissues where telomerase and other neoplastic processes affect telomerase) [3,11,12].

Melanoma susceptibility via an excess of naevi was one of the first tumour to show a link with longer telomeres in white cells [13]. The reason why telomere length was measured in individuals with a high number of naevi was because dermatologists working in high risk melanoma clinics had observed that these high risk patients with multiple atypical naevi showed reduced signs of skin ageing and photoageing compared to their peers. It was therefore suspected that their ageing process may be delayed. In fact, many melanoma patients show far less photoageing compared to patients with squamous cell carcinoma, the latter usually associated with sun damaged skin with solar lentigines and solar keratoses. The link between melanoma susceptibility and longer telomeres suggests that these individuals may have a reduced senescence which is likely to affect many cell types and not just melanocytes. It is yet, unproven, if longer telomeres are causal in the formation of a large numbers of cutaneous naevi in these high risk melanoma patients. However, considering the role of the telomere unit in cell replication, it is highly likely that an excess of naevi is a reflection of the lack of senescence in melanocytes and is linked to their longer telomeres [14]. This has potential implications for other tissues as individuals with high numbers of naevi have been shown to be taller than those with fewer naevi and also have higher bone mineral density [15]. Bone mass and growth early in life is likely to be affected by telomere biology well before melanoma susceptibility is evident later in life. Melanoma case-control studies, family studies and analyses of genetic changes in melanoma tumours have all replicated the link between melanoma susceptibility and telomere biology from 2011 onwards. These studies assessed telomere length in melanoma cases compared to controls or reported on polymorphisms in genes which affect telomere length [16–19].

In contrast, cutaneous squamous cell carcinoma susceptibility is associated with shorter telomeres [20–22]. The cutaneous phenotype associated with squamous cell carcinoma of the skin is characterised by epidermal and dermal atrophy, solar lentigines and solar keratoses which are far less common in melanoma patients (albeit in lentigo maligna in elderly patients) [23]. This photoaged skin phenotype is likely to reflect the altered telomere biology via shorter telomeres and increased senescence in cutaneous squamous cell carcinoma patients.

More recently, large genome wide association studies led to the identification of 10 loci associated with telomere length variation [24–29]. Following the discovery of these loci, a large melanoma case-control study from the Genomel Consortium developed a genetic score to estimate melanoma risk based on these SNPs associated with telomere length variation [26]. The combined SNPs score does predict melanoma risk again confirming the role of the telomere unit in melanoma susceptibility [30]. It is, however, important to note these SNPs only explain 1% of the variance in telomere length so many other genes, environmental factors as well as epigenetics factors may explain the rest of the variance in telomere length. Genetic variants reported in the Codd et al. study [26] may not have an effect solely on telomere length as they may have pleiotropic effects on cancer risk via immune responses and DNA repair as well [30].

In fact, well before these recent telomere studies, dermatologists were well aware that phenotypic risk factors for squamous cell carcinomas and melanoma were different. This may be surprising when there is a tendency to combine all skin cancers together and always assume that they are all caused by excessive sun exposure. Squamous cell carcinoma of the skin is usually associated with chronic sun damage whilst many melanoma patients, especially in the context of familial susceptibility, show very little signs of photoageing. A high number of naevi (high risk of melanoma)

and severe photoageing with solar keratoses (high risk of squamous cell carcinoma) have already been reported to often be mutually exclusive even after adjusting for age which support the fact that melanomas and squamous cell carcinoma do arise from very different “at risk” cutaneous phenotypes. Of the smaller numbers of melanomas associated with a photoageing phenotype these are more likely to be seen in older patients, be a nodular subtype or lentigo maligna and on chronically sun exposed sites [31,32]. This led to the divergent melanoma pathways hypothesis described in the 1990s via the photoageing or alternatively via the naevus phenotype [31,32]. Whilst it is true that squamous cell carcinoma affects older individuals compared to melanoma, telomere studies looking at SCC have adjusted for age and the association remains with shorter telomeres for squamous cell carcinomas [21]. Shorter telomeres in squamous cell carcinomas may also explain the reduced immune surveillance with less robust lymphocytic response in relation to neoplastic keratinocytes. It is common to find a strong lymphocytic response in melanoma tumours whilst for squamous cell carcinomas, this is not commonly reported histologically. While it is not proven that these differences in immune responses in different types of skin cancers are explained by telomere biology, it is quite likely that longer telomeres have a significant impact on lymphocyte functions [33].

Could these differences in the “at risk” skin phenotypes observed for skin cancers shed any light on the association between telomere length and other cancers? In fact, there are already interesting contrasting results regarding telomere length and lung cancer risk where the squamous cell carcinoma type in the lung is more likely to be associated with shorter telomeres whilst adenocarcinoma of the lung (more common in non-smokers) is linked to longer telomeres and this appears to be consistent across studies [34–37]. SNPs in the *TERT* gene have been associated with adenocarcinomas of the lung but not with squamous cell or small cell lung cancers [38]. This again highlights the issue of histological subtypes when looking at associations between cancer and telomere length or telomere SNPs. The association between shorter telomeres and squamous cell carcinoma type tumours is also often reported for squamous cell tumours of other organs such as oro-pharynx [39,40] oesophagus [41–44] or cervix [45]. Adenocarcinomas of the colon, kidney, prostate, uterus and breast are, on the contrary, more likely to be associated with longer telomeres [46–53]. Still, the association with adenocarcinomas of the colon and breast is controversial as other studies have shown inverse relationships with telomere length or no association at all [9]. The association between telomere length and cancer is therefore quite complex as some studies have reported a U shape curve with risk of cancer seen in the extreme groups of telomere length [54–56]. Telomere length can be also be affected by environmental factors and these are often not adjusted for, such as, obesity, smoking (apart from lung cancer) and alcohol abuse. Furthermore, there has been discrepant results between studies looking at telomere length measured by Southern Blot (which gives a measurement of the telomere unit in bp but is more time consuming) and those using qPCR (faster and cheaper but does not give an actual length but mean TL ratio) [57]. Recently, whole genome sequencing has been used to measure telomere length and may, in the future, replace qPCR and Southern blot [58].

In Dyskeratosis Congenita, considered a short telomere syndrome caused by mutations in telomere genes, SCCs of the head and neck and oral leucoplakia are reported frequently and the skin is showing all the signs of premature photoageing with keratinocyte dysplasia so again the skin in this syndrome is very informative as it is a window into telomere biology. These patients are also more prone to cutaneous and non cutaneous SCC type tumours and not adenocarcinomas in line with their very short telomeres [59].

In melanoma families, the risk of other solid tumours is increased and the most commonly reported tumours are brain, pancreas, breast, colon, kidney, lymphomas and sarcomas amongst many others [60–65]. For the brain, gliomas have already been linked to longer telomeres and the same has been observed for soft tissue sarcoma [66–68]. In melanoma families, tumours are also more likely to be of the adenocarcinoma type from the colon, breast, pancreas or prostate cancer whilst the squamous cell carcinoma type such as SCC of the skin, lung, oropharynx or cervix are not as common [61,69,70]. So even in the familial clustering of tumours within melanoma families, the association is more likely to be observed with tumours linked to longer telomeres rather than shorter telomeres.

Telomere biology may also be relevant for cancer survival as shown for colon cancer where telomere length has been used as a predictor of immune function in colon cancer [71,72] as longer telomeres are a favourable prognostic factor in this tumour [73]. It is speculated that this is explained by a more efficient immune response. The implications are that telomere length measurements have a potential as risk stratification enrichment biomarkers for estimating cancer risk but also for prognosis and for individualised therapeutic interventions [14]. Globally shorter telomere length in circulating white cells have been associated with reduced cancer survival and these results have been confirmed in a meta-analysis [10, 47].

In melanoma, high number of naevi offer a survival advantage after adjusting for all melanoma prognostic factors and this again may be related to telomere biology as an excess of naevi is linked to longer telomeres [74]. To date, there are no studies published looking at melanoma survival in relation to telomere length and this is because telomere length is not an easy phenotype to collect and 5 years follow up data is needed to assess melanoma survival. These studies are not yet available.

Conclusion

The skin is a very important window into the body reflecting many changes in cellular functions, especially in relation to the ageing process. Research so far suggests that changes in skin properties with age have good correlations with telomere length especially in skin carcinogenesis. This, in turn, may be highly relevant for cancer biology in general and not just skin so that assessing skin phenotypes in different types of cancers in the future may shed some light on underlying telomere biology and further elucidate the complex relationship between ageing and cancer susceptibility.

Conflict of interest statement

The authors declare no conflict of interest.

References

- [1] Blackburn EH. Switching and signaling at the telomere. *Cell* 2001;106(6):661–73.
- [2] Valdes AM, Andrew T, Gardner JP, et al. Obesity, cigarette smoking, and telomere length in women. *Lancet* 2005;366(9486):662–4.
- [3] Daniali L, Benetos A, Susser E, et al. Telomeres shorten at equivalent rates in somatic tissues of adults. *Nat Commun* 2013;4:1597.
- [4] Hayflick L. Living forever and dying in the attempt. *Exp Gerontol* 2003;38(11–12):1231–41.
- [5] Broer L, Codd V, Nyholt DR, et al. Meta-analysis of telomere length in 19,713 subjects reveals high heritability, stronger maternal inheritance and a paternal age effect. *Eur J Hum Genet* 2013;21(10):1163–8.
- [6] von Zglinicki T, Martin-Ruiz CM. Telomeres as biomarkers for ageing and age-related diseases. *Curr Mol Med* 2005;5(2):197–203.
- [7] Martínez P, Blasco MA. Telomeric and extra-telomeric roles for telomerase and the telomere-binding proteins. *Nat Rev Cancer* 2011;11(3):161–76.
- [8] Sanders JL, Newman AB. Telomere length in epidemiology: a biomarker of aging, age-related disease, both, or neither? *Epidemiol Rev* 2013;35:112–31.
- [9] Wentzensen IM, Mirabello L, Pfeiffer RM, Savage SA. The association of telomere length and cancer: a meta-analysis. *Cancer Epidemiol Biomarkers Prev* 2011;20(6):1238–50.
- [10] Ma H, Zhou Z, Wei S, et al. Shortened telomere length is associated with increased risk of cancer: a meta-analysis. *PLoS One* 2011;6(6).
- [11] Takubo K, Izumiya-Shimomura N, Honma N, et al. Telomere lengths are characteristic in each human individual. *Exp Gerontol* 2002;37(4):523–31.
- [12] Aubert G, Hills M, Lansdorp PM. Telomere length measurement—caveats and a critical assessment of the available technologies and tools. *Mutat Res* 2012;730(1–2):59–67.
- [13] Bataille V, Kato BS, Falchi M, et al. Nevus size and number are associated with telomere length and represent potential markers of a reduced senescence in vivo. *Cancer Epidemiol Biomarkers Prev* 2007;16:1499–502.
- [14] Shay JW. Are short telomeres predictive of advanced cancer? *Cancer Discov* 2013;3(10):1096–8.
- [15] Ribero S, Glass D, Aviv A, et al. Height and bone mineral density are associated with naevus count supporting the importance of growth in melanoma susceptibility. *PLoS One* 2015;10(1):e0116863.
- [16] Nan H, Du M, De Vivo I, et al. Shorter telomeres associate with a reduced risk of melanoma development. *Cancer Res* 2011;71(21):6758–63.
- [17] Bodelon C, Pfeiffer RM, Bollati V, et al. On the interplay of telomeres, nevi and the risk of melanoma. *PLoS One* 2012;7(12):e52466.
- [18] Huang FW, Hodis E, Xu MJ, et al. Highly recurrent TERT promoter mutations in human melanoma. *Science* 2013;339(6122):957–9.
- [19] Horn S, Figl A, Rachakonda PS, et al. TERT promoter mutations in familial and sporadic melanoma. *Science* 2013;339(6122):959–61.
- [20] Han J, Qureshi AA, Prescott J, et al. A prospective study of telomere length and the risk of skin cancer. *J Invest Dermatol* 2009;129:415–21.
- [21] Anic GM, Sondak VK, Messina JL, et al. Telomere length and risk of melanoma, squamous cell carcinoma, and basal cell carcinoma. *Cancer Epidemiol* 2013;37(4):434–9.
- [22] Caini S, Raimondi S, Johansson H, et al. Telomere length and the risk of cutaneous melanoma and non-melanoma skin cancer: a review of the literature and meta-analysis. *J Dermatol Sci* 2015;80(3):168–74.
- [23] Liang X, Qureshi AA, Guo Q, De Vivo I, Han J. No association between telomere length in peripheral blood leukocytes and the risk of non-melanoma skin cancer. *Cancer Epidemiol Biomarkers Prev* 2011;20(5):1043–5.
- [24] Levy D, Neuhausen SL, Hunt SC, et al. Genome-wide association identifies OBF1 as a locus involved in human leukocyte telomere biology. *Proc Natl Acad Sci USA* 2010;107(20):9293–8.
- [25] Codd V, Mangino M, van der Harst P, et al. Common variants near TERC are associated with mean telomere length. *Nat Genet* 2010;42(3):197–9.
- [26] Codd V, Nelson CP, Albrecht E, et al. Identification of seven loci affecting mean telomere length and their association with disease. *Nat Genet* 2013;45(4):422–7.
- [27] Mangino M, Brouillette S, Braund P, et al. A regulatory SNP of the BICD1 gene contributes to telomere length variation in humans. *Hum Mol Genet* 2008;17(16):2518–23.
- [28] Mangino M, Hwang SJ, Spector TD, et al. Genome-wide meta-analysis points to CTC1 and ZNF676 as genes regulating telomere homeostasis in humans. *Hum Mol Genet* 2012;21(24):5385–94.
- [29] Mangino M, Christiansen L, Stone R, et al. DCAF4, a novel gene associated with leukocyte telomere length. *J Med Genet* 2015;52(3):157–62.
- [30] Iles MM, Bishop DT, Taylor JC, Hayward NK, Brossard M, Cust AE, et al. The effect on melanoma risk of genes previously associated with telomere length. *J Natl Cancer Inst* 2014;106(10).
- [31] Bataille V, Sasieni P, Grulich A, et al. Solar keratoses: a risk factor for melanoma but negative association with melanocytic naevi. *Int J Cancer* 1998;78(1):8–12.
- [32] Whiteman DC, Watt P, Purdie DM, et al. Melanocytic nevi, solar keratoses, and divergent pathways to cutaneous melanoma. *J Natl Cancer Inst* 2003;95:806–12.
- [33] Kaszubowska L. Telomere shortening and ageing of the immune system. *J Physiol Pharmacol* 2008;59(Suppl. 9):169–86.
- [34] Seow WJ, Cawthon RM, Purdie MP, et al. Telomere length in white blood cell DNA and lung cancer: a pooled analysis of three prospective cohorts. *Cancer Res* 2014;74(15):4090–8.
- [35] Sanchez-Espiridon B, Chen M, Chang JY, et al. Telomere length in peripheral blood leukocytes and lung cancer risk: a large case-control study in Caucasians. *Cancer Res* 2014;74(9):2476–86.
- [36] Wei R, DeVilbiss FT, Liu W. Genetic polymorphism, telomere biology and non-small lung cancer risk. *J Genet Genomics* 2015;42(10):549–61.
- [37] Yoo SS, Do SK, Choi JE, et al. TERT polymorphism rs2853669 influences on lung cancer risk in the Korean population. *J Korean Med Sci* 2015;30(10):1423–8.
- [38] Walsh KM, Gorlov IP, Hansen HM, et al. Fine-mapping of the 5p15.33, 6p22.1-p21.31, and 15q25.1 regions identifies functional and histology-specific lung cancer susceptibility loci in African-Americans. *Cancer Epidemiol Biomarkers Prev* 2013 Feb;22(2):251–60.
- [39] Bau DT, Lippman SM, Xu E, et al. Short telomere lengths in peripheral blood leukocytes are associated with an increased risk of oral premalignant lesion and oral squamous cell carcinoma. *Cancer* 2013;119(24):4277–83.
- [40] Zhang Y, Sturgis EM, Dahlstrom KR, Wen J, Liu H, Wei Q, Li G, Liu Z. Telomere length in peripheral blood lymphocytes contributes to the development of

- HPV-associated oropharyngeal carcinoma. *Cancer Res* 2013 Oct 1;73 (19):5996–6003.
- [41] Shi J, Sun F, Peng L, et al. Leukocyte telomere length-related genetic variants in 1p34.2 and 14q21 loci contribute to the risk of esophageal squamous cell carcinoma. *Int J Cancer* 2013;132(12):2799–807.
- [42] Yu Q, Yang J, Liu B, et al. Combined effects of leukocyte telomere length, p53 polymorphism and human papillomavirus infection on esophageal squamous cell carcinoma in a Han Chinese population. *Cancer Epidemiol* 2014;38 (5):569–75.
- [43] Lin SW, Abnet CC, Freedman ND, et al. Measuring telomere length for the early detection of precursor lesions of esophageal squamous cell carcinoma. *BMC Cancer* 2013;13:578.
- [44] Du J, Xue W, Ji Y, et al. U-shaped association between telomere length and esophageal squamous cell carcinoma risk: a case-control study in Chinese population. *Front Med* 2015;9(4):478–86.
- [45] Steenbergen RD, Kramer D, Meijer CJ, et al. Telomerase suppression by chromosome 6 in a human papillomavirus type 16-immortalized keratinocyte cell line and in a cervical cancer cell line. *J Natl Cancer Inst* 2001;93 (11):865–72.
- [46] Qu S, Wen W, Shu XO, et al. Association of leukocyte telomere length with breast cancer risk: nested case-control findings from the Shanghai Women's Health Study. *Am J Epidemiol* 2013;177(7):617–24.
- [47] Weischer M, Nordestgaard BG, Cawthon RM, Freiberg JJ, Tybjærg-Hansen A, Bojesen SE. Short telomere length, cancer survival, and cancer risk in 47102 individuals. *J Natl Cancer Inst* 2013;105(7):459–68.
- [48] Lan Q, Cawthon R, Gao Y, et al. Longer telomere length in peripheral white blood cells is associated with risk of lung cancer and the rs2736100 (CLPTM1L-TERT) polymorphism in a prospective cohort study among women in China. *PLoS One* 2013;8(3):e59230.
- [49] Hofmann JN, Lan Q, Cawthon R, et al. A prospective study of leukocyte telomere length and risk of renal cell carcinoma. *Cancer Epidemiol Biomarkers Prev* 2013;22(5):997–1000.
- [50] Pooley KA, Bojesen SE, Weischer M, et al. A genome-wide association scan (GWAS) for mean telomere length within the COGS project: identified loci show little association with hormone-related cancer risk. *Hum Mol Genet* 2013;22(24):5056–64.
- [51] Zee RY, Castonguay AJ, Barton NS, Buring JE. Mean telomere length and risk of incident colorectal carcinoma: a prospective, nested case-control approach. *Cancer Epidemiol Biomarkers Prev* 2009;18(8):2280–2.
- [52] Julin B, Shui I, Heaphy CM, et al. Circulating leukocyte telomere length and risk of overall and aggressive prostate cancer. *Br J Cancer* 2015;112(4):769–76.
- [53] Sun L, Tan R, Xu J, et al. Targeted DNA damage at individual telomeres disrupts their integrity and triggers cell death. *Nucleic Acids Res* 2015;43(13):6334–47.
- [54] Skinner HG, Gangnon RE, Litzelman K, et al. Telomere length and pancreatic cancer: a case-control study. *Cancer Epidemiol Biomarkers Prev* 2012;21 (11):2095–100.
- [55] Cui Y, Cai Q, Qu S, et al. Association of leukocyte telomere length with colorectal cancer risk: nested case-control findings from the Shanghai Women's Health Study. *Cancer Epidemiol Biomarkers Prev* 2012;21 (10):1807–13.
- [56] Wang L, Xiao H, Zhang X, Wang C, Huang H. The role of telomeres and telomerase in hematologic malignancies and hematopoietic stem cell transplantation. *J Hematol Oncol* 2014;7:61.
- [57] Martin-Ruiz CM, Baird D, Roger L, et al. Reproducibility of telomere length assessment – an international collaborative study. *Int J Epidemiol* 2015;44 (5):1749–54.
- [58] Ding Z, Mangino M, Aviv A, Spector T, Durbin R, UK10K Consortium. Estimating telomere length from whole genome sequence data. *Nucleic Acids Res* 2014;42 (9):e75.
- [59] Barbaro P, Ziegler DS, Reddel RR. The wide-ranging clinical implications of the Short Telomere Syndromes. *Intern Med J* 2015.
- [60] Robles-Espinoza CD, Harland M, Ramsay AJ, et al. POT1 loss-of-function variants predispose to familial melanoma. *Nat Genet* 2014;46(5):478–81.
- [61] de Snoo FA, Bishop DT, Bergman W, et al. Increased risk of cancer other than melanoma in CDKN2A founder mutation (p16-Leiden)-positive melanoma families. *Clin Cancer Res* 2008;14(21):7151–7.
- [62] Goldstein AM, Chan M, Harland M, et al. High risk melanoma susceptibility genes and pancreatic cancer, neural system tumors and uveal melanomas across GenoMEL. *Cancer Res* 2006;15:9818–28.
- [63] Maubec E, Chaudru V, Mohamdi H, et al. Characteristics of the coexistence of melanoma and renal cell carcinoma. *Cancer* 2010;116(24):5716–24.
- [64] Monnerat C, Chompret A, Kannengiesser C, Avril MF, Janin N, Spatz A. BRCA1, BRCA2, TP53, and CDKN2A germline mutations in patients with breast cancer and cutaneous melanoma. *Fam Cancer* 2007;6(4):453–61.
- [65] Bradford PT, Freedman DM, Goldstein AM, Tucker MA. Increased risk of second primary cancers after a diagnosis of melanoma. *Arch Dermatol* 2010;146 (3):265–72.
- [66] Walsh KM, Codd V, Smirnov IV, Rice T, et al. Variants near TERT and TERC influencing telomere length are associated with high-grade glioma risk. *Nat Genet* 2014;46(7):731–5.
- [67] Xie H, Wu X, Wang S, et al. Long telomeres in peripheral blood leukocytes are associated with an increased risk of soft tissue sarcoma. *Cancer* 2013;119 (10):1885–91.
- [68] Heidenreich B, Rachakonda PS, Hosen I, et al. TERT promoter mutations and telomere length in adult malignant gliomas and recurrences. *Oncotarget* 2015;6(12):10617–33.
- [69] Goldstein AM, Cahn M, Harland M, et al. Features associated with germline CDKN2A mutations: a GenoMEL study of melanoma prone families from three continents. *J Med Genet* 2007;44:99–106.
- [70] Potrony M, Puig-Butillé JA, Aguilera P, et al. Increased prevalence of lung, breast, and pancreatic cancers in addition to melanoma risk in families bearing the cyclin-dependent kinase inhibitor 2A mutation: implications for genetic counseling. *J Am Acad Dermatol* 2014;71(5):888–95.
- [71] Chen Y, Qu F, He X, Bao G, Liu X, Wan S, Xing J. Short leukocyte telomere length predicts poor prognosis and indicates altered immune functions in colorectal cancer patients. *Ann. Oncol.* 2014;25(4):869–76.
- [72] Jia H, Wang Z. Telomere length as a prognostic factor for overall survival in colorectal cancer patients. *Cell Physiol Biochem* 2016;38(1):122–8.
- [73] Valls C, Piñol C, Reñé JM, Buenestado J, Viñas J. Telomere length is a prognostic factor for overall survival in colorectal cancer. *Colorectal Dis* 2011;13 (11):1265–72.
- [74] Ribero S, Davies JR, Requena C, et al. High nevus counts confer a favorable prognosis in melanoma patients. *Int J Cancer* 2015;137(7):1691–8.