



# UNIVERSITÀ DEGLI STUDI DI TORINO

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12

***This is an author version of the contribution published on:***

*[Biological Conservation, 211: 102-109, 2017,  
<http://dx.doi.org/10.1016/j.biocon.2017.05.013>]*

***The definitive version is available at:***

*[<http://www.sciencedirect.com/science/article/pii/S0006320716310229>]*

13 **Restoration treatments to control *Molinia arundinacea* and woody and alien species**  
14 **encroachment in *Calluna vulgaris* heathlands at the southern edge of their distribution**

15  
16 Massimiliano Probo<sup>a\*</sup>, Davide Ascoli<sup>b</sup>, Michele Lonati<sup>a</sup>, Raffaella Marzano<sup>a</sup> and Giampiero  
17 Lombardi<sup>a</sup>

18

19 <sup>a</sup> Department of Agricultural, Forest and Food Sciences, University of Torino, Largo Braccini 2,  
20 Grugliasco, I 10095, Italy

21 <sup>b</sup> Dipartimento di Agraria, University of Naples Federico II, via Università 100, 80055 Portici,  
22 Napoli, Italy

23 \* Corresponding author: Massimiliano Probo. e-mail: massimiliano.probo@unito.it. Phone number:  
24 +390116708790

25

26 **Abstract**

27 Throughout the last decades, *Calluna vulgaris* (L.) Hull heathlands have declined across Europe  
28 and nowadays their conservation is particularly challenging at the southern edge of their  
29 distribution. In the Nature Reserve of Vauda (north-western Italy), six restoration treatments were  
30 applied (extensive annual goat browsing, one-off mowing, annual mowing, one-off fire without and  
31 with annual browsing, and annual fire) and their effects on plant diversity and the cover of *C.*  
32 *vulgaris*, its competitor grass *Molinia arundinacea* Schrank, woody, and alien species were  
33 monitored between 2005 and 2011. In the short-term, most of the treatments changed the vegetation  
34 community, reducing *C. vulgaris* cover according to a gradient of increasing biomass removal. In  
35 the mid-term, *C. vulgaris*, *M. arundinacea*, woody and alien species cover followed different  
36 trajectories according to the treatment and functional group. Annual fire shifted the vegetation  
37 towards a *M. arundinacea*-dominated community, while extensive annual browsing did not affect  
38 the heathland community and resulted in the lowest increase in *M. arundinacea*, which showed a  
39 remarkable fitness in these environments. Moreover, annual burning and mowing were effective in  
40 reducing woody species encroachment ( $p < 0.05$ ), and fire treatments triggered a peak in alien  
41 species cover (mainly *Panicum acuminatum* Swartz) in the short-term. Six years after treatment,  
42 species richness and Shannon index did not differ between treated and control sites ( $p > 0.05$ ). In  
43 conclusion, these results highlight the need and potential benefit of integrating multiple techniques  
44 to preserve *C. vulgaris* heathlands at their southern edge.

45 **Keywords.** Goat browsing, mowing, *Panicum acuminatum*, plant diversity, prescribed burning.

## 46 1. Introduction

47 The conservation of threatened habitats is particularly challenging near their range edges, where  
48 populations are smaller, fragmented, and more vulnerable to environmental changes (Sexton et al.,  
49 2009). Techniques for habitat restoration, which are effective at the core of target species  
50 distribution, may have unexpected outcomes at the distribution edges. This is the case of *Calluna*  
51 *vulgaris* (L.) Hull heathlands, a key European cultural landscape and habitat (EU Council Directive,  
52 1992), developed after human mediated disturbance regimes, such as grazing, burning, and mowing  
53 (Davies et al., 2016; Fagundez, 2013). Nowadays, heathlands are declining in most European  
54 countries due to different drivers of land use and environmental change. The abandonment of  
55 traditional management has often led to their conversion to woodlands (Pywell et al., 2011).  
56 Moreover, increasing atmospheric nitrogen deposition has favored the replacement of *C. vulgaris*  
57 by grasses such as *Molinia* spp. (Bobbink et al., 2010; Terry et al., 2004).

58 Despite the extensive knowledge on heathland conservation measures in their main oceanic  
59 distribution area (Davies et al., 2016; Littlewood et al., 2014; Pywell et al., 2011), very little work is  
60 available for their southern edges (Fagundez, 2013). Here, these ecosystems often occur under  
61 Continental rather than Atlantic climates and mineral rather than thick organic soils (Lonati et al.,  
62 2009). At these southern margins, heathlands are facing major threats because of increased heath  
63 fragmentation, minor adaptive capacity, and higher pressure by local and exotic grass, shrub and  
64 tree encroachment (Bartolome et al., 2005; Borghesio, 2014). Indeed, woody encroachment happens  
65 at faster rates (Ascoli and Bovio 2010), and *C. vulgaris* competes with vicariant and more  
66 productive grasses, such as *Molinia arundinacea* Schrank rather than *Molinia caerulea* (L.) Moench  
67 (Borghesio et al., 2014; Daněák et al., 2012). Consequently, techniques that are effective in the re-  
68 establishment of the dominance of *C. vulgaris* (e.g. browsing, prescribed burning, mowing) may not  
69 successfully achieve the target of restoring the composition of the whole plant community  
70 (Littlewood et al., 2014) and may promote competitor and alien species (Davies et al., 2016).

71 To address these issues, a heathland restoration experiment was established in a highly  
72 fragmented, continental dry heathland located on Po Plain lowlands, northern Italy (Ascoli et al.,  
73 2009). The study aimed to assess the long-term effects (six years after treatments) of browsing,  
74 prescribed burning, and mowing for the restoration of heathland vegetation, by answering the  
75 following questions: i) what is the effect of single restoration techniques and their combination on  
76 plant diversity and species community assemblage? ii) How does restoration affect the cover of *C.*  
77 *vulgaris*, *M. arundinacea* and encroaching woody species? iii) Is there any restoration treatment  
78 that triggers the invasion of alien species?

79

## 80 2. Materials and Methods

### 81 2.1 Study area, experimental design, and vegetation surveys

82 The study area was located within the Nature Reserve of Vauda, northwest Italy (7°41'17"E,  
83 45°13'13"N), at an altitude ranging from 240 to 480 m a.s.l. The climate is continental, with 81%  
84 of mean annual precipitation (1 000-1 100 mm) falling between April and November and mean  
85 annual temperature about 12°C. The Reserve lies on a fluvio-glacial terrace, characterized by  
86 ancient and leached soils with low pH (4.8), high clay content, and a thin organic layer (Borghesio,  
87 2014). The Reserve was instituted in 1993 to maintain a relict heathland ecosystem. Despite  
88 protection policies, in the last decades the heathland has declined because of *M. arundinacea* and  
89 woody species encroachment (mainly European aspen *Populus tremula* L. and silver birch *Betula*  
90 *pendula* Roth) due to the abandonment of traditional management (i.e. grazing and mowing).  
91 Moreover, large and frequent pastoral uncontrolled fires during the winter dry season, when grasses  
92 dry out, threaten the heathland (Ascoli and Bovio, 2010).

93 The experimental area was composed of *C. vulgaris* stands in the building phase (*sensu*  
94 Watt, 1955) with an advanced encroachment of woody species, i.e. average ( $\pm$ SE) tree density and  
95 basal area were  $22,722 \pm 1518$  stems  $\text{ha}^{-1}$  and  $3.1 \pm 0.4$   $\text{m}^2$   $\text{ha}^{-1}$ , respectively (Ascoli et al., 2013).  
96 Six restoration treatments were applied from 2005 to 2011: 1) annual fire, 2) one-off fire, 3) annual

97 mowing, 4) one-off mowing, 5) extensive annual browsing, and 6) one-off fire + extensive annual  
98 browsing. Annual fire was implemented every winter from 2005 to mimic current pastoral  
99 practices, one-off fire once under prescribed burning conditions in winter 2005 (for details see  
100 Lonati et al., 2009), annual mowing every spring, and one-off mowing once in spring 2005.  
101 Mowing was performed mechanically at 8 cm height and included biomass harvesting. Annual and  
102 one-off fire were carried out over eight 600 m<sup>2</sup> plots each, while annual and one-off mowing over  
103 eight 100 m<sup>2</sup> plots each. A herd of about 100 goats exploited annual browsing and one-off fire +  
104 annual browsing plots (which received a single winter prescribed burn in 2005) for 3.5 h and 3 h  
105 day<sup>-1</sup>, respectively, over a period of four weeks between April and May. Annual browsing and one-  
106 off fire + annual browsing were carried out over 16 plots each (plots were 1250 m<sup>2</sup> and 1000 m<sup>2</sup>,  
107 respectively), with a stocking density of about 135 Animal Units ha<sup>-1</sup> and a stocking rate of 0.05  
108 AU ha<sup>-1</sup> year<sup>-1</sup> (*sensu* Allen et al., 2011). Moreover, eight untreated 300 m<sup>2</sup> plots were used as  
109 control areas. Since we expected a higher variability of vegetation cover and composition after  
110 treatment, due to the more heterogeneous effects produced by the selective feeding behavior of  
111 goats (Iussig et al., 2015), the number of plots for extensive annual browsing and one-off fire +  
112 extensive annual browsing was double compared to other treatments All 72 experimental plots were  
113 fenced and randomly selected within comparable *C. vulgaris* heathland patches, which were chosen  
114 on the basis of similar vegetation cover and composition.

115 In each plot, botanical composition was determined using the vertical point-quadrat method  
116 (Daget and Poissonet, 1971) along one fixed 10 m transect. In each transect, at 20 cm intervals, the  
117 species touching a steel needle were identified and recorded (i.e. 50 points of vegetation  
118 measurement). Since rare species are often missed by this method, a complete list of all other plant  
119 species included within a 1 m buffer around the transect line was also recorded (Orlandi et al.,  
120 2016). Vegetation surveys were conducted during summer 2004 (pre-treatment year), 2007, 2009,  
121 and 2011 (i.e. two, four, and six years after treatments, respectively).  
122

## 123 2.2 Statistical analyses

124 For each species recorded, the frequency of occurrence (number of occurrences/50 points) was  
125 calculated for each transect and converted to percentage cover (%) (Pittarello et al., 2016). In  
126 particular, the percentage cover of *C. vulgaris*, *M. arundinacea*, woody encroaching species, i.e.  
127 species classified as chamaephyte, phanerophyte, or nanophanerophyte according to Raunkiaer  
128 (1937), and alien species (Celesti-Gradow et al., 2009) was computed. Species richness and  
129 Shannon diversity index were also calculated for each survey.

130 A Principal Response Curve (PRC) analysis was performed to visualize the overall effect  
131 produced by treatments on the botanical composition of treated plots compared to that of control  
132 plots over time. The PRC analysis was performed using Canoco 4.5 software (Ter Braak and  
133 Šmilauer, 2009).

134 Generalized Linear Mixed Models (GLMMs) were used to test for differences of each  
135 treatment against control for all the vegetation variables (i.e. species richness, Shannon diversity  
136 index, *C. vulgaris* cover, *M. arundinacea* cover, woody species cover, and alien species cover) for  
137 each of the four years during which vegetation surveys were carried-out. Each treatment was  
138 considered as a fixed effect, with control used as a reference level for all the analyses. Poisson  
139 distribution was specified for count variables and Gaussian or Gamma distributions were specified  
140 for continuous data, depending if normality was met or not, respectively (normality was tested with  
141 Kolmogorov-Smirnoff test). Significance tests were performed using the Wald statistic. The  
142 GLMMs were carried out using R 3.0.3 (R Development Core Team, 2012), with the glmmADMB  
143 package (Fournier et al., 2012).  
144

## 145 3. Results and Discussion

146 A total of 66 plant species was detected in botanical surveys (Appendix 1). Six years after  
147 treatments, species richness did not differ between treated and control sites, underlying the high

148 stability and resistance to treatments of the floristic composition of *C. vulgaris* heathlands at their  
149 southern edge, though inter-annual fluctuation in species richness can occur among years (Figure  
150 1a). In the short-term, most of the treatments changed the heathland community, reducing *C.*  
151 *vulgaris* cover in 2007 (Figures 1c) according to a gradient of increasing biomass removal:  
152 extensive annual browsing (which removed little biomass), one-off mowing, annual mowing, one-  
153 off fire without and with annual browsing, and annual fire (which removed biomass repeatedly). In  
154 the mid-term, we observed changes in *C. vulgaris*, *M. arundinacea*, woody and alien species cover,  
155 which followed different trajectories according to treatment and functional group (Figure 1c-f).

156 *Molinia arundinacea* cover increased proportionally to biomass removal, as it produced  
157 more biomass in southern edge heathlands as compared to *M. caerulea* in the Atlantic European  
158 ones (Marrs et al., 2004) (Figure 1d). Annual fire, which simulated current uncontrolled pastoral  
159 fires, repeatedly removed the heathland and shifted vegetation towards a *M. arundinacea*-  
160 dominated community. Mowing treatments and one-off fire, combined or not with annual browsing,  
161 initially reduced *C. vulgaris* cover (Figures 1c) and increased *M. arundinacea* cover in the short-  
162 term (Figure 1d), but *C. vulgaris* started recovering at all sites after the first growing season mainly  
163 by stump resprouting. However, six years later, one-off fire, combined or not with annual browsing,  
164 displayed a lower *C. vulgaris* and a higher *M. arundinacea* cover in comparison to mowing  
165 treatments. Indeed, graminoids benefited from both litter and crown biomass consumption in fire  
166 treatments. Conversely, mowing did not completely remove the crown and left the litter, which  
167 resulted in a higher *C. vulgaris* cover since 2007. In the following years, the recovery rate of *C.*  
168 *vulgaris* in all these treatments was similar, but graminoids maintained a higher abundance in fire  
169 treatments, in contrast to herbaceous forbs, as evidenced in Figure 2. Notably, one-off mowing did  
170 not show significant differences in both *C. vulgaris* and *M. arundinacea* cover when compared to  
171 control plots. Extensive annual browsing, the treatment with the lowest biomass removal, did not  
172 affect the heathland structure as *C. vulgaris* is barely consumed by goats (Iussig et al., 2015), and it  
173 resulted in the lowest increase of *M. arundinacea*, which was comparable to the one of the control.  
174 Since heathland vegetation was always dominated by a low number of species (namely *C. vulgaris*,  
175 *M. arundinacea*, and a few other graminoids, Figure 2 and Appendix 1), a situation comparable to  
176 that of other heathlands (Hancock and Legg, 2012; Muñoz et al., 2012), Shannon diversity index  
177 was not different between treatments and control at the end of the experiment (Figure 1b).

178 Woody species displayed opposite responses to treatments in comparison to graminoids  
179 (Figure 1e). Annual burning and mowing were effective in reducing woody species encroachment.  
180 One-off fire and mowing top-killed trees, but aspen and birch sprouted vigorously and only  
181 subsequent annual browsing effectively controlled shoot growth. Annual browsing had a delayed  
182 effect and woody species cover reached the same level as one-off fire + annual browsing by the end  
183 of the study. PRC analysis (Figure 2) highlighted significant differences in the botanical  
184 composition between treated and untreated plots ( $p < 0.01$ ), with a marked increase in woody  
185 species cover in unmanaged control plots, as showed by the trend of phanerophytes (*P. tremula*,  
186 *Frangula alnus* Miller, and *B. pendula*).

187 Five alien species were inventoried, but only *Panicum acuminatum* Swartz reached a  
188 noticeable percentage cover (Appendix 1). Interestingly, *P. acuminatum* was triggered only by fire  
189 treatments, displayed a peak in the short-term and decreased to the level of control in 2011 (Figure  
190 1f), remaining higher only in the annual fire treatment (Figure 2). In North America *P. acuminatum*  
191 showed a great fitness after annual fire (Walsh, 1995), a trait maintained also outside its natural  
192 distribution area (Lonati et al., 2009). However, our results confirm that this invasive species  
193 quickly declines when fire frequency is low and it does not become important in terms of density  
194 and biomass (Walsh, 1995). Under a long-term perspective, prescribed burning might have the  
195 effect in rejuvenating the *P. acuminatum* seed bank rather than considerably increasing its  
196 vegetation cover, which is however unfavorable to the control of this alien species.

197

#### 198 4. Conclusions

199 Both frequent fires simulating current uncontrolled pastoral burns and lack of management  
200 promote heathland losses at the southern edge of their distribution, stressing the need for  
201 conservation measures. Moreover, the encroaching grass *Molinia arundinacea*, woody and alien  
202 species, such as *Populus tremula* and *Panicum acuminatum*, appear to have a high resilience to  
203 different restoration treatments. Results evidence the need and potential benefit of integrating  
204 multiple techniques to preserve southern edge fragmented heathlands. The restoration of these  
205 habitats may not be effective with just one of the tested treatments, since all the techniques involve  
206 trade-offs between undesired effects, efficacy and operational difficulties. However, six years after  
207 treatments, extensive goat browsing and annual mowing provided the best results for the  
208 maintenance of *Calluna vulgaris* and kept woody and alien species under a critical level. Likewise  
209 in Atlantic heathlands, prescribed burning may be also valuable for *Calluna* heathlands restoration  
210 at their southern range, but only when applied with long return intervals (i.e. longer than six years,  
211 but further research is needed to establish a suitable return interval). A higher caution in the use of  
212 fire is mandatory because of the presence of encroaching species with marked fire-traits adapted to  
213 a more fire-prone environment when compared to Atlantic regions, which can benefit greatly from  
214 repeated burns at the expense of *Calluna* heaths.

215

## 216 **5. Acknowledgements**

217 We thank Vauda managers (Antonio Aschieri, Andrea Maccioni) for technical support and setting  
218 the experimental design, the State Forestry Corp (Diego Noveri) and Regione Piemonte Fire  
219 Fighting Volunteers for logistic support in prescribed burning, Davide Cugno and Alessandra  
220 Gorlier for fieldwork assistance.

221

## 222 **Role of the funding source**

223 The work was funded by Regione Piemonte, but experimental design, data collection, analysis and  
224 discussion, reporting, and the decision to submit this article for publication are exclusively  
225 attributable to authors.

226


227

228 **6. References**


- 229 Allen, V.G., Batello, C., Berretta, E.J., Hodgson, J., Kotchmann, M., Li, X., McIvor, J., Milne, J.,  
230 Morris, C., Peeters, A., Sanderson, M., 2011. An international terminology for grazing lands  
231 and grazing animals. *Grass Forage Sci.* 66, 2-28.
- 232 Ascoli, D., Beghin, R., Ceccato, R., Gorlier, A., Lombardi, G., Lonati, M., Marzano, R., Bovio, G.,  
233 Cavallero, A., 2009. Developing an Adaptive Management approach to prescribed burning:  
234 a long-term heathland conservation experiment in north-west Italy. *Int J Wildland Fire.* 18,  
235 727-735.
- 236 Ascoli, D., Bovio, G., 2010. Tree encroachment dynamics in heathlands of north-west Italy: the fire  
237 regime hypothesis. *iForest.* 3, 137-143.
- 238 Ascoli, D., Lonati, M., Marzano, R., Bovio, G., Cavallero, A., Lombardi, G., 2013. Prescribed  
239 burning and browsing to control tree encroachment in southern European heathlands. *Forest*  
240 *Ecol Manag.* 289, 68-77.
- 241 Bartolome, J., Plaixats, J., Fanlo, R., Boada, M., 2005. Conservation of isolated Atlantic heathlands  
242 in the Mediterranean region: effects of land-use changes in the Montseny biosphere reserve  
243 (Spain). *Biol. Conserv.* 122, 81-88.
- 244 Bobbink, R., Hicks, K., Galloway, J., Spranger, T., Alkemade, R., Ashmore, M., Bustamante, M.,  
245 Cinderby, S., Davidson, E., Dentener, F., Emmett, B., Erisman, J.W., Fenn, M., Gilliam, F.,  
246 Nordin, A., Pardo, L., De Vries, W., 2010. Global assessment of nitrogen deposition effects  
247 on plant terrestrial biodiversity: a synthesis. *Ecol. Appl.* 20, 30–59.
- 248 Borghesio, L. (2014). Can fire avoid massive and rapid habitat change in Italian heathlands? *J Nat*  
249 *Conserv.* 22(1), 68-74.
- 250 Celesti-Grapow, L., Alessandrini, A., Arrigoni, P.V., Banfi, E., Bernardo, L., Bovio, M., Brundu,  
251 G., Cagiotti, M.R., Camarda, I., Carli, E., Conti, F., Fascetti, S., Galasso, G., Gubellini, L.,  
252 La Valva, V., Lucchese, F., Marchiori, S., Mazzola, P., Peccenini, S., Poldini, L., Pretto, F.,  
253 Prosser, F., Siniscalco, C., Villani, M.C., Viegi, L., Wilhalm, T., Blasi, C., 2009. The  
254 inventory of the non-native flora of Italy. *Plant Biosyst.* 143, 386–430.
- 255 Daget, P., Poissonet, J., 1971. Une méthode d'analyse phytologique des prairies. *Ann. Agron.* 22, 5–  
256 41.
- 257 Daněák, M., Duchoslav, M., Trávníček, B., 2012. Taxonomy and cytogeography of the *Molinea*  
258 *caerulea* complex in central Europe. *Preslia,* 84, 351-374.
- 259 Davies, G. M., Kettridge, N., Stoof, C. R., Gray, A., Ascoli, D., Fernandes, P. M., Marss, R., Allen,  
260 K. A., Doerr, S. H., Clay, G. D., McMorrow, J., 2016. The role of fire in UK peatland and  
261 moorland management: the need for informed, unbiased debate. *Phil. Trans. R. Soc.*  
262 *B.* 371(1696), 20150342.
- 263 EU Council Directive, 1992. Natura 2000 Network on the Conservation of Natural Habitats and of  
264 Wild Fauna. COUNCIL DIRECTIVE 92/43/EEC (1) of 21 May 1992. The Council of the  
265 European Communities.
- 266 Fagundez, J., 2013. Heathlands confronting global change: drivers of biodiversity loss from past to  
267 future scenarios. *Ann Bot-Lonon.* 111, 151-172.
- 268 Fournier, D.A., Skaug, H.J., Ancheta, J., Ianelli, J., Magnusson, A., Maunder, M.N., Nielsen, A.,  
269 Sibert, T.J. 2012. AD Model Builder: using automatic differentiation for statistical inference  
270 of highly parameterized complex nonlinear models. *Optim Method Softw.* 27, 233–249.
- 271 Hancock, M. H., Legg, C. L., 2012. Diversity and stability of ericaceous shrub cover during two  
272 disturbance experiments: one on heathland and one in forest. *Plant Ecol. Diversity.* 5, 275-  
273 287.
- 274 Iussig, G., Lonati, M., Probo, M., Lombardi, G., 2015. Plant species selection by goats foraging on  
275 montane semi-natural grasslands and grazable forestlands in the Italian alps. *Ital J Anim Sci.*  
276 14, 484-494.



277 Littlewood, N.A., Greenwood, S., Quin, S.L.O., Pakeman, R.J., Woodin, S.J., 2014. Long-term  
 278 trends in restored moorland vegetation assemblages. *Community Ecol.* 15, 104-112.  
 279 Lonati, M., Gorlier, A., Ascoli, D., Marzano, R., Lombardi, G., 2009. Response of the alien species  
 280 *Panicum acuminatum* to disturbance in an Italian lowland heathland. *Bot Helv.* 119, 105-  
 281 111.  
 282 Marrs, R. H., Philipps, J. D. P., Todd P. A., Ghorbani J., Le Duc, M. G. 2004. Control of *Molinia*  
 283 *caerulea* on upland moors. *J. Appl. Ecol.* 41, 398-411.  
 284 Muñoz, A., García-Duro, J., Álvarez, R., Pesqueira, X. M., Reyes, O., Casal, M., 2012. Structure  
 285 and diversity of *Erica ciliaris* and *Erica tetralix* heathlands at different successional stages  
 286 after cutting. *J Environ Manage.* 94, 34-40.  
 287 Orlandi, S., Probo, M., Sitzia, T., Trentanovi, G., Garbarino, M., Lombardi, G., Lonati, M., 2016.  
 288 Environmental and land use determinants of grassland patch diversity in the western and  
 289 eastern Alps under agro-pastoral abandonment. *Biodivers. Conserv.* 25(2), 275-293.  
 290 Pittarello, M., Probo, M., Lonati, M., Lombardi, G., 2016. Restoration of sub-alpine shrub-  
 291 encroached grasslands through pastoral practices: effects on vegetation structure and  
 292 botanical composition. *Appl Veg Sci.* 19(3), 381-390.  
 293 Pywell, R.F., Meek, W.R., Webb, N.R., Putwain, P.D., Bullock, J.M., 2011. Long-term heathland  
 294 restoration on former grassland: The results of a 17-year experiment. *Biol. Conserv.* 144,  
 295 1602-1609.  
 296 R Development Core Team, 2012. R Development Core Team, Vienna, Austria.  
 297 Raunkiaer, C., 1937. *Plant Life Forms*. The Clarendon Press, Oxford.  
 298 Sexton, J.P., McIntyre, P.J., Angert, A.L., Rice, K.J. 2009. Evolution and ecology of species range  
 299 limits. *Annu Rev Ecol Syst.* 40, 415-436.  
 300 Ter Braak, C.J.F., Šmilauer, P., 2009. *CANOCO 4.5*. Biometrics - Plant Research International,  
 301 Wageningen, the Netherlands.  
 302 Terry, A.C., Ashmore, M.R., Power, S.A., Allchin, E.A., Heil, G.W., 2004. Modelling the impacts  
 303 of atmospheric nitrogen deposition on *Calluna*-dominated ecosystems in the UK. *J. Appl.*  
 304 *Ecol.* 41, 897-909.  
 305 Walsh, R. A., 1995. *Dichanthelium acuminatum*. In: *Fire Effects Information System*, [Online].  
 306 Watt, A. S., 1955. Bracken versus heather, a study in plant sociology. *J Ecol.* 43(2), 490-506.  
 307



309  
 310 **Fig. 1.** a) Species richness, b) Shannon diversity index, c) *Calluna vulgaris* cover (%), d) *Molinia*  
 311 *arundinacea* cover (%), e) woody species cover (%), and f) alien species cover (%) of *C.*  
 312 *vulgaris* heathlands, untreated (i.e. control plots) or subjected to six restoration treatments. †  
 313 =  $p < 0.1$ ; \* =  $p < 0.05$ ; \*\* =  $p < 0.01$ ; \*\*\* =  $p < 0.001$ . Statistical significance values are  
 314 related to differences of each treatment against control. ■ = annual fire; □ = one-off fire;  
 315 ◆ = one-off fire + annual goat browsing; ▲ = annual mowing; △ = one-off mowing; ● =  
 316 annual goat browsing; ○ = untreated.



317  
 318 **Fig. 2.** PRC (axis 1) showing effects on botanical composition of *C. vulgaris* heathlands produced  
 319 by the implementation of six restoration treatments with respect to untreated control (control  
 320 line) from pre-treatment stage (2004) to 2011. The scores of the most frequent species  
 321 (namely, species present in at least the 14 plots, i.e. the 5% of total plots) are shown on the  
 322 right side of the graph: positive values represent species whose canopy cover increased after  
 323 treatments, whereas negative values represent species whose canopy cover decreased over  
 324 time. Letters within brackets indicate: G = graminoid, F = herbaceous forb, NP =  
 325 nanophanerophyte, CH = chamaephyte, P = phanerophyte. ■ = annual fire; □ = one-off fire;  
 326 ◆ = one-off fire + annual goat browsing; ▲ = annual mowing; △ = one-off mowing; ● =  
 327 annual goat browsing; ○ = untreated.

328

329

## 8. Appendix

330

**Appendix 1.** Species frequency, name, growth form, life form, origin, mean cover under different restoration treatments and years (2004, i.e. pre-treatment year, 2007, 2009, and 2011).

331

Species frequency (%)	Species name	Growth forms (Raunkiaer, 1937)	Life form	Origin	Mean species cover (%)																												
					Annual fire				One-off fire				Fire + browsing				Annual mowing				One-off mowing				Annual browsing				Control				
					2004	2007	2009	2011	2004	2007	2009	2011	2004	2007	2009	2011	2004	2007	2009	2011	2004	2007	2009	2011	2004	2007	2009	2011	2004	2007	2009	2011	
100.0	<i>Molinia arundinacea</i>	Hemicryptophyte	Graminoid	Native	68.8	97.0	91.0	96.8	73.3	94.0	91.0	94.8	77.3	94.8	94.5	94.0	84.3	92.5	86.3	91.3	77.8	89.5	90.5	91.3	72.3	81.4	80.3	85.0	72.8	80.5	83.5	85.3	
96.5	<i>Calluna vulgaris</i>	Chamaephyte	Forb	Native	50.0	0.6	0.2	2.2	70.5	21.3	39.3	45.8	65.8	14.8	23.4	32.3	53.0	38.8	48.8	56.5	65.0	51.0	59.3	64.5	60.3	74.1	72.9	65.8	66.0	77.5	78.3	75.3	
93.1	<i>Potentilla erecta</i>	Hemicryptophyte	Forb	Native	1.4	3.6	5.5	4.0	0.9	4.3	4.1	7.3	1.8	8.1	5.5	6.7	1.0	2.4	1.7	1.4	0.5	3.0	1.8	2.8	2.2	2.0	1.7	2.1	1.8	3.8	2.4	2.1	
92.7	<i>Frangula alnus</i>	Phanerophyte	Forb	Native	0.5	1.1	0.6	1.2	0.9	1.2	0.9	0.9	2.8	2.3	1.5	1.7	2.4	0.3	0.3	0.3	2.9	1.9	0.9	0.5	1.0	2.7	3.0	2.6	1.9	2.6	4.9	4.4	
92.4	<i>Carex panicea</i>	Geophyte	Graminoid	Native	2.8	11.8	18.8	43.8	2.5	16.3	9.8	23.0	3.3	17.9	17.6	38.9	3.6	22.0	30.3	39.3	2.8	15.8	7.5	17.3	4.4	7.6	8.1	23.9	2.8	14.5	7.0	10.5	
86.5	<i>Populus tremula</i>	Phanerophyte	Forb	Native	4.6	9.5	5.0	7.0	4.3	8.5	9.0	11.0	6.4	8.3	9.1	10.5	3.3	5.8	3.5	6.3	3.4	11.0	12.0	16.5	7.0	18.2	10.1	9.6	8.8	17.3	24.0	21.5	
58.0	<i>Serratula tinctoria</i>	Hemicryptophyte	Forb	Native	0.2	0.2	0.2	0.4	0.2	0.4	0.3	0.2	0.2	0.5	0.2	0.2	0.2	0.6	0.4	0.2	0.2	0.7	0.4	0.1	0.6	0.4	0.5	0.3	0.2	1.6	0.4	0.3	
55.6	<i>Salix rosmarinifolia</i>	Nanophanerophyte	Forb	Native	1.4	1.7	1.6	0.6	0.6	7.8	8.8	6.3	2.3	1.8	1.7	3.0	2.1	1.8	2.3	1.0	4.0	3.5	3.8	4.0	0.9	1.0	0.7	1.0	2.3	3.8	2.5	2.8	
54.5	<i>Panicum acuminatum</i>	Therophyte	Graminoid	Alien	0.5	24.3	6.1	7.4	0.5	18.5	0.6	0.1	0.9	29.3	2.8	1.1	-	0.3	4.6	1.5	-	1.0	0.2	-	1.4	2.7	1.1	0.9	0.3	1.0	-	-	
53.1	<i>Danthonia decumbens</i>	Hemicryptophyte	Graminoid	Native	0.1	0.1	6.1	1.0	1.0	0.1	3.6	1.8	0.7	1.1	6.7	3.3	0.4	-	1.2	0.9	-	0.6	2.2	1.3	1.1	0.6	0.5	0.5	0.3	0.5	1.3	0.3	
51.7	<i>Betula pendula</i>	Phanerophyte	Forb	Native	2.7	0.8	0.4	0.3	0.3	0.2	0.4	0.4	1.1	0.7	0.3	0.2	4.0	2.8	0.4	0.1	1.9	2.2	1.1	1.8	0.4	0.7	0.1	1.0	-	0.7	1.3	0.2	
37.2	<i>Carex lumidicarpa</i>	Hemicryptophyte	Graminoid	Native	1.5	1.0	7.5	-	-	3.8	7.8	0.3	0.2	1.5	3.4	-	0.3	1.0	2.8	-	0.3	0.9	5.0	0.8	0.1	0.2	1.2	0.1	1.3	0.5	0.1	0.3	
33.7	<i>Peucedanum cervaria</i>	Hemicryptophyte	Forb	Native	0.3	-	-	-	0.9	0.8	0.3	-	0.9	0.6	0.5	0.3	0.9	0.8	1.0	0.4	0.8	0.6	1.1	0.1	0.4	0.2	0.2	-	1.3	0.2	0.6	-	
32.6	<i>Quercus robur</i>	Phanerophyte	Forb	Native	0.8	0.2	0.2	0.2	0.1	0.2	0.6	0.6	0.4	0.1	0.1	0.1	0.8	0.3	0.3	0.6	0.1	0.1	0.2	0.1	0.5	0.2	0.1	-	0.8	0.3	1.9	2.0	
31.9	<i>Melampyrum pratense</i>	Therophyte	Forb	Native	0.1	0.6	1.5	-	0.1	0.2	-	-	0.1	1.6	0.1	-	0.1	0.2	-	-	0.1	4.8	0.6	-	0.1	2.0	0.1	0.2	-	0.9	0.3	-	
29.9	<i>Agrostis tenuis</i>	Hemicryptophyte	Graminoid	Native	-	1.7	0.7	1.1	-	1.8	0.4	0.4	-	2.1	0.2	0.5	-	-	0.3	0.8	-	0.1	0.5	-	-	0.2	-	0.1	0.3	-	-	-	
28.8	<i>Festuca tenuifolia</i>	Hemicryptophyte	Graminoid	Native	-	1.8	2.8	3.8	-	1.9	1.8	1.8	0.1	2.1	0.1	2.4	-	0.5	0.8	1.8	-	1.3	2.3	2.0	-	0.2	0.4	1.4	0.3	1.3	0.3	1.0	
25.7	<i>Rosa gallica</i>	Nanophanerophyte	Forb	Native	0.1	0.4	0.4	0.2	0.1	-	0.3	0.3	0.6	0.1	0.2	-	-	-	-	-	-	-	-	-	0.3	0.6	0.4	0.7	0.1	0.1	0.1	0.3	
22.9	<i>Genista tinctoria</i>	Chamaephyte	Forb	Native	-	0.5	0.9	-	1.0	1.3	0.8	0.1	0.6	1.3	0.3	0.1	0.5	0.3	0.3	-	0.3	1.0	0.1	0.1	0.1	-	0.3	0.2	1.8	2.5	0.3	0.3	
22.2	<i>Stachys officinalis</i>	Hemicryptophyte	Forb	Native	-	0.9	0.1	0.3	-	0.1	-	-	0.6	0.1	0.1	-	0.6	0.3	-	-	0.1	-	0.1	-	-	0.2	-	-	-	-	-	-	
21.5	<i>Genista germanica</i>	Chamaephyte	Forb	Native	0.1	0.3	0.1	-	0.5	0.1	0.1	0.8	0.1	0.5	0.2	0.1	0.3	0.1	-	-	0.1	0.6	0.3	-	0.2	0.1	-	0.1	0.5	1.0	0.3	0.3	
12.2	<i>Nardus stricta</i>	Hemicryptophyte	Graminoid	Native	2.3	0.3	1.0	-	8.5	-	-	0.1	-	0.1	-	-	0.3	0.3	0.1	-	-	0.1	-	-	4.5	2.3	1.3	0.5	2.5	1.0	-	-	
10.8	<i>Rubus ulmifolius</i>	Nanophanerophyte	Forb	Native	-	0.3	0.3	0.1	-	-	-	-	0.1	-	0.1	-	-	-	-	-	-	-	-	-	0.1	0.1	0.1	-	-	-	-	-	
10.4	<i>Peucedanum oroselinum</i>	Hemicryptophyte	Forb	Native	-	0.1	0.3	-	1.3	-	-	0.1	0.3	-	-	-	-	-	-	-	-	-	-	-	0.2	-	-	1.5	1.8	0.3	0.5	-	
9.0	<i>Plantago serpentina</i>	Hemicryptophyte	Forb	Native	-	-	0.1	-	0.8	-	-	0.3	-	-	-	-	-	-	-	-	-	-	-	0.6	0.3	0.1	-	-	0.8	0.3	-	-	
8.3	<i>Carex umbrosa</i>	Hemicryptophyte	Graminoid	Native	-	-	-	1.5	-	-	-	3.3	-	-	-	1.3	-	-	-	-	-	-	-	0.8	0.1	-	-	0.5	-	1.0	-	-	
8.3	<i>Succisa pratensis</i>	Hemicryptophyte	Forb	Native	-	0.1	0.1	-	0.1	-	0.3	0.1	-	-	-	-	-	0.1	0.1	-	-	-	-	0.1	0.1	-	-	-	-	-	-	-	
6.6	<i>Gentiana pneumonanthes</i>	Hemicryptophyte	Forb	Native	-	0.1	-	-	0.1	-	-	-	-	-	-	-	-	-	-	-	-	-	0.3	-	0.1	-	-	-	0.5	-	-	-	
5.2	<i>Chrysopogon gryllus</i>	Hemicryptophyte	Graminoid	Native	-	-	-	-	-	-	-	-	0.1	0.4	0.3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
4.5	<i>Aristida gracilis</i>	Therophyte	Graminoid	Alien	-	0.5	-	0.3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.1	-	0.4	-	-	-	-	-	
4.2	<i>Lotus corniculatus</i>	Hemicryptophyte	Forb	Native	-	-	-	-	-	-	-	-	0.1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
3.8	<i>Crataegus monogyna</i>	Phanerophyte	Forb	Native	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.3	-	-	-	-	-	-	-	-	
3.8	<i>Glaucium palustris</i>	Geophyte	Forb	Native	-	0.1	-	-	-	-	-	-	0.1	-	-	-	0.3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
3.8	<i>Inula hirta</i>	Hemicryptophyte	Forb	Native	-	-	-	-	-	-	-	0.1	-	-	-	-	-	-	0.3	-	-	-	-	-	-	-	-	0.3	0.1	-	-	-	
3.1	<i>Salix caprea</i>	Phanerophyte	Forb	Native	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
2.8	<i>Juncus conglomeratus</i>	Hemicryptophyte	Graminoid	Native	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.1	-	-	-	-	-	-	-	-	-	
2.1	<i>Leontodon hispidus</i>	Hemicryptophyte	Forb	Native	-	-	-	-	-	-	-	0.4	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.8	0.3	-	0.8	-	
1.7	<i>Carex acutiformis</i>	Geophyte	Graminoid	Native	-	-	-	-	-	-	-	0.5	-	-	-	-	-	-	-	-	-	-	-	0.5	-	-	1.5	-	-	-	-	-	
1.7	<i>Galium verum</i>	Therophyte	Forb	Native	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
1.7	<i>Hieracium umbellatum</i>	Hemicryptophyte	Forb	Native	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.1	-	-	-	-	-	-	-	-	
1.7	<i>Prunella grandiflora</i>	Hemicryptophyte	Forb	Native	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
1.7	<i>Viburnum opulus</i>	Phanerophyte	Forb	Native	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
1.4	<i>Carex pallidiflora</i>	Hemicryptophyte	Graminoid	Native	-	-	-	-	0.3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.3	-	-	-	-	
1.4	<i>Corylus avellana</i>	Phanerophyte	Forb	Native	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
1.4	<i>Eleocharis carniolica</i>	Hemicryptophyte	Graminoid	Native	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.3	-	-	-	-	-	-	-	-	-	-
1.4	<i>Euphorbia flavicoma</i>	Chamaephyte	Forb	Native	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
1.4	<i>Gratiola officinalis</i>	Hemicryptophyte	Forb	Native	1.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1.0	-	-	-	-	-
1.4	<i>Vincetoxicum hircundinaria</i>	Hemicryptophyte	Forb	Native	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1.0	<i>Juncus effusus</i>	Hemicryptophyte	Graminoid	Native	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.1	-	-	-	-	-	-	-	-	-	-
1.0	<i>Lysimachia vulgaris</i>	Hemicryptophyte	Forb	Native	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.8	-	-	-	-	-	-	-	-	-	-
0.7	<i>Carex caryophylla</i>	Hemicryptophyte	Graminoid	Native	0.3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.1	-	-	-	-	-	-	-	-
0.7	<i>Fraxinus excelsior</i>	Phanerophyte	Forb																														

333  
334  
335  
336  
337  
338  
339  
340

## **9. Highlights**

1. Six restoration treatments were applied in southern *Calluna vulgaris* heathlands
2. Annual fire shifted the vegetation towards a *Molinia arundinacea*-dominated community
3. In the short-term, fire treatments triggered a peak in alien species cover
4. Six years after treatments, plant diversity did not differ between treated and control sites
5. Six years after treatments, goat browsing and annual mowing provided the best results