


# Accepted Manuscript

A Prospective International *Aspergillus terreus* Survey: An EFISG, ISHAM and ECMM Joint Study

Brigitte Risslegger, Tamara Zoran, Michaela Lackner, Maria Aigner, Ferran Sánchez-Reus, Antonio Rezusta, Anuradha Chowdhary, Saad Jaber Taj-Aldeen, Maiken C. Arendrup, Salvatore Oliveri, Dimitrios P. Kontoyiannis, Ana Alastruey-Izquierdo, Katrien Lagrou, Giuliana Lo Cascio, Jacques F. Meis, Walter Buzina, Claudio Farina, Miranda Drogari-Apiranthitou, Anna Grancini, Anna Maria Tortorano, Birgit Willinger, Axel Hamprecht, Elizabeth Johnson, Lena Klingspor, Valentina Arsic-Arsenijevic, Oliver A. Cornely, Joseph Meletiadis, Wolfgang Prammer, Vivian Tullio, Jörg-Janne Vehreschild, Laura Trovato, Russell E. Lewis, Esther Segal, Peter-Michael Rath, Petr Hamal, Manuel Rodriguez-Iglesias, Emmanuel Roilides, Sevtap Arikan-Akdagli, Arunaloke Chakrabarti, Arnaldo L. Colombo, Mariana S. Fernández, M. Teresa Martin-Gomez, Hamid Badali, Georgios Petrikkos, Nikolai Klimko, Sebastian M. Heimann, Jos Houbraeken, Omrum Uzun, Michael Edlinger, Sonia de la Fuente, Cornelia Lass-Flörl



PII: S1198-743X(17)30220-3

DOI: [10.1016/j.cmi.2017.04.012](https://doi.org/10.1016/j.cmi.2017.04.012)

Reference: CMI 924

To appear in: *Clinical Microbiology and Infection*

Received Date: 24 February 2017

Revised Date: 7 April 2017

Accepted Date: 9 April 2017

Please cite this article as: Risslegger B, Zoran T, Lackner M, Aigner M, Sánchez-Reus F, Rezusta A, Chowdhary A, Taj-Aldeen SJ, Arendrup MC, Oliveri S, Kontoyiannis DP, Alastruey-Izquierdo A, Lagrou K, Lo Cascio G, Meis JF, Buzina W, Farina C, Drogari-Apiranthitou M, Grancini A, Tortorano AM, Willinger B, Hamprecht A, Johnson E, Klingspor L, Arsic-Arsenijevic V, Cornely OA, Meletiadis J, Prammer W, Tullio V, Vehreschild J-J, Trovato L, Lewis RE, Segal E, Rath P-M, Hamal P, Rodriguez-Iglesias M, Roilides E, Arikan-Akdagli S, Chakrabarti A, Colombo AL, Fernández MS, Martin-Gomez MT, Badali H, Petrikkos G, Klimko N, Heimann SM, Houbraeken J, Uzun O, Edlinger M, Fuente Sdl, Lass-Flörl C, A Prospective International *Aspergillus terreus* Survey: An EFISG, ISHAM and ECMM Joint Study, *Clinical Microbiology and Infection* (2017), doi: 10.1016/j.cmi.2017.04.012.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## 1 RESEARCH NOTE

2

3 A Prospective International *Aspergillus terreus* Survey:

4 An EFISG, ISHAM and ECMM Joint Study

5

6

7 Brigitte Risslegger<sup>1\*</sup>, Tamara Zoran<sup>1\*</sup>, Michaela Lackner<sup>1</sup>,  
8 Maria Aigner<sup>1</sup>, Ferran Sánchez-Reus<sup>2</sup>, Antonio Rezusta<sup>3</sup>,  
9 Anuradha Chowdhary<sup>4</sup>, Saad Jaber Taj-Aldeen<sup>5</sup>, Maiken C.  
10 Arendrup<sup>6</sup>, Salvatore Oliveri<sup>7</sup>, Dimitrios P. Kontoyiannis<sup>8</sup>, Ana  
11 Alastruey-Izquierdo<sup>9</sup>, Katrien Lagrou<sup>10</sup>, Giuliana Lo Cascio<sup>11</sup>,  
12 Jacques F. Meis<sup>12</sup>, Walter Buzina<sup>13</sup>, Claudio Farina<sup>14</sup>, Miranda  
13 Drogari-Apiranthitou<sup>15</sup>, Anna Grancini<sup>16</sup>, Anna Maria  
14 Tortorano<sup>17</sup>, Birgit Willinger<sup>18</sup>, Axel Hamprecht<sup>19</sup>, Elizabeth  
15 Johnson<sup>20</sup>, Lena Klingspor<sup>21</sup>, Valentina Arsic-Arsenijevic<sup>22</sup>,  
16 Oliver A. Cornely<sup>23</sup>, Joseph Meletiadis<sup>24</sup>, Wolfgang Prammer<sup>25</sup>,  
17 Vivian Tullio<sup>26</sup>, Jörg-Janne Vehreschild<sup>27</sup>, Laura Trovato<sup>28</sup>,  
18 Russell E. Lewis<sup>29</sup>, Esther Segal<sup>30</sup>, Peter-Michael Rath<sup>31</sup>, Petr  
19 Hamal<sup>32</sup>, Manuel Rodriguez-Iglesias<sup>33</sup>, Emmanuel Roilides<sup>34</sup>,  
20 Sevtap Arikan-Akdagli<sup>35</sup>, Arunaloke Chakrabarti<sup>36</sup>, Arnaldo L.  
21 Colombo<sup>37</sup>, Mariana S. Fernández<sup>38</sup>, M. Teresa Martin-  
22 Gomez<sup>39</sup>, Hamid Badali<sup>40</sup>, Georgios Petrikos<sup>41</sup>, Nikolai  
23 Klimko<sup>42</sup>, Sebastian M. Heimann<sup>43</sup>, Jos Houbraken<sup>44</sup>, Omrum

24 Uzun<sup>45</sup>, Michael Edlinger<sup>46</sup>, Sonia de la Fuente<sup>47</sup>, Cornelia Lass-

25 Flörl<sup>1</sup>

26

27 \* contributed equally

28

29 Author affiliations:

30 <sup>1</sup>Division of Hygiene and Medical Microbiology, Medical

31 University of Innsbruck, Innsbruck/Austria, <sup>2</sup>Servei de

32 Microbiologia, Hospital de la Santa Creu I Sant Pau,

33 Barcelona/Spain, <sup>3</sup>Microbiologia, Hospital Universitario Miguel

34 Servet, IIS Aragon, Universidad de Zaragoza, Zaragoza/Spain,

35 <sup>4</sup>Department of Medical Mycology, Vallabhbhai Patel Chest

36 Institute, University of Delhi, Delhi/India, <sup>5</sup>Microbiology

37 Division, Department of Laboratory Medicine and Pathology,

38 Hamad Medical Corporation, Doha/Qatar, <sup>6</sup>Statens Serum

39 Institute, Unit of Mycology, Copenhagen/Denmark &

40 Department of Clinical Microbiology, Copenhagen University,

41 Rigshospitalet, Copenhagen/Denmark, <sup>7</sup>Department of

42 Biomedical and Biotechnological Sciences, University of

43 Catania, Catania/Italy, <sup>8</sup>The University of Texas MD Anderson

44 Cancer Center, Houston, Texas/USA, <sup>9</sup>National Centre for

45 Microbiology, Instituto de Salud Carlos III, Madrid/Spain,

46 <sup>10</sup>Department of Microbiology and Immunology, KU Leuven,

47 Leuven/Belgium, <sup>11</sup>Unità Operativa Complessa di Microbiologia

48 e virologia, Dipartimento di Patologia e diagnostica, Azienda  
49 Ospedaliera Universitaria Integrata, Verona/Italy,  
50 <sup>12</sup>Department of Medical Microbiology and Infectious  
51 Diseases, Canisius Wilhelmina Hospital, Nijmegen/The  
52 Netherlands, <sup>13</sup>Institute of Hygiene, Microbiology and  
53 Environmental Medicine, Medical University of Graz,  
54 Graz/Austria, <sup>14</sup>Microbiology Institute, ASST Papa Giovanni  
55 XXIII, Bergamo/Italy, <sup>15</sup>Infectious Diseases Research  
56 Laboratory/4<sup>th</sup> Department of Internal Medicine, ATTIKON  
57 University Hospital, National and Kapodistrian University of  
58 Athens, Athens/Greece, <sup>16</sup>Laboratorio Centrale di Analisi  
59 Chimico Cliniche e Microbiologia, IRCCS Foundation, Cà  
60 Granda Ospedale Maggiore Policlinico, Milan/Italy,  
61 <sup>17</sup>Department of Biomedical Sciences for Health, Università  
62 degli Studi di Milano, Milan/Italy, <sup>18</sup>Department of Laboratory  
63 Medicine, Division of Clinical Microbiology, Medical University  
64 of Vienna, Vienna/Austria, <sup>19</sup>Institute for Medical  
65 Microbiology, Immunology and Hygiene, University of  
66 Cologne, Cologne/Germany, <sup>20</sup>Mycology Reference Laboratory,  
67 Public Health England, Bristol/United Kingdom, <sup>21</sup>Karolinska  
68 Institutet, Department of Laboratory Medicine, F 68,  
69 Karolinska University Hospital, Huddinge, Stockholm/Sweden,  
70 <sup>22</sup>National Reference Medical Mycology Laboratory, Institute  
71 of Microbiology and Immunology, Faculty of Medicine,

72 University of Belgrade, Belgrade/Serbia, <sup>23</sup>Cologne Excellence  
73 Cluster on Cellular Stress Responses in Aging-Associated  
74 Diseases (CECAD), Department I of Internal Medicine, Clinical  
75 Trials Centre Cologne (ZKS Köln), Center for Integrated  
76 Oncology (CIO Köln-Bonn), German Centre for Infection  
77 Research (DZIF), University of Cologne, Cologne/Germany,  
78 <sup>24</sup>Clinical Microbiology Laboratory, National Kapodistrian  
79 University of Athens, ATTIKON University Hospital Athens,  
80 Athens/Greece, <sup>25</sup>Department of Hygiene and Medical  
81 Microbiology, Klinikum Wels-Grieskirchen, Wels/Austria,  
82 <sup>26</sup>Department of Public Health and Pediatrics, Microbiology  
83 Division, Turin/Italy, <sup>27</sup>Department I for Internal Medicine,  
84 University Hospital of Cologne, Cologne/Germany and German  
85 Centre for Infection Research, partner site Bonn-Cologne,  
86 Germany, <sup>28</sup>A.O.U. Policlinico Vittorio Emanuele Catania,  
87 Biometec – University of Catania/Italy, <sup>29</sup>Infectious Diseases  
88 Unit, S. Orsola-Malpighi, Department of Medical and Surgical  
89 Sciences, University of Bologna, Bologna/Italy, <sup>30</sup>Department  
90 of Clinical Microbiology and Immunology, Sackler School of  
91 Medicine, Tel Aviv University, Tel Aviv/Israel, <sup>31</sup>Institute of  
92 Medical Microbiology, University Hospital Essen, University of  
93 Duisburg- Essen, Essen/Germany, <sup>32</sup>Department of of  
94 Microbiology, Faculty of Medicine and Dentistry, Palacky  
95 University Olomouc and University Hospital Olomouc/Czech

96 Republic, <sup>33</sup>Clinical Microbiology, Puerta del Mar University  
97 Hospital, University of Cádiz, Cádiz/Spain, <sup>34</sup>Infectious Diseases  
98 Unit, 3<sup>rd</sup> Department of Pediatrics, Faculty of Medicine,  
99 Aristotle University School of Health Sciences, Hippokration  
100 General Hospital, Thessaloniki/Greece, <sup>35</sup>Department of  
101 Medical Microbiology, Hacettepe University Medical School,  
102 Ankara/Turkey, <sup>36</sup>Division of Mycology, Department of Medical  
103 Microbiology, Chandigarh/India, <sup>37</sup>Escola Paulista de Medicina,  
104 Federal University of São Paulo, São Paulo/Brazil,  
105 <sup>38</sup>Departamento de Micología, Instituto de Medicina Regional,  
106 Universidad Nacional del Nordeste, CONICET,  
107 Resistencia/Argentina, <sup>39</sup> Division of Clinical Mycology,  
108 Department of Microbiology, Vall d'Hebron University  
109 Hospital, Barcelona, <sup>40</sup>Department of Medical Mycology and  
110 Parasitology/Invasive Fungi Research Center, Mazandaran  
111 University of Medical Sciences, Sari/Iran, <sup>41</sup>School of Medicine,  
112 European University Cyprus, Nicosia/Cyprus, <sup>42</sup>Department of  
113 Clinical Mycology, Allergy and Immunology, North Western  
114 State Medical University, Saint Petersburg/Russia,  
115 <sup>43</sup>Department I for Internal Medicine, University Hospital of  
116 Cologne, Cologne/Germany, <sup>44</sup>CBS-KNAW Fungal Biodiversity  
117 Centre, Utrecht/The Netherlands, <sup>45</sup>Hacettepe University  
118 Medical School, Department of Infectious Diseases and Clinical  
119 Microbiology, Ankara/Turkey, <sup>46</sup>Department of Medical

120 Statistics, Informatics and Health Economics, Medical  
121 University of Innsbruck, Innsbruck/Austria, <sup>47</sup>Department of  
122 Dermatology, Hospital Ernest Lluch Martin, Calatayud,  
123 Zaragoza/Spain  
124  
125 Running title: TerrNet Study  
126  
127 Corresponding author: Cornelia Lass-Flörl, Division of Hygiene  
128 and Medical Microbiology. University of Innsbruck,  
129 Schöpfstraße 41, 6020 Innsbruck, Austria  
130 Tel.No. 0043 512 9003-70703, Fax No. 0043 512 9003-73700  
131 email: [cornelia.lass-floerl@i-med.ac.at](mailto:cornelia.lass-floerl@i-med.ac.at)  
132  
133 Length of abstract : 87 words  
134 Length of paper (excluding abstract and ref.) : 1172 words



135 **Abstract**

136 Objectives: A prospective international multicentre

137 surveillance study was conducted to investigate the

138 prevalence and amphotericin B (AMB) susceptibility of

139 *Aspergillus terreus* species complex infections.

140 Methods: Three hundred seventy cases from 21 countries

141 were evaluated.

142 Results: The overall prevalence of *A. terreus* species complex

143 among patients investigated and with mold positive cultures

144 was 5.2% (370/7116). AMB MICs were ranging from 0.125 to

145 32 mg/L, (median 8mg/L).

146 Conclusions: *A. terreus* species complex infections cause a

147 wide spectrum of aspergillosis and the majority of cryptic

148 species display high AMB MICs.

149 Introduction:  
150 *Aspergillus terreus* species complex holds an exceptional  
151 position within the aspergilli, as it appears to be a rare  
152 pathogen of infection and displays polyene resistance [1,2,3].  
153 *A. terreus* is a common cause of invasive aspergillosis (IA) at  
154 the M. D. Anderson Cancer Center in Houston, USA, and the  
155 University Hospital of Innsbruck, Austria [3,4,5]. Almost no  
156 data are available on how frequently this species occurs  
157 elsewhere and whether differences within amphotericin B  
158 (AMB) susceptibility exist. Our objective was to investigate the  
159 global prevalence of *A. terreus* species complex in fungal  
160 diseases and to survey AMB susceptibility.

161

162 Methods:

163 An international surveillance network was established on  
164 behalf of the European Fungal Infection Study Group, the  
165 International Society for Human and Animal Mycology  
166 *Aspergillus terreus* working group, and the European  
167 Confederation of Medical Mycology. 38 centres from 21  
168 countries participated. Each centre collected isolates and  
169 reported the number of *A. terreus* and fungal pathogens  
170 detected for 12 consecutive months (2014-2015). Patient  
171 characteristics', epidemiological data, and antifungal  
172 treatment were documented through an online questionnaire

173 using www.clinicalsurveys.net online platform. Patients were  
174 classified according to the European Organisation for the  
175 Research and Treatment of Cancer/Mycoses Study Group  
176 consensus definitions [6] by the participating centres. Unless  
177 otherwise noted, the isolation of *A. terreus* from sputa of non-  
178 neutropenic patients was categorized as colonisation. Isolates  
179 were sent to the Division of Hygiene and Medical Microbiology  
180 for molecular species identification [7,8] and susceptibility  
181 testing according to EUCAST (European Committee on  
182 Antimicrobial Susceptibility Testing) method [2]. *A. terreus*  
183 strains were identified to the cryptic species level by  
184 sequencing partial beta-tubulin and applying a validated in-  
185 house database owned by Jos Houbraken, CBS Fungal  
186 Biodiversity Center, Utrecht, The Netherlands. An AMB  
187 epidemiological cut-off value of 4mg/L was set for *A. terreus*  
188 [2].

189 This study was approved by the Ethics Commission of the  
190 Medical University of Innsbruck (UN4926).

191 Results:

192 461 cases were enrolled of which 91 were excluded because of  
193 insufficient patient documentation (n=45) or lack of fungal  
194 isolates (n=46) being available. Consequently, this survey  
195 comprises 370 eligible cases with an equal number of  
196 corresponding *A. terreus* isolates. Cases derived from Europe

197 (n=261), followed by Middle East (n=70), India (n=19), South  
198 America (n=10), and North America (n=10) (Figure 1). *A.*  
199 *terreus* sensu stricto (n=315), *A. citrinoterreus* (n=36), *A.*  
200 *alabamensis* (n=6), *A. hortai* (n=10), *A. floccosus* (n=1), and *A.*  
201 *neoafricanus* (n=1) were identified. One isolate (*A. terreus*  
202 1214) was most close to *A. alabamensis* and might represent a  
203 new species. Thus, cryptic species accounted for 14.9%  
204 (55/370) with *A. citrinoterreus* (36/55, 65.5%) being dominant.  
205 AMB MICs ranged from 0.125 to 32 mg/L for *A. terreus* sensu  
206 stricto; MICs for all cryptic species were consistently higher,  
207 ranging from 2 to 32 mg/L, see Table 1. According to the  
208 EUCAST cut-off values, 194 isolates (52.4%) were classified as  
209 non-wild types. A proportion of 6.3% (n=20) of the *A. terreus*  
210 sensu stricto isolates displayed lower MICs, ranging from 0.25  
211 – 0.5 mg/L. Isolates were predominantly acquired from Spain  
212 (n=85) and Austria (n=49), see Figure 1.  
213 Underlying diseases e described in Table 2. Species distribution  
214 did not differ per underlying disease and specimen  
215 investigated (Table 2). Diseases comprised IA (25.1%), allergic  
216 broncho-pulmonary aspergillosis (12.4%), chronic aspergillosis  
217 (11.4%), COPD exacerbation (5.5%), aspergilloma (3.7%), otitis  
218 externa (2.5%), and wound infections (0.7%). 25.1% and 27.3%  
219 of the patients suffered from proven and probable IA, 28.6%  
220 were colonized, 10.1% had onychomycosis, and 8.9% had

221 mycological documented diseases such as otitis externa,  
222 aspergilloma and others.  
223 Using a random effects model the pooled estimated  
224 proportion was 5.6% (95% CI 3.8 to 7.7) with  $I^2 = 92\%$   
225 ( $p < 0.0001$ ) and the proportions ranged from 0.0% to 58.3%.  
226 These calculations were done with MedCalc 16.8.4. Four  
227 reference centres and one centre dealing with onychomycosis  
228 only were excluded from the analysis.  
229 A total of 68 patients received antifungal treatment at the  
230 time of fungal diagnosis, 12 were treated with AMB or  
231 liposomal - AMB. The remaining 56 received combinations of  
232 azoles and echinocandins and improved. Only one patient died  
233 due to the *A. terreus* infection. No information on outcome  
234 was available in 13 patients.

235

236 Discussion:

237 Infections due to *A. terreus* species complex were detected in  
238 21 countries and 38 centres with an overall prevalence of 5.2%  
239 among mold infections. High AMB MICs were frequently  
240 observed and crossed all cryptic species. Infections were  
241 reported from all over the world with three main specific  
242 findings. Firstly, Spain and Austria were the countries with the  
243 highest density of *A. terreus* isolates collected. Secondly, the  
244 number of *A. terreus* cases enrolled varied from centre to

245 centre, and displayed a broad range from zero to several cases  
246 per country. Thirdly, it seems that few susceptible AMB  
247 variants exist within *A. terreus* sensu stricto.  
248 Taking into account the differences on the environmental  
249 conditions, host related characteristics, and the use of  
250 antifungal agents, it is not possible to conclude on the  
251 particular biogeography of *A. terreus* species complex. In  
252 addition, one has to be aware that data collected may depend  
253 on the quality of care, patient demographics, infection control  
254 practices, frequency of specimen collection, and laboratory  
255 methodology. Hence, further studies are needed to determine  
256 whether specific risk and/or environmental factors are  
257 associated with infections by *A. terreus*.  
258 Notable was the fact that *Aspergillus* section *Terrei* was most  
259 commonly isolated from patients suffering from chronic lung  
260 diseases (39.2%). No similar data have been reported [10] and  
261 it remains to be seen whether *A. terreus* reflects an emerging  
262 pathogen of this disease entity.  
263 *A. terreus* is a poor target for AMB and hence is reported as  
264 resistant [2]. The role of isolates with MICs <0.5 mg/L needs  
265 further evaluation. The pharmacodynamic target may be  
266 attained with the standard AMB dose for isolates with MICs  
267  $\leq 0.25$  mg/l [10] and infections were successfully treated with  
268 high dose liposomal-AMB [11].

269 Cryptic species accounted for 14.8%, with *A. citrinoterreus*  
270 being the most prevalent. Although the clinical implications of  
271 sibling species of *A. terreus* are less well understood, our study  
272 confirms that these species are generally resistant to AMB and  
273 are causing a wide spectrum of invasive and non-invasive  
274 aspergillosis. Guinea et al. [12] observed *A. citrinoterreus*  
275 acting mainly as a co-pathogen with *A. fumigatus*.  
276 Our study has some limitations. We do not have a  
277 comprehensive worldwide *A. terreus* survey network and  
278 some countries are missing for a variety of reasons. Also,  
279 generally, the diagnosis of fungal infections is difficult to  
280 obtain and may often be based on detection of biomarkers  
281 rather than on isolation of the infecting organism. Hence,  
282 some cases may have been missed and chronic lung diseases  
283 were not specified in more detail. Further, we have no data  
284 available on co-infections which may complicate diseases. The  
285 centres included represent a convenience sample. However,  
286 this is the largest and geographically most diverse study on the  
287 contemporary epidemiology of *A. terreus* species complex  
288 infections worldwide.

289 Our study shows that *A. terreus* sensu stricto is widely  
290 distributed in climatically divergent countries, and that cryptic  
291 species display high AMB MICs. *A. terreus* species complex was

- 292 most commonly isolated from patients suffering from chronic  
293 lung diseases (39.2%).

ACCEPTED MANUSCRIPT



**294 Funding**

295 This work was supported by ECMM, ISHAM and EFISG and in  
296 part by an unrestricted research grant through the Investigator  
297 Initiated Studies Program of Astellas and Pfizer as well as by  
298 the Austrian Science Fund (W1253-B24 doctoral program  
299 HOROS).

300

301

**302 Transparency Declaration**

303 We declare that we have no conflicts of interest related to this  
304 study.

305 CLF received research grants, travel grants or honorarium as a  
306 speaker or advisor from Gilead Sciences, Pfizer, Schering

307 Plough, MSD and Basilea, VAA received research grants or

308 honorarium as a speaker or advisor from Astellas, Pfizer and

309 Schering Plough, MSD. WB received honorarium as a speaker

310 from Pfizer, Schering Plough and MSD. CF received research

311 grants or honorarium as a speaker or advisor from Astellas,

312 Gilead Sciences, MSD and Basilea. AH received travel grants

313 from Astellas and served on the advisory board of Gilead. EJ

314 received travel grants or honorarium as a speaker or advisor

315 from Bio-Rad, Gilead Sciences, Pfizer, Schering Plough, and

316 MSD. DPK has received research support from Pfizer, Astellas

317 Pharma, honoraria from Astellas, Merck, Pfizer and he serves

318 on the advisory boards of Merck, Amplyx, Cidara and F2G. JFM  
319 received grants from Astellas, Basilea and Merck. He has been  
320 a consultant to Astellas, Basilea and Merck and received  
321 speaker's fees from Merck, United Medical and Gilead. ALC  
322 received educational grants from Pfizer, Gilead and United  
323 Medical and research grant from Pfizer, Astellas. MCA received  
324 research grants/contract work (Paid to SSI) from Amplyx,  
325 Basilea, Cidara, F2G, Gilead, Pfizer & T2Candida, speaker  
326 honoraria from Astellas, Basilea, Gilead, MSD, Novartis, Pfizer  
327 & T2Candida. NK has received research support from Pfizer,  
328 honoraria from Astellas, Merck and Pfizer, and he serves on  
329 the advisory boards of Merck and Pfizer. OAC is supported by  
330 the German Federal Ministry of Research and Education, has  
331 received research grants from Actelion, Aranis, Astellas,  
332 AstraZeneca, Basilea, Bayer, Cidara, Duke University (NIH  
333 UM1AI104681), F2G, Gilead, GSK, Leeds University, MedPace,  
334 Melinta Therapeutics, Merck/MSD, Miltenyi, Pfizer, Rempex,  
335 Roche, Sanofi Pasteur, Scynexis, Seres Therapeutics, The  
336 Medicine Company, is a consultant to Anacor, Amplyx,  
337 Actelion, Astellas, Basilea, Cidara, Da Volterra, F2G, Gilead,  
338 Inositec AG, Janssen Pharmaceuticals, Matinas, Menarini,  
339 Merck/MSD, Paratek Pharmaceuticals, Scynexis, Seres,  
340 Summit, Vical, Vifor, and received lecture honoraria from  
341 Astellas, Basilea, Gilead and Merck/MSD. BW received

342 research support from Pfizer, travel grants or honorarium as a  
343 speaker or advisor from Gilead Sciences, Pfizer, Astellas, MSD  
344 and Basilea. LK grant from Gilead and has been an adviser to  
345 Astellas, Gilead, Schering-Plough, has received research grants  
346 from Gilead, Merck, Sharpe&Dohme, Schering-Plough and has  
347 received honoraria for educational lectures from Gilead,  
348 Pfizer, Merck, Sharpe & Dohme, Schering-Plough and Janssen.  
349 AMT received research support from Gilead Sciences, travel  
350 grants or honorarium as speaker from Gilead Sciences, Pfizer,  
351 Astellas. KL has received research grants from Gilead, MSD and  
352 Pfizer, received travel support from MSD, Pfizer and Gilead  
353 and received lecture honoraria from Gilead, MSD, and Pfizer.  
354 SAA received Investigator Initiated Research grant support  
355 from Pfizer and speaker honoraria or travel grants from  
356 Astellas, Gilead, Merck and Pfizer. SO received research grants  
357 from Pfizer and Astellas, honorarium as advisor from MSD,  
358 Astellas Pharma and Gilead Sciences. ER reports grants, and  
359 non-financial support from Astellas, Pfizer, Gilead, and Merck,  
360 outside the submitted work. AAI received research and/or  
361 travel grants from Gilead Sciences, F2G and Pfizer. OU was  
362 consultant for and has received travel grants from Merck,  
363 Astellas, Pfizer, and Gilead. S.M.H. has received research and  
364 travel grants from Astellas, Gilead, and MSD, research grants  
365 from Basilea, and travel grants from Pfizer; was consultant to

- 366 Basilea; and received honoraria as a speaker from Astellas and  
367 Merck.

ACCEPTED MANUSCRIPT

368 **References**

- 369 1. Perkhofer S, Lass-Flörl C, Hell M, Russ G, Krause R, Hönigl  
370 M, et al. The Nationwide Austrian *Aspergillus* Registry: a  
371 prospective data collection on epidemiology, therapy and  
372 outcome of invasive mould infections in  
373 immunocompromised and/or immunosuppressed patients.  
374 Int J Antimicrob Agents 2010;36:531-6.
- 375 2. Arendrup MC, Cuenca-Estrella M, Lass-Flörl C, Hope WW;  
376 European Committee on Antimicrobial Susceptibility  
377 Testing Subcommittee on Antifungal Susceptibility Testing  
378 (EUCAST-AFST). EUCAST technical note on *Aspergillus* and  
379 amphotericin B, itraconazole, and posaconazole. Clin  
380 Microbiol Infect. 2012;18:E248-50.
- 381 3. Lass-Flörl C, Grif K, Mayr A, Petzer A, Gastl G, Bonatti H et  
382 al. Epidemiology and outcome of infections due to  
383 *Aspergillus terreus*: 10-year single center experience. Br J  
384 Haematol. 2005;131:201-207.
- 385 4. Lass-Flörl C, Grif K, Kontoyiannis DP. Molecular typing of *A.*  
386 *terreus* isolates collected in Houston, Texas and Innsbruck,  
387 Austria: evidence of great genetic diversity. J Clin  
388 Microbiol. 2007;45:2686-90.
- 389 5. Lackner M, Coassin S, Haun M, Binder U, Kronenberg F,  
390 Haas H et al. Geographically predominant genotypes of  
391 *Aspergillus terreus* species complex in Austria: a

- 392        microsatellite typing study. Clin Microbiol Infect.  
393        2016;22:270-276.
- 394    6. De Pauw B, Walsh TJ, Donnelly JP, Stevens DA, Edwards JE,  
395        Calandra T et al. Revised definitions of invasive fungal  
396        disease from the European Organization for Research and  
397        Treatment of Cancer/Invasive Fungal Infections  
398        Cooperative Group and the National Institute of Allergy  
399        and Infectious Diseases Mycoses Study Group  
400        (EORTC/MSG) Consensus Group. Clin Infect Dis  
401        2008;46:1813-21.
- 402    7. Balajee S A, Baddley JW, Peterson SW, Nickle D, Varga J,  
403        Boey A et al., ISHAM Working Group on *A. terreus*.  
404        *Aspergillus alabamensis*, a new clinically relevant species in  
405        the section *Terrei*. Eukaryotic Cell 2009;8:713–22.
- 406    8. Kathuria S, Sharma C, Singh PK, Agarwal P, Agarwal K,  
407        Hagen F et al. Molecular epidemiology and in-vitro  
408        antifungal susceptibility of *Aspergillus terreus* species  
409        complex isolates in Delhi, India: Evidence of genetic  
410        diversity by amplified fragment length polymorphism and  
411        microsatellite typing. PLoS ONE 2015;10:e0118997.
- 412    9. Denning DW, Cadranel J, Beigelman-Aubry C, Ader F,  
413        Chakrabarti A, Blot S et al. Chronic pulmonary aspergillosis:

- 414 rationale and clinical guidelines for diagnosis and  
415 management. Eur Respir J. 2016;47:45-68.
- 416 10. Elefanti A, Mouton JW, Verweij, PE, Zerva L, Meletiadis J.  
417 Susceptibility breakpoints for amphotericin B and  
418 *Aspergillus* species in an in vitro pharmacokinetic-  
419 pharmacodynamic model simulating free-drug  
420 concentrations in human serum. Antimicrob Agents  
421 Chemother. 2014;58:2356-2362.
- 422 11. Lewis RE, Albert NP, Liao G, Wang W, Prince RA,  
423 Kontoyiannis DP. High-dose induction liposomal  
424 amphotericin B followed by de-escalation is effective in  
425 experimental *Aspergillus terreus* pneumonia. J Antimicrob  
426 Chemother. 2013;68:1148-1151.
- 427 12. Guinea J, Sandoval-Denis M, Escibano P, Peláez T, Guarro  
428 J, Bouza E. *Aspergillus citrinoterreus*, a new species of  
429 section Terrei isolated from samples of patients with non-  
430 hematological predisposing conditions. J Clin Microbiol.  
431 2015;53:611-617.

**Table 1.** Distribution of amphotericin B MICs against *Aspergillus terreus* species complex isolates collected during the study period and tested according to EUCAST methodology

Species	Amphotericin B MICs, mg/L								
	0.125	0.25	0.5	1	2	4	8	16	32
<i>A. terreus sensu stricto</i>	3	7	10	14	36	81	86	55	23
<i>A. citrinoterreus</i>					3	13	8	7	5
<i>A. hortai</i>					1	2	5	2	
<i>A. alabamensis</i>					2	3	1		
<i>A. floccosus</i>						1			
<i>A. neoaffricanus</i>									1
Potential new species							1		

**Table 2.** Species distribution of *Aspergillus terreus* species complex isolated from the various human specimens

Species	Specimens, total numbers							Total
	Sputa	Bronchoalveolar lavages and tracheal secretions	Body-fluids	Biopsies	Swabs	Others		
<i>A. terreus sensu stricto</i>	126	65	53	33	17	21	315	
<i>A. citrinoterreus</i>	14	7	3	5	3	4	36	
<i>A. hortai</i>	4	2			1	3	10	
<i>A. alabamensis</i>	3	2			1		6	
<i>A. floccosus</i>					1		1	
<i>A. neoaffricanus</i>						1	1	
Potential new species				1			1	
Total	147	76	56	39	23	29	370	

\* aspirates, wound secretions, nails




**Fig. 1a - c.** Overview of countries and *Aspergillus terreus* species complex isolated numbers collected during the study period:

a) *Aspergillus citrinoterreus*

b) *Aspergillus terreus sensu stricto*


c) *Aspergillus hortai*

ACCEPTED MANUSCRIPT


*Aspergillus citrinoterreus*

**Total=36**

ACCEPTED

*Aspergillus terreus sensu stricto*

ACCEPTED

*Aspergillus hortai*

Total=10

ACCEPTED