



## Research Article

# Barrett's Disease and *Helicobacter Spp.* Detection in Dogs with Gastritis

Simonetta Appino<sup>1\*</sup>, Paola Pregel<sup>2</sup>, Mitzy Mauthe Von Degerfeld<sup>2</sup>  
and Ada Rota<sup>2</sup>

### Abstract

Barrett's disease (BE) is a metaplastic disorder of the esophageal mucosa in which specialized columnar epithelium replaces healthy squamous epithelium; it is the most common cause of esophageal carcinoma in humans. The dog has been used as a model of Barrett's esophagus, although the pathology is very rare in the species. In man, infection with *Helicobacter pylori* resulted inversely associated with BE. The role of *Helicobacter spp.* in dogs gastritis has not been fully elucidated, as in esophageal disease. Aim of this work was to evaluate esophageal and gastric biopsies of symptomatic dogs in association with the isolation of *Helicobacter spp.* and to describe and discuss a case of BE.

### Keywords

Barrett's disease; Dog; *Helicobacter*; Stomach

## Introduction

Bacteria of the Genus *Helicobacter* colonize the gastrointestinal tract of several mammalian and avian species. Some *Helicobacter* species, such as *H. canis*, *H. pullorum*, *H. heilmannii*, *H. rappini* and *H. cinaedi*, may be zoonotic [1-3]. In human beings, *Helicobacter pylori* colonize the stomach of roughly half the world population [4]. Infection with *H. pylori* typically causes a mild, mixed gastritis; however, chronic infections lead to severe clinical outcomes, such as duodenal and gastric ulcers, in approximately 15% of infected individuals. Chronic *H. pylori* infection also contributes to gastric issues that are associated with the development of gastric cancer, leads to loss of acid secretion and the loss of acidic mammalian chitinase expression [5]. The factors leading to these divergent clinical outcomes are not entirely known, but host genetics that regulate the potency of the immune response towards the infection, bacterial genetics, and environmental factors, such as diet and smoking, contribute [6].

Among human esophagogastric diseases, Barrett's disease (or Barrett esophagus= BE) originates from a long-standing gastroesophageal reflux that damages the integrity of esophageal mucosa; it is a metaplastic disorder in which specialized columnar epithelium replaces healthy squamous epithelium. The metaplastic columnar lining comes in three types: two types are similar to groups of cells found in regions of the stomach lining; the third

type is similar to groups of cells found in the small intestine. This intestinal type of metaplasia is important because it can potentially lead to the development of cancer. Barrett's metaplasia is, in fact, the most common cause or precursor of esophageal carcinoma; the metaplasia derives from basal layer of squamous epithelium, but also submucosal or mucosal glands or ducts, and mesenchyme are possibilities. Columnar epithelium of Barrett's may be more resistant to acid, pepsin and bile [7,8]. Barrett's oesophagus has been recently associated to Dog adenocarcinoma by Chambers et al. [9].

Metaplasia is an adaptive response of tissue to various injuries such as the chemical insult to the distal esophagus due to gastroesophageal reflux. Regarding *Helicobacter pylori*, Rubenstein et al. [10] concluded that infection with this bacterium was inversely associated with BE, particularly the cagA+ strain, possibly because the association of *H. pylori* and erosive esophagitis is only valid in patients without gastric atrophy [11].

BE is rarely reported in dogs but the dog has been used as a model, inducing the development of the pathology by mucosectomy and gastric acid augmentation [12].

While examining some dogs showing gastric symptoms for the presence of *Helicobacter spp.*, some histologic profiles of Barrett disease could be identified and were thought worth to be reported.

Ten dogs of different age (3 to 8 years) and breed were included in the study: they had been referred to the Veterinary University Hospital of the University of Turin, Italy, because of unresolved mild to severe signs of gastric disease: vomiting at distance from meals, weight loss, lack-of or decreased appetite, occasional diarrhoea, abdominal pain referable to stomach pain.

The animals were subjected to esophageal and gastric biopsies: besides standard evaluations, the presence of *Helicobacter spp.* was investigated. An informed consent was obtained by the owners: all the procedures had diagnostic purposes and are common clinical actions; none of them was carried out solely for research purposes.

Food was withheld for 12 hr and gastroscopy was carried out on dogs under general isoflurane anesthesia, according to the procedures adopted at the Veterinary University Hospital. The dogs were positioned in left lateral recumbency and tissue specimens were collected from the distal esophagus and from the stomach, both from the cardias and the fundus: two specimens were collected from each site, one was stored in frozen until exam, the other one was fixed in 10% neutral buffered formalin for histological examination. From paraffin-embedded tissue samples three- $\mu$ m sections were cut and stained with May Grünwald-Giemsa and Warthin Starry in order to detect *Helicobacter spp.*

A gastric washing was performed on each animal, by slowly infusing 10 ml of sterile saline solution through a gastric tube (4 mm diameter) and recollecting the fluid by using a syringe; then the fluid was centrifuged for 5 min at 3000 rpm and the pellet was cytocentrifuged and smeared for cytological examination after May Grünwald-Giemsa staining.

DNA was extracted both from each tissue sample and from each cells pellet by means of a commercial kit (Nucleospin® Tissue

\*Corresponding author: Simonetta Appino, Dipartimento di Medicina Veterinaria, Università di Sassari via Vienna 2, 07100 Sassari, Italy, Tel: +39 079229468; E-mail: [simo@uniss.it](mailto:simo@uniss.it)

Received: September 22, 2017 Accepted: October 14, 2017 Published: October 18, 2017

extraction Kit Macherey-Nagel) and amplified by PCR, using specific primers which amplify a fragment of the 16S rRNA gene of *Helicobacter*. PCR was performed in 25 µl tubes using 0.5U Taq DNA polymerase, 10mM Tris HCl, 50mM KCl, 1.5mM MgCl<sub>2</sub>, 200 µM each dNTP and 25 µM each primer (sense primer: innS: gaa cct cta ggc ttg aca ttg, and antisense primer: innR: ggt gag tac aag acc cgg gaa). Samples were submitted to 30 amplifying cycles at the following temperatures: denaturation at 94°C for 30 sec, annealing at 47°C for 30 sec and extension at 72°C for 30 sec. PCR yielded a 420 bps fragment of the 16S rRNA gene which was evaluated by means of 7% polyacrylamide gel electrophoresis, comparing the obtained products to DNA Molecular Weight Marker V (Roche Diagnostics, Basel, Switzerland).

Table 1 reports the summary of the results of the different exams: histology of gastric biopsies did not show any significant lesion in six dogs; from two of them, *Helicobacter* DNA was found both in gastric mucous and in the gastric specimen. In all the positive dogs, *Helicobacter* was never revealed in the oesophagus but only in the stomach and all the exams agreed: it was directly detected in the tissue, in the cells smears and with the molecular analysis. Warthin Starry (Figure 1) staining showed the presence of brown spiral microorganisms in all the *Helicobacter* PCR positive histological samples. The bacteria were mainly present on the surface of the lumen glands. The same findings were shown by Giemsa staining of the gastric biopsies, in which the spiral bacteria stained dark purple. In cells smears from cytocentrifuged gastric fluid, spiral microorganisms presumptive of *Helicobacter* spp. were evidenced in all the *Helicobacter* PCR positive samples. PCR positive samples gave the typical 420 bps fragment evaluated on SDS-PAGE. Positive samples nucleotide sequences revealed a homology range of 99-100% with a partial sequences from *H. heilmanni* 16S.

Some *Helicobacter* positive dogs showed histological lesions mainly represented by chronic gastritis. In a negative *Helicobacter* dog, histology revealed a mild lymphocytic gastritis. In three cases, histology revealed inflammation of the esophagus, but only in one, a Barrett's esophagus could be diagnosed. It is the case of a Poodle, suffering from chronic vomit and acute episodes of diarrhoea, in which esophageal epithelium was replaced by columnar epithelium of intestinal type with goblet cells (distended, mucin-filled cytoplasm with a barrel-shaped configuration) (Figure 1) associated with mild chronic submucosal inflammation. This dog was negative for *Helicobacter* spp.

In general, esophageal mucosa dysfunction has a multifactorial pathogenesis that includes the reflux of gastric or bile acid into the esophagus, hiatal hernia, dysfunction of esophageal sphincter and esophageal motility, impairment of esophageal epithelial resistance and hypersensitivity. Of these factors, prolonged and frequent esophageal reflux of gastric acid with low pH is a critical risk factor for esophageal mucosal injury. As regards *Helicobacter*, its role in dogs esophageal disease has not been clearly defined. In man, it seems to be inversely associated to the presence of the microorganism in the stomach, probably due to the eradication therapy but its role requires further investigations [13].




Figure 1: (a) gastric biopsy, spiralshape microorganisms, *Helicobacter* spp. in gastric glands lumen (Giemsa staining, 1000x) (b) higher magnification (c) gastric biopsy, *Helicobacter* spp. positive (Warthin Starry staining, 1000x) (d) esophageal biopsy from Poodle dog, hyperplasia of glandular mucosa (HE, 200x).

Table 1: Detection of *Helicobacter* spp. DNA in the dogs gastric mucous (Pellet) and in tissue specimens from oesophagus and stomach; presence of *Helicobacter* spp. in cells smears (Cytology) and in histological samples from oesophagus and stomach; histological observations. BE=Barrett's esophagus.

Animal	<i>Helicobacter</i> DNA			Cytology <i>H. spp.</i>	Histology		Stomach <i>H. spp.</i>	
	Pellet	Esophagus	Stomach		Esophagus <i>H. spp.</i>			
1 Poodle	-	-	-	-	-	BE	-	nsl*
2 G. shepherd	+	-	+	+	-	Mild basal hyperplasia	+	nsl
3 G shepherd	+	-	+	+	-	nsl	+	mild chronic gastritis
4 Rottweiler	+	-	+	+	-		+	Moderate chronic gastritis, plasma cells, small lymphocyte
5 Rottweiler	-	-	-	-	-	nsl	-	Mild lymphocytic gastritis
6 Rottweiler	-	-	-	-	-	nsl	-	nsl
7 Mixed-breed	-	-	-	-	-	Mild lymph. infiltration	-	nsl
8 Dogue de B.	+	-	+	+	-	nsl	+	Chronic gastritis
9 Dogue de B.	-	-	-	-	-	nsl	-	nsl
10 S Husky	+	-	+	+	-	nsl	+	nsl

\*nsl (no significant lesions)

Results of different studies on the relationship between *Helicobacter* and BE are still controversial. According to some authors, colonization with cagA-positive *Helicobacter pylori* strains in humans may protect against gastro-esophageal reflux disease and its complications [14], but other works assessed the opposite [15]. Medications that are helpful for BE include H2 receptor antagonists (or H2 blockers) and proton pump inhibitors, which reduce the amount of acid produced by the stomach, and are typically used to counteract *Helicobacter*.

Nevertheless, the full understanding of canine gastric pathology requires further investigations. The results from histology, cytology and molecular biology were in agreement in our study; *Helicobacter* was detected only in gastric samples, never in esophagus specimens and its presence was not always related to gastric inflammation signs. This study also shows the importance of the gastric washing (followed by PCR or cytological examination) in the detection of *Helicobacter* spp. and it confirms that Giemsa staining is as reliable as Warthin Starry stain and has the advantage of being more rapid and less difficult to perform.

## References

1. Fox JG (2002) The non-*H. pylori* helicobacters: their expanding role in gastrointestinal and systemic diseases. *Gut* 50: 273-283.
2. De Groote D, van Doorn LJ, Ducatelle R, Verschuuren A, Haesebrouck F, et al. (1999) 'Candidatus *Helicobacter suis*', a gastric helicobacter from pigs, and its phylogenetic relatedness to other gastrospirilla. *Int. J. Syst. Bacteriol.* 49: 1769-1777.
3. Kawamura Y, Tomida J, Miyoshi-Akiyama T, Okamoto T, Narita M, et al (2016) Proposal of *Helicobacter canicola* sp. nov., previously identified as *Helicobacter cinaedi*, isolated from canines. *Syst. Appl. Microbiol.* 39: 307-312.
4. Brown LM (2000) *Helicobacter pylori*: Epidemiology and routes of transmission. *Epidemiol. Rev.* 22: 283-297.
5. Nookaew I, Thorell K, Worah K, Wang S, Hibberd ML, et al. (2013) Transcriptome signatures in *Helicobacter pylori*-infected mucosa identifies acidic mammalian chitinase loss as a corpus atrophy marker. *BMC Med. Genomics* 6: 41.
6. Ahn HJ, Lee DS (2015) *Helicobacter pylori* in gastric carcinogenesis. *World J. Gastrointest. Oncol.* 7: 455-465.
7. DeMeester SR, Wickramasinghe KS, Lord RV, Friedman A, Balaji NS, et al. (2002) Cytokeratin and DAS-1 immunostaining reveal similarities among cardiac mucosa, CIM, and Barrett's esophagus. *Am. J. Gastroenterol* 97: 2514-2523.
8. Odze RD (2005) Unraveling the mystery of the gastroesophageal junction: a pathologist's perspective. *Am. J. Gastroenterol* 100: 1853-1867.
9. Chambers JK, Saito T, Fukushima K, Kakuta S, Nakayama J, et al. (2017) Adenocarcinoma of Barrett's esophagus in a dog. *J Toxicol Pathol* 3: 239-24.
10. Rubenstein JH, Inadomi JM, Scheiman J, Schoenfeld P, Appelman H, et al. (2014) Association between *Helicobacter pylori* and Barrett's esophagus, erosive esophagitis, and gastroesophageal reflux symptoms. *Clin. Gastroenterol. Hepatol* 12: 239-245.
11. Vasapolli R, Malferteiner P, Kandulski A (2016) *Helicobacter pylori* and non-malignant upper gastrointestinal diseases. *Helicobacter* 1: 30-33.
12. Kapoor H, Lohani KR, Lee TH, Agrawal DK, Mittal SK (2015) Animal models of Barrett's esophagus and esophageal adenocarcinoma-Past, present, and future. *Clin Transl Sci* 8: 841-847.
13. Corley DA, Kubo A, Levin TR (2008) *Helicobacter pylori* infection and the risk of Barrett's esophagus: a community-based study. *Gut* 57: 727-733.
14. Ackermark P, Kuipers EJ, Wolf C, Breumelhof R, Seldenrijk CA, et al. (2003) Colonization with cagA-positive *Helicobacter pylori* strains in intestinal metaplasia of the esophagus and the esophagogastric junction. *Am J Gastroenterol* 98: 1719-1724.
15. Ferrandez A, Benito R, Arenas J, Garcia-Gonzalez MA, Sopena F, et al. (2006) CagA-positive *Helicobacter pylori* infection is not associated with decreased risk of Barrett's esophagus in a population with high *H. pylori* infection rate. *BMC Gastroenterol* 6: 7.

## Author Affiliation

Top

<sup>1</sup>Dipartimento di Medicina Veterinaria, Università di Sassari, Italy

<sup>2</sup>Dipartimento di Scienze Veterinarie, Università di Torino, Italy

## Submit your next manuscript and get advantages of SciTechnol submissions

- ❖ 80 Journals
- ❖ 21 Day rapid review process
- ❖ 3000 Editorial team
- ❖ 5 Million readers
- ❖ More than 5000 
- ❖ Quality and quick review processing through Editorial Manager System

Submit your next manuscript at • [www.scitechnol.com/submission](http://www.scitechnol.com/submission)