



American Journal of Human Biology

**Evaluation of baseline frequency of sister chromatid exchanges in an Italian population according to age, gender, smoking habit and gene polymorphisms.**

Journal:	<i>American Journal of Human Biology</i>
Manuscript ID	AJHB-16-0253.R3
Wiley - Manuscript type:	Original Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	SANTOVITO, Alfredo; University of Turin, Department of Life Sciences and Systems Biology GENDUSA, Claudio; University of Turin, Department of Life Sciences and Systems Biology CERVELLA, Piero; University of Turin, Department of Life Sciences and Systems Biology
Keywords:	SCE, Gene Polymorphisms, Italian Population, Lifestyles

SCHOLARONE™  
Manuscripts

Review

**Title: Evaluation of baseline frequency of sister chromatid exchanges in an Italian population according to age, gender, smoking habit and gene polymorphisms.**

**Short Title: Baseline frequency of sister chromatid exchanges in an Italian population**

Authors: Alfredo SANTOVITO\*, Claudio GENDUSA, Piero CERVELLA

University of Turin, Department of Life Sciences and Systems Biology, Via Accademia Albertina n. 13, 10123 Torino (Italy)

\*Corresponding Author:

Alfredo SANTOVITO

Department of Life Sciences and Systems Biology

Via Accademia Albertina n. 13

10123 – Torino (Italy)

Tel.: +39-0116704554

Fax: +39-0116704508

## Abstract

**Objectives:** Increased SCEs frequencies in human lymphocytes are an indicator of spontaneous chromosome instability and could be influenced by different exogenous and endogenous factors. In this study, we evaluated the influence of age, sex, smoking habit and genetic polymorphisms on the background levels of SCEs in peripheral blood lymphocytes.

**Methods:** Two hundred thirty healthy Italian subjects were recruited. Data about age, gender and smoking habit were recorded. Subjects were also genotyped for *GSTT1*, *GSTM1*, *GSTP1 A/G*, *CYP1A1 Ile/Val*, *CYP2C19 G/A*, *ERCC2/XPD Lys751Gln*, *XRCC1 Arg194ATrp*, *XRCC1 Arg399Gln* and *XRCC1Arg208His* gene polymorphisms.

**Results:** The frequency of SCEs/Cell was  $5.15 \pm 1.87$ , with females showing a significantly higher SCEs value with respect to males ( $5.36 \pm 2.10$  and  $4.82 \pm 1.39$ , respectively). Smokers showed significantly increased levels of SCEs with respect to non-smokers ( $5.93 \pm 1.75$  and  $4.70 \pm 1.79$ , respectively) whereas no differences were observed between heavy and light smokers. The age correlated with the RI value ( $P = 0.01$ ) but not with the SCEs frequency ( $P = 0.07$ ), although **the** 31-40 age-group showed a significantly lower SCEs frequency with respect to the other age-groups. A significant association was also found between *GSTP2C19-AA*, *GSTT1-null*, *GSTM1-null*, *ERCC2/XPD Gln751Gln* and *XRCC1 His208His* genotypes and higher frequencies of SCEs.

**Conclusion:** We describe the association between some phase I, phase II and DNA-repair gene polymorphisms with increased SCEs frequencies, reinforcing the importance of genetic analysis in bio-monitoring studies. Gender and age were found to be important endogenous factors **that** affect the level of genomic damage and the replicative capacity of cells, respectively.

1  
2 **Key words:** SCE, Gene Polymorphisms, Italian Population, Lifestyles  
3  
4

5 **ABBREVIATIONS**  
6  
7

8  
9  
10 BrdUrd = bromodeoxyuridine  
11

12 CAs = chromosomal aberrations  
13

14 CYP = cytochrome P  
15

16 GST = glutathione- S-transferase  
17

18 HFC = high frequency cells  
19

20 HFI = high frequency individuals  
21

22 LT = long time  
23

24 NER = nucleotide excision repair  
25

26 RI = replication index  
27

28 SCEs = Sister chromatid exchanges  
29

30 ST = short time  
31

32 XPD = Xeroderma Pigmentosum complementation group D  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 INTRODUCTION  
3

4 The frequency of sister chromatid exchanges (SCEs) in peripheral blood lymphocytes is extensively  
5 used as a biomarker of chromosomal damage and genome stability in human populations. SCEs  
6 occur as a consequence of interchanges between DNA replication products at homologous  
7 chromosomal loci, and these exchanges involve DNA breakage and reunion (Knudsen and Hansen,  
8 2007). Nevertheless, some compounds are able to form covalent adducts with the DNA or to  
9 interfere with DNA metabolism and repair, with consequent induction of [genomic damage](#).  
10  
11

12 SCEs test is also used in surveillance of work environments with low-dose exposures to mutagens  
13 or carcinogens. Indeed, increased frequencies of SCEs [are](#) found to be associated with [a](#) higher risk  
14 of various types of cancers (Medves et al., 2016; Baltaci et al., 2002) [as well as they represent](#)  
15 [indicators](#) of spontaneous chromosome instability among human populations (Salah et al., 2011). In  
16 this [context](#), one of the objectives of the present study was to evaluate, by means of the SCEs assay,  
17 the level of genomic damage in the peripheral blood lymphocytes of non-occupationally exposed,  
18 healthy subjects. Indeed, it is known that the level of genomic damage can be partly influenced by a  
19 variety of external factors such as chemical and physical agents, life styles (smoking and drinking  
20 habits, nutrition) [or](#) residential and/or working areas, as well as by endogenous factors, including  
21 those of biological origins such as gender and age (Santovito et al., 2015; 2016).  
22  
23

24 [Individual](#) genetic susceptibility was found to play an increasingly important role in determining the  
25 levels of genomic damage. From [a](#) genetic point of view, this susceptibility is due to a battery of  
26 gene polymorphisms, principally those of metabolic genes (such as cytochrome P (*CYP*) 450 and  
27 glutathione- S-transferase (*GST*) family genes) (Autrup, 2000; Wang et al., 2013). In particular,  
28 phase I cytochrome P450 (*CYP1*) gene products are involved in the oxidative metabolism of  
29 xenobiotics, producing compounds subsequently processed by phase II enzymes, such as GSTT1. It  
30 was observed that variants of *CYP1* and *GSTs* metabolic genes were associated with increased  
31 levels of chromosomal aberrations and SCEs (Laczmanska et al., 2006; Kumar et al., 2011;  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Hemminki et al., 2015).

In order to prevent the potentially mutagenic consequences of DNA modifications, cells have evolved different mechanisms of DNA repair, depending on the specific type of DNA damage. These mechanisms include Base Excision Repair (BER) and Nucleotide Excision Repair (NER) that correct non-bulky damage and lesions that distort the DNA double helical structure, respectively (Cleaver et al., 2009; Collins and Azqueta, 2012). Most of the genes encoding DNA-repair enzymes are polymorphic, and some of these polymorphisms are directly related with increased levels of chromosomal aberrations and SCEs (Vodicka et al., 2004; Laczmanska et al., 2007; Toolaram et al., 2014).

We decided to evaluate the relationships between some phase I (*CYP1A1* Ile/Val and *CYP2C19* G/A), phase II (*GSTT1* positive/null, *GSTM1* positive/null and *GSTP1* A/G) and DNA-repair (*ERCC2/XPD* Lys/Gln, *XRCC1* Arg194Tr, *XRCC1* Arg399Gln and *XRCC1* Arg208His) gene polymorphisms, as they are associated with an increase susceptibility to DNA damage (Laczmanska et al., 2006; Toolaram et al., 2014, Santovito et al., 2015), and the levels of genomic damage measured by SCEs assay.

## MATERIALS AND METHODS

### *Study population*

The present work is part of a national study designed to analyse the health of the urban Italian population. Our group is interested in analyzing the relationships between life style, endogenous factors and levels of genomic damage. The demographic characteristics of the studied group are reported in Table 1. The study population comprised 230 blood donors sampled in Turin (Piedmont, North-Western Italy). We recruited subjects without any known exposure except those of the routine household, traffic and/or clerical work. Subjects were randomly recruited among different university departments, hospital workers of the administrative staff and healthy voluntaries enrolled

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

in these structures. The equal representation of both sexes and of all age-groups was the only adopted selection criteria.

Subjects were divided into four groups, from A to D, according to age: 21-30 (group A), 31-40 (group B), 41-50 (group C), and 51-70 (group D). The last group included a wider age range because only three individuals were found to belong to the age-group 61-70. Therefore, we decided to join the classes of age 51-60 and 61-70 in a single age-group for this analysis.

In order to analyse the influence of smoking on the level of genomic damage, the total population sample was subdivided in two groups, smokers and non-smokers. Smokers were then subdivided into four sub-groups based on the number of cigarettes smoked/day (cig/day) [heavy smokers, >10 cig/day; light smokers, ≤10 cig/day] and on the number of years of smoking [LT-smokers = long time smokers, >10 years; ST-smokers = short time smokers, ≤10 years].

The present study was performed in accordance with ethical standards of the University of Turin bioethics committee and with the 1964 Declaration of Helsinki. Blood donors were informed about the aim and the experimental details of the study, gave their informed consent, and volunteered to donate blood for sampling. They were healthy at the moment of blood sampling and interviews. In our sample, we exclusively considered individuals that did not consume drugs or alcohol and were not exposed to X-ray for a period of at least two year prior to the analysis.

#### *Blood sampling and SCEs assay*

Blood samples were obtained by venipuncture (5-10 ml) and collected in heparinised tubes. After collection, all blood samples were coded, cooled (4°C), and processed within 2 h after collection. Approximately 0.4 mL of each sample was cultured using RPMI 1640 medium supplemented with 20% foetal calf serum, 2 mM L-glutamine, 0.2 mL of the mitogenic agent phytohemagglutinin (PHA), and antibiotics (100 IU/ml penicillin, and 100 µg/ml streptomycin). To arrest cells in mitosis, colchicine (0.25 µg/mL) was added at a concentration of 0.06 µg/mL during the last 2 h of

1  
2 culture. The cultures were incubated for 72 h at 37 °C in an atmosphere of 5% CO<sub>2</sub> in the air. To  
3  
4 measure SCEs in second division metaphases, bromodeoxyuridine (BrdUrd, 5 µg/mL) was added at  
5  
6 24 h. BrdUrd closely resembles thymidine and is efficiently incorporated into the elongating DNA  
7  
8 strands during replication. After two cell cycles in BrdUrd medium, the two sister chromatids differ  
9  
10 in the amount of BrdUrd and the chromatid with more BrdUrd is lighter in appearance (a  
11  
12 “bleaching” effect). Chromosome preparation was done following standard procedures, as  
13  
14 described in Santovito et al. (2014). For each subject, we scored 50 well-spread second-division  
15  
16 metaphases containing 46 chromosomes. A total of 100 cells from each donor was scored for the  
17  
18 determination of the replication index (RI), calculated according to the following formula:  $RI = (M_1$   
19  
20  $+ 2M_2 + 3M_3)/N$ , where  $M_1$ ,  $M_2$  and  $M_3$  represent the number of cells undergoing first second and  
21  
22 third mitosis and  $N$  is the total number of metaphase scored.  
23  
24  
25  
26

27  
28 Together with the individual mean value of SCEs per cell, in order to increase the sensitivity of the  
29  
30 assay, we calculated d the high frequency cells (HFC) and the number of high frequency individuals  
31  
32 (HFI). These measures took into account cells with a high frequency of SCEs that represented a  
33  
34 subpopulation of more sensitive cells or of long living lymphocytes which accumulated DNA  
35  
36 lesions *in vivo* (Carrano and Moore, 1982). The evaluation of HFC was performed to assess  
37  
38 individual variability in susceptibility to genotoxic agents. The occurrence of HFC in population  
39  
40 studies is generally evaluated using aggregate measures such as the HFI, i.e., subjects who show a  
41  
42 high proportion of HFC (Bonassi et al., 1999).  
43  
44  
45  
46  
47  
48

#### 49 *DNA extraction and Genotyping*

50  
51 DNA extraction was conducted using a Chelex solution, according to the following protocol: 10 µL  
52  
53 of peripheral blood was diluted in 1 mL of sterile distilled water for 15 min at room temperature.  
54  
55 After centrifugation at 14,000 rpm for 1 min, the pellet was re-suspended in 200 µL of 5% Chelex  
56  
57 solution in Tris-EDTA at pH 8, heated to 56°C for 15 min and, after vortex for 10 sec, at 100°C in  
58  
59  
60



1  
2 boiler water for 8 min. For PCR reactions we used 19  $\mu$ L of this solution containing extracted  
3  
4 DNA, whereas primers and methodologies are described in Pemble et al. (1994), Zhong et al.  
5  
6 (1993), García-Gonzalés et al. (2012), Chen et al. (2001), Bonello et al. (2010), Li et al. (2009),  
7  
8 Matullo et al. (2001) and Wang et al. (2010). PCR reactions were performed in a 25  $\mu$ L volume  
9  
10 containing about 10 ng DNA (template), with a final concentration of 1X Reaction Buffer, 1.5 mM  
11  
12 of  $MgCl_2$ , 5% of DMSO, 250  $\mu$ M of dNTPs, 0.5  $\mu$ M of each primer, and 1 U/sample of Taq DNA  
13  
14 polymerase (Fischer, U.S.). Cycles were set as follows: 35 cycles, 1 min at 95°C, 1 min at 60°C, 1  
15  
16 min at 72°C, and a final extension step 10 min at 72 °C. Amplification products were detected by  
17  
18 ethidium bromide staining after 3% agarose gel electrophoresis. In order to improve our results, all  
19  
20 null genotypes and a random 20% (n = 75) of the subjects were re-genotyped for all analyzed  
21  
22 polymorphisms. Only when the results of the two genotyping efforts were similar, did we include  
23  
24 the subject in our sample.  
25  
26  
27  
28  
29  
30

### 31 *Statistical analysis*

32  
33 Statistical analysis was assessed using the SPSS software statistical package programme (version  
34  
35 23.0 SPSS Inc., Chicago, IL). The Wilcoxon Mann-Whitney *U* test and the ANOVA with Tukey  
36  
37 correction test were used to analyse the differences in the frequency of SCEs between males and  
38  
39 females, smokers and non-smokers, and age groups, as well as to test the influence of the analysed  
40  
41 gene polymorphisms on the level of genomic damage. Multiple regression analysis was also used to  
42  
43 evaluate the influence of age, sex and smoke on SCEs frequency. All *P*-values were two tailed and  
44  
45 the level of statistical significance was set at  $P < 0.05$  for all tests.  
46  
47  
48  
49

## 50 **RESULTS**

### 51 *Study population*

52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 | The general characteristics of the study subjects are reported in Table 1. Ninety subjects were male  
4 (mean age $\pm$ SD 42.01 $\pm$ 9.34, range 22-70) and 140 were female (mean age $\pm$ SD 36.16 $\pm$ 8.94, range  
5  
6  
7 21-58).

8  
9  
10 | Eighty-three subjects were regular smokers (48 females and 35 males) while 147 were not (92  
11  
12 females and 55 males). Among smokers, the heavy smokers were 66 and the light smokers 17, with  
13  
14 an average number of cigarettes smoked per day of 17.83 $\pm$ 6.42 (ranging from 8 to 40). The long  
15  
16 time (LT) and short time (ST) smokers were 47 and 36, respectively, with an average number of  
17  
18 years of smoking of 14.26 $\pm$ 9.04 (ranging from 2 to 38).

19  
20  
21 | Fifty-four subjects fell into age group A (mean age 26.41 $\pm$ 2.84), 91 in age group B (mean age  
22  
23 36.42 $\pm$ 2.60), 59 in age group C (mean age 44.88 $\pm$ 2.81) and 26 in age group D (mean age  
24  
25 55.96 $\pm$ 4.83).

### 26 27 28 29 30 *SCEs analysis*

31  
32  
33 | The results of the SCEs analysis are summarized in Table 2. The mean frequency of SCEs and the  
34  
35 value of RI in the total sample were 5.15 $\pm$ 1.87 and 1.90 $\pm$ 0.24, respectively. According to the  
36  
37 Carrano and Moore (1982) methodology, 62 individuals were classified as HFI with a mean value  
38  
39 of SCEs/Cell of 7.54 $\pm$ 0.99, whereas 168 subjects were classified as Non-HFI, with a mean value of  
40  
41 SCEs/Cell of 4.26 $\pm$ 1.23. We observed significant differences between HFI and Non-HFI in terms of  
42  
43 SCEs frequency ( $P < 0.001$ ) and RI value ( $P = 0.03$ ).

44  
45  
46  
47  
48 | Similarly, sex seemed to play an important role in the determining the amount of genomic damage.  
49  
50 as females showed a significantly ( $P = 0.01$ ) higher level of SCEs with respect to males (5.36 $\pm$ 2.10  
51  
52 and 4.82 $\pm$ 1.39, respectively), as also confirmed by the ANOVA analysis ( $P = 0.03$ , Table 5).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

With regard to smoking habit, smokers showed a significantly ( $P < 0.001$ ) higher frequency of SCEs compared to non-smokers ( $5.93 \pm 1.75$  and  $4.70 \pm 1.79$ , respectively), whereas no statistical differences were found in terms of SCEs between heavy- and light-smokers ( $6.07 \pm 1.54$  and  $5.38 \pm 2.39$ , respectively) as well as between LT- and ST-smokers ( $5.84 \pm 1.63$  and  $6.04 \pm 1.92$ , respectively). However, LT-smokers showed a significantly lower RI value with respect to ST-smokers ( $P = 0.04$ ), indicating a possible cytotoxic effect of smoking over time. Finally, among smokers, the regression analysis indicated a significant correlation ( $P = 0.03$ ) between the number of cigarettes/day and the level of SCEs (Table 6).

Age was found to correlate with the RI value ( $P = 0.01$ ) but not with the SCEs frequency ( $P = 0.07$ ) (Table 6), although the ANOVA indicated significant differences in the level of SCEs among age-groups ( $P = 0.02$ , Table 5). Indeed, the 31-40 age-group showed a significantly lower SCEs frequency with respect to all other age-groups (Table 3).

### Gene polymorphisms

Finally, the influence of some gene polymorphisms on SCEs frequency was also evaluated (Tables 4 and 5). Our results showed that *GSTP2C19-AA*, *GSTT1-null*, *GSTM1-null*, *ERCC2/XPD Lys751Gln* and *XRCC1 Arg208His* gene polymorphisms were associated with significantly higher levels of SCEs, whereas this association was not found for the other analysed genic polymorphisms.

## DISCUSSION

In the present study, we evaluated, by means of the SCEs assay, the influence of some exogenous (smoking habits) and endogenous (age, gender, metabolic and DNA-repair genes polymorphisms) parameters on the level of the cytogenetic damage in the peripheral blood lymphocytes of non-occupationally exposed, healthy subjects living in the city of Turin (North Italy).

Although Turin is one of the most polluted cities in Europe (Santovito et al., 2016), mainly in terms

1  
2 of air fine particular matter whose mutagenic potential has been suggested in a number of studies  
3  
4 (Buschini et al., 2001; Wei and Meng, 2006), we observed a frequency of SCEs generally similar to  
5  
6 that found by Carere et al. (2002) for Rome and lower with respect to that observed by Barale et al.  
7  
8 (1998) for Pisa (Tuscany Region, Central Italy). On the contrary, in our previous published work  
9  
10 (Santovito et al., 2016), we reported a high baseline frequency of chromosomal aberrations (such as  
11  
12 breaks, dicentrics and rearrangements) in a control sample of subjects living in Turin, one of the  
13  
14 highest value reported in literature for European control populations.  
15  
16

17  
18  
19 It is known that spontaneous genomic damage can also be induced by a variety of other endogenous  
20  
21 and exogenous factors. Among them, smoking has been found to increase the level of SCEs and  
22  
23 other cytogenetic biomarkers in peripheral blood lymphocytes of many human populations (Salah  
24  
25 et al., 2011; Bonassi et al., 2011).  
26  
27

28  
29 According to data reported by other authors (Salah et al., 2011; Sebastià et al., 2014), a significant  
30  
31 increase in SCEs frequency was observed among smokers (Tables 2 and 5). Moreover, the  
32  
33 chromosomal damage seemed to correlate with the number of cigarettes/day (Table 6), indicating a  
34  
35 possible cumulative genotoxic effect of the cigarette smoke on human lymphocytes. This increased  
36  
37 level of genomic damage observed among smokers can be explained by the fact that cigarette  
38  
39 smoke contains several genotoxic compounds, most of them also having carcinogenic properties,  
40  
41 such as polycyclic aromatic hydrocarbons, aromatic amines and metals (IARC, 1986).  
42  
43  
44

45  
46 Contrary to what was observed in other reports (Salah et al., 2011; for a review see DeMarini, 2004  
47  
48 and Husgafvel-Pursiainen, 2004), in our work and other published studies, no significant  
49  
50 differences were found between light and heavy smokers and between LT- and ST-smokers in  
51  
52 terms of SCEs frequency. Finally, among smokers, no correlation was found between the number of  
53  
54 years of smoking and SCEs frequency (Table 6). A possible interpretation of this pattern has been  
55  
56 postulated by Costa et al. (2008), who hypothesized that various physiological systems (induction  
57  
58 of metabolizing and detoxifying enzymes, induction of DNA repair processes) had adopted over  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

time. In this sense a few cigarettes per day may stimulate a cell-adaptive response, thereby causing an apparent lowering in SCEs frequency. Moreover, as suggested by Donmez-Altuntas and Bitgen (2012) for other cytogenetic markers, it could be that cells damaged by cigarette smoke may not survive in culture or may not divide because they are more likely to die of necrosis or apoptosis, making it impossible to carry out the SCEs assay with them.

As for the role of sex, although in some studies (Santovito et al., 2015; Sebastia et al., 2014) no sex effect was observed, in our work females showed significantly higher SCE values than males, while sex had no effect on the replicative capacity of the cells, as indicated by the RI values (Tables 2 and 5).

The influence of age on the frequencies of SCEs have been evaluated in many studies (for a review see Bolognesi et al., 1997). In the present work, although the regression analysis indicated that the level of SCEs did not correlate with age (Table 6), we observed a significantly lower frequency of SCEs for the 31-40 age-group with respect to the 41-50 and 51-70 age-groups (Tables 3 and 5).

The increase in the amount of SCEs among subjects belonging to the last two age-groups could be explained by a decreased efficiency in the repair of DNA damage, with consequent accumulation of “aberrant cells”, in peripheral lymphocytes of older individuals. Indeed, it has been well documented that cells from older individuals exhibit increased levels of damaged DNA and chromosomal instability (Donmez-Altuntas and Bitgen, 2012; Milosevic-Djordjevic et al., 2002; Bukvic et al., 2001).

The individuals enrolled in our study were also genotyped for five phase I and phase II metabolic gene polymorphisms, as well as for four DNA-repair gene polymorphisms (Table 4). In agreement with data obtained by Kumar et al. (2011), our results showed that *GSTT1-null* and *GSTM1-null* gene polymorphisms were associated with increased cytogenetic damage. This result was not

1  
2 surprising because the reduced detoxification ability of the null genotypes has been related to an  
3  
4 increased susceptibility to DNA damage (Palma et al., 2007), as well as to an increased cancer risk  
5  
6 (Bajpai et al., 2007; Cha et al., 2007).  
7  
8  
9

10  
11 *CYP2C19 A/A* subjects also showed a frequency of SCEs significantly higher with respect to the  
12  
13 *CYP2C19 G/G* homozygote genotypes. Although other polymorphisms in genes belonging to *CYP2*  
14  
15 family, such as [these in the \*CYP2E1\* c1/c2 gene](#), were found to affect the frequency of SCEs  
16  
17 (Laczmanska et al., 2006), this is the first study showing a possible association of the *CYP2C19 A/A*  
18  
19 genotype with [increased](#) SCEs levels in a control population.  
20  
21  
22  
23

24  
25 Finally, analysing the effect of DNA-repair gene polymorphisms on the amount of genomic  
26  
27 damage, we found, for the first time in a control population, an association between *XPD*  
28  
29 *Gln751Gln* and *XRCC1 His208His* genotypes and increased levels of SCEs. It is known that high  
30  
31 frequencies of SCEs are related to defects in the DNA-repair machinery (Garcia-Sagredo, 2008). At  
32  
33 the same time, the *XPD* codon 751 Gln allele was found to be associated with lower DNA repair  
34  
35 capacity and lower cell viability in *in vitro* systems (Xiao et al., 2016; Zhang et al., 2017), and with  
36  
37 higher levels of DNA adducts (Palli et al., 2001; Matullo et al., 2003) and chromatid aberrations  
38  
39 (Ma et al., 2013). Similarly, the *XRCC1 His* allele results in defective DNA repair capacity, due to  
40  
41 the inefficient localisation of protein to the DNA damage site (Ji et al., 2010). In this scenario, we  
42  
43 can postulate that the reduced DNA repair capacity could [potentially contribute to](#) the higher levels  
44  
45 of SCEs observed among homozygous *XPD Gln751Gln* and *XRCC1 His208His* genotypes.  
46  
47  
48  
49  
50

51 [However, it should be emphasized that data related to the association of \*ERCC2/XPD Lys751Gln\*](#)  
52  
53 [and the cytogenetic damage are contradictory. In previous published studies, the \*XPD751Gln\*](#)  
54  
55 [variant allele was found to be associated with increased micronuclei frequencies \(Pérez-Cadahía,](#)  
56  
57 [2008\) but with decreased chromatid aberration frequencies \(Lunn et al., 2000; Vodicka et al.,](#)  
58  
59  
60

2015). *Vice versa*, in other studies the *Gln* allele failed to influence the levels of SCEs and DNA adducts (Duell et al., 2000). A possible explanation for these conflicting results could be that the accumulation of chromosomal aberrations requires a complex interplay between different DNA repair pathways. Gene-gene interactions in DNA repair genes could also influence enhanced or decreased chromosomal aberration frequencies. For example, Spitz et coll. (2001) observed the best repair activity in cells from wild-type individuals who were homozygous at both Lys751Gln and Asp312Asn loci and the lowest repair capacity in those carrying at least two variant alleles.

## CONCLUSIONS

Beyond the classical endogenous and exogenous factors, such as sex, age and smoking habits, which are already known to have in some cases a stronger effect on the level of genomic damage, we describe a positive association between CYP2C19 A/A, *GSTT-null*, *GSTM1-null*, *XPD 751 CC* and *XRCC1 His208His* genotypes with increased frequencies of SCEs. This finding reinforces, in bio-monitoring studies of human populations, the importance of genetic analysis designed to evaluate more classic endogenous and exogenous factors that could influence the level of the genomic damage. Moreover, our data assume a more important connotation if we consider the fact that, in the present study, we analysed a control population consisting of subjects not exposed for professional reasons to xenobiotics, but living in a city, like Turin, with many problems related to urban pollution (Traversi et al., 2009; Raaschou-Nielsen et al., 2013; Bono et al., 2016; Santovito et al., 2016). In this sense, we hope that the results of this study can be used as a stimulus for future bio-monitoring programs in other Italian and globally distributed cities.

Finally, it should be emphasize that the results of the present work cannot be generalized for all “Caucasians” because this group is heterogeneous, with differences in the distribution of genetic polymorphisms and in life styles among individuals.

## ACKNOWLEDGMENTS

1  
2 This investigation was supported by a grant from University of Turin (grant ex 60%). The authors  
3  
4 declare no conflicts of interest, with respect to the authorship and/or publication of this article.

5  
6 We would like to thank Sergio Castellano for his valuable help in improving the statistical analysis.  
7  
8  
9

#### 10 11 12 13 AUTHOR CONTRIBUTIONS

14  
15 Alfredo SANTOVITO designed the study, performed the laboratory analysis, analysed the data and  
16  
17 drafted the manuscript.

18  
19  
20 Claudio GENDUSA, collected the data, contributed to draft the manuscript and to perform the  
21  
22 laboratory analysis.

23  
24 Piero CERVELLA provided necessary logistical and economical support, analyzed the data and  
25  
26 edited the manuscript for intellectual content providing critical comments on the manuscript.  
27  
28  
29  
30

#### 31 32 LITERATURE CITED

- 33 1. Autrup, H. (2000). Genetic polymorphisms in human xenobiotica metabolizing enzymes as  
34 susceptibility factors in toxic response. *Mutat Res*, 464, 65-76.
- 35  
36  
37 2. Bajpai, P., Tripathi, A.K., & Agrawal, D. (2007). Increased frequencies of glutathione-S-  
38 transferase (GSTM1 and GSTT1) null genotypes in Indian patients with chronic myeloid  
39 leukaemia. *Leukemia Res*, 31, 407-420.
- 40  
41  
42  
43 3. Baltaci, V., Kayikçioğlu, F., Alpas, I., Zeyneloğlu, H. & Haberal, A. (2002). Sister  
44 chromatid exchange rate and alkaline comet assay scores in patients with ovarian cancer.  
45  
46  
47  
48  
49  
50  
51  
52  
53 4. Barale, R., Chelotti, L., Davini, T., Derl Ry, S., Andreassi, M.G., Ballardini, M., Bulleri, M.,  
54 He, J., Baldacci, S., Di Pede, F., Germignani, F., & Landi, S. (1998). Sister chromatid  
55 exchange and micronucleus frequency in human lymphocytes of 1,650 subjects in an Italian  
56 population: II. Contribution of sex, age, and lifestyle. *Environ Mol Mutagen*, 31(3), 228-  
57  
58  
59  
60 242.



- 1  
2  
3  
4 5. Bolognesi, C., Abbondandolo, A., Barale, R., Casalone, R., Dalprà, L., De Ferrari, M.,  
5 Degrassi, F., Forni, A., Lamberti, L., Lando, C., Migliore, L., Padovani, P., Pasquini, R.,  
6 Puntoni, R., Sbrana, I., Stella, M., & Bonassi, S. (1997). Age-related increase of baseline  
7 frequencies of sister chromatid exchanges, chromosome aberrations, and micronuclei in  
8 human lymphocytes. *Cancer Epidemiol Biomarkers Prev*, 6 (4), 249-256.  
9  
10  
11  
12
- 13  
14 6. Bonassi, S., Fontana, V., Ceppi, M., Barale, R., & Biggeri, A. (1999). Analysis of correlated  
15 data in human biomonitoring studies. The case of high sister chromatid exchange frequency  
16 cells. *Mutat Res-Gen Tox En*, 438 (1), 13-21.  
17  
18  
19
- 20  
21 7. Bonassi, S., Coskun, E., Ceppi, M., Lando, C., Bolognesi, C., Burgaz, S., Holland, N.,  
22 Kirsh-Volders, M., Knasmueller, S., Zeiger, E., Carnesoltas, D., Cavallo, D., da Silva, J., de  
23 Andrade, V.M., Dermicigil, G.C., Domínguez Odio, A., Domnez-Altuntas, H., Gattas, G.,  
24 Giri, A., Giri, S., Gómez-Meda, B., Gómez-Arroyo, S., Hadjidekova, V., Haveric, A.,  
25 Kamboj, M., Kurteshi, K., Martino-Roth, M.G., Montero Montoya, R., Nersesyan, A.,  
26 Pastor-Benito, S., Favero Salvadori, D.M., Shaposhnikova, A., Stopper, H., Thomas, P.,  
27 Torres-Bugarín, O., Yadav, A.S., Zúñiga González, G., & Fenech, M. (2011). The HUMAN  
28 MicroNucleus project on exfoliated buccal cells (HUMN(XL): the role of life-style, host  
29 factors, occupational exposures, health status, and assay protocol. *Mutat Res*, 728 (3), 88-  
30 97.  
31  
32  
33  
34  
35  
36  
37  
38
- 39 8. Bonello, L., Armero, S., Mokhtar, O.A., Mancini, J., Aldebert, P., Saut, N., Bonello, N.,  
40 Barragan, P., Arques, S., Giacomoni, M.P., Bonello-Burignat, C., Bartholomei, M.N.,  
41 Dignat-George, F., Camoi-Jau, L., & Paganelli, F. (2010). Clopidogrel Loading Dose  
42 Adjustment According to Platelet Reactivity Monitoring in Patients Carrying the 2C19\*2  
43 Loss of Function Polymorphism. *J Am Coll Cardiol* 56, 1630-1636.  
44  
45  
46  
47  
48
- 49 9. Bono, R., Romanazzi, V., Bellisario, V., Tassinari, R., Trucco, G., Urbino, A., Cassardo, C.,  
50 Siniscalco, C., Marchetti, P., & Marcon, A. (2016). Air pollution, aeroallergens and  
51 admissions to pediatric emergency room for respiratory reasons in Turin, northwestern Italy.  
52 *BMC Public Health*, 16, 722.  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
10. Bukvic, N., Gentile, M., Susca, F., Fanelli, M., Serio, G., Buonadonna, L., Capurso, A., & Guanti, G. (2001). Sex chromosome loss, micronuclei, sister chromatid exchange and aging: a study including 16 centenarians. *Mutat Res*, 498, 159–167.
  11. Buschini, A., Cassoni, F., Anceschi, E., Pasini, L., Poli, P., & Rossi, C. (2001). Urban Airbone particulate: genotoxicity evaluation of different size fractions by mutagenesis tests on microorganisms and comet assay. *Chemosphere*, 44, 1723-1736;
  12. Carere, A., Andreoli, C., Galati, R., Leopardi, P., Marcon, F., Rosati, M.V., Rossi, S., Tomei, F., Verdina, A., Zijno, A., & Crebelli, R. (2002). Biomonitoring of exposure to urban air pollutants: analysis of sister chromatid exchanges and DNA lesions in peripheral lymphocytes of traffic policemen. *Mutat Res*, 518, 215-224.
  13. Carrano, A.V., & Moore II, D.H. (1982). The rationale and methodology for quantifying sister chromatid exchange in humans. In J.A. Heddle (Ed.), *Mutagenicity: New Horizons in Genetic Toxicology*, (pp. 267–304). New York: Academic Press.
  14. Cha, I.H., Park, J.Y., Chung, W.Y., Choi, M.A., Kim, H.J., & Park, K.K. (2007). Polymorphisms of CYP1A1 and GSTM1 genes and susceptibility to oral cancer. *Yonsei Med J*, 48, 233-239.
  15. Chen, S., Xue, K., Xu, L., Ma, G., & Wu, J. (2001). Polymorphisms of the CYP1A1 and GSTM1 genes in relation to individual susceptibility to lung carcinoma in Chinese population. *Mutat Res*, 458, 41-47.
  16. Cleaver, J.E., Lam, E.T., & Revet, I. (2009). Disorders of nucleotide excision repair: the genetic and molecular basis of heterogeneity. *Nature*, 10, 756-768.
  17. Collins, A.R., & Azqueta, A. (2012). DNA repair as a biomarker in human biomonitoring studies: further applications of the comet assay. *Mutat Res*, 736, 122-129.
  18. Costa, S., Coelho, P., Costa, C., Silva, S., Mayan, O., Santos, L.S., Gaspar, J., & Teixeira, J.P. (2008). Genotoxic damage in pathology anatomy laboratory workers exposed to formaldehyde. *Toxicology*, 252 (1–3), 40-48.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
19. DeMarini, D. M. (2004). Genotoxicity of tobacco smoke and tobacco smoke condensate: a review. *Mut Res*, 567, 447-474.
  20. Donmez-Altuntas, H., & Bitgen, N. (2012). Evaluation of the genotoxicity and cytotoxicity in the general population in Turkey by use of the cytokinesis-block micronucleus cytome assay. *Mutat Res-Gen Tox En*, 748 (1-2), 1-7.
  21. Duell, E.J., Wiencke, J.K., Cheng, T.J., Varkonyi, A., Zuo, Z.F., Ashok, T.D., Mark, E.J., Wain, J.C., Christiani, D.C., & Kelsey, K.T. (2000). Polymorphisms in the DNA repair genes XRCC1 and ERCC2 and biomarkers of DNA damage in human blood mononuclear cells. *Carcinogenesis*, 21(5), 965-971.
  22. García-González, M.A., Quintero, E., Bujanda, L., Nicolás, D. Benito, R., Strunk, M., Santolaria, S., Sopeña, F., Badía, M., Hijona, E., Pérez-Aísa, M.A., Méndez-Sánchez, I.M., Thomson, C., Carrera, P., Piazuelo, E., Jiménez, P., Espinel, J., Campo, R., Manzano, M., Geijo, F., Pellisé, M., González-Huix, F., Espinós, J., Titó, L., Zaballa, M., Pazo, R., & Lanás, A. (2012). Relevance of GSTM1, GSTT1, and GSTP1 gene polymorphisms to gastric cancer susceptibility and phenotype. *Mutagenesis*, 27(6), 771-777.
  23. Garcia-Sagredo, J.M. (2008). Fifty years of cytogenetics: a parallel view of the evolution of cytogenetics and genotoxicology. *Biochim Biophys Acta*, 1779 (6-7), 363-375.
  24. Hemminki, K., Frank, C., Försti, A., Musak, L., Kazimirova, A, Barancokova, M., Horska, A., Vymetalkova, V., Smerhovsky, Z., Naccarati, A., Soucek, P., Vodickova, L., Buchancova, J., Smolkova, B., Dusinska, M., & Vodicka, P. (2015). Metabolic gene variants associated with chromosomal aberrations in healthy humans. *Genes Chrom Cancer*, 54(4), 260-266.
  25. Husgafvel-Pursiainen, K. (2004). Genotoxicity of environmental tobacco smoke: a review. *Mutat Res-Rev Mutat*, 567 (2-3), 427-445.
  26. Knudsen, L.E., & Hansen, Å.M. (2007). Biomarkers of intermediate endpoints in environmental and occupational health. *Int J Hyg Environ Health*, 210, 461-470.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
27. Kumar, A., Yadav, A., Giri, S.K., Dev, K., Gautam, S.K., Gupta, R., & Aggarwal, N. (2011). Influence of GSTM1 and GSTT1 genotypes and confounding factors on the frequency of sister chromatid exchange and micronucleus among road construction workers. *Chemosphere*, 84 (5), 564-570.
28. Ji, F., Wang, W., Xia, Z.L., Zheng, Y.J., Qiu, Y.L., Wu, F., Miao, W.B., Jin, R.F., Qian, J., Jin, L., Zhu, Y.L., & Christiani, D.C. (2010). Prevalence and persistence of chromosomal damage, and susceptible genotypes of metabolic and DNA repair genes in Chinese vinyl chloride-exposed workers. *Carcinogenesis* 31, 648-653.
29. International agency for research on cancer tobacco smoking (IARC). Monographs on the Evaluation of Carcinogenic Risks to Humans, vol. 38 IARC, Lyon, France (1986).
30. Laczmanska, I., Gil, J., Karpinski, P., Stembalska, A., Kozłowska, J., Busza, H., Trusewicz, A., Pesz, K., Ramsey, D., Schlade-Bartusiak, K., Blin, N., & Sasiadek, M.M. (2006). Influence of polymorphisms in xenobiotic-metabolizing genes and DNA-repair genes on diepoxybutane-induced SCE frequency. *Environ Mol Mutagen*, 47(9). 666-673.
31. Laczmanska, I., Gil J., Karpinski, P., Stembalska, A., Trusewicz, A., Pesz, K., Ramsey, D., Schlade-Bartusiak, K., Blin, N., Sasiadek, M.M. (2007). Polymorphism in nucleotide excision repair gene XPC correlates with bleomycin-induced chromosomal aberrations. *Environ Mol Mutagen*, 48, 666-671.
32. Li, Y., Marion, M-J, Zipprich, J., Santella, R. M., Freyer, G. & Brandt-Rauf, P. W. (2009). Gene-environment interactions between DNA repair polymorphisms and exposure to the carcinogen vinyl chloride. *Biomarkers*, 14(3), 148-155.
33. Lunn, R. M., Helzlsouer, K. J., Parshad, R., Umbach, D. M., Harris, E. L., Sanford, K.K., & Bell, D. A. (2000). XPD polymorphisms: effects on DNA repair proficiency. *Carcinogenesis*, 21, 551-555.
34. Ma, Q., Qui, C., Tie, C., & Guo, Z. (2013). Genetic polymorphisms of xeroderma pigmentosum group D gene Asp312 Asn and Lys751Gln and susceptibility to prostate cancer: a systematic review and meta-analysis. *Gene*, 530(2), 309-314.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
35. Matullo, G., Palli, D., Peluso, M., Guarrera, S., Carturan, S., Celentano, E., Krogh, V., Munnia, A., Tumino, R., Polidoro, S., Piazza, A., & Vineis, P. (2001). XRCC1, XRCC3, XPD gene polymorphisms, smoking and (32)P-DNA adducts in a sample of healthy subjects. *Carcinogenesis*, 22(9):1437-1445.
36. Matullo, G., Peluso, M., Polidoro, S., Guarrera, S., Munnia, A., Krogh, V., Masala, G., Berrino, F., Panico, S., Tumino, R., Vineis, P., & Palli, D. (2003). Combination of DNA repair gene single nucleotide polymorphisms and increased levels of DNA adducts in a population-based study. *Cancer Epidemiol Biom Prev*, 12, 674-677.
37. Medves, S., Auchter, M., Chambeau, L., Gazzo, S., Poncet, D., Grangier, B., Verney, A., Moussay, E., Ammerlaan, W., Brisou, G., Morjani, H., Géli, V., Palissot, V., Berchem, G., Salles, G., & Wenner, T. (2016) A high rate of telomeric sister chromatid exchange occurs in chronic lymphocytic leukaemia B-cells. *Br J Haematol*, 174(1), 57-70.
38. Milosevic-Djordjevic, O., Grujicic, D., Novakovic, T., Arsenijevic, S., & Marinkovic, D.(2002). Micronuclei and ageing in a sample of Yugoslavian population. *Russ J Genet (Genetika)*, 38, 201-204.
39. Palli, D., Russo, A., Masala, G., Saieva, C., Guarrera, S., Carturan, S., Munnia, A., Matullo, G., Peluso, M. (2001). DNA adduct levels and DNA repair polymorphisms in traffic-exposed workers and a general population sample. *Int J Cancer*, 94(1), 121-127.
40. Palma, S., Cornetta, T., Padua, L., Cozzi, R., Appolloni, M., Ievoli, E., & Testa A (2007). Influence of glutathione S-transferase polymorphisms on genotoxic effects induced by tobacco smoke. *Mut Res*, 633, 1-12.
41. Pemble, S., Schroeder, K. R., Spencer, S. R., Meyer, D. J., Hallier, E., Bolt, H.M., Ketterer, B., & Taylor, J.B. (1994). Human glutathione S-transferase theta (GSTT1): cDNA cloning and the characterization of a genetic polymorphism. *Biochem J*, 300, 271-276.
42. Pérez-Cadahía, B., Laffon, B., Valdiglesias, V., Pásaro, E., & Méndez, J. (2008). Cytogenetic effects induced by Prestige oil on human populations: the role of polymorphisms in genes involved in metabolism and DNA repair. *Mutat Res*, 653(1-2), 117-123.

- 1  
2  
3  
4 43. Raaschou-Nielsen, O., Andersen, Z.J., Beelen, R., Samoli, E., Stafoggia, M., Weinmayr, G.,  
5 Hoffmann, B., Fischer, P., Nieuwenhuijsen, M.J., Brunekreef, B., Xun, W.W., Katsouyanni,  
6 K., Dimakopoulou, K., Sommar, J., Forsberg, B., Modig, L., Oudin, A., Oftedal, B.,  
7 Schwarze, P.E., Nafstad, P., De Faire, U., Pedersen, N.L., Ostenson, C.G., Fratiglioni, L.,  
8 Penell, J., Korek, M., Pershagen, G., Eriksen, K.T., Sørensen, M., Tjønneland, A.,  
9 Ellermann, T., Eeftens, M., Peeters, P.H., Meliefste, K., Wang, M., Bueno-de-Mesquita, B.,  
10 Key, T.J., de Hoogh, K., Concin, H., Nagel, G., Vilier, A., Grioni, S., Krogh, V., Tsai,  
11 M.Y., Ricceri, F., Sacerdote, C., Galassi, C., Migliore, E., Ranzi, A., Cesaroni, G., Badaloni,  
12 C., Forastiere, F., Tamayo, I., Amiano, P., Dorronsoro, M., Trichopoulou, A., Bamia, C.,  
13 Vineis, P., & Hoek, G. (2013). Air pollution and lung cancer incidence in 17 European  
14 cohorts: prospective analyses from the European Study of Cohorts for Air Pollution Effects  
15 (ESCAPE). *Lancet Oncol*, 14(9), 813-822.
- 16 44. Salah, G.B., Kamoun H., Rebai A., Youssef, A.B., Ayadi H., Belghith-Mahfoudh, N.,  
17 Fourati, A., Ayadi, H., & Fakhfakh, F. (2011). Sister chromatid exchange (SCE) and high-  
18 frequency cells (HFC) in peripheral blood lymphocytes of healthy Tunisian smokers. *Mut*  
19 *Res-Gen Tox En*, 719 (1-2), 1-6.
- 20 45. Santovito, A., Cervella, P., & Delpero, M. (2014). Increased frequency of chromosomal  
21 aberrations and sister chromatid exchanges in peripheral lymphocytes of radiology  
22 technicians chronically-exposed to low levels of ionizing radiations. *Environ Toxicol*  
23 *Pharmacol*, 37, 396-403.
- 24 46. Santovito, A., Cervella, P., & DelPero, M. (2015). Evaluation of genomic damage in  
25 peripheral lymphocytes from occupationally exposed anesthetists: assessment of the effects  
26 of age, sex, and GSTT1 Gene Polymorphism. *J Biochem Mol Toxicol*, 29 (5), 234-239.
- 27 47. Santovito, A., Cervella, P., & Delpero, M. (2016). Baseline frequency of chromosomal  
28 aberrations and sister chromatid exchanges in peripheral blood lymphocytes of healthy  
29 individuals living in Turin (North-Western Italy): assessment of the effects of age, sex and  
30 GSTs gene polymorphisms on the levels of genomic damage. *Ann Hum Biol*, 43(3), 269-  
31 278.
- 32 48. Sebastià, N., Hervás, D., Almonacid, M., Villaescusa, J. I., Soriano, J. M., Sahuquillo, V.,  
33 Esteban, V., Barquinero, J. F., Verdú, G., Cervera, J., Such, E., & Montoro, A. (2014).
- 34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2 Sister chromatid exchange (SCE), High-Frequency Cells (HFCs) and SCE distribution  
3 patterns in peripheral blood lymphocytes of Spanish adult smokers compared to non-  
4 smokers. *Food Chem Toxicol*, 66, 107-112.  
5  
6  
7  
8  
9 49. Spitz, M.R., Wu, X., Wang, Y., Wang, L.E., Shete, S., Amos, C.I., Guo, Z., Lei, L.,  
10 Mohrenweiser, H., & Wei, Q. (2001) Modulation of nucleotide excision repair capacity by  
11 XPD polymorphisms in lung cancer patients. *Cancer Res*, 61, 1354-1357.  
12  
13  
14  
15 50. Toolaram, A.P., Kümmerer, K., & Schneider, M. (2014). Environmental risk assessment of  
16 anti-cancer drugs and their transformation products: A focus on their genotoxicity  
17 characterization-state of knowledge and short comings. *Mutat Res-Rev Mutat*, 760, 18-35.  
18  
19  
20  
21 51. Traversi, D., Degan, R., De Marco, R., Gilli, G., Pignata, C., Villani, S., & Bono, R. (2009).  
22 Mutagenic properties of PM2.5 urban pollution in the Northern Italy: The nitro-compounds  
23 contribution. *Environ Int*, 35, 905-910.  
24  
25  
26  
27  
28  
29 52. Vodicka, P., Kumar, R., Stetina, R., Sanyal, S., Soucek, P., Haufroid, V., Dusinska M.,  
30 Kuricova, M., Zamecnikova, M., Musak, L., Buchancova, J., Norppa, H., Hirvonen, A.,  
31 Vodickova L., Naccarati, A., Matousu, Z., & Hemminki, K. (2004). Genetic polymorphisms  
32 in DNA repair genes and possible links with DNA repair rates, chromosomal aberrations  
33 and single-strand breaks in DNA. *Carcinogenesis* 25:757-63.  
34  
35  
36  
37  
38  
39 53. Vodicka, P., Musak, L., Frank, C., Kazimirova, A., Vymetalkova, V., Barancokova, M.,  
40 Smolkova, B., Dzapinkova, Z., Jiraskova, K., Vodenkova, S., Kroupa, M., Osina, O.,  
41 Naccarati, A., Palitti, F., Försti, A., Dusinska, M., Vodickova, L., & Hemminki, K. (2015).  
42 Interactions of DNA repair gene variants modulate chromosomal aberrations in healthy  
43 subjects. *Carcinogenesis*, 36 (11), 1299-1306.  
44  
45  
46  
47  
48  
49 54. Wang, Q., Ji, F., Sun, Y., Qui, Y. L., Wang, W., Wu, F., Miao, W., Li, Y., Brandt-Rauf, P.  
50 W. & Xia, Z. (2010). Genetic polymorphisms of XRCC1, HOGG1 and MGMT and  
51 micronucleus occurrence in Chinese vinyl chloride-exposed workers. *Carcinogenesis*, 31(6),  
52 1068-1073.  
53  
54  
55  
56  
57 55. Wang, Q., Tan, H., Zhang, F., Sun, Y., Feng, N., Zhou, L., Ye, Y., Zhu Y., Li Y., Brandt-  
58 Rauf P. W., Shao, H., & Xia, Z. (2013) Polymorphisms in BER and NER pathway genes:  
59  
60

- 1  
2  
3 Effects on micronucleus frequencies among vinyl chloride-exposed workers in northern  
4 China. *Mut Res-Gen Tox En*, 754 (1-2), 7-14.  
5  
6  
7  
8 56. Wei, A., & Meng, Z. (2006). Induction of chromosome aberrations in cultured human  
9 lymphocytes treated with sand dust storm fine particles (PM2.5). *Toxicol Lett*, 166, 37-43.  
10  
11  
12 57. Xiao, S., Cui, S., Lu, X., Guan, Y., Li, D., Liu, Q., Cai, Y., Jin, C., Yang, J., Wu, S., & van  
13 der Straaten, T. (2016). The ERCC2/XPD Lys751Gln polymorphism affects DNA repair of  
14 benzo[a]pyrene induced damage, tested in an in vitro model. *Toxicol in Vitro*, 34, 300-308.  
15  
16  
17  
18 58. Zhang, G., Guan, Y., Zhao, Y., van der Straaten, T., Xiao, S., Xue, P., Zhu, G., Liu, Q.,  
19 Cai, Y., Jin, C., Yang, J., Wu, S., & Lu, X. (2017). *ERCC2/XPD* Lys751Gln alter DNA  
20 repair efficiency of platinum-induced DNA damage through P53 pathway. *Chem-Biol*  
21 *Interact*, 263, 55-65.  
22  
23  
24  
25  
26  
27 59. Zhong, S., Wyllie, A.H., Barnes, D., Wolf, C. R., & Spurr, N. K. (1993). Relationship  
28 between GSTM1 genetic polymorphism and susceptibility to bladder, breast and colon  
29 cancer. *Carcinogenesis*, 14, 1821-1824.  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



Table 1 – General characteristics of the studied subjects

Subjects	N	Age (Mean ± S.D.)	Age range
<b>Total</b>	<b>230</b>	<b>38.45±9.51</b>	<b>21-70</b>
<b>Sex</b>			
Males	90	42.01±9.34	22-70
Females	140	36.16±8.94	21-58
<b>Smoking Habit</b>			
Smokers	83	39.60±10.65	22-70
Non-Smokers	147	37.80±8.79	21-66
Heavy Smokers	66	40.20±10.61	22-70
Light Smokers	17	37.29±10.81	22-58
LT - Smokers	47	46.92±7.44	33-70
ST - Smokers	36	30.06±5.25	22-42
<b>Age groups</b>			
A	54	26.41±2.84	21-30
B	91	36.42±2.60	31-40
C	59	44.88±2.81	41-50
D	26	55.96±4.83	51-70

N = number of studied subjects; S.D. = Standard Deviation;

Heavy Smokers = >10 cig/die; Light-Smokers = ≤10 cig/die

LT - Smokers = Long Time Smokers = >10 years of smoking habit

ST – Smokers = Short Time Smokers = ≤10 years of smoking habit

Table 2 – Frequency of SCEs in the studied population according to sex and smoking habit

Groups	N	Cells	SCEs	SCEs/Cell± S.D.	M <sub>1</sub>	M <sub>2</sub>	M <sub>3</sub>	RI ± S.D.
<b>Total</b>	<b>230</b>	<b>11500</b>	<b>59172</b>	<b>5.15±1.87</b>	<b>8027</b>	<b>8992</b>	<b>5940</b>	<b>1.90±0.24</b>
HFI	62	3100	23379	7.54±0.99 <sup>a</sup>	2338	2662	1165	1.80±0.26 <sup>d</sup>
Non-HFI	168	8400	35793	4.26±1.23 <sup>a</sup>	5690	6328	4776	1.95±0.22 <sup>d</sup>
<b>Sex</b>								
Males	90	4500	21685	4.82±1.39 <sup>b</sup>	3102	3505	2389	1.92±0.19
Females	140	7000	37487	5.36±2.10 <sup>b</sup>	4926	5485	3552	1.90±0.27
<b>Smoking Habit</b>								
Non-Smokers	147	7350	34573	4.70±1.79 <sup>c</sup>	5022	5763	3902	1.92±0.22
Smokers	83	4150	24599	5.93±1.75 <sup>c</sup>	3006	3227	2039	1.88±0.28
Heavy Smokers	66	3300	20026	6.07±1.54	2404	2545	1623	1.87±0.25
Light Smokers	17	850	4573	5.38±2.39	602	682	416	1.89±0.36
LT Smokers	47	2350	13722	5.84±1.63	1775	1891	997	1.81±0.22 <sup>e</sup>
ST Smokers	36	1800	10877	6.04±1.92	1231	1136	1042	1.96±0.33 <sup>e</sup>

<sup>a,c</sup>  $P < 0.001$ ; <sup>b</sup>  $P = 0.01$ ; <sup>d</sup>  $P = 0.03$ ; <sup>e</sup>  $P = 0.04$  (Wilcoxon Mann-Whitney  $U$  test)

HFI = High Frequency Individuals; LT = Long Time Smokers (>10 years of smoking habit); ST = Short Time smokers ( $\leq 10$  years of smoking habit); N = Number of analysed subjects; SCEs = Sister chromatid exchanges; Metaphases; RI (Replication Index) =  $(M_1 + 2M_2 + 3M_3)/N$ , where  $M_1$ ,  $M_2$  and  $M_3$  represent the number of cells undergoing first second and third mitosis and N is the total number of metaphase scored; S.D. = Standard Deviation.

Table 3 – SCEs frequency according to age groups

Age Groups	N	Cells	SCEs	SCEs/Cell± S.D.	M <sub>1</sub>	M <sub>2</sub>	M <sub>3</sub>	RI ± S.D.
A (21-30)	54	2700	14530	5.38±2.02 <sup>a</sup>	1930	1959	1520	1.93±0.30 <sup>d</sup>
B (31-40)	91	4550	20453	4.50±1.81 <sup>a,b,c</sup>	2971	3543	2582	1.95±0.21 <sup>e f</sup>
C (41-50)	59	2950	16757	5.68±1.77 <sup>b</sup>	2154	2492	1205	1.83±0.24 <sup>e f g</sup>
D (51-70)	26	1300	7432	5.72±1.26 <sup>c</sup>	972	998	633	1.87±0.20 <sup>d g</sup>

<sup>a</sup> P = 0.01; <sup>b</sup> P = <0.001; <sup>c</sup> P = 0.001; <sup>d</sup> P = 0.04; <sup>e</sup> P = 0.01; <sup>f</sup> P = 0.04; <sup>g</sup> P = 0.03 (Wilcoxon Mann-Whitney *U* test)

N = Number of analysed subjects; SCEs = Sister chromatid exchanges; Metaphases; RI (Replication Index) = (M<sub>1</sub> + 2M<sub>2</sub> + 3M<sub>3</sub>)/N, where M<sub>1</sub>, M<sub>2</sub> and M<sub>3</sub> represent the number of cells undergoing first second and third mitosis and N is the total number of metaphase scored; S.D. = Standard Deviation.

For Peer Review

Table 4 – SCEs frequency according to genetic polymorphisms

Gene Polymorphisms	N	Cells	SCEs	SCEs/Cell± S.D.	M <sub>1</sub>	M <sub>2</sub>	M <sub>3</sub>	RI ± S.D.
<b>PHASE I</b>								
<i>CYP1A1 Ile/Ile</i>	177	8850	44395	5.02±1.87	6082	6837	4738	1.92±0.25
<i>CYP1A1 Ile/Val</i>	44	2200	12093	5.50±1.86	1615	1802	983	1.86±0.18
<i>CYP1A1 Val/Val</i>	9	450	2684	5.96±1.69	331	351	220	1.88±0.28
<i>CYP2C19 GG</i>	202	10100	51179	5.07±1.81 <sup>a,b</sup>	7025	7838	5310	1.91±0.24
<i>CYP2C19 G/A</i>	23	1150	6554	5.70±1.97 <sup>a</sup>	845	929	512	1.84±0.24
<i>CYPC192 A/A</i>	5	250	1440	5.76±3.34 <sup>b</sup>	157	225	118	1.92±0.29
<b>PHASE II</b>								
<i>GSTT1+</i>	192	9600	47191	4.92±1.69 <sup>c</sup>	6679	509	4979	1.91±0.24
<i>GSTT1-</i>		1900	11981	6.31±2.28 <sup>c</sup>	1349	1481	962	1.89±0.25
<i>GSTM1+</i>	168	8400	41584	4.95±1.84 <sup>d</sup>	5727	6635	4397	1.92±0.24
<i>GSTM1-</i>		3100	17588	5.67±1.86 <sup>d</sup>	2301	2355	1544	1.88±0.23
<i>GSTT1+/GSTM+</i>	125	6250	31066	4.97±1.71	4379	4971	3117	1.89±0.25
<i>GSTT1-/GSTM1-</i>		1700	9231	5.43±2.06	1151	1359	889	1.93±0.22
<i>GSTT1+/GSTM1-</i>	50	2500	12574	5.03±1.80	1779	1855	1367	1.92±0.25
<i>GSTT1-/GSTM1+</i>		1050	6301	6.00±2.40	718	807	567	1.92±0.23
<i>GSTP1 AA</i>	186	9300	46711	5.02±1.80	6561	7238	4771	1.90±0.25
<i>GSTP1 AG</i>		1800	10088	5.60±2.06	1190	1444	955	1.93±0.23
<i>GSTP1 GG</i>		400	2373	5.93±2.24	276	310	214	1.92±0.24
<b>DNA-REPAIR</b>								
<i>ERCC2/XPD Lys751Lys</i>	185	9250	45375	4.91±1.72 <sup>e,f</sup>	6420	7214	4834	1.91±0.25
<i>ERCC2/XPD Lys751Gln</i>		1450	8423	5.81±2.01 <sup>e</sup>	1063	1161	677	1.87±0.23
<i>ERCC2/XPD Gln751Gln</i>	16	800	5374	6.72±2.31 <sup>f</sup>	544	617	429	1.92±0.26

<i>XRCCI Arg194Arg</i>								
<i>XRCCI Arg194Trp</i>	192	9600	48673	5.07±1.89	6727	7497	4931	1.90±0.25
<i>XRCCI Trp194Trp</i>	31	1550	8504	5.49±1.65	1070	1198	836	1.92±0.25
	7	350	1994	5.70±2.25	230	297	1343	1.92±0.18
<i>XRCCI Arg399Arg</i>								
<i>XRCCI Arg399Gln</i>	178	8900	45469	5.11±1.91	6188	6938	4641	1.91±0.25
<i>XRCCI Gln399Gln</i>	45	2250	11870	5.28±1.73	1592	1759	1141	1.90±0.24
	7	350	1832	5.24±1.91	247	295	158	1.87±0.19
<i>XRCCI Arg208Arg</i>								
<i>XRCCI Arg208His</i>	200	10000	50695	5.07±1.88 <sup>g</sup>	6956	7883	5133	1.90±0.24
<i>XRCCI His208His</i>	26	1300	7062	5.43±1.69 <sup>h</sup>	957	923	707	1.89±0.27
	4	200	1415	7.08±1.22 <sup>g,h</sup>	114	186	100	1.97±0.16

<sup>a,b,c,f,h</sup>  $P < 0.001$ ; <sup>d</sup>  $P = 0.01$ ; <sup>e</sup>  $P = 0.01$ ; <sup>g</sup>  $P = 0.001$  (Wilcoxon Mann-Whitney  $U$  test)

N = Number of analysed subjects; SCEs = Sister chromatid exchanges; Metaphases; RI (Replication Index) =  $(M_1 + 2M_2 + 3M_3)/N$ , where  $M_1$ ,  $M_2$  and  $M_3$  represent the number of cells undergoing first second and third mitosis and N is the total number of metaphase scored; S.D. = Standard Deviation;

Table 5 - Factors affecting Sister Chromatid Exchanges  
analyzed by ANOVA

Factors	F-value	P-value
<b>Demographic characteristics</b>		
Age groups	6.87	0.02
Sex	4.58	0.03
<b>Smoking habit</b>		
Smoke vs Non smokers	25.13	<0.001
Heavy vs Light	2.12	0.15
Long-Time vs Short-Time	0.28	0.60
<b>Gene Polymorphisms</b>		
CYP1A1 Ile/Val	2.08	0.28
CYP2C19 G/A	14.62	<0.001
GSTT1 +/-	18.88	<0.001
GSTM1 +/-	6.93	0.01
GSTP1 A/G	2.22	0.20
ERCC Lys751Glc	9.68	0.03
XRCC Arg194trp	0.97	0.49
XRCC Arg399Glc	1.39	0.22
XRCC Arg208His	12.62	0.00

Table 6. Multiple regression analysis of confounding factors on SCEs and RI values in lymphocytes of the study groups

CF	SCEs frequency			RI value		
	$\beta$ -co	<i>P</i> -value	95% CI (Lower) - (Upper)	$\beta$ -co	<i>P</i> -value	95% CI (Lower) - (Upper)
Age	0.10	0.07	(-0.89 – 24.85)	-0.16	0.01	(-0.01) - (-0.00)
Cig/day	0.23	0.03	(0.35) - (6.20)	0.04	0.79	(-0.62) - (0.88)
Years of smoking	-0.05	0.67	(-2.60) - (1.68)	0.04	0.69	(-0.43) - (0.64)

CF = Confounding Factor;  $\beta$ -co =  $\beta$ -coefficient; SCE = Sister Chromatid Exchanges; RI = Replication Index

For Peer Review