

1 **Plasticizers used in food-contact materials affect adipogenesis** 2 **in 3T3-L1 cells**

3

4 Valentina Pomatto^a, Erika Cottone^a, Paolo Cocci^b, Matteo Mozzicafreddo^b, Gilberto
5 Mosconi^b, Erik Russel Nelson^c, Francesco Alessandro Palermo^b, Patrizia Bovolin^{a*}

6

7 ^aDepartment of Life Sciences and Systems Biology, University of Turin, 10123 Turin, Italy

8 ^bSchool of Biosciences and Veterinary Medicines, University of Camerino, 62032 Camerino,
9 Italy

10 ^cDepartment of Molecular and Integrative Physiology, University of Illinois at Urbana-
11 Champaign, Urbana, IL 61801, USA; University of Illinois Cancer Center, Chicago, IL 60612,
12 USA; Division of Nutritional Sciences, University of Illinois at Urbana-Champaign, Urbana,
13 IL 61801, USA

14

15

16 *Corresponding author:

17 Patrizia Bovolin

18 Department of Life Sciences and Systems Biology, University of Turin

19 Via Accademia Albertina 13, 10123 Turin, Italy

20 Tel. +390116704679

21 Fax +390116704508

22 E-mail: patrizia.bovolin@unito.it

1 **Abstract**

2 Recent studies suggest that exposure to some plasticizers, such as Bisphenol A (BPA), play
3 a role in endocrine/metabolic disruption and can affect lipid accumulation in adipocytes.
4 Here, we investigated the adipogenic activity and nuclear receptor interactions of four
5 plasticizers approved for the manufacturing of food-contact materials (FCMs) and currently
6 considered safer alternatives. Differentiating 3T3-L1 mouse preadipocytes were exposed
7 to scalar concentrations (0.01-25 μ M) of DiNP (Di-iso-nonyl-phthalate), DiDP (Di-iso-decyl-
8 phthalate), DEGDB (Diethylene glycol dibenzoate), or TMCP (Tri-m-cresyl phosphate).
9 Rosiglitazone, a well-known pro-adipogenic peroxisome proliferator activated receptor
10 gamma (PPAR γ) agonist, and the plasticizer BPA were included as reference compounds.
11 All concentrations of plasticizers were able to enhance lipid accumulation, with TMCP being
12 the most effective one. Accordingly, when comparing *in silico* the ligand binding efficiencies
13 to the nuclear receptors PPAR γ and retinoid-X-receptor-alpha (RXR α), TMPC displayed the
14 highest affinity to both receptors. Differently from BPA, the four plasticizers were most
15 effective in enhancing lipid accumulation when added in the mid-late phase of differentiation,
16 thus suggesting the involvement of different intracellular signalling pathways. In line with
17 this, TMCP, DiDP, DiNP and DEGDB were able to activate PPAR γ in transient transfection
18 assays, while previous studies demonstrated that BPA acts mainly through other nuclear
19 receptors. qRT-PCR studies showed that all plasticizers were able to increase the
20 expression of CCAAT/enhancer binding protein β (*Cebp β*) in the early steps of adipogenesis,
21 and the adipogenesis master gene *Ppar γ 2* in the middle phase, with very similar efficacy to
22 that of Rosiglitazone. In addition, TMCP was able to modulate the expression of both Fatty
23 Acid Binding Protein 4/Adipocyte Protein 2 (*Fabp4/Ap2*) and Lipoprotein Lipase (*Lpl*)

1 transcripts in the late phase of adipogenesis. DEGDB increased the expression of *Lpl* only,
2 while the phthalate DiDP did not change the expression of either late-phase marker genes
3 *Fabp4* and *Lpl*. Taken together, our results suggest that exposure to low, environmentally
4 relevant doses of the plasticizers DiNP, DiDP DEGDB and TMCP increase lipid
5 accumulation in 3T3-L1 adipocytes, an effect likely mediated through activation of PPAR γ
6 and interference at different levels with the transcriptional cascade driving adipogenesis.

7

8 **Keywords:** plasticizer; endocrine disruptor; phthalates; adipogenesis; nuclear receptors;
9 lipid accumulation

10

11

12 This work was supported by MIUR-PRIN (Ministero dell'Istruzione, dell'Università e della
13 Ricerca, Progetti di Ricerca di Interesse Nazionale) prot. 2010W87LBJ_005 to PB, and prot.
14 2010W87LBJ_002 to GM; and Fondazione CRT (Cassa di Risparmio di Torino) RF
15 2014.0814 to PB.

16

1 **1. Introduction**

2 Obesity is the fastest growing health problem in Europe and worldwide. In the European
3 Union, overweight affects between 36% and 67.5% of adults, while obesity affect between
4 10% and 28% of adults (last update 2014) [1]. In addition to genetic factors, life style factors
5 such as excessive caloric intake, high fat diets, and low physical activity contribute to
6 obesity. However, there is also increasing evidence that environmental pollutants including
7 endocrine-disrupting chemicals (EDCs) may contribute to the development of obesity and
8 metabolic disorders. A subset of EDCs have been named "obesogens" or "metabolic
9 disruptors" [2–5], because of their ability to promote adiposity by altering fat cell
10 development and increasing energy storage of fat tissue, and because of their implication in
11 metabolic syndrome and obesity [6].

12 The EU regulation (1907/2006 and subsequent updates) regarding the Registration,
13 Evaluation, Authorisation and Restriction of Chemicals (REACH) has identified so far 181
14 substances of very high concern (SVHC) for the environment and human health (last update
15 January 2018). Several SVHC are plasticizers, a class of diverse additives used in plastics
16 production, that are poorly bound or not bound to the polymers. These features facilitate
17 their migration from food-contact materials (FCMs) and several household plastic items, thus
18 coming in contact with humans through food consumption, skin absorption and inhalation
19 [7]. FCMs, including plastic packaging, are not generally perceived to be a chemical health
20 threat when compared to pesticides, veterinary drugs or heavy metals arising from
21 agricultural practices or environmental contamination. However, within the last decade it has
22 been increasingly reported that certain FCMs can act like EDCs [8]; a good example are
23 plastic additives used in food containers like Bisphenol A (BPA), a substance recently

1 included in the SVHC list and whose impact on the endocrine system has been increasingly
2 reported [2,3].

3 The EFSA (European Food Safety Authority) regulation 10/2011 has provided a list of
4 plasticizers permitted in EU for FCMs manufacturing, which has become a useful source of
5 alternatives to currently used SVHC. In the present work, we focused our attention on four
6 plasticizers employed in food packaging: Di-iso-nonyl-phthalate (DiNP), Di-iso-decyl-phthalate
7 (DiDP), Diethylen glycol dibenzoate (DEGDB), and Tri-m-cresyl phosphate (TMCP).
8 Notably, DiNP and DiDP are comprised in the EFSA list of permitted compounds and are
9 indeed among the most used in the plastic market (33% United States; 63% European
10 Union) as substitutes of di(2-ethylhexyl) phthalate (DEHP), a substance classified as SVHC
11 [9,10]. DEGDB is another emerging plasticizer designed to substitute phthalates, since it is
12 considered more eco-friendly due to its biodegradation pathways [11]. Tri-cresyl
13 phosphates, such as tri-m-cresyl phosphate (TMCP), are mainly used as substitutes of the
14 plasticizers polybrominated diphenyl ethers (e.g. BDE-47) [12]. Along with the increased
15 usage of these SVHC substitutes as alternative plasticizers, new biomonitoring data are
16 becoming available associating the exposure to these chemicals with adverse effects in
17 living beings. Notably, DiNP and DiDP have both been associated with increased insulin
18 resistance in adolescent cohorts [13] and in general with several different adverse effects
19 after peri- and post-natal exposure [14]. Interestingly, *in silico* approaches demonstrated
20 that DiNP and DiDP can act as ligands of human peroxisome proliferator activated receptor
21 γ (PPAR γ) and retinoid-X-receptor- α (RXR α), possibly triggering a cascade of intracellular
22 events [15]. DiDP is also a confirmed modulator of PPAR:RXR-dependent gene expression
23 pathways in fish hepatocytes [16]. Similarly, TMCP was found to affect lipid/cholesterol

1 metabolism through a functional interplay between PPARs and liver X receptor (LXR) in a
2 fish *in vitro* system [17]. Also, in fish DEGDB was demonstrated to have high affinities for
3 PPAR α , RXR α and LXR, showing the ability to modulate PPAR α transcriptional pathways
4 [18].

5 The 3T3-L1 preadipocyte cell line has proved to be a useful tool to study *in vitro* mechanisms
6 by which obesogens can affect lipid accumulation and adipocyte differentiation. In 3T3-L1
7 cells, these two processes are regulated by a strict transcriptional activity in which PPAR γ
8 is the master regulator [19]. During adipocyte differentiation, three different time windows
9 can be distinguished, each one characterized by the upregulation/activation of a different
10 set of transcription factors: an early phase of induction, characterized by the upregulation of
11 *Cebp* (CCAAT/enhancer binding protein) β and δ and the activation of *Cebp* β and *Rxrs*; a
12 middle phase, with RXR α and PPAR γ 2 as obligate heterodimers; a late phase, where
13 adipocyte specific genes such as *Fabp4/Ap2* (Fatty Acid Binding Protein 4/Adipocyte Protein
14 2), *Lpl* (Lipoprotein Lipase), *AdipoQ* (adiponectin) and leptin are upregulated [20–22].

15 Several studies have shown how environmental chemicals can perturb this intracellular
16 cascade by targeting transcription factors and consequently enhance or decrease
17 adipogenesis [5,6,22–24]. For example, certain EDCs may target PPAR γ by binding to it
18 directly to activate downstream cascades leading to enhanced lipid accumulation or by
19 increasing PPAR γ expression to favour its activation [24].

20 In the present work we used 3T3-L1 preadipocytes to investigate the possible adipogenic
21 effects of plasticizers considered safe SVHC substitutes and used in FCMs manufacturing.
22 First, we evaluated possible modifications in lipid accumulation following exposure to scalar
23 concentrations of the plasticizers DiNP, DiDP, DEGDB and TMCP. Since adipogenesis

1 occurs in 3T3-L1 with a defined timeline of transcription factors and receptors activity, we
2 also evaluated the possible different effects of plasticizer exposure alternatively during 3T3-
3 L1 early or mid-late differentiation. We then verified, by *in silico* molecular docking analysis
4 and reporter gene assays, the ability of these molecules to bind and activate the major
5 transcription factor involved in adipogenesis, namely PPAR γ . To better understand the
6 intracellular mechanisms underlying the changes in the adipogenic process, we investigated
7 the regulation of the expression of genes belonging to the early, mid and late phase of
8 adipocyte differentiation.

9

10

11 **2. Material and Methods**

12 **2.1. Chemicals/Reagents**

13 All the reagents for cell culture (including medium supplements), Oil Red O (CAS Number
14 1320-06-5), Rosiglitazone (BRL49653; CAS Number 122320-73-4, purity $\geq 98\%$), DiNP (di-
15 iso-nonyl-phtalate; CAS Number 28553-12-0, purity $\geq 99\%$), DiDP (di-iso-decyl-phtalate;
16 CAS Number 26761-40-0, purity $\geq 99\%$), DEGDB (diethylene glycol dibenzoate; CAS
17 Number 120-55-8, purity 90%), TMCP (tri-m-cresyl phosphate; CAS Number 563-04-2) and
18 BPA (Bisphenol A; CAS Number 80-05-7, purity $\geq 99\%$) were obtained from Sigma Aldrich
19 (USA).

20

21 **2.2. 3T3-L1 culture and adipocyte differentiation experiments**

22 3T3-L1 preadipocytes (ATCC[®] CL-173[™]; ATCC, USA) were cultured in Dulbecco's modified

1 Eagle's medium high-glucose (DMEM) supplemented with 10% calf serum, 2 mM L-
2 glutamine, 50 IU/mL penicillin, and 50 µg/mL streptomycin. 2×10^4 cells/well were seeded in
3 24-well plates. Two days after reaching confluence (day 0), cells were exposed to the
4 differentiation medium (MDI; DMEM containing 10% fetal bovine serum, 1 µg/mL insulin, 1
5 µM dexamethasone, 0.5 mM isobutylmethylxanthine). Two days later (day 2), MDI medium
6 was replaced with maintenance medium (MM; DMEM 10% FBS, 1 µg/mL insulin). Fresh
7 medium was provided every two days. Experiments were ended after 10 days from the
8 beginning of the differentiation (day 10).

9 Cells were exposed to the following plasticizers: DiNP, DiDP, DEGDB, TMCP or BPA at
10 concentrations ranging from 0.01 to 25 µM, that were excluded to be toxic by visual analysis.
11 100 nM Rosiglitazone was used as a positive control. All the chemicals were dissolved in
12 100% DMSO as vehicle, and cells were exposed to a final concentration of 0.1% DMSO.
13 Cells were treated with chemicals alternatively from day 0 to day 10 (whole differentiation
14 period treatment), from day 0 to day 2 (early phase treatment), or from day 2 to day 10
15 (middle-late phase treatment). Control cells were kept in MDI plus 0.1% DMSO from day 0
16 to day 2 and in MM plus 0.1% DMSO from day 2 to day 10.

17 Three independent replicates were set in each experiment; experiments were repeated
18 three times at different passage numbers (p8-p11).

19

20 **2.3. Quantification of adipocyte lipid accumulation**

21 Lipid accumulation in 3T3-L1 adipocytes was determined by quantitative Oil Red O (ORO)
22 staining at day 10. Oil Red O was dissolved in isopropanol overnight at a concentration of
23 0.35%, followed by 0.2 µm filtration, dilution in water to a final concentration of 0.2%, and

1 refiltration. Adipocytes were washed twice with PBS, then they were fixed in 10%
2 paraformaldehyde for 10 min at room temperature. Cells were washed with ddH₂O, allowed
3 to dry, and stained with ORO solution for 20 min. Following several washes with ddH₂O,
4 plates were dried at room temperature; ORO was then eluted in 100% isopropanol, and
5 absorbance at 500 nm was measured using a microplate reader (BioRad, USA). The mean
6 of 8 absorbance readings (technical replicates) was calculated for each sample; three
7 independent plate replicates were set in each experiment and experiments were repeated
8 three times. Variations in lipid accumulation were expressed as fold changes of the
9 absorbance of treated cells relative to the absorbance of control cells; controls were
10 assigned a value of 1.

11 Results are expressed as the mean of the values obtained in the three independent
12 experiments \pm standard error of the mean (SEM).

13

14 **2.4. Molecular docking studies**

15 Molecular docking analysis were performed using Autodock Vina 1.1.2 [25] on an Intel Core
16 i7/Mac OS X 10.9 – based platform, setting a docking zone of 24, 26, and 28 points (in the
17 x, y, and z directions) and of 26, 25, and 27 points with a grid spacing of 1 Å over the human
18 PPAR γ and RXR α binding site, respectively.

19 The crystallographic structures of PPAR γ and RXR α receptors were obtained from the
20 Protein Data Bank [26]: PPAR γ 1I7I.pdb [27], RXR α 3DZY.pdb [28]. The molecular
21 structures of ligands were obtained from the PubChem database [29] and minimized (with
22 a universal force field, UFF, and a conjugate gradient algorithm until a ΔE lower than
23 0.001kJ/mol) using the Avogadro software (Version 1.1.0;

1 <http://avogadro.openmolecules.net/>) [30].

2 The affinity constants, expressed as equilibrium dissociation constants (K_d), were
3 determined analysing the 10 best complexes, obtained for each ligand from Autodock Vina,
4 with the NNScore algorithm, version 2.0 [31].

5 All models and images were rendered using UCSF Chimera software, version 1.11 [32],
6 whereas 2D ligand interaction diagrams were obtained using Maestro software, version 10.6
7 (Schrödinger, LLC, USA).

8

9 **2.5. Transfection and reporter gene assays**

10 HepG2 human hepatoblastoma cell line (ATCC® HB-8065™; ATCC, USA) was used for
11 gene reporter assays; cells were plated on a 24 well plate and then transfected with the
12 following constructs [33,34]: (1) 1.5 µg DR1-Luc (containing a direct repeat 1 upstream of
13 luciferase gene), (2) 100 ng pCMV-βgal (pCMV-β-galactosidase normalization plasmid), and
14 (3) 400 ng pcDNA3-PPARγ (an expression vector for human PPARγ) using Lipofectin
15 (Invitrogen). As described previously [35], cells were treated with the indicated ligands 24
16 hrs post transfection and assayed for luciferase activity 24 hrs post-treatment. Luciferase
17 activity was normalized to β-galactosidase activity to control for transfection efficiency.

18

19 **2.6. Gene expression analysis**

20 Cells were exposed from day 0 to 25 µM DiDP, DEGDB, TMCP or 100 nM Rosiglitazone;
21 control cells were treated with 0.1% DMSO. Three independent replicates were set in each
22 experiment; experiments were repeated three times. Total RNA was isolated from control
23 and treated 3T3-L1 cells at day 2, day 4 and day 8. Briefly, cells were washed with PBS and

1 Tri-Reagent (Sigma, USA) was used for RNA extraction following manufacturer guidelines.
2 qReal-Time PCR was performed using Superscript III Platinum One-step qRT-PCR system
3 (Invitrogen, USA) and the thermal cycler Rotor Gene Q (Qiagen, Germany). Intron-spanning
4 primers for representative genes were designed with Primer-BLAST software (NCBI, USA)
5 and are listed in Table 1. Each sample was analysed in three technical replicates containing
6 50 ng of total RNA. The relative quantification of gene expression was done using a standard
7 curve that was built by pooling all the RNA samples and making serial dilutions (range: 200-
8 6.25 ng of total RNA). The amplicon concentrations were expressed in arbitrary units and
9 were normalized for the expression of *β-actin*, a commonly used housekeeping gene, proved
10 to be a suitable reference gene for qRT-PCR expression studies in 3T3-L1 cells [36]. For
11 each gene, the mRNA expression of the samples was reported as fold changes relative to
12 the expression of control cells; controls were assigned a value of 1.

13

14 **2.7. Statistical analysis**

15 Statistical analysis was performed with SPSS software (version 24; IBM, USA). All data were
16 analysed with one-way ANOVA plus Tukey or Bonferroni post-hoc test ($p < 0.05$). Data were
17 expressed as fold changes versus control \pm standard error of the mean (SEM) or \pm standard
18 deviation (SD); controls were assigned a value of 1.

19

20

21 **3. Results**

22 **3.1. The plasticizers DiNP, DiDP, DEGDB and TMCP enhance lipid accumulation in** 23 **3T3-L1 cells**

1 We evaluated the effect of four plasticizers belonging to different chemical categories (the
2 phthalates DiNP and DiDP, the benzoate ester DEGDB and the organophosphate TMCP)
3 on adipocyte differentiation by assessing lipid accumulation using Oil Red O (ORO) staining.
4 The plasticizer BPA (Bisphenol A), whose well-documented pro-adipogenic effects have
5 been ascribed to multiple pathways [37], was included as a reference compound; another
6 reference molecule included in the study was Rosiglitazone (BRL49653), because of its well-
7 defined agonist activity toward PPAR γ [24]. 3T3-L1 preadipocytes were induced to start
8 adipogenic differentiation and were treated throughout differentiation with vehicle only (0.1%
9 DMSO) or with scalar concentrations (0.01-25 μ M) of each plasticizer, while Rosiglitazone
10 was used at a concentration of 100 nM, selected according to published data [6,24,38]. At
11 the end of the experiment (day 10), lipid accumulation was measured by ORO lipid staining
12 and quantification (Fig. 1). As expected, 100 nM Rosiglitazone-exposed cells displayed a
13 strong enhancement (about 7 folds) in lipid accumulation in respect to untreated cells
14 (cultured in MDI-MM medium containing 0.1% DMSO). BPA exerted a clear dose-dependent
15 enhancement of lipid accumulation, the highest concentration (25 μ M) being markedly more
16 effective in inducing lipogenesis than lower concentrations (5 folds for 25 μ M versus 1.2-
17 1.8 folds for 0.01-10 μ M). Interestingly, also DiNP, DiDP, DEGDB, and TMCP led to a
18 significant increase in lipid accumulation at all tested concentrations. Although lower than
19 the maximal effect reached by the highest doses of BPA, the increase induced by
20 plasticizers was about 20-50% compared to control cells, with TMCP being the most
21 effective plasticizer at all concentrations.

22

23 **3.2. Plasticizers are more effective in enhancing lipid accumulation when**

1 **administered during mid-late differentiation**

2 Since lipidogenesis occurs in 3T3-L1 cells with a defined timeline of transcription factors and
3 receptors activity, we tried to identify windows of susceptibility to plasticizer exposure. For
4 this purpose, plasticizers were added at the lowest concentration tested (0.01 μM)
5 alternatively during the early (day 0-2) or the mid-late differentiation (day 2-10) and lipid
6 accumulation was measured by ORO staining at day 10. An increase in lipid accumulation
7 was observed both when 3T3-L1 cells were treated with plasticizers during the early or the
8 mid-late differentiation (Fig. 2). However, the highest effect on lipidogenesis was reached
9 when plasticizer administration was performed during the mid-late differentiation, except for
10 BPA, for which no statistically significant differences were seen between the two phases.
11 Notably, when administered at 0.01 μM during the mid-late differentiation, BPA resulted the
12 least effective molecule in inducing lipidogenesis, while TMCP was the most effective one.
13 As a matter of fact, exposure to TMCP at days 2-10 was 37% more effective than exposure
14 at days 0-2 (2.33 versus 1.70 folds relative to control), indicating that the mid-late
15 differentiation is considerably more sensitive to TMCP.

16

17 **3.3. Computational analysis predicts specific interactions of the plasticizers with** 18 **PPAR γ and RXR α**

19 Metabolic disruptors are known to control lipidogenesis and adipocyte differentiation
20 interacting with transcription regulators of gene networks, the main of which belong to the
21 PPAR and RXR receptor families. Since our above-reported results show that plasticizers
22 can enhance *in vitro* 3T3-L1 preadipocytes lipid accumulation, we evaluated if these
23 plasticizers could potentially act via an interaction with the nuclear receptors PPAR γ and

1 RXR α . *In silico* molecular docking analysis, that consider the affinity and the geometry of
2 binding, actually showed the capability of DiNP, DiDP and TMCP to specifically bind the
3 PPAR γ receptor with affinities ranging in the submicromolar order; as expected, BPA
4 showed a lower affinity for PPAR γ , in respect to the other plasticizers. All the ligands
5 analysed showed a higher binding affinity with RXR α , although their predicted equilibrium
6 dissociation constants for PPAR γ are in the same order of magnitude (Table 2). To validate
7 the molecular docking procedure, we added Rosiglitazone to the ligands set and found a
8 predicted equilibrium dissociation constant for PPAR γ highly comparable to the K_d value
9 already published [39]. Moreover, the molecular docking model of the best predicted
10 Rosiglitazone/PPAR γ complex and the crystallographic structure of this complex
11 (4EMA.pdb) [40] are extremely comparable (data not shown), on the basis of both
12 orientation and average distance of each atom of the ligand (RMSD value = 1.05Å). Among
13 the molecules analysed, TMCP resulted to be the best ligand for PPAR γ and RXR α
14 receptors, showing two equilibrium dissociation constants comparable to those of
15 Rosiglitazone. Molecular docking analysis between TMCP and the two receptors ligand
16 binding domains showed that TMCP is exclusively stabilized by non-polar interactions and,
17 in particular, it could establish a pi-pi stacking interaction with Arg²⁸⁸ of PPAR γ and with
18 Phe³¹³ of RXR α (Fig. 3).

19

20 **3.4. Plasticizers can transactivate PPAR γ**

21 We confirmed the ability of the plasticizers TMCP, DiDP, DiNP and DEGDB to bind and
22 activate PPAR γ by examining their capacity to induce PPAR γ -driven reporter expression
23 following transient transfection of HepG2 cells with pcDNA3-PPAR γ . In this assay, all

1 plasticizers significantly induced PPAR γ -driven reporter activity at a concentration of 25 μ M,
2 with DiNP and TMPC being already active at 10 μ M (Fig. 4). The maximal activity was
3 reached by 25 μ M TMCP, that lead to an induction of 2.5 folds, corresponding to about half
4 of the induction obtained by 10 μ M Rosiglitazone.

5

6 **3.5. Plasticizers modulate the expression of adipogenic marker genes**

7 Differentiation of 3T3-L1 preadipocytes, similarly to what occurs *in vivo*, involves a
8 transcriptional cascade initially activated by an adipogenic cocktail (MDI, see Methods)
9 inducing, among others, the transcription factor *Cebp β* (*early phase* of differentiation).
10 CEPB β is a direct activator of *Ppar γ* transcription (*mid phase*), and 7-12)PPAR γ in turn
11 binds as an obligate heterodimer with the nuclear receptor RXR to numerous promoter sites
12 of adipocyte specific genes (*late phase*), including *Fabp4/Ap2* and *Lpl*. In the effort to further
13 elucidate the mechanisms of plasticizer action on preadipocyte differentiation, we analysed
14 by qReal-Time PCR the expression of *Cebp β* , *Rxra*, *Ppar γ 2*, *Fabp4/Ap2* and *Lpl* transcripts
15 at day 2, day 4 or day 8 post-induction. 3T3-L1 cells were exposed to 100 nM Rosiglitazone
16 or to 25 μ M TMCP, DiDP and DEGDB, a concentration able to induce the highest lipid
17 accumulation in the absence of cytotoxic effects.

18 At day 2 (Fig. 5, upper panel), corresponding to the early phase of adipogenic differentiation,
19 all the tested molecules were able to enhance the expression of the *Cebp β* transcript,
20 suggesting that Rosiglitazone, TMCP, DiDP and DEGDB can influence the first steps of
21 differentiation by regulating the expression of this early gene. On the other hand, at day 4
22 (Fig. 5, lower panel) only DEGDB still enhanced *Cebp β* mRNA expression. The expression
23 of *Rxra* was selectively modified only by Rosiglitazone administration both during the early

1 (day 2; Fig. 5, upper panel) and mid phase of differentiation (day 4; Fig. 5, lower panel). The
2 expression of *Ppar γ 2*, the adipogenesis master gene, was markedly increased at day 2 (Fig.
3 5, upper panel) by DiDP and DEGDB, while Rosiglitazone and TMCP did not exert any
4 effect. At day 4 (Fig. 5, lower panel) all the analysed plasticizers were able to increase
5 *Ppar γ 2* mRNA expression. Overall, the plasticizer-induced regulation of *Ppar γ 2* expression
6 on day 2 and 4 was quite similar to the one exerted by Rosiglitazone.

7 As expected, in the late phase of differentiation (day 8) (Fig. 6), the levels of the *Fabp4*
8 transcript were highly increased by Rosiglitazone. The plasticizers TMCP and DEGDB had
9 also a positive effect (4 and 3.5 folds respectively compared to untreated cells) on the
10 expression of this transcript. *Lpl*, another adipogenesis marker gene belonging to the late
11 phase, was modulated by Rosiglitazone and TMCP at comparable levels (about 8 and 6
12 folds respectively). Conversely, the expression of both *Fabp4* and *Lpl* was not modified by
13 exposure to the phthalate DiDP.

14

15 **4. Discussion**

16 Plasticizers and their metabolites are a frequent finding in human biomonitoring data of
17 industrialized countries [41–46]. Published datasets in national surveys referring to the last
18 decade track the coexistence of both dismissed compounds, still present in relevant
19 amounts, and new plasticizers and their metabolites [45,46] that are slowly substituting the
20 former ones. Some of the new plasticizers could represent an emerging class of
21 contaminants, therefore evaluation of their potential biological effects is needed [47].

22 The results of the present study suggest that plasticizers considered safer alternatives to
23 SVHC may actually affect metabolic processes, such as adipogenesis. We demonstrate

1 that low nanomolar concentrations of four plasticizers currently used in FCMs manufacturing
2 (namely DiNP, DiDP, DEGDB and TMCP) enhance the ability of 3T3-L1 cells to differentiate
3 into mature adipocytes, as shown by a 1.2-2.3 fold increase in lipid accumulation, depending
4 on the chemical and time window of exposure. Computational analysis shows the capability
5 of these compounds to bind to PPAR γ and RXR α , two nuclear receptors specifically involved
6 in the adipogenic transcriptional cascade. Each plasticizer was able to transactivate PPAR γ
7 and to modulate the expression of adipogenic marker genes to various extents. By
8 analysing the regulation of *Ppar γ 2* gene expression exerted by test plasticizers we found a
9 certain similarity to the one exerted by Rosiglitazone, a PPAR γ agonist, suggesting some
10 degree of overlapping in the cellular mechanisms involved.

11 Besides Rosiglitazone, we included also BPA as a useful reference compound in all our
12 experiments, since considerable amount of knowledge has been accumulated from *in vitro*
13 and *in vivo* studies on this plasticizer. While some controversy exists in epidemiological data
14 associating BPA exposure and development of obesity and/or metabolic syndrome in human
15 populations [2,3,48] , several animal studies demonstrate that exposure to BPA can affect
16 adipogenesis [2,3,37]. In addition, a number of studies on 3T3-L1 cells have shown that
17 BPA administered during adipocyte differentiation increases lipid accumulation, generally
18 from 2 to 5 folds compared to control, depending on protocol and dosage [24,49–53]. Our
19 results regarding BPA are in line with most previous literature data.

20 Phthalate pro-obesogenic effects in the human population are still under investigation. Some
21 studies relate the presence of phthalates in blood samples and urine with an increased risk
22 of obesity and metabolic syndrome [54–57], however in a context of a larger dataset these
23 links seem to have some uncertainty [58]. Differently from epidemiological data, there is

1 extensive knowledge that phthalates exposure, particularly DEHP and its metabolite mono
2 (2-ethylhexyl) phthalate (MEHP), have negative outcomes on glucose and lipid homeostasis
3 in cellular and animal models [47,58,59]. However, there are scarce or no studies on the
4 emerging phthalate substitutes DiNP and DiDP. We show that low nanomolar
5 concentrations of DiNP and DiDP are able to enhance lipid accumulation from 20% to 80%,
6 depending on the time-frame of administration. While this is the first report showing that
7 DiDP can increase lipodogenesis in 3T3-L1 adipocytes, a previous study reported small
8 statistically significant effects of DiNP on lipid accumulation [24]. Human biomonitoring
9 studies employing metabolites of DiNP and DiDP as biomarkers of exposure, reported
10 median values of 5.10 $\mu\text{g/L}$ (16 nM) for MCiOP (mono carboxy-isoctyl phthalate, a DiNP
11 metabolite) and 2.7 $\mu\text{g/L}$ (7.9 nM) for MCiNP (mono-carboxy-isononyl phthalate, a DiDP
12 metabolite) in the urine of the United States general population (>6 years, 2005-2006 survey,
13 Calafat et al. [60]). These levels found in urine are comparable to the 10 nM concentration
14 used in our experiments. Additional studies compared the levels of phthalate metabolites in
15 urine among mother-child pairs [61,62] showing that children's DiNP and DiDP metabolite
16 excretion was higher than that of the mothers, indicating a possible higher children
17 exposure. In addition, multiple studies [62,63] found a significant temporal decline over the
18 last 15-20 years in urinary levels of metabolites of strictly regulated phthalates (such as
19 DHEP), paralleled by a marked increase in urinary metabolite concentrations of DiNP and
20 DiDP. Given the existing biomonitoring data and the results of our study, further research
21 on the adverse health effects of DiNP and DiDP, including obesity and metabolic
22 dysfunctions, is warranted.

1 DEGDB is defined by many as a “green plasticizer”. There are currently no published human
2 biomonitoring studies on this chemical and there is only some preliminary evidence of the
3 potential impact of DEGDB on tissue-specific regulation of genes involved in lipid
4 metabolism and energy balance *in vivo* [18]. In our *in vitro* experiments, DEGDB actually
5 showed a lipidogenic effect similar to that of the phthalates DiNP and DiDP. To the best of
6 our knowledge, this is the first report linking this compound to *in vitro*-induced adipogenesis.
7 TMCP belongs to the class of organophosphates, a group of compounds that range from
8 slightly to highly toxic depending on chemical structure, dose and route of exposure [64,65].
9 Epidemiological data on organophosphates indicate that a prenatal exposure may lead to
10 adverse effects on glucose metabolism at birth [66], but little is known about the outcomes
11 of long-term exposure and adult datasets often report controversial results [67]. Concerning
12 animal studies, recent data show that a chronic or subchronic dietary or perinatal exposure
13 to organophosphates alters metabolic functions causing an obese-like phenotype and a
14 diabetic profile in mice [68–70] and rat models [71,72]. Although there are no data available
15 on TMCP effects in 3T3-L1 adipocytes, recent results on the organophosphate triphenyl
16 phosphate (TPhP) indicate that this molecule is able to increase 3T3-L1 preadipocyte
17 proliferation and subsequent adipocyte differentiation, as well as glucose uptake and
18 lipolysis [73]. TMCP was the most effective compound in enhancing lipid accumulation
19 among the plasticizers we tested. This evidence, together with the fact that this
20 organophosphate showed the highest computational binding affinity and capability to
21 transactivate PPAR γ , potentially make TMCP the most obesogenic of the four plasticizers
22 that we tested. Clearly, *in vivo* studies are needed to confirm the plasticizer obesogenic
23 potentials defined *in vitro* in the present study.

1 To elucidate the mechanism by which DiNP, DiDP, DEGDB and TMCP enhance lipid
2 accumulation in 3T3-L1 cells, we evaluated by computational analysis their interaction with
3 PPAR γ and RXR α . *In silico* binding affinity of plasticizers for PPAR γ and RXR α receptors
4 was highly indicative of *in vivo* interactions, particularly for TMCP whose K_d values were
5 similar to those calculated for Rosiglitazone. BPA showed a higher K_d value for PPAR γ ,
6 suggesting an action mainly through other nuclear receptors. The predicted interaction of
7 the plasticizers with PPAR γ was confirmed by transient transfection studies and is in line
8 with the significant increase found in the expression of the PPAR γ target gene *Fabp4* after
9 exposure to plasticizers during differentiation. These data are in agreement with previous
10 studies suggesting that phthalates and TMCP or TPhP can modulate the regulatory
11 mechanism of lipid metabolism pathways through PPARs and RXRs [15–17,74]. Another
12 significant result from our data is that both DiNP and TMCP show high binding affinity for
13 RXR α . This finding, which is a common trend for other potential obesogens [16,59,75],
14 suggests that binding of these plasticizers to RXR α may independently increase PPAR γ
15 transcriptional activity. This possibility, although needing experimental confirmation, is in line
16 with the “permissive” features of the PPAR γ /RXR α heterodimer [76], meaning that also RXR
17 ligands can activate it, amplifying the effects on downstream genes. Taken together, *in silico*
18 predictions and transactivation experiments suggest that the mechanisms through which
19 DiNP, DiDP, DEGDB and TMCP increase lipid accumulation involve the direct activation of
20 the PPAR γ /RXR α complex.

21 It is expected that the effect of plasticizers on lipid accumulation in 3T3-L1 cells is linked to
22 and, perhaps, promoted by modifications in the pro-adipogenic transcription factor cascade.
23 In order to test this hypothesis, we analysed a set of transcripts that play a key role in the

1 adipogenic process: *Cebpb*, *Rxra*, *Ppar* γ 2, *Fabp*4 and *Lpl*. In the early phase of adipocyte
2 differentiation, all plasticizers were able to increase *Cebpb* expression, a transcription factor
3 playing a crucial role in the induction of 3T3-L1 differentiation and required for the binding
4 to genomic adipogenic hotspots of other adipogenic transcription factors [19,77]. CEPB β is
5 a direct activator of *Ppar* γ transcription, therefore an increase in *Cebpb* expression is
6 expected to reverberate on *Ppar* γ expression [77,78]. In line with this, we found that all
7 plasticizers induced also a significant enhancement in *Ppar* γ 2 transcript levels in the middle-
8 late phase of differentiation. Exposure to plasticizers only in the early phase (days 0-2),
9 corresponding to the enhancement of *Cebpb* expression, was enough to induce a significant
10 increase in lipid accumulation measured at the end of differentiation (day 10). This result
11 suggests that any molecule able to modify the expression and therefore the activity of
12 CEPB β can have profound consequences on adipocyte differentiation. We can hypothesize
13 that the plasticizers could increase the expression of *Cebpb* through the activation of the
14 cAMP response element-binding protein (CREB) and the glucocorticoid receptor (GR),
15 however recently additional transcription factors have been found to regulate *Cebpb*
16 transcription as a consequence of different adipogenic stimuli [21,79]. We observed no
17 changes in the expression of the *Rxra* gene, except for a moderate increase exerted by
18 Rosiglitazone. Absence of regulation of *Rxra* expression was somehow expected, since
19 previous studies showed that the human *Rxra* gene displays features of a housekeeping
20 gene [80]. Additional studies report that RXR α activity is modulated by extensive
21 posttranslational modifications and proteasomal degradation [76], suggesting that RXR α is
22 mainly regulated at the protein level. Like Rosiglitazone, in the late phase of 3T3-L1 cell
23 differentiation TMCP was able to modulate the expression of the adipogenesis marker genes

1 *Lpl* and *Fabp4*. Similarly to our result, a recent study [73] reported that 25 μ M of the
2 organophosphate TPhP is able to increase 3T3-L1 differentiation by upregulating the
3 expression of *Cebp β* , *Ppar γ* and *Lpl* during early and mid-late differentiation, respectively.
4 Activation of PPAR γ and increased differentiation of 3T3-L1 cells into adipocytes by
5 phthalates (i.e. MHEP and DHEP) has been previously reported [81–83]. Nonetheless, this
6 effects not always correlated with a modulation in late genes involved in lipidogenesis [84].
7 Similarly, we also observed that DiDP and DEGDB were both unable to modify *Lpl* transcript
8 levels. It is possible that other late genes, not considered in our study, are regulated by these
9 plasticizers. In addition, both DiDP and DEGDB were able to activate PPAR γ in transient
10 transfection studies only at the highest concentration (25 μ M), therefore showing a lower
11 capability to interact with PPAR γ compared to TMCP.

12 The plasticizer-mediated enhancement of lipid accumulation in 3T3-L1 cells was present
13 when exposing cells both in the early or in the mid-late phase of adipogenic differentiation.
14 However, plasticizers were more effective when added during mid-late differentiation. We
15 can postulate that when plasticizers are delivered in the early phase, they positively
16 modulate *Cebp β* transcription, leading to enhanced PPAR γ expression and receptor
17 availability in the subsequent steps of the lipidogenic process. On the other hand, if
18 plasticizers are added in the mid-late phase, when PPAR γ is highly expressed, they can
19 interact directly with this receptor. As a result, lipid accumulation increases even further
20 compared to the early phase treatment. We observed that cells exposed to BPA did not
21 behave differently in the two phases, possibly because of the low BPA binding affinity for
22 PPAR γ . Multiple intracellular pathways involved in the induction of adipogenesis by BPA
23 have been described [85,86], mostly characterized by PPAR γ /RXR α independent

1 mechanisms [37,86,87]. The wide range of 3T3-L1 cells responses observed after plasticizer
2 treatments probably reflects not only the multiple pathways engaged by each type of
3 chemical compound [88], but also the high complexity of the cell processes leading to the
4 differentiation into mature adipocytes [20,89].

5

6 **5. Conclusions**

7 Our study demonstrates that the plasticizers DiNP, DiDP, DEGDB and TMCP, used as safer
8 alternatives to SVHC chemicals, are able to interfere with the adipogenic process in 3T3-L1
9 cells at low nanomolar concentrations. Our results suggest that the observed increase in
10 lipid accumulation is at least partly mediated by direct binding to the transcription factors
11 PPAR γ and RXR α and through regulation of several genes involved in the adipogenic
12 transcriptional cascade. The effect of single chemicals on lipid accumulation was moderate,
13 however it should be considered that multiple plasticizers often occur in the same FCM,
14 therefore the global effect of singularly active plasticizers could be significantly higher in
15 mixtures. For this reason, future studies should address the metabolic effects of mixtures
16 containing TMCP, DiNP, DiDP and DEGDB. Our findings also suggest that these four
17 plasticizers may not be harmless substitute of currently restricted compounds. Given the
18 growing exposure of humans to these plasticizers, further *in vivo* investigation of their effects
19 is warranted.

20

21 **Acknowledgments**

22 The Authors wish to thank Dr. Fabio Penna and Dr. Claudio Dati for 3T3-L1 cell culture set
23 up and Dr. Stefania Rapelli for the initial assistance with qRealTime PCR.

1 This work was supported by MIUR-PRIN (Ministero dell'Istruzione, dell'Università e della
2 Ricerca, Progetti di Ricerca di Interesse Nazionale) prot. 2010W87LBJ_005 to PB, and prot.
3 2010W87LBJ_002 to GM; and Fondazione CRT (Cassa di Risparmio di Torino) RF
4 2014.0814 to PB.

1 **References**

- 2 [1] Eurostat, Overweight and obesity - BMI statistics, (2014).
3 [http://ec.europa.eu/eurostat/statistics-](http://ec.europa.eu/eurostat/statistics-explained/index.php/Overweight_and_obesity_-_BMI_statistics)
4 [explained/index.php/Overweight_and_obesity_-_BMI_statistics.](http://ec.europa.eu/eurostat/statistics-explained/index.php/Overweight_and_obesity_-_BMI_statistics)
- 5 [2] J.J. Heindel, B. Blumberg, M. Cave, R. Machtinger, A. Mantovani, M.A. Mendez, A.
6 Nadal, P. Palanza, G. Panzica, R. Sargis, L.N. Vandenberg, F. vom Saal,
7 Metabolism disrupting chemicals and metabolic disorders, *Reprod. Toxicol.* 68
8 (2017) 3–33. doi:10.1016/j.reprotox.2016.10.001.
- 9 [3] J.J. Heindel, F.S. Vom Saal, B. Blumberg, P. Bovolini, G. Calamandrei, G. Ceresini,
10 B.A. Cohn, E. Fabbri, L. Gioiosa, C. Kassotis, J. Legler, M. La Merrill, L. Rizzir, R.
11 Machtinger, A. Mantovani, M.A. Mendez, L. Montanini, L. Molteni, S.C. Nagel, S.
12 Parmigiani, G. Panzica, S. Paterlini, V. Pomatto, J. Ruzzin, G. Sartor, T.T. Schug,
13 M.E. Street, A. Suvorov, R. Volpi, R.T. Zoeller, P. Palanza, Parma consensus
14 statement on metabolic disruptors, *Environ. Heal. A Glob. Access Sci. Source.* 14
15 (2015). doi:10.1186/s12940-015-0042-7.
- 16 [4] F. Grun, B. Blumberg, Environmental obesogens: Organotins and endocrine
17 disruption via nuclear receptor signaling, *Endocrinology.* 147 (2006) 50–55.
18 doi:10.1210/en.2005-1129.
- 19 [5] F. Grün, B. Blumberg, Endocrine disruptors as obesogens, *Mol. Cell. Endocrinol.*
20 304 (2009) 19–29. doi:10.1016/j.mce.2009.02.018.
- 21 [6] A.S. Janesick, B. Blumberg, Obesogens: An emerging threat to public health, *Am. J.*
22 *Obstet. Gynecol.* 214 (2016) 559–565. doi:10.1016/j.ajog.2016.01.182.
- 23 [7] E. Diamanti-Kandarakis, J.-P. Bourguignon, L.C. Giudice, R. Hauser, G.S. Prins,

- 1 A.M. Soto, R.T. Zoeller, A.C. Gore, Endocrine-Disrupting Chemicals: An Endocrine
2 Society Scientific Statement, *Endocr. Rev.* 30 (2009) 293–342. doi:10.1210/er.2009-
3 0002.
- 4 [8] K. Cwiek-Ludwicka, J.K. Ludwicki, Endocrine disruptors in food contact materials; is
5 there a health threat?, *Rocz. Państwowego Zakładu Hig.* 65 (2014) 169–177.
- 6 [9] Calvin E, Plasticizer market update. BASF corporation. Presentation at the 22nd
7 Annual Vinyl Compounding Conference, Burlington, July 10–13 2011, in: 2011.
8 <http://www.cpsc.gov/PageFiles/126090/spi.pdf>.
- 9 [10] ECHA (European Chemicals Agency), Evaluation of new scientific evidence
10 concerning DINP and DIDP in relation to entry 52 of annex XVII to reach regulation
11 (EC) No 1907/2006 (Final Review Report), 2013.
12 <http://echa.europa.eu/documents/10162/31b4067e-de40-4044-93e8-9c9ff1960715>.
- 13 [11] A. Kermanshahi pour, D.G. Cooper, O.A. Mamer, M. Maric, J.A. Nicell, Mechanisms
14 of biodegradation of dibenzoate plasticizers, *Chemosphere.* 77 (2009) 258–263.
15 doi:10.1016/j.chemosphere.2009.06.048.
- 16 [12] S. Posner, Guidance on alternative flame retardants to the use of commercial
17 pentabromodiphenylether (c-PentaBDE). Norwegian Pollution control Authority
18 (SFT) on behalf of UNEP, United Nations Oslo, 2009.
- 19 [13] T.M. Attina, L. Trasande, Association of exposure to Di-2-ethylhexylphthalate
20 replacements with increased insulin resistance in adolescents from NHANES 2009-
21 2012, *J. Clin. Endocrinol. Metab.* 100 (2015) 2640–2650. doi:10.1210/jc.2015-1686.
- 22 [14] Å. Bergman, J. Heindel, S. Jobling, K. Kidd, R. Zoeller, S. Jobling, State of the
23 science of endocrine disrupting chemicals 2012: an assessment of the state of the

- 1 science of endocrine disruptors prepared by a group of experts for the United
2 Nations Environment Program and World Health Organization. WHO, 2013.
- 3 [15] M.K. Sarath Josh, S. Pradeep, K.S. Vijayalekshmi Amma, S. Balachandran, U.C.
4 Abdul Jaleel, M. Doble, F. Spener, S. Benjamin, Phthalates efficiently bind to human
5 peroxisome proliferator activated receptor and retinoid X receptor α , β , γ subtypes:
6 An in silico approach, *J. Appl. Toxicol.* 34 (2014) 754–765. doi:10.1002/jat.2902.
- 7 [16] P. Cocci, G. Mosconi, A. Arukwe, M. Mozzicafreddo, M. Angeletti, G. Aretusi, F.A.
8 Palermo, Effects of Diisodecyl Phthalate on PPAR:RXR-Dependent Gene
9 Expression Pathways in Sea Bream Hepatocytes, *Chem. Res. Toxicol.* 28 (2015)
10 935–947. doi:10.1021/tx500529x.
- 11 [17] F.A. Palermo, P. Cocci, M. Mozzicafreddo, A. Arukwe, M. Angeletti, G. Aretusi, G.
12 Mosconi, Tri-m-cresyl phosphate and PPAR/LXR interactions in seabream
13 hepatocytes: Revealed by computational modeling (docking) and transcriptional
14 regulation of signaling pathways, *Toxicol. Res. (Camb)*. 5 (2016) 471–481.
15 doi:10.1039/c5tx00314h.
- 16 [18] P. Cocci, M. Mozzicafreddo, M. Angeletti, G. Mosconi, F.A. Palermo, Differential
17 tissue regulation of peroxisome proliferator-activated receptor α (PPAR α) and
18 cannabinoid receptor 1 (CB1) gene transcription pathways by diethylene glycol
19 dibenzoate (DEGB): preliminary observations in a seabream (*Sparus aurata*) in vivo
20 mode, *Environ. Toxicol. Pharmacol.* 55 (2017) 87–93.
21 doi:10.1016/j.etap.2017.08.015.
- 22 [19] A.G. Cristancho, M.A. Lazar, Forming functional fat: A growing understanding of
23 adipocyte differentiation, *Nat. Rev. Mol. Cell Biol.* 12 (2011) 722–734.

- 1 doi:10.1038/nrm3198.
- 2 [20] R. Siersbæk, R. Nielsen, S. Mandrup, PPAR γ in adipocyte differentiation and
3 metabolism - Novel insights from genome-wide studies, *FEBS Lett.* 584 (2010)
4 3242–3249. doi:10.1016/j.febslet.2010.06.010.
- 5 [21] R. Siersbæk, R. Nielsen, S. Mandrup, Transcriptional networks and chromatin
6 remodeling controlling adipogenesis, *Trends Endocrinol. Metab.* 23 (2012) 56–64.
7 doi:10.1016/j.tem.2011.10.001.
- 8 [22] J.H. Kamstra, E. Hrubá, B. Blumberg, A. Janesick, S. Mandrup, T. Hamers, J.
9 Legler, Transcriptional and Epigenetic Mechanisms Underlying Enhanced in Vitro
10 Adipocyte Differentiation by the Brominated Flame Retardant BDE-47, *Environ. Sci.*
11 *Technol.* 48 (2014) 4110–4119.
- 12 [23] E.W.Y. Tung, A. Boudreau, M.G. Wade, E. Atlas, Induction of Adipocyte
13 Differentiation by Polybrominated Diphenyl Ethers (PBDEs) in 3T3-L1 Cells, *PLoS*
14 *One.* 9 (2014) e94583. doi:10.1371/journal.pone.0094583.
- 15 [24] A. Pereira-Fernandes, H. Demaegdt, K. Vandermeiren, T.L.M. Hectors, P.G. Jorens,
16 R. Blust, C. Vanparys, Evaluation of a Screening System for Obesogenic
17 Compounds: Screening of Endocrine Disrupting Compounds and Evaluation of the
18 PPAR Dependency of the Effect, *PLoS One.* 8 (2013) 1–17.
19 doi:10.1371/journal.pone.0077481.
- 20 [25] O. Trott, A.J. Olson, AutoDock Vina: Improving the Speed and Accuracy of Docking
21 with a New Scoring Function, Efficient Optimization, and Multithreading, *J. Comput.*
22 *Chem.* 31 (2010) 455–61. doi:10.1002/jcc.
- 23 [26] H.M. Berman, J. Westbrook, Z. Feng, G. Gilliland, T.N. Bhat, H. Weissig, I.N.

- 1 Shindyalov, P.E. Bourne, The Protein Data Bank, *Nucl. Acids Res.* 28 (2000) 235–
2 242. doi:10.1093/nar/28.1.235.
- 3 [27] P. Cronet, J.F.W. Petersen, R. Folmer, N. Blomberg, K. Sjöblom, U. Karlsson, E.L.
4 Lindstedt, K. Bamberg, Structure of the PPAR α and γ ligand binding domain in
5 complex with AZ 242; ligand selectivity and agonist activation in the PPAR family,
6 *Structure*. 9 (2001) 699–706. doi:10.1016/S0969-2126(01)00634-7.
- 7 [28] V. Chandra, P. Huang, Y. Hamuro, S. Raghuram, Y. Wang, T.P. Burris, F.
8 Rastinejad, Structure of the intact PPAR- γ -RXR- nuclear receptor complex on
9 DNA., *Nature*. 456 (2008) 350–356. doi:10.1038/nature07413.
- 10 [29] S. Kim, P.A. Thiessen, E.E. Bolton, J. Chen, G. Fu, A. Gindulyte, L. Han, J. He, S.
11 He, B.A. Shoemaker, J. Wang, B. Yu, J. Zhang, S.H. Bryant, PubChem substance
12 and compound databases, *Nucleic Acids Res.* 44 (2016) D1202–D1213.
13 doi:10.1093/nar/gkv951.
- 14 [30] M.D. Hanwell, D.E. Curtis, D.C. Lonie, T. Vandermeersch, E. Zurek, G.R.
15 Hutchison, Avogadro: An advanced semantic chemical editor, visualization, and
16 analysis platform, *J. Cheminform.* 4 (2012). doi:10.1186/1758-2946-4-17.
- 17 [31] J.D. Durrant, J.A. McCammon, NNScore: A neural-network-based scoring function
18 for the characterization of protein-ligand complexes, *J. Chem. Inf. Model.* 50 (2010)
19 1865–1871. doi:10.1021/ci100244v.
- 20 [32] E.F. Pettersen, T.D. Goddard, C.C. Huang, G.S. Couch, D.M. Greenblatt, E.C.
21 Meng, T.E. Ferrin, UCSF Chimera - A visualization system for exploratory research
22 and analysis, *J. Comput. Chem.* 25 (2004) 1605–1612. doi:10.1002/jcc.20084.
- 23 [33] J.M. Hall, D.P. McDonnell, The Molecular Mechanisms Underlying the

- 1 Proinflammatory Actions of Thiazolidinediones in Human Macrophages, *Mol.*
2 *Endocrinol.* 21 (2007) 1756–1768. doi:10.1210/me.2007-0060.
- 3 [34] R. Safi, A. Kovacic, S. Gaillard, Y. Murata, E.R. Simpson, D.P. McDonnell, C.D.
4 Clyne, Coactivation of liver receptor homologue-1 by peroxisome proliferator-
5 activated receptor gamma coactivator-1 alpha on aromatase promoter II and its
6 inhibition by activated retinoid X receptor suggest a novel target for breast-specific
7 antiestrogen therapy, *Cancer Res.* 65 (2005) 11762–11770. doi:10.1158/0008-
8 5472.CAN-05-2792.
- 9 [35] E.R. Nelson, C.D. DuSell, X. Wang, M.K. Howe, G. Evans, R.D. Michalek, M.
10 Umetani, J.C. Rathmell, S. Khosla, D. Gesty-Palmer, D.P. McDonnell, The oxysterol,
11 27-hydroxycholesterol, links cholesterol metabolism to bone homeostasis through its
12 actions on the estrogen and liver X receptors, *Endocrinology.* 152 (2011) 4691–
13 4705. doi:10.1210/en.2011-1298.
- 14 [36] J. Zhang, H. Tang, Y. Zhang, R. Deng, L. Shao, Y. Liu, F. Li, X. Wang, L. Zhou,
15 Identification of suitable reference genes for quantitative RT-PCR during 3T3-L1
16 adipocyte differentiation, *Int. J. Mol. Med.* 33 (2014) 1209–1218.
17 doi:10.3892/ijmm.2014.1695.
- 18 [37] L. Le Corre, P. Besnard, M.C. Chagnon, BPA, an Energy Balance Disruptor, *Crit.*
19 *Rev. Food Sci. Nutr.* 55 (2015) 769–777. doi:10.1080/10408398.2012.678421.
- 20 [38] X. Li, J. Ycaza, B. Blumberg, The environmental obesogen tributyltin chloride acts
21 via peroxisome proliferator activated receptor gamma to induce adipogenesis in
22 murine 3T3-L1 preadipocytes, *J. Steroid Biochem. Mol. Biol.* 127 (2011) 9–15.
23 doi:10.1016/j.jsbmb.2011.03.012.

- 1 [39] J.M. Lehmann, L.B. Moore, T. a Smith-Oliver, W.O. Wilkison, T.M. Willson, S. a
2 Kliewer, An antidiabetic thiazolidinedione is a high affinity ligand for peroxisome
3 proliferator-activated receptor gamma (PPAR gamma)., *J. Biol. Chem.* 270 (1995)
4 12953–12956. doi:10.1074/jbc.270.22.12953.
- 5 [40] M.V. Liberato, A.S. Nascimento, S.D. Ayers, J.Z. Lin, A. Cvorov, R.L. Silveira, L.
6 Martínez, P.C.T. Souza, D. Saidenberg, T. Deng, A.A. Amato, M. Togashi, W.A.
7 Hsueh, K. Phillips, M.S. Palma, F.A.R. Neves, M.S. Skaf, P. Webb, I. Polikarpov,
8 Medium chain fatty acids are selective peroxisome proliferator activated receptor
9 (PPAR) γ activators and Pan-PPAR partial agonists, *PLoS One.* 7 (2012).
10 doi:10.1371/journal.pone.0036297.
- 11 [41] CDC, Fourth National Report on Human Exposure to Environmental Chemicals
12 Centers for Disease Control and Prevention, Department of Health and Human
13 Services, Atlanta, GA, 2009.
14 <https://www.cdc.gov/exposurereport/pdf/fourthreport.pdf>.
- 15 [42] N. Fréry, A. Saoudi, R. Garnier, A. Zeghnoun, G. Falq, Exposition De La Population
16 Française Aux Substances Chimiques De l'Environnement. Saint-Maurice, France:
17 Institut de veille sanitaire, (2011) 151. <http://invs.santepubliquefrance.fr>.
- 18 [43] M. Kolossa-Gehring, K. Becker, A. Conrad, C. Schröter-Kermani, C. Schulz, M.
19 Seiwert, Environmental surveys, specimen bank and health related environmental
20 monitoring in Germany, *Int. J. Hyg. Environ. Health.* 215 (2012) 120–126.
21 doi:10.1016/j.ijheh.2011.10.013.
- 22 [44] J.W. Lee, C.K. Lee, C.S. Moon, I.J. Choi, K.J. Lee, S.M. Yi, B.K. Jang, B. jun Yoon,
23 D.S. Kim, D. Peak, D. Sul, E. Oh, H. Im, H.S. Kang, J. Kim, J.T. Lee, K. Kim, K.L.

- 1 Park, R. Ahn, S.H. Park, S.C. Kim, C.H. Park, J.H. Lee, Korea National Survey for
2 Environmental Pollutants in the Human Body 2008: Heavy metals in the blood or
3 urine of the Korean population, *Int. J. Hyg. Environ. Health.* 215 (2012) 449–457.
4 doi:10.1016/j.ijheh.2012.01.002.
- 5 [45] WHO Regional Office for Europe, Human biomonitoring: facts and figures, (2015)
6 88. [http://www.euro.who.int/__data/assets/pdf_file/0020/276311/Human-](http://www.euro.who.int/__data/assets/pdf_file/0020/276311/Human-biomonitoring-facts-figures-en.pdf)
7 [biomonitoring-facts-figures-en.pdf](http://www.euro.who.int/__data/assets/pdf_file/0020/276311/Human-biomonitoring-facts-figures-en.pdf).
- 8 [46] D.A. Haines, G. Saravanabhavan, K. Werry, C. Khoury, An overview of human
9 biomonitoring of environmental chemicals in the Canadian Health Measures Survey:
10 2007-2019, *Int. J. Hyg. Environ. Health.* (2016) 2007–2019.
11 doi:10.1016/j.ijheh.2016.08.002.
- 12 [47] A.C. Gore, V.A. Chappell, S.E. Fenton, J.A. Flaws, A. Nadal, G.S. Prins, J. Toppari,
13 R.T. Zoeller, Executive Summary to EDC-2: The Endocrine Society's second
14 Scientific Statement on endocrine-disrupting chemicals, *Endocr. Rev.* 36 (2015)
15 593–602. doi:10.1210/er.2015-1093.
- 16 [48] N. Chevalier, P. Fénelichel, Bisphenol A: Targeting metabolic tissues, *Rev. Endocr.*
17 *Metab. Disord.* 16 (2015) 299–309. doi:10.1007/s11154-016-9333-8.
- 18 [49] H. Masuno, T. Kidani, K. Sekiya, K. Sakayama, T. Shiosaka, H. Yamamoto, K.
19 Honda, Bisphenol A in combination with insulin can accelerate the conversion of
20 3T3-L1 fibroblasts to adipocytes., *J. Lipid Res.* 43 (2002) 676–84.
21 <http://www.ncbi.nlm.nih.gov/pubmed/11971937>.
- 22 [50] R.M. Sargis, D.N. Johnson, R.A. Choudhury, M.J. Brady, Environmental endocrine
23 disruptors promote adipogenesis in the 3T3-L1 cell line through glucocorticoid

- 1 receptor activation, *Obesity*. 18 (2010) 1283–1288. doi:10.1038/oby.2009.419.
- 2 [51] S. Ahmed, E. Atlas, Bisphenol S- and bisphenol A-induced adipogenesis of murine
3 preadipocytes occurs through direct peroxisome proliferator-activated receptor
4 gamma activation, *Int. J. Obes.* 40 (2016) 1566–1573. doi:10.1038/ijo.2016.95.
- 5 [52] G. Biasiotto, I. Zanella, A. Masserdotti, R. Pedrazzani, M. Papa, L. Caimi, D. Di
6 Lorenzo, Municipal wastewater affects adipose deposition in male mice and
7 increases 3T3-L1 cell differentiation, *Toxicol. Appl. Pharmacol.* 297 (2016) 32–40.
8 doi:10.1016/j.taap.2016.02.023.
- 9 [53] X. Xie, J. Song, G. Li, MiR-21a-5p suppresses bisphenol A-induced pre-adipocyte
10 differentiation by targeting map2k3 through MKK3/p38/MAPK, *Biochem. Biophys.*
11 *Res. Commun.* 473 (2016) 140–146. doi:10.1016/j.bbrc.2016.03.066.
- 12 [54] R.W. Stahlhut, E. van Wijngaarden, T.D. Dye, S. Cook, S.H. Swan, Concentrations
13 of urinary phthalate metabolites are associated with increased waist circumference
14 and insulin resistance in adult U.S. males, *Environ. Health Perspect.* 115 (2007)
15 876–882. doi:10.1289/ehp.9882.
- 16 [55] E.E. Hatch, J.W. Nelson, R.W. Stahlhut, T.F. Webster, Association of endocrine
17 disruptors and obesity: Perspectives from epidemiological studies, *Int. J. Androl.* 33
18 (2010) 324–331. doi:10.1111/j.1365-2605.2009.01035.x.
- 19 [56] L. Trasande, T.M. Attina, S. Sathyanarayana, A.J. Spanier, J. Blustein,
20 Race/ethnicity-specific associations of urinary phthalates with childhood body mass
21 in a nationally representative sample, *Environ. Health Perspect.* 121 (2013) 501–
22 506. doi:10.1289/ehp.1205526.
- 23 [57] E.E. Hatch, J.W. Nelson, M.M. Qureshi, J. Weinberg, L.L. Moore, M. Singer, T.F.

- 1 Webster, Association of urinary phthalate metabolite concentrations with body mass
2 index and waist circumference: A cross-sectional study of NHANES data, 1999-
3 2002, *Environ. Heal. A Glob. Access Sci. Source.* 7 (2008) 1–15. doi:10.1186/1476-
4 069X-7-27.
- 5 [58] D.B. Martinez-Arguelles, E. Campioli, M. Culty, B.R. Zirkin, V. Papadopoulos, Fetal
6 origin of endocrine dysfunction in the adult: The phthalate model, *J. Steroid*
7 *Biochem. Mol. Biol.* 137 (2013) 5–17. doi:10.1016/j.jsbmb.2013.01.007.
- 8 [59] B. Desvergne, J.N. Feige, C. Casals-Casas, PPAR-mediated activity of phthalates:
9 A link to the obesity epidemic?, *Mol. Cell. Endocrinol.* 304 (2009) 43–48.
10 doi:10.1016/j.mce.2009.02.017.
- 11 [60] A.M. Calafat, L.Y. Wong, M.J. Silva, E. Samandar, J.L. Preau, L.T. Jia, L.L.
12 Needham, Selecting adequate exposure biomarkers of diisononyl and diisodecyl
13 phthalates: Data from the 2005-2006 national health and nutrition examination
14 survey, *Environ. Health Perspect.* 119 (2011) 50–55. doi:10.1289/ehp.1002316.
- 15 [61] M. Kasper-Sonnenberg, H.M. Koch, J. Wittsiepe, M. Wilhelm, Levels of phthalate
16 metabolites in urine among mother-child-pairs - Results from the Duisburg birth
17 cohort study, Germany, *Int. J. Hyg. Environ. Health.* 215 (2012) 373–382.
18 doi:10.1016/j.ijheh.2011.09.004.
- 19 [62] K. Larsson, C.H. Lindh, B.A. Jönsson, G. Giovanoulis, M. Bibi, M. Bottai, A.
20 Bergström, M. Berglund, Phthalates, non-phthalate plasticizers and bisphenols in
21 Swedish preschool dust in relation to children’s exposure, *Environ. Int.* 102 (2017)
22 114–124. doi:10.1016/j.envint.2017.02.006.
- 23 [63] A.R. Zota, A.M. Calafat, T.J. Woodruff, Temporal Trends in Phthalate Exposures:

- 1 Findings from the National Health and Nutrition Examination Survey, 2001-2010,
2 Environ. Health Perspect. 122 (2014) 235–241. doi:10.1289/ehp.1306681.
- 3 [64] C.J. Burns, L.J. McIntosh, P.J. Mink, A.M. Jurek, A.A. Li, Pesticide Exposure and
4 Neurodevelopmental Outcomes: Review of the Epidemiologic and Animal Studies, J.
5 Toxicol. Environ. Heal. Part B. 16 (2013) 127–283.
6 doi:10.1080/10937404.2013.783383.
- 7 [65] L. V. Dishaw, L.J. Macaulay, S.C. Roberts, H.M. Stapleton, Exposures,
8 mechanisms, and impacts of endocrine-active flame retardants, Curr. Opin.
9 Pharmacol. 19 (2014) 125–133. doi:10.1016/j.coph.2014.09.018.
- 10 [66] A. Debost-Legend, C. Warembourg, C. Massart, C. Chevrier, N. Bonvallot, C.
11 Monfort, F. Rouget, F. Bonnet, S. Cordier, Prenatal exposure to persistent organic
12 pollutants and organophosphate pesticides, and markers of glucose metabolism at
13 birth, Environ. Res. 146 (2016) 207–217. doi:10.1016/j.envres.2016.01.005.
- 14 [67] M. Ranjbar, M.A. Rotondi, C.I. Ardern, J.L. Kuk, The influence of urinary
15 concentrations of organophosphate metabolites on the relationship between BMI
16 and cardiometabolic health risk, J. Obes. 2015 (2015). doi:10.1155/2015/687914.
- 17 [68] T.A. Slotkin, Does early-life exposure to organophosphate insecticides lead to
18 prediabetes and obesity?, Reprod. Toxicol. 31 (2011) 297–301.
19 doi:10.1016/j.reprotox.2010.07.012.
- 20 [69] F. Peris-Sampedro, P. Basaure, I. Reverte, M. Cabré, J.L. Domingo, M.T. Colomina,
21 Chronic exposure to chlorpyrifos triggered body weight increase and memory
22 impairment depending on human apoE polymorphisms in a targeted replacement
23 mouse model, Physiol. Behav. 144 (2015) 37–45.

- 1 doi:10.1016/j.physbeh.2015.03.006.
- 2 [70] F. Peris-Sampedro, M. Cabré, P. Basaure, I. Reverte, J.L. Domingo, M. Teresa
3 Colomina, Adulthood dietary exposure to a common pesticide leads to an obese-like
4 phenotype and a diabetic profile in apoE3 mice, *Environ. Res.* 142 (2015) 169–176.
5 doi:10.1016/j.envres.2015.06.036.
- 6 [71] T.L. Lassiter, S. Brimijoin, Rats gain excess weight after developmental exposure to
7 the organophosphorothionate pesticide, chlorpyrifos, *Neurotoxicol. Teratol.* 30
8 (2008) 125–130. doi:10.1016/j.ntt.2007.10.004.
- 9 [72] A.J. Green, J.L. Graham, E.A. Gonzalez, M.R. La Frano, S.S.E. Petropoulou, J.S.
10 Park, J.W. Newman, K.L. Stanhope, P.J. Havel, M.A. La Merrill, Perinatal triphenyl
11 phosphate exposure accelerates type 2 diabetes onset and increases adipose
12 accumulation in UCD-type 2 diabetes mellitus rats, *Reprod. Toxicol.* 68 (2017) 119–
13 129. doi:10.1016/j.reprotox.2016.07.009.
- 14 [73] G. Cano-Sancho, A. Smith, M.A. La Merrill, Triphenyl phosphate enhances
15 adipogenic differentiation, glucose uptake and lipolysis via endocrine and
16 noradrenergic mechanisms, *Toxicol. Vit.* 40 (2017) 280–288.
17 doi:10.1016/j.tiv.2017.01.021.
- 18 [74] H.K. Pillai, M. Fang, D. Beglov, D. Kozakov, S. Vajda, H.M. Stapleton, T.F. Webster,
19 J.J. Schlezinger, Ligand binding and activation of PPARgamma by Firemaster(R)
20 550: effects on adipogenesis and osteogenesis in vitro, *Env. Heal. Perspect.* 122
21 (2014) 1225–1232. doi:10.1289/ehp.1408111.
- 22 [75] J.C. Caldwell, DEHP: Genotoxicity and potential carcinogenic mechanisms-A review,
23 *Mutat. Res. - Rev. Mutat. Res.* 751 (2012) 82–157. doi:10.1016/j.mrrev.2012.03.001.

- 1 [76] P. Lefebvre, Y. Benomar, B. Staels, Retinoid X receptors: Common
2 heterodimerization partners with distinct functions, *Trends Endocrinol. Metab.* 21
3 (2010) 676–683. doi:10.1016/j.tem.2010.06.009.
- 4 [77] L. Guo, X. Li, Q.Q. Tang, Transcriptional regulation of adipocyte differentiation: A
5 central role for CCAAT/ enhancer-binding protein (C/EBP) β , *J. Biol. Chem.* 290
6 (2015) 755–761. doi:10.1074/jbc.R114.619957.
- 7 [78] E.D. Rosen, O.A. MacDougald, Adipocyte differentiation from the inside out, *Nat.*
8 *Rev. Mol. Cell Biol.* 7 (2006) 885–896. doi:10.1038/nrm2066.
- 9 [79] H. Kimura, K. Fujimori, Activation of early phase of adipogenesis through Kruppel-
10 like factor KLF9-mediated, enhanced expression of CCAAT/enhancer-binding
11 protein beta in 3T3-L1 cells, *Gene.* 534 (2014) 169–176.
12 doi:10.1016/j.gene.2013.10.065.
- 13 [80] G. Li, W. Yin, R. Chamberlain, D. Hewett-Emmett, J.N. Roberts, X. Yang, S.M.
14 Lippman, J.L. Clifford, Identification and characterization of the human retinoid X
15 receptor alpha gene promoter, *Gene.* 372 (2006) 118–127.
16 doi:10.1016/j.gene.2005.12.027.
- 17 [81] C.H. Hurst, D.J. Waxman, Activation of PPAR α and PPAR γ by environmental
18 phthalate monoesters, *Toxicol. Sci.* 74 (2003) 297–308. doi:10.1093/toxsci/kfg145.
- 19 [82] J.N. Feige, L. Gelman, D. Rossi, V. Zoete, R. M \acute{e} tivier, C. Tudor, S.I. Anghel, A.
20 Grosdidier, C. Lathion, Y. Engelborghs, O. Michielin, W. Wahli, B. Desvergne, The
21 endocrine disruptor monoethyl-hexyl-phthalate is a selective peroxisome proliferator-
22 activated receptor γ modulator that promotes adipogenesis, *J. Biol. Chem.* 282
23 (2007) 19152–19166. doi:10.1074/jbc.M702724200.

- 1 [83] R. Biemann, A. Navarrete Santos, A. Navarrete Santos, D. Riemann, J. Knelangen,
2 M. Bluher, H. Koch, B. Fischer, Endocrine disrupting chemicals affect the adipogenic
3 differentiation of mesenchymal stem cells in distinct ontogenetic windows, *Biochem.*
4 *Biophys. Res. Commun.* 417 (2012) 747–752. doi:10.1016/j.bbrc.2011.12.028.
- 5 [84] H. chih Chiang, Y.T. Kuo, C.C. Shen, Y.H. Lin, S.L. Wang, T.C. Tsou, Mono(2-
6 ethylhexyl)phthalate accumulation disturbs energy metabolism of fat cells, *Arch.*
7 *Toxicol.* 90 (2016) 589–601. doi:10.1007/s00204-014-1446-9.
- 8 [85] E. Atlas, L. Pope, M.G. Wade, A. Kawata, A. Boudreau, J.G. Boucher, Bisphenol A
9 increases aP2 expression in 3T3L1 by enhancing the transcriptional activity of
10 nuclear receptors at the promoter, *Adipocyte.* 3 (2014) 170–179.
11 doi:10.4161/adip.28436.
- 12 [86] F. Acconcia, V. Pallottini, M. Marino, Molecular mechanisms of action of BPA, *Dose-*
13 *Response.* 13 (2015) 1–9. doi:10.1177/1559325815610582.
- 14 [87] R. Chamorro-García, S. Kirchner, X. Li, A. Janesick, S. C. Casey, C. Chow, B.
15 Blumberg, Bisphenol A Diplycidyl Ether Induces Adipogenic Differentiation of
16 Multipotent Stomal Stem Cells through a Peroxisome Proliferator-Activated Receptor
17 Gamma-Independent Mechanism, *Environ. Health Perspect.* 120 (2012) 984–9.
18 doi:10.1289/ehp.1205063.
- 19 [88] S.M. Regnier, R.M. Sargis, Adipocytes under assault: Environmental disruption of
20 adipose physiology, *Biochim. Biophys. Acta - Mol. Basis Dis.* 1842 (2014) 520–533.
21 doi:10.1016/j.bbadis.2013.05.028.
- 22 [89] D. Moseti, A. Regassa, W.K. Kim, Molecular regulation of adipogenesis and
23 potential anti-adipogenic bioactive molecules, *Int. J. Mol. Sci.* 17 (2016) 1–24.

2

3

4 **Figure and Table captions**

5 **Fig. 1. Scalar concentrations of BPA, DiNP, DiDP, DEGDB and TMCP enhance lipid**
6 **accumulation in differentiated 3T3-L1 cells.**

7 Upper left panel: schematic representation of the experimental protocol (for details, see
8 Materials and Methods section). The blue line indicates the presence of plasticizers (or
9 Rosiglitazone) in the cell culture medium. MDI: differentiation medium; MM: maintenance
10 medium. Graphs show quantification of lipid accumulation by Oil Red O (ORO) staining,
11 elution and absorbance reading. Three independent experiments (n=3) with 3 biological
12 replicates each were carried out. Variations in lipid accumulation were expressed as fold
13 changes of the absorbance of treated cells relative to the absorbance of control cells (=1) ±
14 SEM; *** p<0.001.

15

16 **Fig. 2. Low nanomolar concentrations of plasticizers are more effective in enhancing**
17 **lipid accumulation when administered during mid-late differentiation.**

18 Upper left panel: schematic representation of the experimental protocol (for details, see
19 Materials and Methods section). The blue lines indicate the presence of plasticizers (0.01
20 µM) in the cell culture medium. MDI: differentiation medium; MM: maintenance medium.
21 Graphs show quantification of lipid accumulation by Oil Red O (ORO) staining, elution and
22 absorbance reading. 3T3-L1 preadipocytes were treated with plasticizers alternatively from
23 day 0 to day 2 (early differentiation) or from day 2 to day 10 (mid-late differentiation). Three

1 independent experiments (n=3) were carried out with 3 biological replicates each. Variations
2 in lipid accumulation were expressed as fold changes of the absorbance of treated cells
3 relative to the absorbance of control cells (=1) \pm SEM. * differences versus control; #
4 differences between early and mid-late differentiation; * p<0.05; ** p<0.01; *** p<0.001; ##
5 p<0.01; ### p<0.001.

6

7 **Fig. 3. TMCP is predicted to interact with both PPAR γ and RXR α ligand binding**
8 **domains.**

9 3D (left side) and 2D (right side) predicted models of the TMCP/PPAR γ (A) and TMCP/RXR α
10 (B) complexes obtained by molecular docking. In the 3D representations, the receptor is
11 shown in cartoon mode, whereas TMCP is shown as stick. Predicted non-polar interactions
12 between TMCP and PPAR γ Arg²⁸⁸ / RXR α Phe³¹³ are reported in the 2D schemes. See the
13 Material & Methods section for methodological details.

14

15 **Fig. 4. Plasticizers can transactivate PPAR γ .**

16 HepG2 cells were transfected with pcDNA3-PPAR γ , DR1-Luc, and pCMV- β -galactosidase
17 vectors, then were treated with scalar concentrations of Rosiglitazone or plasticizers as
18 described under Material & Methods. Luciferase activities are reported as fold changes of
19 luminescence of treated cells versus control (=1) \pm SEM (n=3). * p<0.05.

20

21 **Fig. 5. All plasticizers modulate the expression of *Cebp β* and *Ppar γ 2* in the early**
22 **and/or mid phase of 3T3-L1 pre-adipocyte differentiation.**

23 Left panels: schematic representations of the experimental protocol. The blue line indicates

1 the presence of plasticizers (25 μ M) or Rosiglitazone (100 nM) in the cell culture medium.
2 MDI: differentiation medium; MM: maintenance medium. mRNA expression was evaluated
3 by qReal-Time PCR at day 2 (upper panel) or at day 4 (lower panel). Data are expressed
4 as fold changes in mRNA expression versus control (=1) \pm SD. * $p < 0.05$; ** $p < 0.01$; ***
5 $p < 0.001$. Graphs are representative of three independent experiments.

6

7 **Fig. 6. Among plasticizers, TMCP shows the highest similarity to Rosiglitazone in**
8 **modulating the expression of late differentiation genes.**

9 Upper left panel: schematic representation of the experimental protocol. The blue line
10 indicates the presence of plasticizers (25 μ M) or Rosiglitazone (100 nM) in the cell culture
11 medium. MDI: differentiation medium; MM: maintenance medium. mRNA expression was
12 evaluated by qReal-Time PCR at day 8. Data are expressed as fold changes in mRNA
13 expression versus control (=1) \pm SD. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$. Graphs are
14 representative of three independent experiments.

15

16 **Table 1. Primer sequences used for gene expression analysis**

17

18 **Table 2. Predicted equilibrium dissociation constants ($K_{d,pred}$) between PPAR γ , RXR α**
19 **and a set of plasticizers**

20

1 **Table 1**

Primers	Sequences
<i>Cebpβ</i>	Forward 5' – CCTGAGTAATCACTTAAAGATGT – 3' Reverse 5' – TTTAATGCTCGAAACGGAAA – 3'
<i>Rxra</i>	Forward 5' – CGGAACAGCGCTCACAGT – 3' Reverse 5' – AGCTCCGTCTTGTCCATCTG – 3'
<i>Pparγ2</i>	Forward 5' – CTGTTATGGGTGAAACTCTG – 3' Reverse 5' – ATGGCATCTCTGTGTCAA – 3'
<i>Fabp4</i>	Forward 5' – GAATTCGATGAAATCACCGCA – 3' Reverse 5' – CTCTTTATTGTGGTCGACTTTCCA – 3'
<i>Lpl</i>	Forward 5' – GATCCGAGTGAAAGCCGGAG – 3' Reverse 5' – TTGTTTGTCCAGTGTTCAGCCA – 3'
<i>β-actin</i>	Forward 5' – TCTTTGCAGCTCCTTCGTTG – 3' Reverse 5' – ACGATGGAGGGGAATACAGC – 3'

2

3

4 **Table 2**








5

6

	$K_{d,pred}$ (M) vs PPAR γ	$K_{d,pred}$ (M) vs RXR α
BPA, Bisphenol A	$1.40 \pm 0.34 \times 10^{-6}$	$8.02 \pm 1.38 \times 10^{-7}$
DiNP, Di-isononyl-phthalate	$1.34 \pm 0.24 \times 10^{-7}$	$6.09 \pm 0.98 \times 10^{-8}$
DiDP, Di-isodecyl-phthalate	$1.39 \pm 0.31 \times 10^{-7}$	$1.72 \pm 0.40 \times 10^{-7}$
DEGDB, Diethylene glycol dibenzoate	$5.55 \pm 1.24 \times 10^{-7}$	$3.74 \pm 0.79 \times 10^{-7}$
TMCP, Tri-m-cresyl phosphate	$4.27 \pm 1.26 \times 10^{-8}$	$2.56 \pm 0.40 \times 10^{-8}$
Rosiglitazone, BRL49653	$4.92 \pm 1.43 \times 10^{-8}$	$3.84 \pm 0.72 \times 10^{-8}$




7

1 Fig.1



2
 3
 4
 5
 6

1 Fig.2





- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15



1 Fig. 3

2

A



B




3

4

1 Fig. 4

2












3

4

1 Fig. 5

2





3

4

5

1 Fig. 6

2



3

