

Evaluation of the efficacy of insecticidal coatings based on teflutrin and chlorpyrifos against *Rhynchophorus ferrugineus*

Pugliese Massimo*, Rettori Andrea Alberto***, Martinis Roberto***, Al-Rohily Khalid***, Al-Maashi Ali***

* AgriNewTech srl, Via G. Quarello 15/A, 10135 Turin, Italy;

** Studio Associato Planta, Via Chiesa 19, 10090 Rosta (TO), Italy;

*** Saudi Basic Industries Corporation, PO Box 5101, KSA-11422 Riyadh, Kingdom of Saudi Arabia

Summary

The date palm (*Phoenix dactylifera* L.) is an important economic resource for many nations worldwide, recently threatened by the presence of different insect pests, like the red palm weevil (RPW) *Rhynchophorus ferrugineus*. Different strategies to control the insect have been used so far worldwide, including insecticides, biological control, endotherapy and pheromones, but it is still causing serious damages to palms in several Countries. In the present study, a new approach has been used to control the insect, based on a non-phytotoxic and non-toxic coating material applied on palm trees to prevent the red palm weevil attacks. Two products, a glue (polyvinyl acetate) and an oil (raw linseed oil), have been used as coatings and applied together with a repellent and two insecticides (teflutrin and chlorpyrifos) at different dosages on 2 species of palm (*Phoenix dactylifera* and *Phoenix canariensis*). Phytotoxic effects of the treatments have been evaluated in greenhouse on 260 potted palms (130 *Phoenix dactylifera* and 130 *P. canariensis*) and no negative effects have been observed. Afterwards, a trial lasting 400 days has been carried out in a nursery located in Sicily (South Italy), treating 572 potted palm trees (286 *P. dactylifera* and 286 *P. canariensis*) with an average diameter at base of 18-20 cm. After 400 days, 48% of the untreated palms were infested, while only 3% of date palms and 7% of Canary palms treated with insecticide at lower dosages have been infested. The application of an insecticide based coating is a good strategy to control and prevent the red palm weevil infestation, in particular on date palms.

Keywords: *Rhynchophorus ferrugineus*; teflutrin; chlorpyrifos; polyvinyl acetate; coating.

35 **Introduction**

36

37 *Rhynchophorus ferrugineus* Olivier (Red Palm Weevil - RPW), a native pest of South and South
38 East Asia, causes serious and important crop and landscape damage attacking date palms and other
39 palm species in many parts of the world (EPPO, 2007). The male and female adults of this insect,
40 member of *Coleoptera: Curculionidae*, are large reddish brown beetles about 3 cm long and with a
41 characteristic long curved rostrum. Due to their strong wings, they are capable of undertaking long
42 flights (Salama *et al.*, 2009).

43 Damage to palms is produced mainly by the larvae. Adult females lay about 200 eggs at the base of
44 young leaves or in wounds to the leaves and trunks; the grubs feed on the soft fibers and terminal
45 bud tissues. They reach a size of more than 5 cm before pupation. Except just before pupating, they
46 move towards inner tissues of the palm tree making tunnels and large cavities damaging its internal
47 tissues. The larvae can be found anywhere within the palm, even in the very base of the trunk where
48 the roots emerge. This burrowing weakens and eventually kills the tree. Although adult weevils can
49 damage trees by feeding on them, larvae can cause greater damage by burrowing.

50 The complete life cycle of the insect takes an average of 82 days (Murphy and Briscoe, 1999), up to
51 93 (Salama *et al.*, 2009) and 124 (Kaakeh, 2005), depending on different feeding substrate. They
52 can be found in any place within the palm (Alkhalil *et al.*, 2009) from the base of the trunk where
53 the roots emerge up to the apical bud. Vertical distribution of infestations along the trunk of date
54 palm trees generally showed that 70% of the attack occurred from the ground up to 1 -1.5 m. On
55 other palm trees (i.e. *Phoenix canariensis*) the attacks occur principally near the apical bud (80-
56 90%).

57 In 1994, this pest had been captured in the south of Spain and it was observed in Italy later in 2004
58 (Vacante, 2013). Today, it can be found in almost all Mediterranean countries (EPPO, 2008). The
59 spread of this pest occurs due to transporting infested young or adult date palm trees and offshoots
60 from contaminated to uninfested areas.

61 The external symptoms on infested palm trees are a progressive yellowing of the leaf area,
62 destruction of the rising leaf and necrosis in the flowers. Leaves begin to dry in ascending order in
63 the crown; the apical leaf bends and eventually drops. Affected plant tissue turns foul, producing
64 strong characteristic odours (EPPO, 2007).

65 Infested palms generally do not show any obvious early symptoms, and usually the weevil is
66 detected only after most damage has been caused on the palm, and preventive measures are mostly
67 indicated to control it (Kehat, 1999). Preventive and curative measures include: trunk injection with
68 systemic insecticides; treatment of wounds with repellents; drenching of the crown of infested trees

69 with insecticides (EPPO, 2008). Control of the red palm weevil is problematic and particularly
70 challenging for several reasons, even with pesticides. Adults are mobile and easily bypass or evade
71 containment barriers thereby expanding infestation outbreaks. Moreover, the pest is concealed deep
72 inside palm trunks and consequently insecticides have to be applied frequently and over a long
73 period (Faleiro, 2006; Llácer *et al.*, 2010; Murphy and Briscoe, 1999). In addition, a restriction in
74 the use of insecticides has occurred recently due to concerns about the side effects of chemical
75 pesticides on the environment. Systemic insecticides may be the best chemicals available for
76 controlling the RPW, also once palms are already infested (Llácer *et al.*, 2012) and are also applied
77 successfully with endoterapy and trunk injection, with concerns about residues on dates.
78 Natural enemies do not play an important part in controlling *R. ferrugineus* (Reginald, 1973) and at
79 present there is no practical biological control. However, pheromones are increasingly used against
80 *R. ferrugineus* and protocols for pheromone-based mass trapping of the weevil have been developed
81 (Hallett *et al.*, 1999).
82 Only in few cases, like in Israel, the eradication of the insect, detected in 1999, was achieved in 4
83 years by implementing several measures such as mass trapping, chemical treatment of infested
84 palms, destruction of heavily infested palms and preventive measures (Hamburger *et al.*, 2003).
85 Similarly in Morocco and Tunisia (Vacante, 2013).
86 Considering the widespread presence of RPW and the difficulties to control it, a new approach has
87 been applied to prevent palms infestation based on the use of non-phytotoxic and non-toxic coating
88 material applied on palm trees to prevent the attacks.

89

90 **Materials and methods**

91 Treatments

92 Two products, a glue (polyvinyl acetate, Vinavil 59, Vinavil Spa, Italy) and an oil (raw linseed oil,
93 Linoil, Chimica C.B.R. Spa, Italy), have been used as coatings and applied together with a repellent
94 (camphor) and two insecticides, tefluthrin (Teflutar, 0.2% p.a.) and chlorpyrifos (Zelig GR, 7.5%
95 p.a.) at different dosages on 2 species of palm (*Phoenix dactylifera* and *Phoenix canariensis*).

96 Tefluthrin is one of the most toxic pyrethroids, while chlorpyrifos is a toxic crystalline
97 organophosphate insecticide that inhibits acetylcholinesterase.

98 The mixtures of products applied in the trials are listed in Table 1.

99

100 Phytotoxicity evaluation

101 Seven years old *P. dactylifera* and 5 years old *P. canariensis* have been used and totally 260 potted
102 palms (n. 130 *P. dactylifera* and n. 130 *P. canariensis*) have been treated and placed in greenhouse

103 to assess the phytotoxicity of the selected products. The average diameter at the base was
104 respectively 9.5 ± 2.6 and 8.5 ± 1.6 cm. Ten palm trees per species have been treated for each product
105 mix. Products have been applied by painting, to avoid the drift of pesticides in an enclosed place
106 (greenhouse).

107 Phytotoxic effects have been evaluated with a monthly visual assessment observing the
108 presence/absence of leaves discoloration or necrosis (scale: 0= no symptoms; 1= light leaf chlorosis
109 or necrosis; 2= leaf chlorosis or necrosis up to 25%; 3= 26-50% leaf chlorosis or necrosis; 4= 51-
110 90% leaf chlorosis or necrosis; 5= plant death),.

111 At the same time of visual assessment other measurements have been carried out:

- 112 • Thickness measurements of the films applied (3 measurements/leaf on three leaves coated
113 portion for each plant) with Positector 200-ultrasonic coating thickness gage.
- 114 • Indirect measurements of chlorophyll content (CCI) with Chlorophyll Content Meter 200
115 (Opti-Sciences). Chlorophyll has distinct optical absorbance characteristics that the CCM-
116 200 plus exploits to non-destructively measure relative chlorophyll concentrations. Strong
117 absorbance bands are present in the blue and red regions but not the green or infrared bands,
118 hence the green appearance of a leaf. By measuring the amount of energy absorbed in the
119 red band an estimate of the amount of chlorophyll present in the tissue is possible.
120 Absorbance in the infrared band can be used to quantify and account for leaf thickness, so
121 providing a more accurate CCI value. This device has optical absorbance in two different
122 wavebands: 653 nm (Chlorophyll) and 931 nm (Near Infra-Red) providing CCI value.

123

124 Efficacy evaluation

125 The nursery activities took place between August 2013 and the end of November 2014.

126 After two months of acclimatization, treatments were carried out in October 2013 and repeated in
127 November 2013 on 572 palm trees (286 *P. dactylifera* and 286 *P. canariensis*). The two species
128 have been positioned in two separate blocks. The distance of each palm from the others was 1
129 meter. Date palms were from 9 to 13 years old, with average diameter at base 20.1 ± 5.4 cm; Canary
130 palms were from 7 to 8 years old with 17.8 ± 1.8 cm diameter. Treatments have been carried out
131 randomly inside the two blocks of palms at 14-16 October 2013, the coatings have been applied by
132 brush and by spray (Black & Decker - SmartSelect HVLP Sprayer mod. BDPH400).

133 During the period November 2013 - November 2014 measurements of palms in open field (in
134 Sicily) were carried out to characterize plant physiology and coating application, in order to assess
135 the efficacy of the products to prevent RPW infestation and to test the efficacy of the curative
136 effect.

137 Forty-four palm trees not treated (22 *P. dactylifera* and 22 *P. canariensis*) served as control. In the
138 months of August and September 2013 irrigations were made every 2 days, in October every 4
139 days, from November 2013 to March 2014 every 7 days and from April to September 2014 every 3
140 days. Fertilizations have been made in October 2013, in May 2014 and in October 2014 using a
141 ternary fertilizer (COMPO NPK Original Gold®, 5-10 g per pot).

142 The average temperature of whole period has been 18°C. Maximum temperatures were recorded in
143 the months of July, August and September and were about 40 degrees.

144 At the end of November 2014 all the palms involved have been sectioned to verify the real
145 infestation. Each palm tree has been cut by chainsaw at the base about 2-4 cm height.

146

147 Data analysis

148 About the phytotoxicity evaluation ANOVA with Tukey's HSD ($p \leq 0.05$) has been applied.

149 About the efficacy, infested palms were calculated as percentage, where healthy palm corresponds
150 to 0 and infested palm to 100. ANOVA with Tukey's HSD ($p \leq 0.05$) has been performed.

151

152

153 **Results and discussion**

154 The average thickness at the end of trials was 0.257 ± 0.088 mm for vinyl acetate and 0.091 ± 0.023
155 mm for raw linseed oil in the greenhouse trial, 0.135 ± 0.038 mm for vinyl acetate and 0.072 ± 0.032
156 mm for raw linseed oil in the nursery trial.

157 At the end of phytotoxicity test, after 7 months, the absence of damages on treated potted palms
158 were confirmed: no phytotoxic effects have been observed (Table 2). For each treatment and for
159 each parameter no significant differences were observed during the months. In particular, no
160 statistical difference have been observed comparing the treatments to the untreated control in terms
161 of chlorophyll content (CCI), leaf discolouration and necrosis.

162 Regarding the trial in Sicily, 71 palms out of 572 were infested (36 date palms and 35 Canary
163 palms) after 400 days from the treatments:

- 164 • 21 controls (not treated): 13 Date palms and 8 Canary palms;
- 165 • 32 palms treated with liquid suspensions insecticide-free;
- 166 • 11 with vinyl acetate based products (5 with Camphor and 6 without Camphor);
- 167 • 21 with raw linseed oil based products (10 with Camphor and 11 without Camphor) ;
- 168 • 3 palm treated with vinyl acetate solution containing insecticide (Teflutrin) at low
169 concentration;

- 170 • 2 palm treated with vinyl acetate solution containing insecticide (Teflutrin) at high
171 concentration;
- 172 • 4 palm treated with oil solution containing insecticide (Teflutrin) at low concentration;
- 173 • 2 palm treated with oil solution containing insecticide (Teflutrin) at high concentration;
- 174 • 3 palm treated with oil solution containing insecticide (Chlorpyrifos) at low concentration;
- 175 • 4 palm treated with oil solution containing insecticide (Chlorpyrifos) at high concentration.

176 Some infested palms were completely decayed or hollow and all stages of RPW were found (young
177 and old larvae, adults).

178 The treatments containing insecticides reduced the infestations, but after 400 days they were losing
179 their efficacy. The best results have been achieved with vinyl acetate and chlorpyrifos at both
180 dosages (T5 and T6) and no infestation have been observed after 400 days. In the case of linseed oil
181 and teflutrin or chlorpyrifos at high dosage (T7 and T9), no infestation have been observed after
182 323 days. For the combinations vinyl acetate and teflutrin at both dosages (T2 and T3), and linseed
183 oil and teflutrin or chlorpyrifos at low dosage (T8 and T10) no infestation have been observed after
184 245 days.

185 After 400 days, 36 date palms were infested by RPW: 13 controls, 4 treated with vinyl acetate
186 products (without insecticide), 1 treated with vinyl acetate product (with insecticide), 13 treated
187 with raw linseed oil (without insecticide) and 5 treated with raw linseed oil (with insecticide) (Table
188 3). Vinyl acetate with teflutrin at higher dosage, vinyl acetate with chlorpyrifos at both dosages and
189 raw linseed oil with teflutrin at higher dosage were able to prevent the infestations on date palms
190 (Table 3).

191 After 400 days, 35 Canary palms were infested by RPW: 8 controls, 7 treated with vinyl acetate
192 product (without insecticide), 4 treated with vinyl acetate product with insecticide, 8 treated with
193 raw linseed oil with insecticide and 8 treated with raw linseed oil (without insecticide) (Table 3).

194 In the case of Canary palms, only the application of vinyl acetate with chlorpyrifos at both dosages
195 was able to prevent the attacks on date palms (Table 3).

196 Comparing the results among the palms treated with linseed oil (with or without insecticides), vinyl
197 acetate (with or without insecticides) and untreated control, vinyl acetate generally reduced the
198 infestation better than raw linseed oil (Tab. 4). Moreover, both vinyl acetate and linseed oil applied
199 without insecticide were able to reduce the infestation (Tab. 4).

200 It seems that the vinyl coating without insecticide has a minimum repellent action, certainly greater
201 than the coating with raw linseed oil. In fact, at the end of the trial , infested palms treated with raw
202 linseed oil without insecticide are double respect those treated with vinyl (Tab. 5).

203 The vinyl acetate coating with insecticide had the best efficacy. Regarding the insecticides,
204 Chlorpyrifos was more effective, but only if combined with the vinyl acetate. Tefluthrin also seems
205 more effective combined with vinyl, especially in the long period (Tab. 6).

206

207 **Conclusions**

208 The treatments carried out had a positive effect on the health of the palms. After 400 days, 71 palm
209 trees were infested by RPW: 21 controls and 50 treated. Only 9.4% of the treated palms were
210 infested, compared to 47.7% of the untreated.

211 The first infestations on insecticide treated palms have been detected after 245 days, after that it
212 begins to lose effectiveness. The efficacy of the treatment is variable and depending on the time, the
213 presence and the type of active principle. The coatings with vinyl acetate and chlorpyrifos (T5 and
214 T6) seem to be the most effective (100% up to 400 days in this test).

215 These products could be easily applied by airsprayer and the solution have been prepared and
216 immediately used. Treatments on adult palms could leave small areas not covered (especially with
217 not well pruned petioles) and therefore the possibility for RPW to penetrate would be higher
218 compared to small plants. Based on information collected during the test, it can be assumed that a
219 vinyl-based coating with an insecticide like chlorpyrifos could be used to significantly reduce the
220 possibility of RPW infestation, providing a treatment every six months.

221 Similar results have been achieved by the use of the insecticidal paint Inesfly IGR FITO (Industrias
222 Químicas Inesba S.L., Paiporta, Spain), having as active ingredients 3.0% chlorpyrifos and
223 0.063% pyriproxyfen in a microencapsulated formulation (Llàcer *et al.*, 2010). The coating
224 application confers the advantage of releasing the active ingredients slowly, so the treatments do not
225 need to be applied frequently and the effect can last for a longer time (López *et al.*, 1999,
226 Mosqueira *et al.* 2005, Amelotti *et al.* 2009). In our case, the coating was able to last 400 days,
227 while the insecticidal paint Inesfly IGR FITO has been tested for 180 days (Llàcer *et al.*, 2010).

228 According to our results, the coating without chlorpyrifos was also initially effective, but of course
229 this active ingredient is the primary cause of the efficacy of the product. Chlorpyrifos is widespread
230 used to control insect pests and it is recommended to control *R. ferrugineus* in several countries
231 within the Europe. However, it is not possible to apply the coating as curative effect on already
232 infested palms, because the target pest is protected inside the palm, and other products are
233 recommended, such as imidacloprid (Kaakeh, 2006), or biological control agents (Llàcer *et al.*,
234 2009).The product can be applied within an integrated pest management strategy for *R. ferrugineus*
235 which includes surveillance, early detection, trapping using pheromones, protecting wounds,
236 physical methods, sanitation, use of attractants and other chemicals (Hoddle *et al.*, 2013; Hussain *et*

237 *al.*, 2013). In particular the adoption of early detection method (Rettori *et al.*, 2015) integrated with
238 vinyl acetate coating containing chlorpyrifos can be considered an optimal combination of methods
239 for IPM.

240
241

242 REFERENCES

243

- 244 • Alkhazal, M.H.; Youssef L.A.; Abdel-Wahaed M.S.; Kassab A.S.; Saleh M.M.E. (2009).
245 Factors affecting infestation pattern of the red palm weevil, *Rhynchophorus ferrugineus* Oliv.
246 In date palm farm in Qatif, Saudi Arabia. Arab. Univ. J. Agric. Sci. Ain Shams Univ.,
247 Cairo, 17 (1) 177-183.
- 248 • Amelotti I., Català S. S., and Gorla D. E. (2009). Experimental evaluation of insecticidal
249 paints against *Triatoma infestans* (Hemiptera: Reduviidae), under natural climatic
250 conditions. Parasites Vectors 2, 30 (<http://www.parasitesandvectors.com/content/2/1/30>).
- 251 • EPPO (2007). *Rhynchophorus ferrugineus* and *Rhynchophorus palmarum*. Bulletin
252 OEPP/EPPO Bulletin, 37(3), 571-579. [http://www.blackwell-](http://www.blackwell-synergy.com/doi/full/10.1111/j.1365-2338.2007.01165.x)
253 [synergy.com/doi/full/10.1111/j.1365-2338.2007.01165.x](http://www.blackwell-synergy.com/doi/full/10.1111/j.1365-2338.2007.01165.x)
- 254 • EPPO (2008). Data sheets on quarantine pests *Rhynchophorus ferrugineus*. EPPO Bull. 38,
255 55-59.
- 256 • Faleiro J. R. (2006). A review of the issues and management of the red palm weevil
257 *Rhynchophorus ferrugineus* (Coleoptera: Rhynchophoridae) in coconut and date palm
258 during the last one hundred years. Int. J. Trop. Insect Sci. 26, 135-154.
- 259 • Hallett R. H., Oehlschlager A. C., Borden J. H. (1999). Pheromone-trapping protocols for
260 *Rhynchophorus ferrugineus*. International Journal of Pest Management 45, 231–237.
- 261 • Hamburger M., Bitton S., Nakache J. (2003). Control of *Rhynchophorus ferrugineus*, a
262 quarantine pest in Israel. Phytoparasitica, 3, 299 –300.
- 263 • Hoddle M. S., Al-Abbad A. H., El-Shafie H.A.F., Faleiro J. R., Sallam A. A., Hoddle C. D.
264 (2013). Assessing the impact of areawide pheromone trapping, pesticide applications, and
265 eradication of infested date palms for *Rhynchophorus ferrugineus* (Coleoptera:
266 Curculionidae) management in Al Ghowaybah, Saudi Arabia. Crop Protection, 53, 152-160.
- 267 • Hussain A., Rizwan-ul-Haq M., Al-Jabr A. M., Al-Ayied H. Y. (2013) – Managing invasive
268 populations of red palm weevil: a worldwide perspective. Journal of Food Agriculture and
269 Environment, 11 (2), 456-463.

- 270 • Kaakeh W. (2005). Longevity, fecundity, and fertility of the red palm weevil,
271 *Rhynchophorus ferrugineus* Olivier (Coleoptera: Curculionidae) on natural and artificial
272 diets. Emir J agric sciences 17: 23-33
- 273 • Kaakeh W. (2006). Toxicity of imidacloprid to developmental stages of *Rhynchophorus*
274 *ferrugineus* (Curculionidae: Coleoptera): laboratory and field tests. Crop Prot. 25, 432-439.
- 275 • Kehat M. (1999). Threat to date palms in Israel, Jordan and the Palestinian authority by the
276 red palm weevil, *Rhynchophorus ferrugineus*. Phytoparasitica 27, 241-242.
- 277 • Llàcer E., Martínez de Altube M. M., Jacas J. A. (2009). Evaluation of the efficacy of
278 *Steinernema carpocapsae* in a chitosan formulation against the red palm weevil,
279 *Rhynchophorus ferrugineus* in *Phoenix canariensis*. Biocontrol 54, 559-565.
- 280 • Llàcer E., Dembilio O., Jacas J. A. (2010). Evaluation of the efficacy of an insecticidal paint
281 based on chlorpyrifos and pyriproxyfen in a microencapsulated formulation against
282 *Rhynchophorus ferrugineus* (Coleoptera: Curculionidae). Journal of Economic Entomology,
283 103 (2), 402-408.
- 284 • Llàcer E., Negre M., Jacas J.A. (2012). Evaluation of an oil dispersion formulation of
285 imidacloprid as a drench against *Rhynchophorus ferrugineus* (Coleoptera: Curculionidae) in
286 young palm trees. Pest Manag. Sci. 68, 878-882.
- 287 • Lòpez J., Jimènez R., Mateo M. P., Moreno J., Oltra M. T. (1999). Efficacy of Inesfly 5A
288 IGR paint against *Periplaneta americana* in sewers in Canary Islands (Spain). In:
289 Proceedings of the 3rd International Conference on Urban Pests, 19-22 July 1999, Prague,
290 Czech Republic (W. H. Robinson, F. Rettich, and G. W. Rambo coord.). Czech University
291 of Agriculture, Prague, Czech Republic, p. 623.
- 292 • Mosqueira B., Finot L., Chabi J., Akogbeto M., Chandre F., Hougard J. M., Carnevale P.,
293 and Mas-Coma S. (2005). Evaluation of the efficacy of a new insecticide paint for Malaria
294 control. In: Proceedings of the Medicine and Health in the Tropics Congress, 11-15
295 September 2004, Marseille, France, pp. 107-108, (<http://www.festmih.eu/document/1146>).
- 296 • Murphy S. T., Briscoe B. R. (1999). The red palm weevil as an alien invasive: biology and
297 the prospects for biological control as a component of IPM. Biocontrol News Inf. 20, 35-46.
- 298 • Reginald C. (1973). Principal Insect Pests. In: Coconuts. Tropical Agriculture Series,
299 London, UK, 68.
- 300 • Rettori A. A., Martinis R., Pugliese M., Al-Rohily K., Al-Maashi Al, Occhiello E., Gullino
301 M. L. (2015). Use of tree stability assessment devices to detect red palm weevil infestation,
302 submitted.

- 303 • Salama, H. S.; Zaki, F. N.; Abdel-Razek, A. S. (2009). Ecological and biological studies on
304 the red palm weevil *Rhynchophorus ferrugineus* (Olivier). Archives of Phytopathology and
305 Plant Protection Vol. 42 No. 4 pp. 392-399.
- 306 • Vacante V. (2013) - Punteruolo rosso delle palme: aggiornamenti sulle tecniche di lotta.
307 Protezione delle Colture, 6 (5), 38-51.

308

309

310

311

312

313 Table 1 – Different treatments applied in the trials.

314

Thesis	Coating	Solvent	Insecticide	Repellent
T1	Vinyl acetate 59 (100 g)	38.7 g H ₂ O		5 g Camphor White oil
T2	Vinyl acetate 59 (100 g)	109.3 g H ₂ O	40 g Teflutar (0.08 g teflutrin)	5 g Camphor White oil
T3	Vinyl acetate 59 (100 g)	109.3 g H ₂ O	20 g Teflutar (0.04 g teflutrin)	5 g Camphor White oil
T4	Vinyl acetate 59 (100 g)	38.7 g H ₂ O		
T5	Vinyl acetate 59 (100 g)	82.7 g H ₂ O	30 g Zelig GR (2.25 g chlorpyrifos)	5 g Camphor White oil
T6	Vinyl acetate 59 (100 g)	46.7 g H ₂ O	15 g Zelig GR (1.125 g chlorpyrifos)	5 g Camphor White oil
T7	Raw Linseed oil (100 g)	25 g Limonene	40 g Teflutar (0.08 g teflutrin)	5 g Camphor White oil
T8	Raw Linseed oil (100 g)	25 g Limonene	20 g Teflutar (0.04 g teflutrin)	5 g Camphor White oil
T9	Raw Linseed oil (100 g)	25 g Limonene	30 g Zelig GR (2.25 g chlorpyrifos)	5 g Camphor White oil
T10	Raw Linseed oil (100 g)	25 g Limonene	15 g Zelig GR (2.25 g 1.125 g chlorpyrifos)	5 g Camphor White oil
T11	Raw Linseed oil (100 g)	25 g Limonene		
T12	Raw Linseed oil (100 g)	25 g Limonene		5 g Camphor White oil
T13	Untreated control			

315

316

317

318 Table 2 – Evaluation of phytotoxicity of coating treatments on date and canary palms.

319

Treatment	CCI		Discolouration**		Necrosis**	
	<i>Phoenix dactylifera</i>	<i>Phoenix canariensis</i>	<i>Phoenix dactylifera</i>	<i>Phoenix canariensis</i>	<i>Phoenix dactylifera</i>	<i>Phoenix canariensis</i>
T1	28.42±4.61*	53.37±15.52a	0.00±0.00a	0.00±0.00a	0.20±0.42a	0.50±0.53a
T2	35.13±11.11a	53.00±11.95a	0.00±0.00a	0.10±0.00a	0.10±0.32a	0.70±0.67a
T3	34.53±8.78a	45.37±10.46a	0.00±0.00a	0.10±0.32a	0.00±0.00a	0.40±0.52a
T4	30.41±9.49a	51.80±8.77a	0.00±0.00a	0.00±0.00a	0.10±0.32a	0.60±0.70a
T5	38.27±9.53a	44.85±15.13a	0.00±0.00a	0.00±0.00a	0.00±0.00a	0.10±0.32a
T6	36.55±5.95a	54.96±8.18a	0.00±0.00a	0.10±0.32a	0.10±0.32a	0.40±0.52a
T7	39.43±7.54a	46.26±11.01a	0.00±0.00a	0.10±0.32a	0.00±0.00a	0.40±0.70a
T8	41.78±8.57a	52.38±9.97a	0.00±0.00a	0.10±0.32a	0.00±0.00a	0.40±0.52a
T9	38.09±10.17a	52.32±10.64a	0.00±0.00a	0.00±0.00a	0.00±0.00a	0.10±0.32a
T10	43.29±10.83a	53.82±13.16a	0.00±0.00a	0.30±0.42a	0.10±0.32a	0.70±0.67a
T11	39.79±10.22a	48.25±12.70a	0.00±0.00a	0.00±0.00a	0.00±0.00a	0.50±0.53a
T12	41.73±8.45a	50.54±12.81a	0.00±0.00a	0.10±0.32a	0.00±0.00a	0.50±0.53a
Control	32.65±7.01a	58.11±15.33a	0.00±0.00a	0.00±0.00a	0.10±0.32a	0.30±0.42a
Total average	36.93±9.48	51.16±12.21	0.00±0.00	0.07±0.23	0.0538±0.22	0.43±0.55

320

321 * Tukey's HSD ($p \leq 0.05$)

322 **Discolouration/necrosis of leaves, scale: 0= no symptoms; 1= light leaf chlorosis/necrosis; 2=

323 leaf chlorosis/necrosis up to 25%; 3= 26-50% leaf chlorosis/necrosis; 4= 51-90% leaf

324 chlorosis/necrosis; 5= plant death.

325

326

327

328 Table 3 – Efficacy evaluation of coating treatments on date and canary palms, after 400 days.
 329 Infested palms were calculated as percentage, where healthy palm corresponds to 0 and infested
 330 palm to 100.

331

Treatment	Valid cases	<i>Phoenix dactylifera</i>	<i>Phoenix canariensis</i>	Average
		% of healthy palms		
T1	22	13.63±7.49ab*	9.09±6.27ab	11.36±4.84bc
T2	22	0±0a	9.09±6.27ab	4.54±3.17ab
T3	22	4.54±4.54ab	9.09±6.27ab	4.83±3.84ab
T4	22	4.54±4.54ab	22.73±9.14cd	13.64±5.23bc
T5	22	0±0a	0±0a	0±0a
T6	22	0±0a	0±0a	0±0a
T7	22	0±0a	9.09±6.27a	4.54±3.18ab
T8	22	9.09±6.27ab	9.09±6.27a	9.09±4.38bc
T9	22	9.09±6.27ab	9.09±6.27a	9.09±4.38bc
T10	22	4.54±4.54ab	9.09±6.27ab	6.82±3.84ab
T11	22	36.36±10.50c	13.64±7.49bc	25.00±6.60cd
T12	22	22.73±9.14bc	22.73±9.14cd	22.73±6.39cd
CONTROL	22	59.09±10.73c	36.36±10.50d	47.73±7.62e

332

333 * Tukey's HSD ($p \leq 0.05$)

334

335 Table 4 – Efficacy evaluation of treatments 1-4 (vinyl acetate without insecticide), 2-3-5-6 (vinyl
 336 acetate with insecticide), 7-8-9-10 (Raw Linseed oil with insecticide), 11-12 (Raw Linseed oil
 337 without insecticide) and control comparison on date and canary palms, after 400 days. Infested
 338 palms were calculated as percentage, where healthy palm corresponds to 0 and infested palm to
 339 100.

Treatment	<i>Phoenix dactylifera</i>	<i>Phoenix canariensis</i>	Average
	% of healthy palms		
T2-T3-T5-T6 (vynil acetate based products with insecticide)	1.14±1.14a*	4.54±2.23ab	2.84±1.25a
T1-T4 (vynil acetate based products without insecticide)	9.09±4.38a	15.91±5.58ac	12.50±3.54b
T7-T8-T9-T10 (Raw Linseed based products with insecticide)	5.68±2.48a	9.09±3.08ab	7.39±1.98ab
T11-T12 (Raw Linseed based products without insecticide)	29.54±6.95b	18.18±5.88c	23.86±4.57c
Control	59.09±10.73c	36.36±10.49d	47.73±7.62d

340

341 * Tukey's HSD ($p \leq 0.05$)

342
343 Table 5 - Percentage of infested palms related to products and insecticides
344

		110 days	152 days	180 days	210 days	245 days	285 days	323 days	400 days
Vinyl acetate	with insecticide	0,0%	0,0%	0,0%	0,0%	0,0%	1,1%	1,7%	2,8%
	without insecticide	2,3%	2,3%	3,4%	3,4%	4,5%	5,7%	10,2%	12,5%
Raw Linseed Oil	with insecticide	0,0%	0,0%	0,0%	0,0%	0,0%	1,1%	1,1%	7,4%
	without insecticide	3,4%	5,7%	8,0%	9,1%	12,5%	15,9%	18,2%	23,9%
Control		6,8%	6,8%	9,1%	15,9%	22,7%	29,5%	38,6%	47,7%

345
346 Table 6 - Percentage of infested palms related to insecticides.
347
348

		110 days	152 days	180 days	210 days	245 days	285 days	323 days	400 days
Teflutrin	Vinyl acetate	0,0%	0,0%	0,0%	0,0%	0,0%	2,3%	3,4%	5,7%
	Raw Linseed Oil	0,0%	0,0%	0,0%	0,0%	0,0%	1,1%	1,1%	6,8%
Chlorpyrifos	Vinyl acetate	0,0%	0,0%	0,0%	0,0%	0,0%	0,0%	0,0%	0,0%
	Raw Linseed Oil	0,0%	0,0%	0,0%	0,0%	0,0%	1,1%	1,1%	8,0%

349

350

351