



18th European Weed Research Society Symposium

EWRS 2018

17-21 June 2018
Ljubljana, Slovenia

**New approaches for
smarter weed management**

Book of Abstracts

Organiser

Kmetijski inštitut Slovenije (KIS) – Agricultural Institute of Slovenia, Hacquetova ulica 17, 1000 Ljubljana, Slovenija

Programme Committee

Dr. Jukka Salonen, Chairman / EWRS Vice-President; Prof. Dr. Hüsrev Mennan, Scientific Secretary; Prof. Dr. Paul Neve, EWRS President

EWRS Scientific Committee

Dr. Theo Verwijst, Department of Crop Production Ecology, Swedish University of Agricultural Sciences, Sweden; Prof. Dr. Garifalia Economou-Antonaka, Faculty of Crop Production Science, Agricultural University of Athens, Greece; Dr. Kirsten Torresen, Norwegian Institute of Bioeconomy Research (NIBIO), Norway; Dr. Roland Beffa, Bayer Crop Science AG, Integrated Weed Management & Resistance Biology, Germany; Dr. Ivo O. Brants, Monsanto Europe S.A, Belgium; Jan Petersen, University of Applied Science Bingen, Germany; Christian Bohren, Research Station Agroscope Changins-Wädenswil (ACW), Switzerland; Dr. Per Kudsk, Professor & Head of Section, Dept. of Agroecology, University of Aarhus, Denmark; Maurizio Vurro, Istituto di Scienze delle Produzioni Alimentari – CNR, Italy; Dr. Marleen Riemens, Wageningen University and Research Centre, Netherlands; Prof. Dr. Svend Christensen, University of Copenhagen, Faculty of Life Sciences, Denmark; Dr. Hanan Eizenberg, Newe Ya'ar Research Center, Israel; Dr. Euro Pannacci, Dept. of Agricultural, Food and Environmental Sciences – University of Perugia, Italy; Dr. Paula Westerman, Group Crop Health, Faculty of Agricultural and Environmental Science, University of Rostock, Germany

Local Organising Committee

Assoc. Prof. Dr. Andrej Simončič, President, Agricultural Institute of Slovenia; Dr. Robert Leskovšek, Agricultural Institute of Slovenia; Dr. Gregor Urek, Agricultural Institute of Slovenia; Prof. Dr. Stanislav Trdan, Biotechnical Faculty, University of Ljubljana; Prof. Dr. Mario Lešnik, Faculty of Agriculture and Life Sciences, University of Maribor; Ela Žilič, M. Sc., Agricultural Institute of Slovenia; Marjeta Urbančič Zemljič, M. Sc., Agricultural Institute of Slovenia

Editor

Andrej Simončič

Published by

Kmetijski inštitut Slovenije, 2018

The publication is published e-only – <http://www.ewrs.org>

Cover crops as mulching to manage weeds in organic rice cultivation

Francesco Vidotto, Silvia Fogliatto, Fernando De Palo, Marco Milan, Aldo Ferrero
University of Torino, GRUGLIASCO, Italy

Cover crops sowing after rice harvesting has begun to spread in organic rice cultivation in Italy as a practice to improve soil fertility and weed management. A study aimed at evaluating the use of cover crops during winter in organic rice production and to estimate the effect of the technique on weeds and rice yield was carried out at two sites in NW Italy. In both experiments, the following cover crops were compared: *Vicia villosa*, *Lolium multiflorum*, and a mixture of *V. villosa* (40%) and *L. multiflorum* (60%). A plot without cover crops was added as a reference. Cover crops were sown in October and let grow until spring. In May, rice was broadcasted in the standing cover crops and the cover crops were terminated immediately after. Roller-crimping and shredding were compared as termination techniques. After cover crop termination, the plots were flooded for about 10 days in order to start cover crop decay and stimulate the production of phytotoxic compounds that may hamper weed infestation. Afterwards, rice field was dried to favor rice rooting. Rice was then grown organically with continuous flooding. Cover crop density was assessed after establishment, while weed density was assessed before cover crop termination and during rice growing season. Rice yield was also recorded for each plot. Weed density assessed in June showed that in both experiments the control plots were the most infested (> 900 plants m⁻²), while the plots with *L. multiflorum* both crimped or shredded were the less infested (< 60 plants m⁻²), followed by the plots hosting *V. villosa*. Rice yield was higher for *V. villosa* (3.4 t ha⁻¹) plots, while lowest for mix plots (1.1 t ha⁻¹), regardless the cover crop termination technique.