

1 **Evaluation of the short term effect of nursery treatments with phosphite-based products,**
2 **acibenzolar-S-methyl, pelleted *Brassica carinata* and biocontrol agents, against lettuce and**
3 **cultivated rocket *Fusarium* wilt under artificial inoculation and greenhouse conditions**

4

5 **Giovanna Gilardi ^a, Stefano Demarchi ^a, Maria Lodovica Gullino ^{a,b} and Angelo Garibaldi ^a**

6 ^a Centre for Innovation in the Agro-Environmental Sector, AGROINNOVA, University of Torino, Largo
7 Braccini 2, 10095 Grugliasco (TO), Italy

8 ^b Department of Agricultural, Forest and Food Sciences (DISAFA), University of Torino, Largo Braccini 2,
9 10095 Grugliasco (TO), Italy

10 *Corresponding author. Tel.: +39 0116708539; fax:+39 0116709307 E-mail address:

11 marialodovica.gullino@unito.it

12

13 **ABSTRACT**

14

15 Experimental trials have been carried out in order to evaluate the efficacy of preventative
16 treatments based on plant defense activator products, biocontrol agents, a microbial complex with
17 arbuscular mycorrhizal fungi, and *Brassica carinata* pellets against *Fusarium oxysporum* f. sp.
18 *lactucae* race 1 on lettuce and *Fusarium oxysporum* f. sp. *raphani* on cultivated rocket under
19 greenhouse conditions. These products were compared with fungicides known for their ability to
20 induce host resistance (phosethyl-Al and acibenzolar-S-methyl), and with azoxystrobin. Three and
21 four applications of the tested products were carried out on lettuce and rocket seedlings grown in
22 nursery conditions. Treated and untreated plants were transplanted into soil infested with *Fusarium*
23 wilt agents to obtain an average disease severity (DS) of 65.6-69.2 and of 56.9-62.1 on the untreated
24 lettuce and rocket plants, respectively. The best *Fusarium* wilt biocontrol was obtained after four

25 applications of *Bacillus subtilis* Qst713 and with the Glomas microbial complex (42 and 46.7%,
26 efficacy, respectively). *Brassica carinata* pellets provided a consistent control when applied 14 days
27 before the rocket and lettuce were transplanted into the infested soil. Acibenzolar-S-methyl, applied
28 at 0.025 g/liter, showed a DS reduction in *F. oxysporum* f. sp. *lactucae* from 36 to 61% and of *F.*
29 *oxysporum* f. sp. *raphani* from 54 to 73%, thus showing statistically similar results to those of
30 azoxystrobin, which was used as a reference (DS reduction from 59 to 65%). Although the
31 Fusarium wilt control provided by such products was not complete in the present experimental
32 conditions, these products can be considered interesting components for an integrated pest
33 management of the Fusarium wilt of leafy vegetables, starting from nursery applications. Moreover,
34 the tested BCAs could become potentially useful, especially for plant monocultures. This study has
35 been produced new information on the effects of potassium phosphite, applied at the nursery level,
36 on reducing lettuce and rocket fusarium wilt. An average efficacy of 69.5% was observed for
37 lettuce, while an average efficacy of 65.2% was observed for cultivated rocket. The good fungicidal
38 activity of the phosphite-based product, coupled with the positive effect on plant biomass, is of special
39 interest.

40

41 **Keywords:** *Lactuca sativa*, *Eruca vesicaria*, *Fusarium oxysporum* f. p. *lactucae*; *Fusarium*
42 *oxysporum* f. sp. *raphani*, integrated control

43

44 **1. Introduction**

45 In recent years, the economic relevance of lettuce (*Lactuca sativa* L.) and cultivated rocket [*Eruca*
46 *vesicaria* (L.) Cav.] has increased, since many farms produce fresh cut and ready-to-eat salads. In
47 these intensive systems, where leafy vegetables are continuously grown in the same soil, the
48 phytopathological situation is constantly in evolution, as a consequence of the dynamism and
49 specialization of such crops (Garibaldi and Gullino, 2010; Garibaldi et al., 2014). Fusarium wilt

50 incited by *Fusarium oxysporum* f. sp. *lactucae* on lettuce and by *Fusarium oxysporum*: ff. spp.
51 *raphani* and *conglutinans* on cultivated rocket, can lead to serious losses (Matheron and Gullino,
52 2012). The *Fusarium* wilt of lettuce was detected for the first time in Europe, in northern Italy, in
53 2002 (Garibaldi and Gullino, 2010). The two *formae speciales* that affect cultivated rocket, that is,
54 *Fusarium oxysporum* ff.spp. *raphani* and *conglutinans*, the first of which is more frequently
55 detected (Garibaldi et al., 2006; Srinivasan et al., 2012), also affect other genera belonging to
56 Brassicaceae, such as cabbage, brussel sprouts, broccoli, turnip, radish and stock (Garibaldi et al.,
57 2006). The *Fusarium* wilt of lettuce and rocket are easily and frequently seed-transmitted (Gullino
58 et al., 2014), thus suggesting the importance of preventative disease management strategies.

59 Lettuce varieties that are resistant or at least tolerant to *Fusarium* wilt are available (Scott et al.,
60 2010; Matheron and Gullino, 2012; Gilardi et al., 2014b), but their effective use is complicated by
61 the presence of three races of the pathogen (Fujinaga, 2005). In the case of rocket, the use of
62 resistant varieties is still very limited (Gilardi et al., 2007).

63 In general, the management of soil-borne pathogens is complicated by the limited number of
64 registered chemicals and by the restrictions in the use of pre-plant fumigants, including metam
65 sodium and dazomet (Colla et al., 2012). Several approaches to soil-borne disease management
66 have been investigated intensively in an attempt to find an answer to the many practical problems
67 associated with the loss or limitation of use of effective fumigants encountered by growers.
68 Moreover, more emphasis is now given to crop and soil health instead of disease control (Barrière
69 et al., 2014). Among the exploited strategies for disease management, systemic acquired resistance
70 (SAR) and induced systemic resistance (ISR), which are mainly triggered by microorganisms, such
71 as plant growth-promoting rhizobacteria, by the metabolic products of affected plants, or by
72 chemicals (Sticher et al., 1997; Oostendorp et al., 2001; Vallad and Goodman, 2004; Shores et al.,
73 2005) are at present attracting a great deal of interest. Induced systemic and localized resistance to
74 soil-borne pathogens, by means of *Trichoderma* treatments, has been well documented (Shores et

75 al., 2005; Vinale et al., 2008). Kloepper et al., (2004) proved the ability of *Bacillus* spp. to induce
76 systemic resistance, and in most cases, to also elicit plant growth promotion. Moreover, arbuscular
77 mycorrhizal fungi have been reported to be implicated in protecting plants against soil-borne
78 pathogens, through different mechanisms, including an improvement in plant nutrition, damage
79 compensation, competition, changes in the root system and activation of plant defense signaling
80 (Whipps, 2004; Pozo and Azcón-Aguliar, 2007). Moreover, among the chemical resistance
81 inducers, the phosphite-based fertilizers and acibenzolar-S-methyl have been shown to lead to a
82 reduction in disease against soil-borne pathogens, on vegetable and ornamental crops (Eikemo et
83 al., 2003; Elmer 2004; 2006; Hyeon et al., 2009; Bubici et al., 2006; Walters, 2012; 2013; Gilardi et
84 al., 2014a), but no data are available concerning their efficacy against the *Fusarium* wilts of lettuce
85 and rocket.

86 This study has been carried out in order to evaluate the efficacy of preventative treatments,
87 including SAR and IRS activator products, biocontrol agents, a microbial complex based on
88 arbuscular mycorrhizal fungi, and *Brassica carinata* pellets against *Fusarium oxysporum* f. sp.
89 *lactucae* race 1 on lettuce and *Fusarium oxysporum* f. sp. *raphani* on cultivated rocket under
90 simulated nursery conditions, in a greenhouse.

91

92 **2. Material and methods**

93 *2.1. Plant material and experimental layout*

94 Experimental trials (Table 1) were carried out in 2012 and 2013 under greenhouse conditions in
95 order to test the efficacy of different products against the *Fusarium* wilts of lettuce and cultivated
96 rocket. Lettuce (cv. Crispilla) and cultivated rocket seeds (cv. Coltivata), which are very susceptible
97 to *Fusarium* wilt (Garibaldi et al., 2004; Gilardi et al., 2014,), were sown in 100-plug trays (2.5 cm
98 Ø per pot, 4-L of soil capacity) filled with a steamed (90°C for 30 minutes) peat mix substrate

99 (blond peat:black peat 15:85, pH 5.5-6.0, 1,100 g m⁻³ of N:P:K and traces of molybdenum, Brill
100 Type 5, Georgsdorf, Germany).

101 Fifteen-day-old lettuce and cultivated rocket seedlings were transplanted into the same substrate in
102 12-L plastic pots. Ten plants/pot of each tested crop were kept in a greenhouse on benches with air
103 temperatures ranging from 26 to 34°C during the day and from 20 to 25°C during the night (Table
104 1). The substrate was artificially infested with the Fusarium wilt agents, as described hereafter.
105 Forty plants per treatment were arranged in a complete randomized block design in each trial, which
106 represented the experimental unit, with three (Protocol 1) and four replicated trials (Protocol 2) as
107 reported in Table 1.

108

109 2.2. Fungal strains and artificial inoculation

110 A highly virulent strain of *F. oxysporum* f. sp. *lactucae*, isolated in 2002 from infected lettuce plants
111 in north-western Italy, ATCCMYA3040, belonging to Race 1 of the pathogen (Garibaldi et al.,
112 2002), and the FusRuc 13/03 strain, isolated from rocket grown in a commercial plastic greenhouse
113 in northern-Italy in 2003 and identified as *F. oxysporum* f. sp. *raphani* (Garibaldi et al., 2006), were
114 used throughout the experiments. The single-spore culture of each isolate was stored in glycerol at -
115 80°C.

116 These strains were grown in potato dextrose broth (Sigma-Aldrich, St. Luis, USA) and kept in a
117 rotatory shaker working at 90-100 rpm for 10 days at 25°C. The biomass produced after
118 centrifugation (9,600 g at 4°C) was prepared as a dry talc-based powder (biomass: talc 2:1 w/w), as
119 described by Locke and Colhoun (1974). After 20 days at 22-25°C, the number of chlamydospores
120 per gram of talc was assessed by serial plating on potato dextrose agar, PDA (Merck, Darmstadt,
121 Germany), which contained 25 mg L⁻¹ streptomycin sulphate. The talc formulations of *F.*
122 *oxysporum* f.sp. *lactucae* (strain MYA3040) and of *F. oxysporum* f.sp. *raphani*, (strain

123 FusRuc13/03) at 2 and 5×10^7 chlamidospores/g, respectively, were used. These formulated
124 pathogens (2 and 1 g liter/respectively) were mixed into the steamed substrate as chlamydo-
125 spores dispersed in talc to achieve a final concentration of 5×10^4 chlamydo-
126 spores ml⁻¹ of substrate. A non-infested substrate was used as a control (Table 1).

127

128 2.3. Products used in the test

129 Different compounds known for their capability to induce resistance in the host, that is, phosphite-
130 based fertilizers, organic amendments, biocontrol agents (BCAs) and fungicides were tested (Table
131 2).

132 Among the considered phosphite-based products and organic amendments, a phosphite-based
133 glucohumate complex (Glucoinductor + GlucoActivator, N 4%, P₂O₅ 18%, International patent
134 PCT, IB2004/001905, Fertirev, Torino, Italy), a mineral fertilizer based on potassium phosphite
135 (Alexin 95PS, P₂O₅ 52%, K₂O 42%, Massò, Spain), and a patented formulation of *Brassica*
136 *carinata* defatted seed meal (Biofence, N organic 3%, P 2.2%, K 2%, organic C 52%, Triumph,
137 Spain) were tested.

138 Among the BCAs, *Bacillus subtilis* QST 713 (Serenade, 14.6 % a.i., BayerCropScience, Italy),
139 *Bacillus velezensis* (Cilus Plus IT45, 95%, Massò, Spain), *Trichoderma asperellum* +*T. gamsii*
140 (Remedier WP, Isagro Ricerca, Milano, Italy), a product based on arbuscular mycorrhizal fungi
141 combined with a microbial complex of *Trichoderma* and *Bacillus* (Rizocore, *Glomus* spp.
142 5%+*Bacillus megaterium* 10⁴ UFCg⁻¹ +*Trichoderma* 10¹⁰ UFCg⁻¹, Biogard, division of CBCEurope,
143 Italy) were tested.

144 The chemicals known for their ability to induce resistance, that is, acibenzolar-S-methyl (Bion
145 50WG, 50% a.i., Syngenta Crop Protection, Italy) and phosethyl-Al (Alliette, 80% a.i, Bayer Crop

146 Science, Italy) were tested, while azoxystrobin (Ortiva, 23.2% a. i., Syngenta Crop Protection, Italy)
147 was used as traditional chemical control (Gullino et al., 2002).

148

149 2.4. Type and timing of the treatments

150 The biocontrol agents, acibenzolar-S-methyl, the phosphite based products, as well as the tested
151 fungicides phosethyl-Al and azoxystrobin were applied as a leaf spray with a high volume of water
152 (1,500 L ha⁻¹), using a 1 L capacity hand sprayer.

153 The temporal organization of the experimental trials, as well as, the timing of the artificial
154 infestation of the soil and treatments are summarized in Table 1. The timing and the application
155 dosages of the tested products, which were based on the manufacturer's suggestions, are given in
156 table 2. The lettuce and cultivated rocket seedlings grown in each tray were treated by leaf spraying
157 with three applications at 6-7 day intervals according to protocol 1 (Trial block 1 to 3), or with one
158 more spray applications in the 12 L plastic pots, as in protocol 2 (Trial block 4 to 7), seven days
159 after transplanting the lettuce and cultivated rocket seedlings. Azoxystrobin and *B. carinata* pellets
160 were only applied once (Tables 1 and 2).

161 The first treatment (T0) was carried out in a greenhouse, at a temperature of 22-24°C, on plants still
162 in the plug tray, at the second true leaf stage, 7-10 days after sowing.

163 The product based on arbuscular mycorrhizal fungi and the microbial complex (Rizocore) was
164 mixed with 4 L of the substrate that was used in the plug tray at T0, than leaf sprayed at 6-7 day
165 intervals (Table1). The patented formulation of *B. carinata* defatted seed meals was mixed with the
166 substrate used to fill the 12 L plastic pots at T0 (14 days before the crops transplant) and at the same
167 time as the artificial infestation of the soil at T7.

168

169

170 2.5. Data collection and analysis

171

172 The plants were monitored weekly and the data were recorded, starting 10-14 days after the lettuce
173 and rocket had been transplanted into the infested substrate, at the appearance of the first yellow
174 leaf symptoms and reduced growth. The number of infected plants showing wilting and stem
175 necrosis was counted to assess disease incidence. The totally wilted (dead) plants were removed.
176 The final disease rating was made four weeks after the soil infestation by dissecting each plant. The
177 used disease severity (DS) index was: 0 = healthy plant, 25 = initial leaf chlorosis, 50 = severe leaf
178 chlorosis and initial symptoms of wilting during the hottest hours of the day, 75 = severe wilting
179 and severe symptoms of leaf chlorosis; 100 = plant totally wilted, leaves completely necrotic. At the
180 end of the trials, the total biomass was weighed in order to evaluate the effect of the tested
181 treatments on the yield. DS data were arcsine transformed in order to normalize the distribution of
182 variance. The data from the non-inoculated and untreated controls were not included in the
183 statistical evaluation of the DS data (Figure 1).

184 The efficacy of the different treatments in controlling the Fusarium wilt of lettuce and rocket,
185 corresponding to the percentage of DS reduction, was calculated as:

186

187
$$\% \text{ efficacy} = 100 - (DS_t \times 100 / DS_i \text{ control})$$

188

189 where,

190 i = inoculated and untreated control

191 t = tested treatments.

192

193 All the data were analysed by univariate ANOVA in SPSS 22.0, and the means were separated by
194 means of Tukey's multiple comparison test ($p=0.05$). The standard errors are marked with error
195 bars in all the figures.

196 3. Results

197

198 Effect of pre-planting treatment on lettuce fusarium wilt

199 The average disease severity (DS) at the end of the lettuce trials in the inoculated non-treated
200 control plots was 62.1 and 56.9, respectively (Figures 1a and 2a). When at least three treatments
201 were carried out, the best control of Fusarium wilt on lettuce was provided by the phosphite-based
202 product Alexin (69.2% efficacy), and this was followed by acibezolar-S-methyl at 0.0125 (59.9%
203 efficacy), by phosethyl-Al (57.8% efficacy) and the phosphite-based glucohumate complex (52.4%
204 efficacy). These treatments were effective as one application of azoxystrobin, which was found to
205 protect the lettuce from Fusarium wilt with an efficacy 55.7%. The biocontrol agents *Trichoderma*
206 *asperellum* + *T. gamsii* and *B. subtilis* provided a partial disease reduction, with statistically
207 different results compared to the inoculated and untreated control (32.8 and 30.7% efficacy,
208 respectively), while *B. velenzensis* and the microbial *Glomus* spp. + *B. megaterium* + *Trichoderma*
209 complex (Rizocore) were similar to the inoculated and untreated control. The *Brassica carinata*
210 pellets, applied once seven days before transplanting, were not effective (Figure 1a). The higher
211 fresh weight of lettuce plants obtained with the first protocol reflected the greatest efficacy in
212 disease reduction, which was provided by phosphite-based products, by acibenzolar-S-methyl at
213 the lower tested dosages, and by phosethyl-Al (Figure 1b).

214 In general, when an extra treatment was applied to the potted plants after transplanting, according
215 to protocol 2 (Figure 2a), all the tested biocontrol products differed significantly from the untreated
216 control, and an improved efficacy for *Bacillus subtilis* (42.7% efficacy), *T. asperellum* + *T. gamsii*
217 (38.3%, efficacy), and the microbial complex based *Glomus* spp complex (38.3%, efficacy) was
218 observed. On the other hand, the extra treatment with acibenzolar-S-methyl and the phosphite-
219 based products did not improve their efficacy. The application of *B. carinata* pellets led to a better
220 control of Fusarium wilt when they were applied 14 days before transplanting at T0, with an

221 average efficacy of 56% (Figure 2a). As far as the fresh weight of the lettuce plants at the end of
222 the trials is concerned, the best results, in terms of production, were observed in the plots in which
223 the best disease control was achieved (Figure 2b).

224

225 **Effect of the pre-planting treatment on rocket fusarium wilt**

226 Similar results, in terms of disease control, to those observed on lettuce, were observed in the trials
227 performed to evaluate the effects of different pre-planting treatments against *F. oxysporum* f. sp.
228 *raphani* on cultivated rocket, with an average disease severity at the end of the trials in the infested
229 and untreated control of 62.1 and 56.9, respectively (Figures 3a and 4a). When at least three
230 treatments were carried out, according to protocol 1, that is, acibenzolar-S-methyl at 0.0125 g L⁻¹,
231 the phosphite based products (Alexin and the Glucohumate complex), provided similar results to
232 one application of azoxystrobin, which reduced Fusarium wilt by 65%. All the tested biocontrol
233 agents were ineffective, and showed similar results to the inoculated and untreated control (Figure
234 3a). In general, there were no significant differences in plant fresh weight between treatments and
235 the inoculated and non-treated control plants, with the exception of the results shown for *Bacillus*
236 *subtilis*, *T. asperellum* + *T. gamsii* and the *Glomus* spp. + *Bacillus megaterium* + *Trichoderma*
237 product, in one out of four trials (Figure 3b).

238 The tested products, which were applied in four treatments (protocol 2), as previously described,
239 significantly reduced the Fusarium wilt of rocket compared to the infested and untreated control
240 (Figure 4a). The best rocket fusarium wilt control result was provided by potassium phosphite
241 (65%, efficacy), and this was followed by acibenzolar-S-methyl (57.5%, efficacy) and by the
242 Glucohumate complex (56.8%, efficacy), which were statistically similar to azoxystrobin, applied at
243 transplanting as a chemical reference, which showed 56.3% efficacy.

244 The *B. carinata* pellets also significantly reduced disease severity when they were applied 14 days
245 before transplanting (60.3%, efficacy). In general, the four treatments carried out with the

246 biocontrol agents *B. subtilis*, *B. velezensis*, *T. asperellum* +*T. gamsii*, and by the microbial
247 *Glomus* spp. + *B. megaterium* + *Trichoderma* complex, significantly reduced the disease, compared
248 to the untreated control (42.7%, 42.3%, 34.5%, 46.7% of efficacy, respectively).

249 The *Brassica carinata* pellets led to a positive effect on rocket fresh weigh, compared to the non-
250 inoculated and non-treated control, when applied 7-14 days before transplanting. The phosphite-
251 based products led to an improvement in fresh weight after four treatments were applied (Figures
252 3b and 4b).

253 None of the products tested under the present experimental conditions affected the fresh weight of
254 the lettuce or rocket plants, thus confirming the absence of phytotoxicity.

255

256 **4. Discussion**

257

258 According to several authors, Fusarium wilt is a devastating disease that affects many economically
259 important vegetables and it is responsible for severe losses to growers worldwide (Elmer, 2006;
260 Katan et al., 2012; Matheron and Gullino, 2012; Walters, 2012; Garibaldi et al., 2014).

261 The tactics adopted to control this plant disease mainly involve the use of preventative measures,
262 such as the minimization of dissemination in the field, the use of plant resistant cultivars whenever
263 possible, the adoption of a proper crop rotation, and the removal and destruction of infested plant
264 material. Like other soil-borne pathogens, Fusarium wilt is difficult to manage with a single
265 approach and/or with a single product (Katan et al., 2012). Moreover, different measures are needed
266 for different cropping systems. Gordon and Koike (2015) have reviewed different ways of
267 managing lettuce fusarium wilt and they have pointed out the importance of the adoption of proper
268 practices in order to reduce the inoculum level. Among these measures, it has been found that 1-
269 month of solarization, with an average temperature of 47 to 49°C, at a depth of 5 cm, provides a

270 reduction of 41 to 92% of Fusarium wilt on lettuce in naturally infested fields (Matheron and
271 Porchas, 2012). Even in sub-optimal temperature regimes (50 to 48°C for 6 h, 45 to 43°C for 8 h
272 and 40 to 38°C for 10 h/day), 14 days of thermal treatment under controlled conditions has provided
273 very valuable results, in terms of disease control on rocket (Gilardi et al., 2014). Moreover, recent
274 investigations on the antagonistic microorganisms *Trichoderma asperellum* and *T. gamsii*, with a
275 *Glomus* spp. +*Bacillus megaterium* +*Trichoderma* product used as a dressing on lettuce seeds, have
276 shown promising results in the control of fusarium wilt on lettuce (Lopez et al., 2014). On the
277 contrary, crop rotation is not an effective measure, because *F. oxysporum* f.sp. *lactucae* can
278 colonise the roots of other crops and, there is evidence of an expanded host range of *Fusarium*
279 *oxysporum* f.sp. *raphani* (Garibaldi et al., 2006; Scott et al., 2014).

280 Among the various alternatives to chemicals that have been tested so far, it has here been shown
281 that any solution on its own is able to provide a viable level of disease control of the Fusarium wilt
282 of lettuce and rocket, under short-cycles and for high levels of infestation. The success of the
283 biological control strategies used against Fusarium wilt agents, through the adoption of various
284 antagonist, organic amendments and biofumigation, in fact depends on many different factors,
285 including the pathogen infestation level in the soil and the type of inoculum (Termorshuizen and
286 Jeger, 2014). Soil temperature also has an important effect on the disease expression of both of
287 these Fusarium wilt agents, and it has been shown to be a key factor that can influence the success
288 of the control measures (Bosland et al. 1988; Scott et al., 2010a).

289 The present results provide evidence of the capability of commercial BCAs to reduce lettuce and
290 rocket Fusarium wilt by 42 to 47%, respectively, under high disease pressure, when applied four
291 times, starting from the nursery. The limited efficacy of lettuce and rocket Fusarium wilt control,
292 after three applications of *Glomus* spp. +*Bacillus megaterium* +*Trichoderma*, could be due to an
293 unsuccessful establishment of the mycorrhizal symbiosis. However, the mechanism involved in the
294 control of Fusarium wilt by *Glomus* is complex and seems to be host specific. In the present study,

295 four applications of the *Glomus* spp. +*Bacillus megaterium* +*Trichoderma* product provide a better
296 Fusarium wilt reduction on rocket with 47% of efficacy. Martinez-Medina et al., (2010) have
297 suggested that the mechanism involved in *Fusarium oxysporum* f.sp. *melonis* control by *Glomus*
298 *intraradices* is independent of the SA and JA pathways, while the Fusarium wilt control of tomato
299 by means of root colonization with mycorrhizae has been attributed to a possible plant-mediated
300 phenomenon (Kapoor, 2008).

301 The present study provides new information on the effect of resistance inducers, based on either
302 phosphites or acibenzolar-S-methyl, applied as a pre-plant treatment in the nursery, against *F.*
303 *oxysporum* f. sp. *lactucae* and *F. oxysporum* f. sp. *raphani* on lettuce and cultivated rocket,
304 respectively. Among the tested chemical resistance inducers, acibenzolar-S-methyl has been found
305 to be effective in controlling the Fusarium wilt of both lettuce and rocket at the lowest tested
306 dosage, with a positive effect on the yield. Encouraging results pertaining to the control of
307 Verticillium wilt on eggplants and Fusarium wilt on ornamental plants by means of acibenzolar-S-
308 methyl have already been reported by Elmer (2004; 2006) and by Bubici (2006).

309 The good fungicidal activity of the phosphite-based product, coupled with the positive effect on
310 plant biomass, is of special interest. In the present study, the phosphite-based products have
311 generally caused a consistent disease control of *F. oxysporum* f. sp. *lactucae*, that is, from 61 to
312 69%, and of *F. oxysporum* f. sp. *raphani*, from 54 to 65%, and have shown statistically similar
313 results to those of azoxystrobin, which has been used as a reference (59 to 65% efficacy), with a
314 significant effect on the yield. A previous study carried out on lettuce reported conflicting data on
315 plant growth for the use of phosphite-phosphate as a fertilizer. However, this may have been due to
316 the different cultivars that were examined, which could have different capabilities of absorbing
317 phosphite (Thao et al., 2009). The rate and timing of application of resistance inducers are
318 considered critical factors that can affect both the level of disease control and the yield (Walters,
319 2012).

320

321 The effectiveness of soil applications of the patented formulation of *Brassica carinata* defatted seed
322 meals has shown conflicting results against the Fusarium wilt of lettuce and rocket. In the present
323 study, *Brassica carinata* pellets have provided a consistent control, when applied 14 days before
324 transplanting the rocket and lettuce into the infested soil, with an efficacy that ranged from 56 to
325 60%. The present results are consistent with previous research, because organic amendments need
326 protracted periods of time to become effective, since their activity is due to decomposition and the
327 release of volatiles. (Mazzola et al., 2007; Bonanomi et al. 2010). The combination of *Brassica*
328 *carinata* pellets with solarization can be effective, as already observed, in infested soils, against the
329 Fusarium wilt of basil, lettuce and rocket (Garibaldi et al., 2011; Gilardi et al., 2014b).

330 Although the Fusarium wilt control provided by such products was not complete in the present
331 experimental conditions, it should be considered that disease severity is generally lower in the field.
332 Since the BCA treatments have not offered a complete Fusarium control, they could be integrated
333 with other strategies, and their contribution to disease management could be interesting, because
334 they can be used to complement other control measures. For instance, in the case of lettuce, the use
335 of resistant or partially resistant commercial cultivars (Garibaldi et al., 2004; Matheron et al., 2005;
336 Scott et al., 2010b; Gilardi et al., 2014b) could be combined with pre-plant treatments.

337 The four application of *Bacillus subtilis* Qs713 and the microbial *Glomus* spp. + *B. megaterium* +
338 *Trichoderma* complex, as well as the three phosphite-based commercial products, which may be
339 technically effective, especially at the nursery level, as well as economically feasible. The positive
340 effect of these products against leaf pathogens should also be considered. *Bacillus* spp., from the
341 rhizosphere of different vegetable plants, has resulted in induced resistance to *B. cinerea* (Levy et
342 al., 2015). In a preventative application, phosphite and acibenzolar-S-methyl induced resistance to
343 downy mildew on basil (Gilardi et al., 2013).

344 Such products are easy and safe to apply under nursery conditions, and could represent an effective
345 tool for the management of foliar and soil-borne pathogens. Moreover, the use of these products
346 before they are applied at a farm level could help to standardize their applications in an integrated
347 pest management model.

348

349 **Acknowledgements**

350 The present study has received funding from the European Union's Horizon 2020 research and
351 innovation programme EUCLID EU-CHINA Lever for IPM Demonstration. Grant agreement no
352 633999. The authors kindly thank Marguerite Jones for the language revision.

353

354 **References**

355

356 Barrière, V., Lecompte, F., Nicot, F.C., Maisonneuve, B., Tchamitchian, M., Lescourret, F., 2014.

357 Lettuce cropping with less pesticides. A review. *Agron. Sustain. Dev.* 34,175–198.

358 Bonanomi, G., Antignani, V., Capodilupo, M., Scala, F., 2010. Identifying the characteristics of

359 organic soil amendments that suppress soilborne plant diseases. *Soil Biol. Biochem.* 42,

360 136-144.

361 Bosland, P.-W., Williams, P.H., Morrison, R.H., 1988. Influence of soil temperature on the

362 expression of yellows and wilt of crucifers by *Fusarium oxysporum*. *Plant Dis.* 72, 777-780.

363 Bubici, G., Amenduni, M., Colella, C., D'Amico, M., Cirulli, M., 2006. Efficacy of acibenzolar-S-

364 methyl and two strobilurins, azoxystrobin and trifloxystrobin, for the control of corky root of

365 tomato and *Verticillium* wilt of eggplant. *Crop Prot.* 25, 814-820.

366 Colla, P., Gilardi, G., Gullino, M.L., 2012. A review and critical analysis of the European situation

367 of soilborne disease management in the vegetable sector. *Phytoparasitica* 40, 515-523.

368 Eikemo, H., A. Stensvand, A. M. Tronsmo, 2003. Induced resistance as a possible means to control
369 diseases of strawberry caused by *Phytophthora* sp.. *Plant Disease* 87, 345-350.

370 Elmer, W.H., 2004. Effect of acinenzolat-S-methyl on the suppression of *Fusarium* wilt of
371 cyclamen. *Crop Prot.* 25,671-676.

372 Elmer, WH., 2006. Efficacy of pre-plant treatment of gladiolus corns with combinations of
373 acibenzolar-S-methyl and biological pr chemical fungicides for suppression of *Fusarium*
374 corn rot (*Fusarium oxysporum* f. sp. *gladioli*). *Can. J. Plant. Pathol.* 28, 609-614.

375 Fujinaga, M., 2005. Studies on physiological races and phylogenetic analysis of lettuce root rot
376 pathogen, *Fusarium oxysporum* f. sp. *lactucae*. *J. Gen. Plant Pathol.* 71, 457.

377 Garibaldi, A., Gilardi, G., Gullino, M. L., 2002. First Report of *Fusarium oxysporum* on lettuce in
378 Europe. *Plant Disease*, 86 (9), 1052–1052.

379 Garibaldi, A., Gilardi, G., Gullino, M.L., 2004. Varietal resistance of lettuce to *Fusarium*
380 *oxysporum* f. sp. *lactucae*. *Crop Prot.* 23, 845-851.

381 Garibaldi, A., Gilardi, G., Gullino, M.L., 2006. Evidence for an expanded host range of *Fusarium*
382 *oxysporum* f. sp. *raphani*. *Phytoparasitica* 34, 115-121.

383 Garibaldi, A., Gilardi, G., Lazzeri, L., Malaguti, L., Clematis, F., Gullino, M.L., 2011. Effect of
384 green *Brassica* manure and *Brassica* defatted seed meals in combination with grafting and
385 soil solarization against *Verticillium* wilt of eggplant and *Fusarium* wilt of lettuce and basil.
386 *Acta Hortic.* 883, 295-302.

387 Garibaldi, A., Giulardi, G., Gullino, M.L., 2014. Critical aspects in disease management as a
388 consequence of the evolutionof soil-borne pathogens. *Acta Hortic.* 1044, 43-52.

389 Garibaldi, A., Gullino, M.L., 2010. Emerging soilborne diseases of horticultural crops and new
390 trends in their management. *Acta Hortic.* 883, 37-47.

391 Gilardi, G., Chen, G., Garibaldi, A., Zhiping, C., Gullino, M.L., 2007. Resistance of different rocket
392 cultivars to wilt caused by strains of *Fusarium oxysporum* under artificial inoculation
393 conditions. J. of Plant Pathol. 89,127-131.

394 Gilardi, G., Demarchi, S., Gullino, M.L., Garibaldi, A., 2013. Management of downy mildew of
395 sweet basil (*Ocimum basilicum*) caused by *Peronospora belbahrii* by means of resistance
396 inducers, fungicides, biocontrol agents and natural products. Phytoparasitica 41, 59-72.

397 Gilardi, G., Demarchi, S., Gullino, M.L., Garibaldi, A., 2014a. Managing Phytophthora crown and
398 root rot of tomato by pre-plant treatments with biocontrol agents, resistance inducers,
399 organic and mineral fertilize under nursery conditions. Phytopathol. Mediterr. 53,313-323.

400 Gilardi, G., Demarchi, S., Gullino, M.L., Garibaldi, A., 2014b. Varietal resistance to control
401 *Fusarium* wilts of leafy vegetables under greenhouse. Comm. Appl. Biol. Sci, 79(2), 21-27.

402 Gordon, T.R., Koike, S.T., 2015. Management of *Fusarium* wilt of lettuce. Crop Protection 73, 45-
403 49.

404 Gullino, M.L., Gilardi, G., Garibaldi, A., 2014. Seed-borne pathogens of leafy vegetable crops. In:
405 M.L. Gullino and G. Munkvold (eds.). Global perspectives on the health of seeds and plant
406 propagation material. Springer, Dordrecht, pp. 47-53.

407 Gullino, M.L., Minuto, A., Gilardi, G., Garibaldi, A., 2002. Efficacy of azoxystrobin and other
408 strobilurins against *Fusarium* wilts of carnation, cyclamen and Paris daisy. Crop Prot. 21,
409 57–61.

410 Hyeon, M.N., Park, M.S., Kim, H.G., Yoo, J.S., 2009. Biological control of strawberry *Fusarium*
411 wilt caused by *Fusarium oxysporum* f.sp. *fragariae* using *Bacillus velenzensis* BS87 and
412 RK1 formulation. J. Microbiol. Biotechnol. 19, 520-524.

413 Kapoor, R., 2008. Induced resistance in mycorrhizal tomato is correlated to concentration of
414 jasmonic acid. Online Journal of Biological Sciences 8, 49-56.

415 Katan, J., Gullino, M.L., Garibaldi, A., 2012. Strategies and tactics, in Fusarium wilt management.
416 In: Gullino ML, Katan J, Garibaldi A. (eds). Fusarium Wilts of Greenhouse Vegetable and
417 Ornamental Crops. St. Paul, MN, USA, APS Press, pp. 61-66.

418 Kloepper, J.W., Ryu, C.M., Zhang, S., 2004. Induced systemic resistance and promotion of plant
419 growth by *Bacillus* spp.. *Phytopathology* 94, 1259-1266.

420 Levey, N.O., Harel, M., Haile, Z.,M.,Elad., Y., Rav-David, E., Jurkevitch, E.,Katan., J., 2015.
421 Induced resistance to foliar diseases by soil solarization and *Trichoderma harzianum*. *Plant*
422 *Pathology* 64, 365-374.

423 Locke, T., Colhoun, J., 1974. Contribution to a method of testing oil palm seedlings for resistance
424 to *Fusarium oxysporum* f. sp. *elaedis* Toovey. *Phytopathologische Zeitschrift* 79, 77-92.

425 Lopez-Reyes, G.G., Gilardi, G., Garibaldi, A., Gullino, M.L., 2014. Efficacy of bacterial and fungal
426 biocontrol agents as seed treatments against *Fusarium oxysporum* f. sp. *lactucae* on lettuce.
427 *Journal of Plant Pathology* 96, 535-539.

428 Martínez-Medina, A., Pascual, J.A., Pérez-Alfocea, F., Albacete, A., Roldán, A., 2010.
429 *Trichoderma harzianum* and *Glomus intraradices* modify the hormone distribution induced
430 by *Fusarium oxysporum* infection in melon plants. *Phytopathology* 100, 682-688.

431 Matheron M.E., Porchas M., 2012. Evaluation of soil solarization and flooding as management
432 tools for Fusarium Wilt of lettuce. *Plant Dis.* 94, 1323-1328.

433 Matheron, M., Gullino, M.L., 2012. Fusarium wilt of lettuce and other salad crops. In: Gullino,
434 M.L., Katan, J., Garibaldi, A. (eds.), *Fusarium wilt of greenhouse vegetable and ornamental*
435 *crops*. St. Paul, MN, USA: APS Press.

436 Matheron, M.E., McCreight, J.D., Tickes, B.R., Porchas, M., 2005. Effect of planting date, cultivar,
437 and stage of plant development on incidence of Fusarium wilt of lettuce in desert production
438 fields. *Plant Dis.* 89, 565-570.

439 Mazzola, M., Brown, J., Izzo, A.D., Cohen, M.F., 2007. Mechanism of action and efficacy of seed
440 meal-induced pathogen suppression differ in a Brassicaceae species and time-dependent
441 manner. *Phytopathology* 97, 454–460.

442 Oostendorp, M., Kunz, W., Dietrich, B., Staub, T., 2001. Induced disease resistance in plants by
443 chemicals. *Eur. J. Plant Pathol.*, 107,19-28.

444 Pozo, M.J., Azcón-Aguilar C., 2007. Unraveling mycorrhiza-induced resistance. *Curr. Opin. Plant*
445 *Biol.* 10, 393-398.

446 Scott, J.C., Gordon, T.R., Shaw, D.V., Koike, S.T., 2010a. Effect of temperature on severity of
447 *Fusarium* wilt of lettuce caused by *Fusarium oxysporum* f. sp. *lactucae*. *Plant Dis.* 94, 13-
448 17.

449 Scott, J.C., Kirkpatrick, S.C., Gordon, T.R., 2010b. Variation in susceptibility of lettuce cultivars to
450 *Fusarium* wilt caused by *Fusarium oxysporum* f. sp. *lactucae*. *Plant Pathol* 59, 139-146.

451 Scott, J.C., McRoberts, D.N., Gordon, T.R., 2014. Colonization of lettuce cultivars and rotation
452 crops by *Fusarium oxysporum* f. sp. *lactucae*, the cause of *Fusarium* wilt of lettuce. *Plant*
453 *Pathol.* 63, 548-553.

454 Shores, M., Yedidia, I., Chet, I., 2005. Involvement of jasmonic acid/ ethylene signaling pathway
455 in the systemic resistance induced in cucumber by *Trichoderma*
456 *asperellum* T203. *Phytopathology* 95, 76–84.

457 Srinivasan, K., Spadaro, D., Poli, A., Gilardi, G., Gullino, M.L., Garibaldi, A., 2012. Genetic
458 diversity and pathogenicity of *Fusarium oxysporum* isolated from wilted rocket plants in
459 Italy. *Phytoparasitica* 40,157-170.

460 Sticher, L., Mauch-Mani, B., Métraux, J.P., 1997. Systemic acquired resistance. *Annu. Rev.*
461 *Phytopathol.* 35, 235-270.

462 Termorshuizen, A.J., Jeger, M.J. 2014. Assessing inoculum of soilborne plant pathogens: theory
463 and practice in decision-making for soil disinfestation. *Acta Horticulturae*, 1044,75-80.

464 Thao, B.T.H., Yamakawa, T., Shibata, K., 2009. Effect of phosphite-phosphate interaction on
465 growth and quality of hydroponic lettuce (*Lactuca sativa*). J. Plant Nutr. Soil Sci. 172, 378-
466 384.

467 Vallad, G. E., Goodman, R.M., 2004. Systemic acquired resistance and induced systemic resistance
468 in conventional agriculture. Crop Sci. 44, 1920-1934.

469 Vinale, F., Sivasithamparam, K., Ghisalberti, E.L., Marra, R., Woo, S.I., Lorito M., 2008.
470 *Trichoderma* plant pathogen interaction. Soil Biol. Biochem 40, 1-10.

471 Walters, D.R., 2012. Induced resistance as a management tool. In: Gullino, M.L., Katan, J.,
472 Garibaldi, A., (eds.), Fusarium wilt of greenhouse vegetable and ornamental crops. St. Paul,
473 MN, USA: APS Press.

474 Whipps, J.M., 2004. Prospect and limitation of mycorrhizas in biocontrol of root pathogens. Can
475 Journal Bot. 82, 1198-1227

476

477

478

479

480

481

482

483

484

485

486

487

488

489 Tables

490 Table 1. General information on the trials and timing of the operations carried out on lettuce and rocket.

Protocol, (trial)	Sowing	Tray treatment (Day)	Artificial inoculation	Transplanting	Plot Treatment	End of the trial
1(1)	6/04/2012	11/04 (T0) ^a ; 17/04 (T7); 24/04 (T14)	17/04//2012	25/04/2012	-	6/6/2012
1(2)	14/06/2012	21/06(T0); 28/06(T7); 5/07 (T14)	28/06/2012	6/07/2012	-	4/09/2012
1(3)	10/01/2013	21/01(T0); 28/01(T7); 4/02(T14)	28/01/2013	5/02/2013	-	18/03/2013
2 (4)	2/04/2013	10/04(T0); 17/04(T7); 24/04 (T14)	17/04/2013	25/04/2013	30/04/2013	28/05/2013
2 (5)	7/06/2013	14/06 (T0);21/06 (T7); 27/06 (T14)	21/06/2013	28/06/2013	4/07 /2013	23/07/2013
2 (6)	25/07/2013	1/08 (T0); 8/08 (T7); 12/08 (T14)	12/08/2/2013	13/08/2013	19/08/2013	17/09/2013
2 (7)	9/09/2013	20/09 (T0); 27/09 (T7); 3/10 (T14)	27/09/2013	4/10/2013	10/10/2013	28/10/2013

491 ^aT0 corresponding to the first treatment carried out as leaf spraying at the development stage of 1-2 true leaves and soil
 492 mixing application.

493

494 Table 2. List of the tested products and of the experimental protocol.

BCA or active ingredient	Commercial formulation	Dosage a. i. g L ⁻¹	Time (days) of application in tray conditions and type of application	Time (days) of application in plastic pot (2-L and 12-L) and type of application
<i>Bacillus subtilis</i> QST713	Serenade Max	0.58	T0 ^a ,T7,T14 (T21), leaf spray	-
<i>Bacillus velezensis</i>	Cilus Plus ^b	0.4 ^b	T0,T7,T14, (T21)leaf spray	-
<i>T. asperellum</i> + <i>T. gamsii</i>	Remedier	0.04	T0,T7,T14, (T21)leaf spray	-
Acibenzolar-S-methyl	Bion 50 WG	0.025 0.0125	T0,T7,T14, (T21) leaf spray	-
Phosethyl-Al	Aliette	1.6	T0,T7,T14, (T21)leaf spray	-
<i>Glomus</i> spp. + <i>Bacillus megaterium</i> + <i>Trichoderma</i>	Rizocore ^b	0.08 ^b	T0,T7,T14, (T21) leaf spray	-
Potassium phosphite P:K 52:42	Alexin	1.3+1.06	T0,T7,T14, (T21) leaf spray	-
Phosphite based glucohumate N:P 4:18	Glucohumate complex	1.6+0.72	T0,T7,T14, (T21)leaf spray	-
Azoxystrobin	Ortiva	0.19	T14, (T21) leaf spray	-
<i>Brassica carinata</i> pellet N:P:K: C organic	Biofence	0.15+0.055+0.0 5+1.13	-	T0 - T7, soil mixing ^c

495 ^aT0 corresponds to the leaf spraying carried out, starting at the second true leaf stage, and at seven-day intervals (T7
496 and T14).

497 ^b Corresponds to the dosage (g L⁻¹) of the commercial formulation.

498 ^c Corresponds to the treatment carried out at 14 day before transplanting (T0) and 7 days before transplanting the crops.

499

500

501

502

503


504

505

506

507 Figure 1a. Effect of three tray-treatments on the Fusarium wilt of lettuce (cv. Crispilla). The data
508 are expressed as DS at the end of the trials and compared with azoxystrobin and *Brassica carinata*
509 pellets applied once.

510




511

512

513

514

515 Figure 1b. Effect of three tray-treatments on the yield of lettuce plants in the presence of *F. oxysporum* f. sp.
 516 *lactucae*. The data are expressed as fresh weight and compared with azoxystrobin and *Brassica carinata*
 517 pellets applied once (g).



518

519

520

521

522

523


524

525

526


527

528 Figure 2a. Effect of four treatments on Fusarium wilt of lettuce (cv. Crispilla). The data are
 529 expressed as DS at the end of the trials and compared with azoxystrobin and *Brassica carinata*
 530 pellets applied at transplanting (T7) and 14 days before transplanting the lettuce (T0).
 531



532
 533
 534
 535
 536
 537
 538
 539

540 Figure 2b. Effect of four treatments on the yield of lettuce plants in the presence of *F. oxysporum* f. sp.
 541 *lactucae*. The data are expressed as fresh weight and compared with azoxystrobin and *Brassica carinata*
 542 pellets applied once (g).




543
 544
 545
 546
 547
 548
 549
 550
 551
 552
 553

554

555 Figure 3a. Effect of three tray-treatments against the Fusarium wilt of rocket (cv. Coltivata). The data are

556 expressed as DS at the end of the trials and compared with azoxystrobin and *Brassica carinata*

557 pellets applied at transplanting (T7).



558

559

560

561

562

563

564


565

566

567

568

569 Figure 3b. Effect of three treatments on the yield of rocket plants in the presence of *F. oxysporum* f. sp.
570 *raphani*. The data are expressed as fresh weight and compared with azoxystrobin and *Brassica carinata*
571 pellets applied once (g).



572

573

574

575


576

577

578

579

580 Figure 4a. Effect of four treatments on the Fusarium wilt of rocket. The data are expressed as DS at
 581 the end of the trials and compared with azoxystrobin and *Brassica carinata* pellets applied at
 582 transplanting (T7) and 14 days before transplanting rocket, at the same time as the artificial
 583 infestation (T0).



584

585

586

587

588

589

590


591

592

593

594 Figure 4b. Effect of four treatments on the yield of rocket plants in the presence of *F. oxysporum* f. sp.
 595 *raphani*. The data are expressed as fresh weight and compared with azoxystrobin and *Brassica carinata*
 596 pellets applied once (g).

597



598

599

600