

Manuscript Number:

Title: Short chain chito-oligosaccharides promote arbuscular mycorrhizal colonization in *Medicago truncatula*

Article Type: Research Paper

Keywords: chito-oligosaccharides; arbuscular mycorrhiza; sustainable agriculture; *Medicago truncatula*; *Funnelliformis mosseae*; symbiosis

Corresponding Author: Professor Andrea Genre,

Corresponding Author's Institution:

First Author: Veronica Volpe, PhD

Order of Authors: Veronica Volpe, PhD; Gennaro Carotenuto, PhD; Carlotta Berzero; Lavinia Cagnina; Virginie Puech-Pagès; Andrea Genre

Abstract: During the establishment of arbuscular mycorrhizal (AM) symbiosis, the fungus and the host plant exchange chemical signals that are crucial to reciprocal recognition. Short-chain chitin oligomers (CO) released by AM fungi are known to trigger symbiotic signaling in all host plant species tested. Here we applied exogenous CO, derived from crustacean exoskeleton, to pot-grown *Medicago truncatula* inoculated with the AM fungus *Funnelliformis mosseae* and investigated root colonization, plant gene regulation and biomass production. CO treatment strongly promoted AM colonization with significant increases in arbuscule development, biomass production and photosynthetic surface compared to untreated mycorrhizal plants. Gene expression analyses indicated that CO treatment anticipated the expression of MtBCP and MtPT4 plant symbiotic markers, during the first two weeks post inoculation. Altogether, our results provide evidence that plant treatment with symbiotic fungal elicitors, anticipated and enhanced AM development, encouraging the use of CO to promote AM establishment in sustainable agricultural practices.

Suggested Reviewers: Cristina Prandi Prof
Full Professor, Chemistry, University of Turin
cristina.prandi@unito.it
Expert in chemical characterization of symbiotic signaling molecules

Marco Giovannetti PhD
Gregor Mendel Institute of Molecular Plant Biology, Austrian Academy of Sciences
marco.giovannetti@gmi.oeaw.ac.at
Expert in plant molecular responses to short-chain chito-oligosaccharides

Thomas Ott Prof
Full professor, Faculty of Biology, University of Freiburg
thomas.ott@biologie.uni-freiburg.de
Expert in symbiotic plant-microbe signaling

Torino, 30.7.2019

Dear Editor,

We are submitting the manuscript entitled «*Short chain chito-oligosaccharides promote arbuscular mycorrhizal colonization in *Medicago truncatula**» to be considered for publication in Carbohydrate Polymers.

Over 80% of plants, including most crops, live in a mutualistic symbiotic association, called arbuscular mycorrhiza (AM), with an ancient group of soil fungi that provide their hosts with soil water and mineral nutrients in change for sugars and lipids. For this key role in plant nutrition, the introduction of AM as biofertilizers in **sustainable agricultural practices** has been proposed for decades as a promising natural solution to the increasing worldwide demand of food safety. Nevertheless, field applications tests have delivered contrasting results, and experimental data indicate a partial recalcitrance of cultivated species toward AM interaction, likely due to decades of excessive use of fertilizers and to farming practices that have been unaware of the plant underground interactions.

We now know that plant-fungus recognition is an essential step in AM establishment and is based on an exchange of chemical signals: root-exudates alerts the AM fungus of host proximity, while AM fungi release **short-chain chito-oligosaccharides** (CO) eliciting the plant symbiotic responses, such as gene regulation, starch reallocation and cell rearrangement for fungal accommodation.

In the light of this role for CO as AM fungal signals, we decided to **test whether their exogenous application could have a positive impact on symbiosis establishment**. This would on the one hand demonstrate unambiguously the biological role of CO as symbiosis elicitors and, on the other hand, open the way to their use as promoters of AM colonization in crops for sustainable, food safety-oriented agricultural applications.


To this aim, we treated the model legume *Medicago truncatula* with a mixture of short-chain CO (ranging between 2 and 5 N-acetyl-glucosamine residues) and observed how their application influenced plant mycorrhizal status and development.

Indeed, our analyses of root colonization intensity, plant gene expression and biomass development consistently showed that **CO treatments strongly enhance and anticipate AM colonization**. This represents a **breakthrough** on the way to the large-scale introduction of AM in agricultural practices, by positively acting on the mycorrhizal aptitude of the host plants.

For these reasons we consider that the findings in this article are of primary interest in the field of carbohydrate polymer applications to plant-microbe interactions, and appropriate for timely publication in your Journal.

Yours sincerely

Andrea Genre



- Arbuscular mycorrhizal (AM) symbiosis supports life of most crop plants.
- The fungus is recognized as a symbiont via water-soluble short-chain chitin oligomers (CO)
- We tested whether exogenous CO application to *Medicago truncatula* could impact on the symbiosis.
- CO treatment strongly promoted AM colonization with significant increases in arbuscule development, biomass production and photosynthetic surface compared to untreated mycorrhizal plants.
- Plant treatment with symbiotic fungal elicitors anticipated and enhanced AM development, encouraging the use of CO to promote AM establishment in sustainable agricultural practices

1 **Short chain chito-oligosaccharides promote arbuscular mycorrhizal colonization in *Medicago***
2 ***truncatula***

3

4 **Authors**

5 **Veronica Volpe^a, Gennaro Carotenuto^a, Carlotta Berzero^a, Lavinia Cagnina^a, Virginie Puech-**
6 **Pagès^b, Andrea Genre^a**

7 ^a*Department of Life Science and Systems Biology, University of Turin, Viale Mattioli 25, 10125*
8 *Torino, Italy*

9 ^b*Laboratoire de Recherche en Sciences Végétales, Université de Toulouse, CNRS, UPS, Castanet-*
10 *Tolosan, France*

11

12 **Abstract**

13 During the establishment of arbuscular mycorrhizal (AM) symbiosis, the fungus and the host plant
14 exchange chemical signals that are crucial to reciprocal recognition. Short-chain chitin oligomers
15 (CO) released by AM fungi are known to trigger symbiotic signaling in all host plant species tested.
16 Here we applied exogenous CO, derived from crustacean exoskeleton, to pot-grown *Medicago*
17 *truncatula* inoculated with the AM fungus *Funneliformis mosseae* and investigated root colonization,
18 plant gene regulation and biomass production. CO treatment strongly promoted AM colonization with
19 significant increases in arbuscule development, biomass production and photosynthetic surface
20 compared to untreated mycorrhizal plants. Gene expression analyses indicated that CO treatment
21 anticipated the expression of *MtBCP* and *MtPT4* plant symbiotic markers, during the first two weeks
22 post inoculation. Altogether, our results provide evidence that plant treatment with symbiotic fungal
23 elicitors, anticipated and enhanced AM development, encouraging the use of CO to promote AM
24 establishment in sustainable agricultural practices.

25

26 **1. Introduction**

27 The life of most plants, including the majority of crops, is supported by a mutualistic root symbiosis
28 with Glomeromycotina, an ancient group of soil fungi (Spatafora et al., 2016) that provide their hosts
29 with a preferential access to soil water and mineral nutrients, while taking advantage of plant-
30 photosynthesized sugars and lipids (Smith & Read, 2008; Wewer et al. 2014; Keymer et al., 2017).

31 Plant-fungus recognition is essential for AM establishment and is based on an exchange of chemical
32 signals (Bonfante & Requena, 2011; Zipfel & Oldroyd, 2017). Root-exuded strigolactone alerts the
33 AM fungus of host proximity (Akiyama et al., 2005); its perception activates spore germination,
34 hyphal metabolism and branching (Besserer et al., 2006). In turn, AM fungi release diffusible
35 molecules eliciting the plant symbiotic response (Maillet et al, 2011; Genre et al., 2013). These
36 include local and systemic changes in gene expression (Kosuta et al., 2003), root-directed starch
37 reallocation (Gutjahr et al., 2009), and the activation of a signal transduction pathway, partially shared
38 with other symbiotic interactions (Oldroyd, 2013; Delaux et al., 2015; Barker et al., 2017; Genre and
39 Russo, 2016). This pathway includes the triggering of repeated oscillations (spiking) in nuclear
40 calcium concentration, making the detection of nuclear Ca²⁺ spiking an acknowledged benchmark
41 for the induction of symbiotic responses in AM hosts.

42 Indeed, Ca²⁺ spiking is observed in root epidermal nuclei upon adhesion of AM fungal hyphopodia
43 (Chabaud et al., 2011), application of AM fungal exudate (Navazio et al, 2007; Chabaud et al., 2011;
44 Genre et al., 2013) or purified molecules isolated from fungal exudate, such as short-chain chito-
45 oligosaccharides (CO) and lipo-chito-oligosaccharides (LCO) (Maillet et al., 2011; Genre et al., 2013).
46 In particular, fungal release of tetrameric and pentameric CO is boosted upon strigolactone treatment
47 (Genre et al., 2013). Furthermore, their activity has been demonstrated in legumes (Genre et al., 2013),
48 carrot (Genre et al., 2008), rice (Sun et al., 2015; Carotenuto et al., 2017), *Discaria trinervis* and
49 *Casuarina glauca* (Chabaud et al., in press).

50 In the light of this universal role for CO in eliciting early plant responses to AM fungi (Genre &

51 Russo, 2016), we decided to test whether their exogenous application could have a positive impact
52 on symbiosis establishment. This would on the one hand demonstrate the biological role of CO as
53 symbiosis elicitors and, on the other hand, open the way to their use as promoters of AM colonization
54 in crops for sustainable, food safety-oriented agricultural applications (Berruti et al., 2015).

55 To this aim, we treated the model legume *Medicago truncatula* with a mixture of short CO (ranging
56 below 6 N-acetyl-glucosamine residues), obtained from crustacean manufacturing industry, and
57 observed how their application influenced plant development and mycorrhizal status.

58

59 **2. Materials and Methods**

60 *2.1. Plant growth and mycorrhization.*

61 Seeds of *M. truncatula* (cv. Jemalong, genotype A17) were scarified for 3 min in H₂SO₄, rinsed
62 several times in distilled water, sterilized for 2 min in diluted commercial bleach (1:5), rinsed 4 times
63 in sterile distilled water and pre-germinated for 48 hours on agar plates (0.6% w/v) at 23° C in the
64 dark. Seedlings were grown for ten days in a photoperiod of 16h day (23°C) and 8h night (21°C)
65 before transferring them to 10x10x12cm pots containing quartz sand.

66 The AM fungus *Funneliformis mosseae* (strain BEG 12) was inoculated in mycorrhizal pots by
67 replacing 10% of the pot substrate with a commercial inoculum (MycAgroLab, Bretenière, France)
68 and mixing. All plants were watered once a week with half-strength Long-Ashton nutrient solution
69 containing 3,2 µM KH₂PO₄ as P source (Hewitt, 1966).

70

71 *2.2. Chito-oligosaccharides*

72 We used a commercial mixture of short chain CO purified from crustacean manufacturing side-
73 products (Zhengzhou Sigma Chemical Co., Ltd., Henan, China). *HPLC-MS/MS* analysis of the CO
74 mixture (File S1) confirmed the presence of deacetylated, mono-acetylated and di-acetylated CO
75 molecules composed of 2 to 5 N-acetyl-glucosamine residues (Fig. S1).

76

77 2.3. Chito-oligosaccharide solutions

78 1g/L and 1mg/L CO solutions in sterile distilled water were tested for the triggering of Ca²⁺ spiking
79 in the root epidermis. Only the 1g/L solution was chosen for pot treatments, to counteract molecule
80 dilution/absorption and compensate for the presence of biologically inactive CO₂ and CO₃ in the mix,
81 alongside the AM signals CO₄ and CO₅ (Fig. S1). CO application to pot-grown plants was done
82 either by irrigating the substrate with the CO solution, or by spraying the solution on plants and soil
83 surface.

84

85 2.4. Analysis of CO-induced Ca²⁺ signals

86 Confocal microscopy was performed on 2cm-long lateral roots of *Agrobacterium rhizogenes*-
87 generated root organ cultures (ROC) of *M. truncatula* expressing NUP-YC2.1, a nuclear localized
88 Ca²⁺-sensing yellow cameleon construct (Chabaud et al., 2011). Detection and plotting of relative
89 changes in Ca²⁺ concentration in epidermal atrichoblasts was done according to Genre et al. (2013),
90 as described in File S1. At least 100 atrichoblasts from seven independent root samples were analyzed
91 for each condition and the t-Student test with a probability level of $P < 0.05$ was used for statistical
92 validation.

93

94 2.5. Plant treatment with CO

95 In order to evaluate CO effect on plant development and mycorrhization, four experimental
96 conditions were set up: control (Ctr), lacking both CO treatment and AM inoculation; CO-treated
97 control (Ctr+CO); mycorrhizal (Myc), where AM inoculum was added in the absence of CO
98 treatments; and CO-treated mycorrhizal (Myc+CO), where plants were both inoculated and exposed
99 to the CO solution.

100 For CO application by irrigation, residual water was carefully removed from the pot saucers before

101 watering each Ctr+CO and Myc+CO pot with 30 mL of CO solution. Distilled water was used for Ctr
102 and Myc pots. Treatments were applied weekly, and plants were sampled 48 days post inoculation
103 (dpi), after six CO treatments (Fig. S2), for morphometric analyses and mycorrhizal intensity
104 determination.

105 For spray treatments, 5 mL of CO solution containing 0.005% Tween20 as a surfactant, was sprayed
106 weekly over Ctr+CO and Myc+CO pots from a 10 cm distance, using plastic shields to confine spray
107 diffusion. Ctr and Myc pots were sprayed with 5 mL of 0.005% Tween20 in sterile distilled water.
108 Plants treated with CO spray were divided in two groups (Fig. S2): one was grown for 28 dpi (with
109 two CO treatments) and the other for 48 dpi (with six CO treatments) before measuring their
110 morphological and physiological parameters and quantifying AM colonization.

111

112 2.6. *Analyses of plant development after CO treatment*

113 Irrigation-treated plants were harvested at 48 dpi; root and shoot fresh weight was immediately
114 measured. To determine the leaf area, green tissues were carefully isolated and placed on an Epson
115 Perfection V800 Photo scanner, to obtain black and white scanned images of the leaf surface at 600
116 dpi resolution. The resulting PNG images were then analyzed using Fiji/ImageJ
117 (<http://imagej.nih.gov/>; <http://fiji.sc/Fiji>) to calculate total leaf surface in mm². Following image
118 acquisition, shoot dry weight was measured after exsiccation at 60°C for 24h.

119 Spray-treated plant growth was monitored (Fig. S3) by photography at 7 dpi (prior to CO treatment),
120 28 dpi (two treatments) and 48 dpi (six treatments). Five Ctr, five Ctr+CO, ten Myc and ten Myc+CO
121 plants were harvested at 28 dpi and 48 dpi, to measure their fresh and dry weight and photosynthetic
122 surface, as described above.

123

124 2.7. *Nitrogen balance index*

125 The physiological status of spray-treated plants was estimated before sampling using a Dualex 4

126 (FORCE-A, Orsay, France) to determine flavonoid and chlorophyll content by measuring leaf UV
127 absorbance and fluorescence, respectively, in 10 randomly chosen leaves from 5 Ctr and 5 Ctr+CO
128 plants, and 20 leaves from 10 Myc and 10 Myc+CO plants. Measures were done on both adaxial and
129 abaxial leaf surface and the two values were averaged (Cartelat et al., 2005). The
130 chlorophyll/flavonoid ratio, or Nitrogen Balance Index (NBI), is considered an indicator of the plant
131 nitrogen nutritional status (Cerovic et al., 2012).

132

133 *2.8. Quantification of mycorrhizal colonization*

134 Following fresh weight measure, each root system was stained with 0.1% cotton blue in lactic acid
135 and quantitative parameters of mycorrhizal intensity were calculated as described by Trouvelot et al.
136 (1986), using the online tool MYCOCALC (<http://www2.dijon.inra.fr/mychintec/Mycocalc-prg/download.html>).

138

139 *2.9. Gene expression analysis following CO treatment*

140 To investigate the ability of CO treatment to stimulate host receptiveness for AM fungi, a time-
141 course experiment was set up where plants received two rounds of spray treatment before AM
142 inoculation, 21 and 23 days post transplant (dpt). The inoculum was added at 25 dpt, and plants were
143 sampled at 7, 10, 14, 21 and 28 dpi. Five leaf replicates per treatment were sampled to analyse
144 phosphorus (P) content and five root replicates from Myc and Myc+CO were sampled and frozen in
145 liquid nitrogen for gene expression analyses.

146

147 *2.10. RNA isolation and quantitative RT-PCR analysis*

148 Total RNA was isolated from roots, using the RNeasy™ Plant Mini kit (Qiagen, Hilden, Germany).
149 Samples were disrupted with a TissueLyser (Qiagen Retsch GmbH, Hannover, Germany) and treated
150 following manufacturer instructions. RNA quality and quantity were evaluated using a Nanodrop and

151 agarose gel electrophoresis.

152 To remove genomic DNA contamination, RNA samples were digested with TURBO™ DNase
153 (Ambion) according to manufacturer instructions and checked for DNA traces by conventional PCR
154 using *MtTEF* primers, before cDNA synthesis and qRT-PCR analyses (File S1). The sequences of all
155 genes studied are listed in Table S1.

156

157 *2.11. Assessment of arbuscule morphology*

158 Root colonized segments from Myc and Myc+CO plants were embedded in agarose (8%) and cut
159 into 200 µm-thick sections using a Vibratome (Oxford Vibratome®). Sections were collected on
160 microscope slides and stained with fluorescein isothiocyanate-conjugated wheat germ agglutinin
161 (WGA-FITC; Sigma-Aldrich, Milan, Italy), as described in Volpe et al. (2016). Twenty root segments
162 from five biological replicates were analyzed for each treatment.

163

164 *2.12. Phosphorus quantification*

165 Phosphorus quantification was done in 5 Ctr, 5 Ctr+CO, 10 Myc and 10 Myc+CO plants, by
166 collecting the third leaf from the tip of the main stem. Collected leaves were lyophilized, weighed
167 and digested in 1 mL of 6M HNO₃ for one hour at 95°C. The digestion product was diluted in 6 mL
168 of sterile distilled water. A control solution, without any sample, was treated with the same procedure.
169 All solutions were analysed by Inductively Coupled Plasma Atomic Emission Spectrometry (ICP-
170 AES) to determine P content.

171

172 *2.13. GenBank accession numbers*

173 AY116211 (*MtPT4*); XM_003592405.2 (*MtBCP*); XM_013595882.1 (*MtTEF*); XM_013606824
174 (*MtPRI0*); XM_003604044.2 (*MtChit1*); XM_003618769.2 (*MtChitIII.1*); XM_003597718.2
175 (*MtChitIII.2*); AY238969.2 (*MtChitIII.3*); AY238970.1 (*MtChitIII.4*)

176

177 3. Results

178 3.1. Crustacean-derived CO triggered Ca^{2+} spiking in *M. truncatula* atrichoblasts

179 In order to assess the commercial CO effectiveness as symbiotic elicitors, we recorded nuclear Ca^{2+}
180 signals in atrichoblasts of NUP-YC2.1 *M. truncatula* ROCs treated with 1g/L and 1mg/L CO solutions
181 (Fig. 1). Both solutions triggered comparable spiking patterns, with frequent, well-defined peaks, in
182 analogy with those induced by analogous concentrations of purified CO4, CO5 or raw AM spore
183 exudate (Chabaud et al., 2011; Genre et al., 2013). As expected, no spiking was recorded upon control
184 treatment with sterile distilled water. The percentage of responding atrichoblasts and average peak
185 numbers confirmed the lack of statistically significant differences in the cell response to either CO
186 concentration (Fig. 1B).

187 In conclusion, our bioassay confirmed that both 1g/L and 1mg/L solutions of the commercial CO
188 mix was able to mimic AM fungal signals in triggering Ca^{2+} -mediated symbiotic signaling.




Fig. 1. Nuclear Ca^{2+} spiking in response to crustacean-derived CO. A) Representative plots of nuclear Ca^{2+} variations over 30 minutes in atrichoblast nuclei of *M. truncatula* root organ culters (ROCs) treated with water (control), 1mg/L and 1g/L CO. Both CO concentrations induced a series of Ca^{2+} peaks (spiking), characteristic of AM fungal recognition. B) Statistical analyses of the spiking response (t-Student test with $P < 0.05$) did not highlight significant differences between CO treatments, in terms of either the percentage of responding atrichoblasts (top panel) or average peak number in responding cells (bottom). Bars represent standard errors. At least 100 atrichoblasts from seven independent root samples were analyzed for each experimental condition.

189

190 *3.2. CO application by irrigation boosted AM colonization*

191 In a first attempt to investigate the effect of exogenous CO supply, we grew inoculated and non-
192 inoculated *M. truncatula* in pots, with or without a weekly supplement of CO in the watering solution,
193 and analyzed their biomass, photosynthetic surface and mycorrhizal colonization at 48 dpi. While
194 root development was comparable in Ctr, Myc and Myc+CO plants, statistically significant
195 differences were observed in the shoot fresh and dry biomass between Ctr and both mycorrhizal
196 conditions, Myc and Myc+CO (Fig. S4). The same trend was confirmed for photosynthetic surface,
197 with similar top values for Myc and Myc+CO plants and a significantly smaller area for Ctr and
198 Ctr+CO plants. Remarkably, Ctr+CO plants showed a significant reduction in biomass and
199 photosynthetic surface, when compared to all other treatments, suggesting that CO perception in the
200 absence of fungal inoculation could induce a stress condition inhibiting plant growth.

201 AM colonization was then assessed in Myc and Myc+CO plants through optical microscopy.
202 Isolated groups of arbuscules were distributed along the root axis of Myc plants, spaced by
203 uncolonized cortical cells (Fig. S5). By contrast, Myc+CO plants displayed a more extensive and
204 homogeneous distribution of arbuscules, with a continuous mass of arbusculated cells, which also
205 involved multiple layers of the root cortex (Fig. S5). This was confirmed by our quantitative analysis,
206 with significantly higher frequency (F), intensity of mycorrhization (M) and arbuscule abundance (a,
207 A) in Myc+CO compared to Myc plants (Fig. S5).

208 In conclusion, beside a partial growth inhibition in uninoculated plants, exogenous CO application
209 by irrigation strongly improved the mycorrhizal status of AM inoculated plants at 48 dpi.

210


211 *3.3. CO spray treatment enhanced AM colonization*

212 These encouraging results, coupled with the possible induction of a stress response in Ctr+CO,
213 prompted us to test an alternative treatment. We chose spray application as an amenable method of
214 CO supply in view of possible field treatments. For these experiments, we sampled our plants at 28

215 and 48 dpi to obtain a clearer picture of symbiosis development.

216 Our first focus was to check whether the spray treatment could be as efficient in promoting AM
217 development as the irrigation method. Indeed, F, M and A values were all significantly higher in
218 Myc+CO compared to Myc plants at 28 dpi (Fig. 2). Also in this case, CO treatment caused arbuscule
219 proliferation across multiple cortical layers, compared to the fewer arbuscules, with spotted
220 distribution, found in Myc plants. At 48 dpi, all mycorrhization parameters surged in both conditions,
221 reducing the difference between Myc and Myc+CO plants, which anyway remained significant (Fig.
222 2).

223 In short, the quantification of AM colonization based on microscopic observations confirmed a
224 positive effect on mycorrhizal intensity also for CO spray treatment.



225 **Fig. 2.** Effect of spray CO treatment on mycorrhizal plants at 28 dpi (A) and 48 dpi (B). Images of root sections of mycorrhizal plants
226 untreated (Myc) and treated with CO (Myc+CO). A strong proliferation of arbuscules (white dots) was evident in Myc+CO plants, with the
227 colonization of multiple cortical layers at both time points, as evident in the magnified areas (red square) in the lower panels. This was
confirmed by the quantitative analysis of root colonization, shown in the bottom graphs: mycorrhization frequency (F), mycorrhization
intensity (M), arbuscule abundance within colonized root fragments (a) and arbuscule abundance in the whole root system (A) all reported
significantly higher values in Myc+CO compared to Myc plants. The effect of CO on root colonization was anyway more evident at 28 dpi.
Mean values and SDs of ten biological replicates for each treatment are shown. Bars = 100 μ m.

228 3.4. CO spray treatment and AM inoculation synergistically increase plant growth

229 Photographic monitoring of shoot growth showed that prior to any CO treatment - at 7 dpi - all
 230 plants had a comparable development (Fig. S3). By contrast, significant differences became evident
 231 at 28 dpi (two CO treatments) and even more pronounced at 48 dpi (six CO treatments): at both time
 232 points, Myc and Myc+CO plants were larger than Ctr plants. Interestingly, CO application by spray
 233 to uninoculated plants (Ctr+CO) did not affect their growth at any time point (Fig. S3), in contrast
 234 with the results of CO treatment by irrigation.

235 Such macroscopical observations were partially confirmed by biometric measurements. At 28 dpi
 236 (Fig. 3A), the photosynthetic surface was significantly more extensive in mycorrhizal treatments
 237 compared to both Ctr and Ctr+CO, with the highest value for Myc+CO plants, albeit not significantly
 238 larger than in Myc plants. An analogous trend was recorded for shoot dry biomass, with the highest
 239 values for the Myc and Myc + CO plants, even if differences were not statistically significant.




Fig. 3. Aboveground development in response to spray CO treatment. A) Average photosynthetic surface (including representative scans of the plant epigeral parts) and shoot dry biomass at 28 dpi: the photosynthetic surface was significantly more developed in Myc and Myc+CO plants compared to Ctr and Ctr+CO; a similar – albeit statistically non significant - trend was observed for shoot dry weight. B) The photosynthetic surface was significantly more development in Myc+CO plants at 48 dpi, when both mycorrhizal plants showed a significant increase of shoot dry biomass. Mean values and SDs are presented for five and ten biological replicates from control and mycorrhizal plants, respectively.

240

241 At 48 dpi (Fig. 3B), both mycorrhizal conditions (Myc and Myc+CO) showed a significantly larger
 242 photosynthetic surface than both Ctr and Ctr+CO, but in addition, Myc+CO significantly outclassed
 243 Myc plants at this later time point. An analogous significant increase was observed in shoot dry
 244 biomass between mycorrhizal and uninoculated plants, albeit Myc and Myc+CO had very similar
 245 values.

246 In conclusion, two CO spray treatments were sufficient to induce a significant increase in AM
 247 colonization at 28 dpi, while a CO-dependent increase in plant development was evident at 48 dpi,
 248 after six CO treatments.




Fig. 4. Leaf nitrogen (N) status at 28 and 48 dpi. The chlorophyll/flavonoid ratio (Chl/Flav), or Nitrogen Balance Index (NBI), is presented here as an indicator of leaf N content. NBI value was significantly higher in both mycorrhizal lines (Myc and Myc+CO) than controls (Ctr and Ctr+CO) at 28 dpi. By contrast, NBI was significantly higher in all CO treated plants at 48 dpi. Mean values and SDs are presented for five and ten biological replicates from control and mycorrhizal plants, respectively.

249

250

251 3.5. Effect of CO spray on plant nitrogen status

252 We measured the nitrogen balance index (NBI) to indirectly monitor plant nitrogen status and
 253 overall stress. At 28 dpi, the NBI was significantly higher for both mycorrhizal conditions, indicating
 254 a better metabolic performance compared to Ctr and Ctr+CO plants (Fig. 4). No significant difference
 255 was observed between the latter, confirming that CO spray alone did not cause significant stress in
 256 uninoculated plants. By contrast, at 48 dpi, the NBI was significantly higher in CO-treated plants
 257 (Ctr+CO and Myc+CO) compared to Ctr and Myc.


258 Altogether, NBI analysis indicated a comparable increase in leaf nitrogen content for both

259 mycorrhizal conditions at 28 dpi, and a further increase in Myc+CO at 48 dpi, in line with CO effects
260 on plant development.

261
262 *3.6. Plant AM marker genes are regulated in response to CO treatment*

263 Since CO supply had effectively enhanced root colonization at 28 and 48 dpi, we decided to
264 investigate in more detail the effect of CO with a time-course experiment that included earlier time
265 points. In addition, CO ability to stimulate host plant receptiveness for AM fungi was assessed by
266 spraying plants twice with the CO, before fungal inoculation (Fig. S2). We consequently analyzed
267 gene expression for two established AM markers: *MtBCP* (*Blue Copper-binding Protein*), expressed
268 during arbuscule development, and *MtPT4* (*Phosphate Transporter 4*), encoding an AM-specific P
269 transporter localized on the periarbuscular membrane of active, mature arbuscules (Pimprikar &
270 Gutjahr, 2018).

271 *MtBCP* expression was more strongly induced in Myc+CO plants at 7, 10 and 14 dpi, compared to
272 a later activation in Myc plants (Fig. 5), with a significant difference at 7 dpi. A drastic
273 downregulation was recorded at 21 and 28 dpi, in line with the expected reduction of *MtBCP*
274 expression in terminal stages of root colonization. Similarly, *MtPT4* was also activated earlier in
275 Myc+CO plants, but its expression surprisingly decreased at 21 dpi, apparently anticipating the
276 analogous decrease observed in Myc plants at 28 dpi.



277 **Fig. 5.** Time-course analysis of *MtBCP* and *MtPT4* expression. Asterisks indicate the significant early induction of *MtBCP* at 7
278 dpi and the significant downregulation of both genes at 21 and 28 dpi in CO-treated plants (Myc+CO). Average values from at
least 5 biological replicates are presented.

278 To better investigate this unexpected difference in AM marker gene expression, we analyzed the

279 regulation of three AM-induced chitinases (Salzer et al, 2000; Kremer et al., 2013; Malolepszy et al,
 280 2018) in fully developed mycorrhizal roots, at 28 dpi: *MtChitIII.2*, *MtChitIII.3* and *MtChitIII.4*. In
 281 our hands, only *MtChitIII.2* and *MtChitIII.4* were upregulated in inoculated plants compared to
 282 controls (Fig. 6), but both chitinases were less expressed in Myc+CO compared to Myc plants. In
 283 conclusion, two additional markers of arbuscule maturity displayed the same downregulation as
 284 *MtPT4* in Myc+CO plants.




Fig. 6. Molecular analysis on roots at 28 dpi after two CO treatments. Two of the three AM-induced chitinases are strong induced in the mycorrhizal plants, but the transcripts levels for *MtChitIII.2* and *MtChitIII.4* are significant reduced in the Myc+CO compared to the Myc untreated plants, in line with the expression profile of *MtPT4*. Mean values and SDs in three biological replicates for each treatment are shown.

285
 286 On this basis, we wondered whether this pattern of gene regulation could be indicative of anticipated
 287 arbuscule degeneration in CO-treated mycorrhizal plants - an aspect that could not be highlighted by
 288 our quantitative analyses of mycorrhizal colonization. We therefore compared arbuscule morphology
 289 in Myc and Myc+CO plants (Fig. S6). No obvious difference was observed in arbuscule shape and
 290 general organisation: the expected arbuscule morphology, with a large trunk and a mass of fine
 291 branches occupying most of the host cell volume, was consistently recorded for both conditions,
 292 suggesting that the vast majority of arbuscules was fully developed, independent of CO application.

293 Altogether, CO treatment on the one hand accelerated root colonization by the AM fungus,
 294 providing a convincing demonstration of CO role as elicitors of the host plant symbiotic response.
 295 Exogenous CO application led to an anticipated expression of early AM markers such as *MtBCP* and
 296 the development of a higher number of arbuscules that appeared functional, based on their
 297 morphological features. On the other hand, CO treatment reduced the late expression of AM marker
 298 genes that are normally expressed in active arbusculated cells (Harrison et al., 2002; Elfstrand et al.,


299 2005; Pumplin & Harrison, 2009). This unexpected result prompted us to compare the nutritional
300 status of Myc and Myc+CO plants.

301

302 3.7. Leaf phosphorus content was not affected by CO treatment

303 We compared P content in leaf samples from both mycorrhizal conditions. Surprisingly, no
304 significant difference was detected for any time point: plants reached their highest P content at 14 dpi
305 under both conditions (Fig. 7), with a progressive decrease in the following time points.

306 In conclusion, the increase in arbuscule abundance in Myc+CO plants was not reflected by an
307 improvement in plant P uptake under our experimental conditions.




308 **Fig. 7.** Phosphorous (P) content level in untreated (Myc) and treated (Myc+ CO) mycorrhizal plants. Both lines
reached the highest leaf P content at 14 dpi but no significant difference was detected in any time point. Mean
values for five biological replicates are shown. Bars represent standard errors.

309

309 3.8. CO spray does not upregulate plant defense markers

310 To have a more comprehensive view of CO effects on the plant, we also checked whether CO
311 perception could activate plant defense responses, by analyzing the expression of the pathogenesis
312 related gene *MtPR10* and two pathogenesis-related chitinases, *MtChitI* and *MtChitIII.1* (Salzer et al.,
313 2000; Lipka et al., 2005). None of these defense markers was induced by CO treatment (Fig. 8), in
314 line with literature data about the scarce activity of short chain CO as defense elicitors (Shibuya &
315 Minami, 2001). Moreover, the significant reduction of *MtChitI* expression in Ctr+CO, Myc and
316 Myc+CO plants compared to Ctr, suggests a possible partial inhibition of defense responses in the

317 presence of either fungal or externally supplied (or both) signals.



318 **Fig. 8.** Expression of pathogenesis-related genes in roots at 28 dpi after two CO treatments. The three analysed pathogenesis-induced genes, *MtChitI*, *MtChitIII.1* and *MtPR10*, were not upregulated upon treatment with crustacean-derived CO, suggesting that they were not perceived as a pathogenesis signal. Mean values and SDs in three biological replicates for each treatment are shown.

318

319 4. Discussion

320 In spite of several consistent reports of CO role as elicitors of early AM signaling and a few
321 investigations of CO-regulated gene expression (Weidmann et al, 2004; Giovannetti et al, 2015; Sun
322 et al., 2015), surprisingly few studies have investigated the effect of CO on symbiosis development
323 (Ramírez et al., 2010). In the light of our results, CO now appear particularly promising for large
324 scale agricultural applications for several reasons: among currently characterized AM fungal signals,
325 CO can be obtained at a fraction of the cost of the more complex LCO (lipo-chito-oligosaccharides,
326 similar to rhizobial Nod factors), from waste products of crustacean fishing industry. As natural and
327 easily bio-degradable products, CO are considered environmentally friendly and the use of chitin-
328 derived molecules has a relatively long history in agriculture, for their stimulation of plant growth
329 and defense (Benhamou et al., 1998; Sharp, 2003; Zhang et al., 2016; Winkler et al., 2017).

330

331 4.1. Exogenous CO application promotes AM development

332 The most striking result of this study was the observation that CO treatment induced a remarkable
333 increase in arbuscule abundance. This evidence represents a crucial advance in support of the role of
334 CO as positive elicitors of symbiotic responses in the host plant, demonstrating their effect - beyond
335 early signaling and gene expression – on the development of arbuscules, the functional core of AM
336 symbiosis.

337 In addition, our results highlighted that spraying the plants with CO prior to AM fungal inoculation
338 anticipated the upregulation of *MtBCP* and *MtPT4* at 7 and 10 dpi, suggesting the ability of CO to
339 accelerate AM establishment: a very promising aspect in view of their application under agricultural
340 conditions. Interestingly, CO did not anticipate symbiosis senescence, as suggested by the arbuscule
341 regular morphology in both treated and untreated samples. The fact that the longer time-span of AM
342 colonization in CO-treated plants did not correlate with a higher P content could appear surprising.
343 Nevertheless, unchanged P contents are also reported in studies on hypermycorrhizal mutants of *M.*
344 *truncatula* (Truong et al., 2014) and *Pisum sativum* (Jones et al., 2015), where arbuscule proliferation
345 appears analogous to what we have observed upon CO treatment. A few hypotheses can be made to
346 explain this, each of which deserves further investigation. Firstly, P uptake might not be the main
347 function of the morphologically active arbuscules that we have observed at 28 dpi. Under our standard
348 growth conditions, fewer arbuscules - as in untreated mycorrhizal plants - might be sufficient to grant
349 *M. truncatula* an optimal P nutrition. The plant might instead take advantage of such an abundant
350 number of arbuscules for other functions, such as water uptake. Challenging CO-treated and untreated
351 plants with different levels of drought will shed light on this possibility, but our observation of a
352 larger biomass in CO-treated mycorrhizal plants at 48 dpi - when the root system had extended to the
353 whole pot volume – hints at a better performance of this experimental line under incipient stress
354 conditions.

355 Secondly, prolonged CO elicitation could stimulate sustained fungal accommodation, while
356 arbuscule functioning and senescence could be regulated by different - and yet unexplored - signaling
357 processes. Lastly, one cannot exclude that such an abundant root colonization in a mature stage of the
358 symbiosis is favoring the AM fungus absorption of plant-derived organic compounds, with CO
359 elicitation possibly weakening the plant control mechanisms on over-colonization. Under this respect,
360 it will be extremely important to investigate C flux toward the fungus, as well as extraradical
361 mycelium development or spore production.

362

363 *4.2. Plant responses to CO treatments*

364 The use of different chitin-derived molecules in agricultural practices or laboratory conditions is
365 not new. Benefits to the plant have been described in terms of better growth and resistance to biotic
366 or abiotic stresses. Nevertheless, a clear picture of their multiple effects on plants is far from being
367 understood and most agricultural applications rely on the heuristic observation of a correlation
368 between chitin-based treatments and beneficial effects. In this frame, our results provide a possible
369 explanation to such positive effects on crops: if such mixes of compounds include short-chain CO,
370 then part of the benefits to the plant may derive from its improved symbiotic status, which in turn is
371 well known to support plant development, defense and stress resistance.

372 In the field of plant-microbe interactions, the length of CO molecules is acknowledged as a major
373 determinant of plant responses, with the following - likely simplistic - scenario: chito-biose and -
374 triose are reported to produce little or no stimulation in plant cells; -tetraose and -pentaose have been
375 characterized as symbiotic signals in all AM host plants (Genre et al., 2013; Giovannetti et al, 2015;
376 Sun et al., 2015); whereas longer CO progressively shift toward defense elicitation (Liu et al., 2012;
377 Hayafune et al., 2014), with chito-octamers being the best characterized chitin-based pathogen-
378 associated molecular pattern (Zhang et al., 2002; Cao et al., 2017). In this frame, CO have been
379 proposed as primers of plant defenses in agriculture, even if limited attention has been put on the
380 oligomer composition of the mixture. The picture is even less clear about the effects of CO on plant
381 development. A stimulating effect of chito-pentaose was demonstrated on Arabidopsis growth (Khan
382 et al., 2011). Similarly, Winkler and colleagues (2017) demonstrated that a mixture of short-chain
383 CO, increased Arabidopsis and poplar shoot explant development, and chito-tetraose treatment of
384 Arabidopsis seedlings upregulated development, cell organization, biogenesis and transport-related
385 genes. By contrast, no stimulating effect on plant growth was observed for CO by Chatelain et al.
386 (2014), who rather reported a decrease in *Phaseolus vulgaris* biomass. Zhang et al (2016) indicated

387 CO length as a critical determinant of their effect also on plant development, with chito-hexaose and
388 -heptaose inducing the strongest elicitation of biomass development in wheat.

389 This complex scenario is difficult to compare with our experimental system, based on CO treatment
390 of mycorrhizal plants. Nevertheless, a few parallels may be drawn. The reduction of uninoculated
391 plant development observed in CO irrigated plants, appears in line with the inhibition of plant growth
392 described by Zhang et al. (2016), since we were using a mix of CO that did not include the growth-
393 promoting hexa- and heptamers. By contrast, a synergistic positive effect of CO and inoculation on
394 plant development was observed near the end of our experimental time frame. In this case, fungal
395 presence is introducing an additional level of complexity to the experimental system, complicating
396 the interpretation of direct and combined effects of CO, symbiotic status and stress conditions
397 possibly induced by the space limitation of the pot culture.

398

399 **5. Conclusion**

400 The introduction of AM as biofertilizers in sustainable agricultural practices has been proposed for
401 decades as a promising natural solution to the increasing worldwide demand of food safety. Results
402 have anyway been contrasting, with two major obstacles emerging. First, even if AM fungi have low
403 host specificity, studies quickly highlighted that not all fungal inocula are suitable for all agricultural
404 applications: local climate, soil characteristics, as well as the species (if not the variety) of the selected
405 crop generate a matrix of variables that only customized inocula could address, with the time- and
406 resource-consuming selection of performing fungal strains for each condition. Second, a partial
407 recalcitrance has been observed in cultivated species toward AM interaction, likely due to decades of
408 excessive use of fertilizers (Duhamel & Vandenkoornhuyse, 2013) and to farming practices that have
409 always been attentive to the aerial part of the plant but unaware of its underground interplay.

410 Our study showed that CO treatments strongly enhance AM colonization, anticipating the
411 upregulation of AM marker genes, but not arbuscule senescence. Even if several aspects of CO effects

412 on plants remain to be explored, our demonstration of their positive impact on AM establishment
413 represents a breakthrough on the way to the large scale introduction of AM in agricultural practices,
414 by positively acting on the mycorrhizal aptitude of the host plants.

415

416 **Acknowledgements**

417 This research was funded by Fondazione Cassa di Risparmio di Cuneo (Bando Ricerca Scientifica
418 2015 - Project AMforQuality). Mass spectrometry analyses were supported by the ICT-Mass
419 spectrometry and MetaToul facilities and the MetaboHUB-ANR-11-INBS-0010 network.

420

421 **References**

422 Akiyama, K., Matsuzaki, K., & Hayashi, H. (2005). Plant sesquiterpenes induce hyphal branching in
423 arbuscular mycorrhizal fungi. *Nature*, 435, 824–827.

424 Barker, D.G., Chabaud, M., Russo, G., & Genre, A. (2017). Nuclear Ca²⁺ signalling in arbuscular
425 mycorrhizal and actinorhizal endosymbioses: on the trail of novel underground signals. *New*
426 *Phytologist*, 214, 533-538.

427 Benhamou, N., Kloepper, J.W., & Tuzun S. (1998). Induction of resistance against *Fusarium* wilt of
428 tomato by combination of chitosan with an endophytic bacterial strain: ultrastructure and
429 cytochemistry of the host response. *Planta*, 204, 153-168.

430 Berruti, A., Lumini, E., Balestrini, R., & Bianciotto, V. (2015). Arbuscular Mycorrhizal Fungi as
431 Natural Biofertilizers: Let's Benefit from Past Successes. *Frontiers in Microbiology*, 6, 1559.

432 Besserer, A., Puech-Pages, V., Kiefer P., et al. (2006). Strigolactones stimulate arbuscular
433 mycorrhizal fungi by activating mitochondria. *PLoS Biology*, 4, 1239–1247.

434 Bonfante, P., & Requena, N. (2011). Dating in the dark: how roots respond to fungal signals to
435 establish arbuscular mycorrhizal symbiosis. *Current Opinion in Plant Biology*, 14, 451-7.

436 Carotenuto, G., Chabaud, M., Miyata, K., Capozzi, M., Takeda, N., Kaku, H., et al (2017). The rice

437 LysM receptor-like kinase OsCERK1 is required for the perception of short-chain chitin oligomers
438 in arbuscular mycorrhizal signaling. *New Phytologist*, 214, 1440-1446.

439 Cartelat, A., Cerovic, Z.G., Goulas, Y., Meyer, S., Lelarge, C., Prioul J.L., et al. (2005). Optically
440 assessed contents of leaf polyphenolics and chlorophyll as indicators of nitrogen deficiency in
441 wheat (*Triticum aestivum* L.). *Field Crops Research*, 91, 35-49.

442 Cerovic, Z.G., Masdoumier, G., Ghazlen, N.B., & Latouche, G. (2012). A new optical leaf-clip meter
443 for simultaneous non-destructive assessment of leaf chlorophyll and epidermal flavonoids.
444 *Physiologia Plantarum*, 146(3), 251-60.

445 Chabaud, M., Genre, A., Sieberer, B.J., Faccio, A., Fournier, J., Novero, M., et al. (2011). Arbuscular
446 mycorrhizal hyphopodia and germinated spore exudates trigger Ca²⁺ spiking in the legume and
447 nonlegume root epidermis. *New Phytologist*, 189, 347-355.

448 Chatelain, P.G., Pintado, M.E., & Vasconcelos, M.W. (2014). Evaluation of chitoooligosaccharide
449 application on mineral accumulation and plant growth in *Phaseolus vulgaris*. *Plant Science*, 215-
450 216, 135-140.

451 Cao, Y., Halane, M.K., Gassmann, W., & Stacey, G. (2017). The role of plant innate immunity in the
452 legume-rhizobium symbiosis. *Annual Review of Plant Biology*, 68, 535-561.

453 Delaux, P.M., Radhakrishnan, G.V., Jayaraman, D., Cheema, J., Malbreil, M., Volkening, J.D.,
454 Sekimoto, H., Nishiyama, T., Melkonian, M., Pokorny, L. et al. (2015). Algal ancestor of land plants
455 was preadapted for symbiosis. *Proceedings of the National Academy of Sciences, USA*, 112, 13390-
456 13395.

457 Duhamel, M., & Vandenkoornhuyse, P. (2013). Sustainable agriculture: possible trajectories from
458 mutualistic symbiosis and plant neodomestication. *Trends in Plant Science*, 18, 597-600.

459 Elfstrand, M., Feddermann, N., Ineichen, K., Nagaraj, V.J., Wiemken, A., Boller, T., & Salzer, P.
460 (2005). Ectopic expression of the mycorrhiza-specific chitinase gene *Mtchit 3-3* in *Medicago*
461 *truncatula* root-organ cultures stimulates spore germination of glomalean fungi. *New Phytologist*,

462 167, 557-570.

463 Genre, A., Chabaud, M., Faccio, A., Barker, D.G., & Bonfante, P. (2008). Prepenetration apparatus
464 assembly precedes and predicts the colonization patterns of arbuscular mycorrhizal fungi within the
465 root cortex of both *Medicago truncatula* and *Daucus carota*. *Plant Cell*, 20, 1407-1420.

466 Genre, A., Chabaud M., Belzergue, C., Puech-Pages, V., Novero, M., Rey, T., Fournier, J., Rochange,
467 S., Becard, G., Bonfante, P., & Barker, D. (2013). Short-chain chitin oligomers from arbuscular
468 mycorrhizal fungi trigger nuclear Ca²⁺ spiking in *Medicago truncatula* roots and their production is
469 enhanced by strigolactone. *New Phytologist*, 198, 190-202.

470 Genre, A., & Russo, G. (2016). Does a Common Pathway Transduce Symbiotic Signals in Plant–
471 Microbe Interactions? *Frontiers in Plant Science*, 7, 96.

472 Giovannetti, M., Mari, A., Novero, M. & Bonfante, P. (2015) Early *Lotus japonicus* root
473 transcriptomic responses to symbiotic and pathogenic fungal exudates. *Frontiers in Plant Science*,
474 6, 480.

475 Gutjahr, C., Novero, M., Guether, M., Montanari, O., Udvardi, M., & Bonfante, P. (2009).
476 Presymbiotic factors released by the arbuscular mycorrhizal fungus *Gigaspora margarita* induce
477 starch accumulation in *Lotus japonicus* roots. *New Phytologist*, 183(1), 53-61.

478 Harrison, M.J., Dewbre, G.R., & Liu, J. (2002) A phosphate transporter from *Medicago truncatula*
479 involved in the acquisition of phosphate released by arbuscular mycorrhizal fungi. *The Plant Cell*,
480 14, 2413-2429.

481 Hayafune, M., Berisio, R., Marchetti, R., Silipo, A., Kayama, M., Desaki, Y., et al. (2014). Chitin-
482 induced activation of immune signaling by the rice receptor CEBiP relies on a unique sandwich-
483 type dimerization. *Proceedings of the National Academy of Sciences, USA*, 111, E404-E413.

484 Hewitt, E.J. (1966). Sand and water culture methods used in the study of plant nutrition. *Farnham*
485 *Royal, UK: Common wealth Agriculture Bureaux*

486 Jones, J.M.C., Clairmont, L., Macdonald, E.S., Weiner, C.A., Emery, N.R.J., & Guinel, F.C. (2015).

487 E151 (*sym15*), a pleiotropic mutant of pea (*Pisum sativum* L.), displays low nodule number,
488 enhanced mycorrhizae, delayed lateral root emergence, and high root cytokinin levels. *Journal of*
489 *Experimental Botany*, 66, 4047-4059.

490 Lipka, V., Dittgen, J., Bednarek, P., Bhat, R., Wiermer, M., Stein, M., Landtag, J., et al. (2005). Pre-
491 and postinvasion defenses both contribute to nonhost resistance in *Arabidopsis*. *Science*, 310, 1180-
492 1183.

493 Liu, T., Liu, Z., Song, C., Hu, Y., Han, Z., She, J., Fan, F., Wang, J., Jin, C., Chang, J., et al. (2012).
494 Chitin-induced dimerization activates a plant immune receptor. *Science*, 336, 1160-1164.

495 CKeymer A, Pimprikar P, Wewer V et al. (2017). Lipid transfer from plants to arbuscular mycorrhizal
496 fungi. *eLife*: 6, e29107.

497 Khan, W., Costa, C., Souleimanov, A., Prithiviraj, B., & Smith D.L. (2011). Response of *Arabidopsis*
498 *thaliana* roots to lipo-chitooligosaccharide from *Bradyrhizobium japonicum* and other chitin-like
499 compounds. *Plant Growth Regulation*, 63, 243.

500 Kosuta, S., Chabaud, M., Loughon, G., Gough, C., Denarie, J., Barker, D.G., & Becard, G. (2003). A
501 diffusible factor from arbuscular mycorrhizal fungi induces symbiosis-specific *MtENOD11*
502 expression in roots of *Medicago truncatula*. *Plant Physiology*, 131, 952-962.

503 Kosuta, S., Hazledine, S., Sun, J., Miwa, H., Morris, R.J., Downie, J.A., & Oldroyd, G.E. (2008)
504 Differential and chaotic calcium signatures in the symbiosis signaling pathway of legumes.
505 *Proceedings of the National Academy of Sciences, USA*, 105, 9823-9828.

506 Kremer, N., Philipp E.E., Carpentier, M.C., Brennan, C.A., Kraemer, L., Altura, M.A., et al. (2013).
507 Initial symbiont contact orchestrates host-organ-wide transcriptional changes that prime tissue
508 colonization. *Cell Host Microbe*, 14, 183-194.

509 Maillet, F., Poinot, V., André, O., Puech-Pagès, V., Haouy, A., Gueunier, M., et al. (2011). Fungal
510 lipochitooligosaccharide symbiotic signals in arbuscular mycorrhiza. *Nature*, 469, 58-63.

511 Malolepszy, A., Kelly, S., Sørensen, K.K., James, E.K., Kalisch, C., Bozsoki, Z., et al. (2018). A

512 plant chitinase controls cortical infection thread progression and nitrogen-fixing symbiosis. *eLife*,
513 7, e38874

514 Miyawaki, A., Llopis, J., Heim, R., McCaffery, J.M., Adams, J.A., Ikura, M., & Tsien, R.Y. (1997).
515 Fluorescent indicators for Ca²⁺ based on green fluorescent proteins and calmodulin. *Nature*, 388,
516 882-887.

517 Navazio, L., Moscatiello, R., Genre, A., Novero, M., Baldan, B., Bonfante, P., & Mariani, P. (2007).
518 A diffusible signal from arbuscular mycorrhizal fungi elicits a transient cytosolic calcium elevation
519 in host plant cells. *Plant Physiology*, 144, 673-681.

520 Oldroyd, G.E.D. (2013). Speak, friend, and enter: signalling systems that promote beneficial
521 symbiotic associations in plants. *Nature Reviews Microbiology*, 11, 252-263.

522 Pimprikar, P., & Gutjahr, C. (2018). Transcriptional regulation of arbuscular mycorrhiza
523 development. *Plant Cell Physiology*, 0, 1-18

524 Pumplin, N., & Harrison, M.J. (2009) Live-cell imaging reveals periarbuscular membrane domains
525 and organelle location in *Medicago truncatula* roots during arbuscular mycorrhizal symbiosis. *Plant*
526 *Physiology*, 151, 809-819.

527 Ramírez, M.A., Rodríguez, A.T., Alfonso, L., & Peniche, C. (2010). Chitin and its derivatives as
528 biopolymers with potential agricultural applications. *Biotechnología Aplicada*, 27, 270-276.

529 Rasmussen, R. (2001). Quantification on the LightCycler instrument. In: Meuer, S., Wittwer, C.,
530 Nakagawara, K., eds. *Rapid cycle real-time PCR: methods and applications*. Heidelberg, Germany,
531 Springer-Verlag, 21–34.

532 Salzer, P., Bonanomi, A., Beyer, K., Vögeli-Lange, R., Aeschbacher, Roger A., & Lange, J. (2000)
533 Differential Expression of Eight Chitinase Genes in *Medicago truncatula* Roots During Mycorrhiza
534 Formation, Nodulation, and Pathogen Infection. *Molecular Plant-Microbe Interaction*, 13, 763-777.

535 Sharp Russell, G. (2003). A Review of the Applications of Chitin and Its Derivatives in Agriculture
536 to Modify Plant-Microbial Interactions and Improve Crop Yields. *Agronomy*, 3, 757-793

537 Shibuya, N., & Minami, E. (2001) Oligosaccharide signaling for defence responses in plants. *Plant*
538 *Cell Physiology*, 59, 223-233.

539 Smith, S.E., & Read, D.J. (2008). Mycorrhizal symbiosis. *Academic Press, Inc., San Diego, CA*

540 Spatafora, J.W., Benny, G.L., Berbee, M., Bonito, G., Corradi, N., et al. (2016). A phylum-level
541 phylogenetic classification of zygomycete fungi based on genome-scale data. *Mycologia*, 108,
542 1028-1046.

543 Sun, J., Miller, J.B., Granqvist, E., Wiley-Kalil, A., Gobbato, E., Maillet, F., et al. (2015). Activation
544 of Symbiosis Signaling by Arbuscular Mycorrhizal Fungi in Legumes and Rice. *The Plant Cell*, 27,
545 823-838.

546 Truong, H.N., Thalineau, E., Bonneau, L., Fournier, C., Potin, S., Balzergue, S., van Tuinen, D.,
547 Jeandroz, S., & Morandi, D. (2014). The *Medicago truncatula* hypermycorrhizal B9 mutant
548 displays an altered response to phosphate and is more susceptible to *Aphanomyces euteiches*. *Plant,*
549 *Cell & Environment*, 38, 73-88.

550 Volpe, V., Giovannetti, M., Sun, X.G., Fiorilli, V., & Bonfante, P. (2016). The phosphate transporters
551 *LjPT4* and *MtPT4* mediate early root responses to phosphate status in non mycorrhizal roots. *Plant,*
552 *Cell & Environment*, 39, 660-671.

553 Weidmann, S., Sanchez, L., Descombin, J., Chatagnier, O., Gianinazzi, S., & Gianinazzi-Pearson, V.
554 (2004). Fungal elicitation of signal transduction-related plant genes precedes mycorrhiza
555 establishment and requires the *dmi3* gene in *Medicago truncatula*. *Molecular Plant Microbe*
556 *Interaction*, 17, 1385-1393.

557 Wewer, V., Brands, M., & Dörmann, P. (2014). Fatty acid synthesis and lipid metabolism in the
558 obligate biotrophic fungus *Rhizophagus irregularis* during mycorrhization of *Lotus japonicus*.
559 *Plant Journal*, 79, 398-412.

560 Winkler, A.J. Dominguez-Nuñez, J.A., Aranaz, I., Poza-Carrión, C., Ramonell, K., Somerville, S., &
561 Berrocal-Lobo, M. (2017). Short-Chain Chitin Oligomers: Promoters of Plant Growth. *Marine*

562 *Drugs*, 15, 40.

563 Zhang, B., Ramonell, K., Somerville, S., & Stacey, G. (2002). Characterization of early, chitin-
564 induced gene expression in *Arabidopsis*. *Molecular Plant Microbe Interaction*, 15, 963-70.

565 Zhang, X., Li, K., Liu, S., Xing, R., Yu, H., Chen, X., & Li, P. (2016). Size effects of chitooligomers
566 on the growth and photosynthetic characteristics of wheat seedlings. *Carbohydrate Polymers*, 138,
567 27-33.

568 Zipfel, C., & Oldroyd, G.E.D. (2017) Plant signalling in symbiosis and immunity. *Nature*, 543, 328.

569

570

571 **Figure legends.**

572 **Fig. 1.** Nuclear Ca^{2+} spiking in response to crustacean-derived CO. A) Representative plots of nuclear
573 Ca^{2+} variations over 30 minutes in atrichoblast nuclei of *M. truncatula* root organ culters (ROCs)
574 treated with water (control), 1mg/L and 1g/L CO. Both CO concentrations induced a series of Ca^{2+}
575 peaks (spiking), characteristic of AM fungal recognition. B) Statistical analyses of the spiking
576 response (t-Student test with $P < 0.05$) did not highlight significant differences between CO
577 treatments, in terms of either the percentage of responding atrichoblasts (top panel) or average peak
578 number in responding cells (bottom). Bars represent standard errors. At least 100 atrichoblasts from
579 seven independent root samples were analyzed for each experimental condition.

580

581 **Fig. 2.** Effect of spray CO treatment on mycorrhizal plants at 28 dpi (A) and 48 dpi (B). Images of
582 root sections of mycorrhizal plants untreated (Myc) and treated with CO (Myc+CO). A strong
583 proliferation of arbuscules (white dots) was evident in Myc+CO plants, with the colonization of
584 multiple cortical layers at both time points, as evident in the magnified areas (red square) in the lower
585 panels. This was confirmed by the quantitative analysis of root colonization, shown in the bottom
586 graphs: mycorrhization frequency (F), mycorrhization intensity (M), arbuscule abundance within

587 colonized root fragments (a) and arbuscule abundance in the whole root system (A) all reported
588 significantly higher values in Myc+CO compared to Myc plants. The effect of CO on root
589 colonization was anyway more evident at 28 dpi. Mean values and SDs of ten biological replicates
590 for each treatment are shown. Bars = 100 μ m.

591

592 **Fig. 3.** Aboveground development in response to spray CO treatment. A) Average photosynthetic
593 surface (including representative scans of the plant epigeral parts) and shoot dry biomass at 28 dpi:
594 the photosynthetic surface was significantly more developed in Myc and Myc+CO plants compared
595 to Ctr and Ctr+CO; a similar – albeit statistically non significant - trend was observed for shoot dry
596 weight. B) The photosynthetic surface was significantly more development in Myc+CO plants at 48
597 dpi, when both mycorrhizal plants showed a significant increase of shoot dry biomass. Mean values
598 and SDs are presented for five and ten biological replicates from control and mycorrhizal plants,
599 respectively.

600

601 **Fig. 4.** Leaf nitrogen (N) status at 28 and 48 dpi. The chlorophyll/flavonoid ratio (Chl/Flav), or
602 Nitrogen Balance Index (NBI), is presented here as an indicator of leaf N content. NBI value was
603 significantly higher in both mycorrhizal lines (Myc and Myc+CO) than controls (Ctr and Ctr+CO) at
604 28 dpi. By contrast, NBI was significantly higher in all CO treated plants at 48 dpi. Mean values and
605 SDs are presented for five and ten biological replicates from control and mycorrhizal plants,
606 respectively.

607

608 **Fig. 5.** Time-course analysis of *MtBCP* and *MtPT4* expression. Asterisks indicate the significant early
609 induction of *MtBCP* at 7 dpi and the significant downregulation of both genes at 21 and 28 dpi in
610 CO-treated plants (Myc+CO). Average values from at least 5 biological replicates are presented.

611

612 **Fig. 6.** Molecular analysis on roots at 28 dpi after two CO treatments. Two of the three AM-induced
613 chitinases are strong induced in the mycorrhizal plants, but the transcripts levels for *MtChitIII.2* and
614 *MtChitIII.4* are significant reduced in the Myc+CO compared to the Myc untreated plants, in line
615 with the expression profile of *MtPT4*. Mean values and SDs in three biological replicates for each
616 treatment are shown.

617

618 **Fig. 7.** Phosphorous (P) content level in untreated (Myc) and treated (Myc+CO) mycorrhizal plants.
619 Both lines reached the highest leaf P content at 14 dpi but no significant difference was detected in
620 any time point. Mean values for five biological replicates are shown. Bars represent standard errors.

621

622 **Fig. 8.** Expression of pathogenesis-related genes in roots at 28 dpi after two CO treatments. The three
623 analysed pathogenesis-induced genes, *MtChitI*, *MtChitIII.1* and *MtPRI0*, were not upregulated upon
624 treatment with crustacean-derived CO, suggesting that they were not perceived as a pathogenesis
625 signal. Mean values and SDs in three biological replicates for each treatment are shown.

626

627 **Supplementary data**

628 **Fig. S1.** Composition of crustacean-derived CO identified by HPLC-MS/MS. CO molecules were
629 detected with a chain length ranging between 2 and 5 residues and different degrees of deacetylation.

630 A) Since CO detection threshold raises with increasing chain length, only the presence (+) or absence
631 (-) of each type of molecule is presented here. NS = not searched. B) Structure of mono-deacetylated
632 (beta anomer) CO4 and theoretical precursor/product ions forming in LC-MS/MS the positive mode.
633 C) LC-MS/MS chromatogram in the MRM mode (M+H⁺). The two peaks correspond to alpha and
634 beta anomers of the molecule.

635

636 **Fig. S2.** Schemes representing the experimental set-up for CO treatment and sampling used for this

637 work. The camera symbol indicates photographic recording of plant development; dpi, days post
638 inoculation; dpt, days post transplant.

639

640 **Fig. S3.** Monitoring of plant growth during CO spray treatment. The growth of control (Ctr), CO-
641 treated control (Ctr+CO), mycorrhizal (Myc) and CO-treated mycorrhizal plants (Myc+CO) was
642 monitored at 7, 28 and 48 dpi. AM inoculation alone increased plant growth at both 28 and 48 dpi,
643 but aerial part development at both time points was further enhanced when AM inoculation was
644 combined with CO application (Myc+CO) at 14, 21, 28, 34, 40 and 46 dpi as summarized in Fig. S2.
645

646 **Fig. S4.** Effect of CO treatment by irrigation on plant development at 48 dpi. Shoot and root biomass
647 production was monitored in control (Ctr), CO-treated control (Ctr+CO), mycorrhizal (Myc) and CO-
648 treated mycorrhizal plants (Myc+CO). A reduced development of both shoots and roots was observed
649 in Ctr+CO compared to Ctr plants. While root biomass was not affected by AM inoculation, Myc and
650 Myc+CO plants displayed an increased shoot biomass production compared to Ctr, with consistent
651 results for fresh weight, dry weight and photosynthetic surface. Mean values and SDs of ten biological
652 replicates for each treatment are shown.

653

654 **Fig. S5.** Effect of CO treatment by irrigation on mycorrhizal development at 48 dpi. Images of root
655 sections from untreated (Myc) and CO-treated mycorrhizal plants (Myc+CO). Arbuscules
656 proliferation was strongly stimulated in Myc+CO plants, with the involvement of multiple cortical
657 layers, compared to a more limited colonization, surrounding the vascular cylinder (vc), in Myc plants.
658 This observation was confirmed by the quantitative analysis of root colonization, reported in the
659 bottom graph: mycorrhization frequency (F), mycorrhization intensity (M), arbuscule abundance
660 within colonized root fragments (a) and arbuscule abundance in the whole root system (A) all reported
661 significantly higher values in Myc+CO compared to Myc plants. Mean values and SDs of ten

662 biological replicates for each treatment are shown. Bars = 200 μm

663

664 **Fig. S6.** Arbuscule phenotype in Myc and Myc + CO plants. In both cases, fully developed and highly
665 branched arbuscules were observed at 28 dpi. Twenty root colonized sections from five biological
666 replicates were used for each treatment. Bars = 30 μm .

667

668 **Table S1.** Primers used for qRT-PCR analysis

Figure 1
[Click here to download high resolution image](#)




Figure 2
[Click here to download high resolution image](#)







Figure 3
[Click here to download high resolution image](#)





Figure 4
[Click here to download high resolution image](#)





Figure 5
[Click here to download high resolution image](#)




Figure 6
[Click here to download high resolution image](#)




Figure 7
[Click here to download high resolution image](#)




Figure 8
[Click here to download high resolution image](#)




Table S1

[Click here to download Supplementary data: TableS1.pdf](#)

Supplementary figures

[Click here to download Supplementary data: SupplFig.pdf](#)

File S1

[Click here to download Supplementary data: File S1.pdf](#)