

**Effects of clogging on stream macroinvertebrates:
an experimental approach**

Tiziano Bo, Stefano Fenoglio¹, Giorgio Malacarne, Massimo Pessino, Francesca Sgariboldi

University of Piemonte Orientale, Di.S.A.V., Via Bellini n. 25, 15100 Alessandria, Italy

¹ Corresponding author: fenoglio@unipmn.it

Short running title: Effect of clogging on macroinvertebrates

Key words: sedimentation, interstitial zone, macroinvertebrates, functional groups, NW Italy, Piemonte

**Effects of clogging on stream macroinvertebrates:
an experimental approach**

Abstract

The influence of stream bed sediment clogging on macroinvertebrate communities was investigated in the Lemme creek (NW Italy). To assess how fine sediment accumulation can influence the colonisation process and community composition of macroinvertebrates, we placed 48 traps in the riverbed. The traps consisted of boxes built with metal net (mesh 1 cm, height 15 cm, sides 5 cm) covered with nylon net except for the apex, allowing access exclusively from the top. We created 4 trap types filled with: 100% gravel, 30% sand and 70% gravel, 70% sand and 30% gravel and 100% sand. After 20 and 40 days, we removed 6 traps/type. Macroinvertebrates rapidly colonised the traps, as we found no significant community differences between the two removal dates. Among the 4 trap types, we found significant differences in taxa number and abundance, which both decreased with increasing clogging. Thus, our study supports the hypothesis that clogging and the accumulation of fine substratum elements strongly affects benthic stream communities.

1. Introduction

The interstitial zone plays an important role in the ecological processes of lotic ecosystem. Thus studying the interstitial areas in bed sediments (*sensu* Bretschko, 1992) is of major importance for lotic ecologists to better understand the physical, chemical and biological processes that take place in the stream bed (Vervier et al., 1992). For example, within the stream bed interstices, organic matter and nutrients can be retained, transformed and stored (Gibert et al., 1990). Many ecological processes are based on complex exchanges between the streambed surface and subsurface. In this context hydraulic gradients at the streambed surface, depth of the hyporheic zone, and sediment characteristics, such as bed porosity and hydraulic conductivity, play a major role (Harvey & Bencala, 1993; White, 1993).

Pioneer studies underlined the importance of bed sediments for the distribution of stream insects (Wene, 1940). In fact, macroinvertebrates use substrata for deposition and incubation of eggs, for feeding (Minshall, 1984), as a shelter from predation (Brusven & Rose, 1981), refuge from physical disturbances (Palmer et al., 1992; Dole-Olivier et al., 1997; Gayraud et al., 2000) and during extreme drought events (Fenoglio et al., 2006).

Sediment size and depth affect the size of available interstitial space (Maridet *et al.*, 1995), thus they are important parameters in studying macroinvertebrate colonisation and distribution (Rae, 1987). It is likely that fine sediment accumulations may clog the interstices, reducing interstitial water exchange, lowering the concentration of dissolved oxygen among the sediments, and finally constraining the movement of some invertebrates in the substrata.

In the last decades, human-induced alterations of the natural morpho-hydrological characteristics have altered the transport-deposition cycle in many rivers, so that clogging has become an important ecological problem (Weigelhofer & Waringer, 2003).

Therefore the aims of this experimental study were: a) to assess how different amounts of fine sediments (sand) in a coarse substratum (gravel) could influence the colonisation process of macroinvertebrates; b) to investigate the effect of clogging on macroinvertebrate taxonomic richness, abundance and community composition.

2. Methods

We experimented in the Lemme creek, a small tributary (2nd order stream with moderate slope) of the Orba river, NW Italy (44° 35' 45'', 8° 51' 41''; altitude 430 m a.s.l.). Some abiotic parameters were measured using portable instruments (Eijkelkamp 13.14 and 18.28) (Table 1).

The studied stream has a good environmental quality, reaching the first class in the Extended Biotic Index system (Ghetti, 1997), which corresponds to an environment without traces of human-induced alteration. To assess how different amount of fine sediments in the substratum can influence colonisation processes and community composition of macroinvertebrates, we placed 48 traps in a large and uniform riffle of the Lemme creek riverbed, using a random distribution (Fig. 1). Each trap consisted of a box built with metal net (5 cm long, 5 cm wide and 15 cm high, mesh width 1 cm, total volume = 0.37 dm³) that was covered with nylon net (mesh width 250 µm) except for the apex, allowing access for macroinvertebrates exclusively from the top of the trap. Traps were individually numbered: traps from 1 to 12 contained only gravel, traps from 13 to 24 contained 30 % sand and 70 % gravel, traps from 25 to 36 contained 30 % gravel and 70 % sand and traps from 37 to 48 contained only sand. The average dimension of the gravel was 45.0 x 23.0 x 21.0 mm, while the sand measured meanly 0.25 - 0.50 mm (sensu Ghetti, 1997).

Traps were positioned in the stream on 28 June 2005 in a uniform, 100 m-long riffle, with the apex orientated upward at the same level of the natural stream substratum, so the depth reached by the traps in the substratum was the same as their height (15 cm). Traps were placed

randomly in the stream, thus the composition of traps substratum was not related with the one of the surrounding natural stream bed. After 20 and 40 days, we removed 6 traps of each type. In order to analyse abundance and composition of the natural bottom communities, we collected 30 Surber samples, once a week during the study period, in the same riffle area where we had placed the traps, using a 0.06 m² Surber sampler (250 µm mesh).

In the laboratory, all organisms were counted and identified to the species or genus level, except for Annelida, early instars of some Trichoptera and Diptera, and others that were identified to the family level. Each taxon was also assigned to a Functional Feeding Group (FFG: scrapers - Sc, shredders - Sh, collector-gatherers - Cg, filterers - F and predators - P) according to Merritt & Cummins (1996). Moreover, a classification of taxa into seven biological and seven ecological groups was conducted according to the Usseglio-Polatera et al. (2000) species traits approach.

3. Results

In total we collected 14793 organisms: 7916 macroinvertebrates belonging to 66 taxa in the riverbed and 6877 macroinvertebrates belonging to 55 taxa in the traps (Table 2). Mean abundance in the substratum was 4221.8 organisms/m² (\pm 624.8). Highest values of invertebrate densities were found in coarse substrata, while sandy and rocky substrata showed lowest densities. In the functional composition of the invertebrate assemblages, collectors-gatherers were the most represented FFG (33.4 % of all organisms), followed by shredders (30,1 %), predators (19,2 %), filterers (14.4 %) and scrapers (2.74 %). According to the Usseglio-Polatera et al. (2000) classification, the most represented biological groups in the substratum were the '*e group*' (small or medium sized, short-lived, crawlers with aquatic respiration and cemented eggs: 54.8 %), and the '*f group*' (medium sized, crawlers, shredders with aquatic respiration: 39.4 %). Considering ecological groups, the two most abundant groups were '*B*' (organisms living in

rhithronic rheophilous environments: 37.1 %) and 'C' (oligotrophic organisms living in banks and channels of rhithronic and epipotamic, at slow and medium current velocities: 31.4 %).

After exposure, interstitial traps were rapidly colonised, regarding both taxa richness and organism abundance, as we detected no significant differences between the two sampling dates (taxa richness: Mann-Whitney U test = 263.5, P = 0.61; invertebrate abundance: U = 243.5, P = 0.36).

However, comparing the invertebrate assemblages among the four trap types, we detected evident differences in both taxa and invertebrate numbers. Interestingly, the abundance of organisms (Fig. 2a) and the number of taxa (Fig. 2b) were inversely related to the total amount of sand. Differences among traps were significant considering both organism abundance (Kruskal-Wallis test = 14.67, P = 0.002) and taxa richness (Kruskal-Wallis test = 7.61, P = 0.05).

Collectors-gatherers was the most represented Functional Feeding Groups (FFG) in the traps (50.4 % of total), followed by Predators (21.8 %) and Filterers (14.3 %), while Shredders (12.4 %) and Scrapers (0.98 %) were less represented.

Comparing 100% gravel filled and 100% sand filled traps, densities of Filterers and Scrapers feeding groups were reduced by 44.0%, and 12.6 %, respectively. In particular the importance of Filterers was inversely related with the total amount of sand (Kruskal-Wallis Test = 12.3, P = 0.006). The most represented biological group in the traps was the 'e group' (66.8 %), followed by 'f group' (23.3 %), 'c group' (7.50 %), 'h group' (1.64 %), 'g group' (0.39 %) and 'd group' (0.33 %). In the different trap types, the importance of some biological groups, such as 'e' and 'f' decreased with the increase of sand, while on the contrary sandy substrata were positively related with the abundance of 'h' organisms (multivoltine, burrowers or interstitial microphagous). Considering the ecological group composition of the traps, we detected no significant differences, with the exception of 'C' organisms (Fig. 3), that were

significantly more abundant in substrata with more than 70 % of gravel (Kruskal-Wallis test = 9.84, $P = 0.02$).

Analysing the taxonomic composition of the traps, we evidenced some interesting differences among trap types: for example large predators, such as *Perla marginata*, were found only in traps filled with 70 or 100% gravel, avoiding more clogged substrata. Furthermore, rheostenic and lithophilous elements, such as Ephemeroptera Heptageniidae were almost exclusively found in gravel-filled traps (75 % of specimens found in traps with more than 70 % of gravel), or occurred there in highest abundances (Fig. 4). On the contrary, the number of Oligochaeta tend to increase in the traps filled with conspicuous amounts of fine elements.

4. Discussion

The Lemme is a typical Apenninic creek, with almost no human-derived impacts and with high levels of invertebrate biodiversity. Taxonomic richness, invertebrate abundances in the substratum were in the range of other Apenninic environments (Fenoglio et al., 2002), such as functional, biological and ecological composition of the assemblages (Bo et al., 2006).

The composition of the bed sediment plays a key role in the colonisation mechanism, composition and abundance of benthic community. Some studies found that in streams with high sediment porosity, the number of invertebrates beneath the top sediment layer may exceed that in the sediment surface by far (Bretschko, 1981; Maridet & Philippe, 1995). Others (Strommer & Smock, 1989; Wagner et al., 1993) found extreme low organism densities in interstitial habitats if those had high quantities of sand and low porosity. Walling and Amos (1999) evidenced that the increase of turbidity and sedimentation can represent a serious damage for salmonids in chalk streams of southern England, by reducing the supply of oxygen to the hatching eggs, as well as degrading other characteristics of the channel habitat.

Correspondingly, our traps filled with 100% of sand, i.e. the most clogged ones, provided always the poorest interstitial habitat quality, probably because tight packing of sand grains reduced the trapping of organic detritus, limited the availability of oxygen and lowered the accessible pore space for most of the macroinvertebrates.

In our study, the effect of clogging is very evident and affected organisms belonging to several functional, ecological and biological groups. In particular, the abundance of filterers is strongly reduced by the accumulation of fine elements, probably because of the reduction of water circulation but also large, semi-voltine, crawlers organisms were not able to colonise clogging environments. The taxonomical, ecological and functional composition of the interstitial assemblages was significantly changed by the accumulation of sand in the substratum of studied stream: only few groups, such as Oligochaeta benefited from the clogging process.

Some recent studies have pointed out that the macroinvertebrate fauna of lotic systems can be affected by unnatural, human-induced sedimentation by different indirect and direct effects. Direct effects are for example the loss of microhabitat (Rae, 1987), loss of access to trophic resources (Lenat et al., 1981), and the damage to respiratory systems of organisms (Lemly, 1982). Indirect effects include for example the change of the ecological processes that are on the basis of autochthonous and allochthonous sources of energy of the system (Quinn et al., 1992).

Alteration of stream morphology and human activities in stream beds can increase the clogging and sedimentation process, altering the granulometry of large parts of stream beds and reducing the biological diversity of stream communities. In Apenninic streams of North-Western Italy, there was a dramatic increase in the regulation and alteration of many lotic systems. After the floods of 1994 and 2000 (Fenoglio et al., 2003), a lot of rivers and creeks in this area were morphologically modified, with unnatural increases of sedimentation also in low order streams.

This study supports the hypothesis that clogging and the increase of sedimentation of fine substratum elements strongly affects the benthic stream invertebrate community, and that this process can diminish the biological diversity of Apenninic lotic environments. Future research should investigate the importance of the organic components in the clogging of the interstitial spaces of streambed substrata.

5. Acknowledgements

We thanks R. Castellaro for his help during field work and G. Ferro for the determination of Coleoptera Hydraenidae.

6. References

- Bo, T., Cucco, M., Fenoglio, S., & Malacarne, G. (2006). Colonisation patterns and vertical movements of stream invertebrates in the interstitial zone: a case study in the Apennine, NW Italy. *Hydrobiologia*, *568*, 67-78.
- Bretschko, G. (1981). Vertical distribution of zoobenthos in an Alpine brook of the Ritrodat-Lunz study area. *Verhandlungen Internationale Vereinigung für Theoretische und Angewandte Limnologie*, *21*, 873-876.
- Bretschko, G. (1992). Differentiation between epigeic and hypogeic fauna in gravel streams. *Regulated Rivers*, *7*, 17-22.
- Brusven, M. A., & Rose, S. T. (1981). Influence of substrate composition and suspended sediment on insect predation by the torrent sculpin, *Cottus rhotenus*. *Canadian Journal of Fisheries and Aquatic Sciences*, *38*, 1444-1448.
- Dole-Olivier, M. J., Marmonier, P., & Befy, J. L. (1997). Response of invertebrates to lotic disturbance: is the hyporheic zone a patchy refugium? *Freshwater Biology*, *37*, 257-276.
- Fenoglio, S., Agosta, P., Bo, T., & Cucco, M. (2002). Field experiments on colonisation and movements of stream invertebrates in an Apennine river (Visone, NW Italy). *Hydrobiologia*, *474*, 125-130.
- Fenoglio, S., Battezzore, M., & Morisi, A. (2003). Alluvioni e comunità macrobentoniche: una panoramica complessiva ed alcuni casi di studio in Piemonte. *Pianura*, *17*, 199-206.
- Fenoglio, S., Bo, T. & Bosi, G. (2006). Deep interstitial habitat as refuge for *Agabus paludosus* (Fabricius, 1801) (Coleoptera: Dytiscidae) during summer droughts. *The Coleopterists Bulletin*, *60*, 37-41.
- Gayraud, S., Philippe, M., & Maridet, L. (2000). The response of benthic macroinvertebrates to artificial disturbances: Drift or vertical movements in the gravel bed of two sub-alpine streams? *Archiv für Hydrobiologie*, *47*, 431-446.

- Ghetti, P. F. (1997). *Manuale di applicazione Indice Biotico Esteso (I.B.E.), i macroinvertebrati nel controllo della qualità degli ambienti di acque correnti*. Provincia Autonoma di Trento, Trento.
- Gibert, J., Dole-Olivier, M. J., Marmonier, P., & Vervier, P. (1990). Groundwater ecotones. In: R. J., Naiman, & Décamps, H. (Eds.), *Ecology and Management of Aquatic Terrestrial Ecotones. Man and the Biosphere* (pp:199–225). Paris-London: Series UNESCO and Parthenon Publishing, Carnforth.
- Harvey, J. W., & Bencala, K. E. (1993). The effect of streambed topography on surface-subsurface water exchange in mountain catchments. *Water Resource Research*, 29, 89-98.
- Lemly, A. D. (1982). Modification of benthic insect communities in polluted streams: combined effects of sedimentation and nutrient enrichment. *Hydrobiologia* 87: 229–245.
- Lenat, D. R., Penrose, D. L., & Eagleson, K.W. (1981). Variable effects of sediment addition on stream benthos. *Hydrobiologia*, 79, 187–194.
- Maridet, L., & Philippe, M., (1995). Influence of substrate characteristics on the vertical distribution of stream macroinvertebrates in the hyporheic zone. *Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biologia*, 91, 101-105.
- Merritt, R. W., & Cummins K. W. (1996). *An introduction to the aquatic insects of North America*. 3^o Edition, Kendall Hunt, Dubuque, IO.
- Minshall, G. W. (1984). Aquatic insect-substratum relationships. In: Resh, V. H., & Rosenberg, D. M. (Eds.), *The ecology of aquatic insects*. (pp: 358-400). New York: Praeger Publisher.
- Palmer, M. A., Bely, A. E., & Berg, K. E. (1992). Response of invertebrates to lotic disturbance: a test of the hyporheic refuge hypothesis. *Oecologia*, 89, 182-194.
- Quinn, J. M., Davies-Colley, R. J., Hickey, C. W., Vickers, M. L., & Ryan, P.A. (1992). Effects of clay discharges on streams. 2. Benthic invertebrates. *Hydrobiologia*, 248, 235–247.

- Rae, J. G. (1987). The effects of flooding and sediments on the structure of a stream midge assemblage. *Hydrobiologia*, *144*, 3-10.
- Strommer, J. L., & Smock, L. A. (1989). Vertical distribution and abundance of invertebrates within the sandy substrate of a low-gradient headwater stream. *Freshwater Biology*, *22*, 263-274.
- Usseglio-Polatera, P., Bournaud, M., Richoux, P., & Tachet, H. (2000). Biological and ecological traits of benthic fresh-water macroinvertebrates: relationships and definition of groups with similar traits. *Freshwater Biology*, *43*, 175–205.
- Vervier, P., Gibert, J., Marmonier, P., & Dole-Olivier, M. J. (1992). A perspective on the permeability of the surface freshwater-groundwater ecotone. *Journal of North American Benthological Society*, *11*, 93–102.
- Wagner, R., Schmidt, H. H., & Marxsen, J. (1993). The hyporheic habitat of the Breitenbach, spatial structure and physicochemical condition as a basis for benthic life. *Limnologica*, *23*, 285-294.
- Walling, D. E., & Amos, C. M. (1999). Source, storage and mobilisation of fine sediment in a chalk stream system. *Hydrological Processes*, *Hydrological Processes*, *13*, 323-340.
- Weigelhofer, G., & Waringer, J. (2003). Vertical distribution of benthic macroinvertebrates in riffles versus deep runs with differing contents of fine sediments (Weidlingbach, Austria). *International Review of Hydrobiology*, *88*, 304–313.
- Wene, G. (1940). The soil as an ecological factor in the abundance of aquatic chironomid larvae. *Ohio Journal of Sciences*, *40*, 193-199.
- White, D. S. (1993). Perspectives on defining and delineating hyporheic zone. *Journal of North American Benthological Society*, *12*, 61-69.

Caption to figures

Fig. 1: Lemme Creek, traps displacement and scheme of the four trap types.

Fig. 2: a) Abundance of invertebrates in the traps with different granulometry (mean \pm SE); b) taxa richness in the traps with different granulometry (mean \pm SE).

Fig. 3: Total abundance of oligotrophic, rhitronic and epipotamic (sensu Usseglio-Polatera et al., 2000) organisms with the increase of clogging in the artificial substrata (mean \pm SE).

Fig. 4: Abundance of some selected taxa (mean \pm SE) in the four clogged trap types: *Cheumatopsyche lepida* (filterer Trichoptera - right Y axis), *Ecdyonurus* sp. (rheophilous, scraper Ephemeroptera), Elmidae larvae (collector-gatherer Coleoptera).

Table 1 – Environmental characteristics (mean \pm SD) in the stream reach during the study period.

Table 2 – Percent relative abundances of macroinvertebrates collected in the natural stream bed and from traps.