



UNIVERSITÀ DEGLI STUDI DI TORINO

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17

This is an author version of the contribution published on:

Questa è la versione dell'autore dell'opera:

[Avocetta 43: 139-147 (2019), DOI: <https://doi.org/10.30456/AVO.2019204>]

The definitive version is available at:

La versione definitiva è disponibile alla URL:

<http://www.avocetta.org/articles/vol-43-2-ems-a-new-alliance-between-a-riparian-species-and-gravel-diggers-might-counteract-population-declines/>

18 **Proper gravel management may counteract population decline of the Collared Sand**
19 **Martin *Riparia riparia***

20

21 Giulia Masoero¹, Giovanni Boano², Alberto Tamietti³, Enrico Caprio⁴

22 ¹ Section of Ecology, Department of Biology, FI-20014 University of Turku, Finland

23 ² Museo Civico di Storia Naturale di Carmagnola, Via San Francesco di Sales 188,
24 Carmagnola (TO), Italy

25 ³ Aree Protette del Po Torinese, Corso Trieste 98, Moncalieri (TO), Italy

26 ⁴ Università degli Studi di Torino, Dipartimento di Scienze della Vita e Biologia dei
27 Sistemi, Via Accademia Albertina 13, 10123 Torino, Italy

28

29 Corresponding author: Giulia Masoero giulia.masoero@gmail.com

30 ORCID:

31 Giulia Masoero 0000-0003-4429-7726

32 Giovanni Boano 0000-0002-2670-4351

33 Enrico Caprio 0000-0002-5997-5959

34

35 Short title: Sand martins, rivers and quarries

36

37 **ABSTRACT**

38 Riparian habitats have gone through major structural changes, and related bird
39 populations had to suffer the consequences or adapt to the newfound conditions. Here,
40 we present the results of the analysis of on the river and quarry evolution, in relation to a
41 long-term monitoring (1970-2016) of a Sand Martins population nesting along the River

42 Po in northern Italy. During the course of the study, the population changed breeding site,
43 preferring more anthropogenic sites in the surrounding quarries to the natural river banks.
44 The alteration of the river dynamic and linearization of the course, alongside the
45 development of the sand quarries, may have caused this change. We conclude with a
46 consideration on the ways to support the survival of populations of riparian birds.
47 Functional nesting habitat along the river should be better preserved, and potential nesting
48 areas in the surrounding quarries should be protected with long term restoration projects.
49 Sand quarries are not free of threats, and cooperation among stakeholders has proven to
50 be of the utmost importance to ensure the success of the population breeding in the area.

51

52 **Keywords:** nest site selection; river dynamic; *Riparia riparia*; quarry restoration.

53

54

55 INTRODUCTION

56 Habitat loss, due to destruction and alteration, is one of the most important
57 anthropogenic impacts on animal populations (Newton 1998, Baillie et al. 2004, Curry
58 Lindahl 1972). Among the most affected habitats, riparian ecosystems have been deeply
59 altered with radical changes to their courses, flow and dynamics. The regulation of the
60 flow and course forces rivers to become more and more linear, losing their meanders and
61 changing the erosion of river banks (Brookes 1988). Changes in the riparian system can
62 affect many aquatic animals, and have proven to affect the structure of breeding bird
63 community (Figarski & Kajtoch 2015, Girvetz 2010).

64 The Collared Sand Martin (*Riparia riparia*; hereafter “Sand Martin”) is riparian
65 species particularly specialised in nesting habitat (Moffatt et al. 2005). This colonial

66 species usually excavates nest holes in sandy banks, normally in the vicinity of water
67 bodies, such as the eroding banks of meandering rivers and streams, but also in artificial
68 sand deposits (Cramp 1988). The most recent IUCN assessment of the global trend
69 categorized Sand Martin populations as Decreasing (BirdLife International 2016).
70 According to the most recent population assessment published by BirdLife International
71 in 2015, Sand Martin populations are decreasing in many European countries, including
72 Italy (Brichetti & Fracasso 2007, Campedelli et al. 2012). This trend appears even more
73 concerning since the species has also experienced large scale population declines since
74 the early 1960s (Cowley 1979, Jones 1987a).

75 Climate change has often been considered as the major driver behind Sand Martin
76 population changes and this trans-Saharan migrant has been well studied in terms of the
77 influence that climatic factors in the African wintering quarters have on its survival
78 (Cowley 1979, Svensson 1986, Persson 1987a, Szép 1995, Norman & Peach 2013). In
79 more recent years, there has not been any strong drought in the Sahel region (Evan et al.
80 2014, Sanogo et al. 2015) that might have driven the global decrease in Sand Martin
81 population sizes. Habitat change might therefore be the cause behind more recent
82 declines. However, despite it being recognised as a cause of population decline (Garrison
83 1999, del Hoyo et al. 2004), few papers have analysed habitat alteration on the breeding
84 grounds. In California, where the species population is declining mainly due to habitat
85 alteration, Moffatt and colleagues (2005) studied the importance of riparian systems and
86 the effects of different restoration strategies on the species. Sand martins were reported
87 to breed in sandy quarry complexes all over Europe (Alves 1997, Cowley & Siriwardena
88 2005, Heneberg 2007, Jones 1987b, Krištofik et al. 1994, Kitowski et al. 2015, Morgan
89 1979, Norman & Peach 2013, Persson 1987b, 1987a). Some of the eastern populations of

90 Sand Martin, however, were still recently found breeding in natural sites in Poland
91 (Figarski & Kajtoch 2015) and, in large numbers, along the banks of the Tisza River in
92 Hungary (Szép 1995, Szép et al. 2003). Despite this important change in the selection of
93 breeding sites in some areas, to our knowledge there are no studies that have analysed the
94 causes behind this phenomenon.

95 In this paper, we have analysed survey data, collected over 45 years, of a Sand Martin
96 population in north-western Italy. The monitored population did not present signs of
97 decline during the study, and its fluctuations in numbers appeared to have no correlation
98 with climate (Masoero et al. 2016). We have also studied the river dynamic of the past 68
99 years and the evolution of the quarry complexes, from 1960 until the present, in the
100 nesting area of the Sand Martins. Our aim was to understand the relationship between
101 Sand Martins and their nesting habitat and how the modifications of the river and of the
102 quarries could affect them.

103

104 **METHODS**

105 **Study area and the quarries**

106 The study area was included about 25 km of the River Po South of the city of Turin,
107 Italy (from 44.971 N 7.693 E to 44.832 N 7.636 E) and all the 15 surrounding quarries,
108 nine active during the study and six already restored to a more natural state (Fig. 1). The
109 area is located in the River Po Torinese Regional Park. The River Po and its tributaries
110 transported and deposited materials during the Middle Pleistocene and the Holocene
111 creating the river channel deposits that currently constitute this area of the Po plain. The
112 sediments at the top layer are 25-100 m thick and are exploited by quarries complexes
113 (Castiglioni et al. 1999). The raw materials, such as sand and gravel, are excavated from

114 the ground to be used as construction aggregates. After the extraction from the ground,
115 sand and gravel are separated in a range of sizes and stored into stockpiles, creating heaps
116 and banks in which the Sand Martins can build their nests (Fig. 2). The quarries are
117 usually worked in progressive phases, in order to minimise the exposed areas. This
118 method also ensures timely restoration of the already excavated sites while the work is
119 still ongoing. The Park oversees the work of the quarries under its jurisdiction and
120 controls the restoration of the excavated banks and surfaces.

121 We georeferenced seven different maps of the River Po (from Casalgrasso 44.830 N
122 7.634 E to Moncalieri 45.008 N 7.679 E) from the years 1922, 1948, 1955, 1963, 1978,
123 1991 and 2008. To assess river dynamism, we calculated the sinuosity index, i.e. the
124 measures of the degree of sinuosity of meandering rivers calculated as the ratio of the
125 length of the midline of the channel and the air-line distance (Friend & Sinha 1993,
126 Alabyan & Chalov 1998). Values of the sinuosity index close to or equal to 1 correspond
127 to a linear river, whereas higher values correspond to a more winding river.

128 Data concerning the changes in the quarries were obtained from the databases on
129 extraction activities of the Ministry of Industry and of the Piedmont Region. From 1960
130 to 1985, the Italian Ministry of Industry collected the data on the amount of extracted
131 construction aggregates for the whole region of Piedmont. After 1982, the Piedmont
132 Region started collecting the data regarding the quarries, obtaining therefore more precise
133 data and we could retrieve data concerning the situation in the 15 quarries located in the
134 study area. We decided to use the amount of extracted material as an index of the quarry
135 activity, and therefore of the amount of potential available nesting habitat because of two
136 main reasons. First, the studied quarries are authorised to extract materials also going
137 below the level of the aquifer, i.e. sand and gravel can still be extracted with the surface

138 of the active quarry remaining stable. Second, sand martins often colonised the sand heaps
139 of extracted materials and seldom the excavated cliffs.

140

141 **Census of breeding colonies and data analysis**

142 From 1970 up to 2016 the riverbanks and the sand quarries were surveyed each year
143 to identify breeding sites and their location. The census of the colonies with an exhaustive
144 count of all active nests started in 2000, whereas in earlier years (i.e, since 1970), the
145 monitoring of the river was not conducted quantitatively, due to the difficulty to reach
146 every area of the river. From the end of April to July all the possible breeding sites were
147 checked during multiple occasions, both along the river Po and in active and abandoned
148 quarries, and all active nests were counted. Almost every year there was a change in the
149 nesting location chosen by the population, which sometimes split among two or more
150 quarries if the nesting cliffs were not wide enough.

151 We used generalized additive models (GAMs; Hastie & Tibshirani 1990) to test for
152 the effects of year on the number of nests in riverbanks assuming a Poisson data
153 distribution. GAMs were performed using R 3.3.1 (R Core Team 2017), with the package
154 mgcv (Wood 2011). The default cubic smoothing spline method with four degrees of
155 freedom was used to smooth the year component. We also evaluated Collared Sand
156 Martin population trends between 2002 and 2016 (years with a complete census of the
157 population) using the software TRIM (Trend & Indices for Monitoring data, TRIM 3.54;
158 Pannekoek & Van Strien 2006). TRIM estimates annual indices and evaluates trends in
159 these indices implementing log-linear models, an approach commonly employed in
160 temporal series analysis. We also tested for a linear relationship between the number of

161 nests in riverbanks and quarries and the amount of gravel and sand extracted from gravel
162 pits using a generalized linear model, specifying a Poisson data distribution.

163

164 **RESULTS**

165 **River dynamic and quarry evolution**

166 The River Po course changed markedly in the study area from 1922 until 1978 (Fig.
167 3), after which it remained largely unchanged. The sinuosity index of the River Po
168 decreased from a value of 2.19 in 1922, to a minimum of 1.59 calculated in 2008 (Fig.
169 4), showing that the river had become increasingly linear. Two large floods happened in
170 2000 and 2016, and created a few suitable nesting areas that were not colonised by Sand
171 Martins. The new nesting cliffs were nevertheless very limited compared to the previous
172 floods, the bank defences did not allow more substantial changes.

173 From the 1960-1985 data from the Italian Minister of Industry it emerged that until
174 1966 the production of construction materials in Piedmont region was low, and always
175 below 3 Mm³. In the subsequent 10 years it grew steeply, reaching a maximum of 14
176 Mm³ in 1976. It started then to decrease, but always maintaining values above the 6 Mm³
177 (Fig. 5a). After 1982, more precise data from the Piedmont Region are available and it
178 was possible to look in detail at the situation in the 15 quarries present in the study area
179 (Fig. 5b). During the years from 1982 to 2007, the amount of extracted construction
180 aggregates presented a fluctuating trend, ranging from a minimum of 110000 m³ in 1995
181 and a maximum of 1.71 Mm³ in 1983. After 2007, the production almost constantly
182 declined.

183

184 **Census of breeding colonies**

185 The size of colonies along the riverbanks varied from a maximum of 244 active nests
186 recorded in 1973 to a minimum of 30 nests in 1983 (Fig. 4). No active nests were recorded
187 after 2002. A Generalized Additive Model of the number of nests in riverbanks against
188 year revealed a significant non-linear negative relationship ($\chi^2_{7.058} = 276.5$, $p < 0.001$,
189 deviance explained = 79%). We cannot assert that the population breeding along the river
190 was smaller than the actual population, due to incomplete surveys carried on in the last
191 century, but we can assert that after 2002 all the colonies are breeding in the quarries.
192 Colonies in sand quarries were detected the first time in 1983 with 50 active nests and
193 were found always in active quarries. The number of active nests varied from a minimum
194 of 20 nests in 1993 to a maximum of 934 nests in 2007 (Fig. 4). A Generalized Additive
195 Model of the number of nests in sand quarries against year revealed a significant non-
196 linear positive relationship ($\chi^2_{8.108} = 3960$, $p < 0.001$, deviance explained = 93%). Using
197 TRIM, the population trend from 2002-2016 is uncertain (multiplicative overall slope
198 model: 1.0067, SE: +/- 0.0349). The number of nests counted in sand quarries were
199 positively related with the amount of construction aggregates extracted in the study area
200 (with the standardised variable, intercept: 6.02 ± 0.01 , beta: 0.40 ± 0.01 ; $F_{1,19}$: 6.07, $p =$
201 0.023).

202

203 **DISCUSSION**

204 In this paper we analysed breeding observations of Sand Martins along a section of
205 the river Po and the surrounding quarries. At the same time, we looked into detailed
206 data regarding the river course and the development of the quarries. Overall, we think
207 this provides a reasonable explanation to the change in breeding sites and to the lack of
208 decrease in size previously shown by the studied population (Masoero et al. 2016).

209

210 River and quarries

211 By looking at map of the River Po courses along the years and at the values of the
212 sinuosity index we can notice a few things. First, a decrease in the river dynamism in the
213 study area, i.e. the river changed its course markedly until 1978 and after that, the course
214 remained largely unchanged, even after two flooding events in the last 15 years. Second,
215 the low values of the index in the later years show a linearization of the river when
216 compared to the earlier values. The fixed course of the river and the lower sinuosity index
217 indicates a decrease in river dynamism, and as a consequence, a decrease in river
218 functionality (Fehér et al. 2012, Yu et al. 2015). The history behind these changes, both
219 regarding the river and the surrounding quarries, is quite complex and has to be told
220 alongside. Between the 1960s and the 1980s, Italy experienced an economic boom. As
221 the surrounding plain became more anthropized, towns grew. An increasing number of
222 flood protections were built around the surrounding cities, houses, crops and poplar
223 plantations. Without any space left to change its course, the river assumed a more fixed
224 and linear path. In the meanwhile, the amount of construction aggregates excavated was
225 also following the market needs. Up until 1960 the extraction took place mainly inside
226 the river with limited volumes. In the early 1960s, the demand for aggregates slowly
227 started to grow and the quantity of sand and gravel extracted from the riverbed was no
228 longer sufficient and the quarry companies started exploiting the sediments present along
229 the river course and the first lakes created by gravel excavations appeared in the alluvial
230 plain. After 1966, the economic boom lead to a strong increase in the volumes of
231 excavated materials. In the 1970s, the excavation of the riverbed started to decrease, due
232 to more severe environmental laws, until it came to a stop. From the 1980s the extraction

233 of construction aggregates continued along the river in quarries outside of the riverbed.
234 The existing quarries expanded both in area and depth and the amount of extracted
235 construction aggregates presented a fluctuating trend, following the marked needs.
236 Around 2007, to the economic crisis hit the construction market. Some quarries were
237 closed and the active ones started extract less sand and gravel. Inactive quarries are
238 currently under a restoration process coordinated by the River Po Torinese Regional Park.
239 A small increase in the extracted material in 2016 might be the sign of a mild economic
240 recovery.

241

242 **Sand martins and breeding habitat**

243 The evolution of the river habitat and of its quarries had a deep impact on Sand
244 Martins. The population investigated in our study, as other studied populations in many
245 European countries (Cowley & Siriwardena 2005, Norman & Peach 2013), has shown a
246 change in the selection of breeding sites, although the general area remained unchanged.
247 It was not possible to obtain overall population trends for the study area because the
248 census on the riverbanks were not always performed with the same methodology. Despite
249 we reckon this may be a limit for previous years, but we still believe that we were able
250 to identify (qualitatively) the magnitude of the reduction of the nests in the riverbanks,
251 and correctly censused the nests since 2000. In the early years, the population was
252 exclusively breeding in the banks of the River Po. It then began to progressively colonize
253 the adjacent sand quarry complexes, and in 2003 the change had become permanent as
254 the traditional riverbank sites were definitively abandoned. This could be explained by
255 considering the decrease in river dynamism (Moffatt et al. 2005). Lower dynamism
256 means less bank erosion, so new sand cliffs are not created and the old ones, remaining

257 undisturbed, start to become covered in vegetation. At the same time, the river became
258 more linear, determining a change in the river regime, and as a consequence, faster and
259 more destructive floods that could have had an impact on the population survival and
260 breeding success. In the meanwhile, the need for construction aggregates grew because
261 of the economic boom, and new sand and gravel pits began to spread along the course of
262 the river. Sand quarries offered clear sand cliffs, free of vegetation, apparently the perfect
263 substitute to the traditional breeding sites. Nevertheless, nesting in an active sand quarry
264 presents some potential threats. If the workers are careless or not aware of the presence
265 of a colony, they could destroy it. In Italy, active nests are protected by the law (L.
266 157/92), but people are often unaware of the law or of the presence of nests. Quarries can
267 therefore turn into an ecological trap, i.e. Sand Martins can start nesting in a good sand
268 cliff and the nests can be destroyed. Another problem can follow from the quarry
269 management which can change following the market needs. In the quarry environment, a
270 good management, alongside the protection and the monitoring of the colonies, is crucial,
271 because such quarries act as surrogates for riverbanks not only for the Sand Martin, but
272 also for other species such as the European Bee-eater (*Merops apiaster*) and the Common
273 Kingfisher (*Alcedo atthis*). With simple initiatives (i.e. leaving undisturbed the sand
274 heaps with active nests during May, June and July) that have very low impact on quarry
275 management, it is possible to combine economic interests with nature conservation. All
276 of the above has been possible in the study area thanks to a positive interaction between
277 the people in charge of wildlife protection and those working in the quarries. The River
278 Po Torinese Regional Park is actively protecting the colonies having these sand quarries
279 under its jurisdiction, and it is also monitoring them. In 2000, the Park staff have also
280 started to raise awareness among the gravel diggers about the presence and the importance

281 of Sand Martins, and thus, the nests in the gravel pits have been left undisturbed allowing
282 Sand Martins to successfully fledge.

283 After the quarry exploitation, it is important that the restoration plan allows the Sand
284 Martins to nest for many years to come. Quarry restoration is indeed necessary because
285 of the extensive land disturbances and negative impacts in terms of both safety and
286 environment (Milgrom 2008, Zuquette et al. 2002). Reclamation techniques often try to
287 address this issues by creating a safe environment with more natural slopes and by
288 creating of appropriate surfaces for the establishment of vegetation (Down & Stocks
289 1978, Haywood 1979, Gunn & Bailey 1993, Legwaila et al. 2015, Muzzi & Mongardi
290 2016). Nowadays, the fact that colonies were never found in inactive quarries suggests
291 that the sites are not restored in a way that provides suitable nesting habitat. In fact, the
292 sand cliffs and heaps of active quarries used as breeding sites are no longer available after
293 the restoration process.

294

295 **Concluding remarks**

296 Diversity and abundance of riparian birds is usually increased by a renaturalisation of
297 the river channels often favoured by floods (Kajtoch and Figarski 2013). In most
298 industrialised countries, rivers have been showing signs of decrease in dynamism and
299 functionality, which has led to loss in suitable nesting habitat for the Sand Martins
300 (Moffatt et al. 2005). Not finding suitable habitat along the river, the studied population
301 of Sand Martins probably started nesting in the adjacent sand quarries, which have
302 increased their activity around the same years (Figs. 4 and 5). After 2007, fewer and fewer
303 building materials were needed by the building industry, and many of the quarries present
304 in the area began to close and being reclaimed. The population already lost the traditional

305 nesting sites along the river: what will happen when all the currently active quarries will
306 be closed and restored?

307 A thorough analysis of the available literature lead to the conclusion that there are two
308 possible alternatives: to leave sand cliffs close to the water available to the Sand Martins
309 to build their nests, or to build artificial breeding cliffs (Gulickx et al. 2007, Bachmann
310 et al. 2008). The first solution can last a few years, but the cliff can collapse or the
311 vegetation can soon invade it. In any case, it has to be managed throughout the years. The
312 second seems a more stable solution, but comes with another type of problem:
313 ectoparasites. Parasites that infest nest holes include different species, including the tick
314 *Ixodes lividus* (Szép & Møller 1999). Newly formed nest sites do not present high levels
315 of infestation by ectoparasites, but as colonies get older the prevalence can approach
316 100% (e.g. Szép and Møller 1999). In the traditional breeding sites and in the quarries the
317 colony is usually used for only a year since it is destroyed after the nesting season by
318 floods or by quarry activities, and the parasite load does not approach such high levels.
319 Artificial nests are used from one year to another and can therefore develop very high
320 concentrations of ectoparasites. In a colony of nests artificially excavated in the limestone
321 in the UK, all the nest holes were washed out using a water hose (Gulickx et al. 2007),
322 but the effectiveness of this measure remain to be proven.

323 A better management of rivers and of the riverbanks in particular should be promoted
324 to help the Sand Martins in their breeding grounds, and it would be beneficial also for
325 many other species of birds (Jankowiak and Ławicki 2014; Figarski and Kajtoch 2015).
326 In areas where the Sand Martins are present and already nesting in quarries, restoration
327 projects should consider the creation of artificial colonies (as in Bachmann et al. 2008).
328 Moreover, active quarries hosting Sand Martin colonies should provide cliffs that should

329 remain untouched during the whole breeding season, and create new ones every year.
330 This management could help preserving a large number of nests over a long period of
331 time as happened in our study.

332

333

334 **Acknowledgements**

335 We thank all the people that helped collecting the data and the River Po Torinese
336 Regional Park. We are grateful to all the Park staff (in particular E. Castello, F. Nobili
337 and S. Zaghi), and to S. Blasutta, I. Ellena, G. Ramello, S. Tozzi and G. Vaschetti. We
338 acknowledge L. Mancuso and M. Moccagatta for their help in recovering the Piemonte
339 Region data from the database. Thanks also to D.E. Chamberlain, his comments and
340 corrections on an early version of the manuscript were greatly appreciated.

341

342 **References**

343 Alabyan A.M. & Chalov R.S., 1998. Types of river channel patterns and their natural
344 controls. *Earth Surf. Process. Landf.* 23: 467–474.

345 Alves M.A.S., 1997. Effects of ectoparasites on the Sand Martin *Riparia riparia*
346 nestlings. *Ibis*. 139: 494–496.

347 Bachmann S., Haller B., Lötscher R., Rehsteiner U., Spaar R. & Voge C., 2008.

348 Guide de promotion de l'hirondelle de rivage en Suisse. Conseils pratiques pour la
349 gestion des colonies dans les carrières et la construction de parois de nidification.

350 Fondation Paysage et Gravier, Uttigen, Association Suisse de l'industrie des Graviers et

351 du Béton, Berne, Association Suisse pour la Protection des Oiseaux ASPO/BirdLife

352 Suisse, Cudrefin, Station ornithologique suisse, Sempach.

- 353 Baillie J., Hilton-Taylor C. & Stuart S.N., 2004. 2004 IUCN red list of threatened
354 species: a global species assessment Gland, Switzerland: IUCN, pp. 191.
- 355 BirdLife International, 2016. *Riparia riparia*. The IUCN Red List of
356 Threatened Species 2016: e.T103815961A87438023.
357 [http://dx.doi.org/10.2305/IUCN.UK.2016-](http://dx.doi.org/10.2305/IUCN.UK.2016-3.RLTS.T103815961A87438023.en)
358 [3.RLTS.T103815961A87438023.en](http://dx.doi.org/10.2305/IUCN.UK.2016-3.RLTS.T103815961A87438023.en). Downloaded on 05 November 2019.
- 359 BirdLife International, 2017. European birds of conservation concern: populations,
360 trends and national responsibilities. BirdLife International, Cambridge.
- 361 Brichetti P. & Fracasso G., 2007. Ornitologia italiana. Vol. 4 Apodidae –
362 Prunellidae. Alberto Perdisa Editore. Bologna.
- 363 Brookes A., 1988. Channelized Rivers: Perspectives for Environmental
364 Management. Wiley, Chichester.
- 365 Campedelli T., Buvoli L., Bonazzi P., Calabrese L., Calvi G., Celada C., Cutini S.,
366 De Carli E., Fornasari L., Fulco E., La Gioia G., Londi G., Rossi P., Silva L. & Tellini
367 Florenzano G., 2012. Andamenti di popolazione delle specie comuni nidificanti in
368 Italia: 2000-2011. Avocetta 36: 121–143.
- 369 Castiglioni G.B., Biancotti A., Bondesan M., Cortemiglia G.C., Elmi C., Favero V.,
370 Gasperi G., Marchetti G., Orombelli G., Pellegrini G.B. & Tellini C., 1999.
371 Geomorphological map of the Po Plain, Italy, at a scale of 1:250 000. Earth Surf.
372 Process. Landf. 24: 115–1120.
- 373 Cowley E., 1979. Sand Martin population trends in Britain, 1965–1978. Bird Study
374 26: 113–116.
- 375 Cowley E. & Siriwardena G.M., 2005. Long-term variation in survival rates of Sand
376 Martins *Riparia riparia*: dependence on breeding and wintering ground weather, age

- 377 and sex, and their population consequences. *Bird Study* 52: 237–251.
- 378 Cramp S., 1988. *Handbook of the Birds of Europe, the Middle East and North*
379 *Africa: the birds of the Western Palearctic. Volume V. Tyrant Flycatchers to Thrushes.*
380 Oxford University Press, Oxford.
- 381 Curry-Lindahl K., 1972. *Conservation for survival. An ecological strategy.* William
382 Morrow, London.
- 383 Down C.G. & Stocks J., 1978. *Environmental impact of mining.* Applied Science
384 Publishers, London.
- 385 Evan A.T., Flamant C., Lavaysse C., Kocha C. & Saci A., 2014. Water Vapor–
386 Forced Greenhouse Warming over the Sahara Desert and the Recent Recovery from the
387 Sahelian Drought. *Journal of Climate*, 28, 108–123.
- 388 Fehér J., Gáspár J., Veres K.S., Kiss A., Globevnik L., Peterlin M., Kirn T., Stein U.,
389 Prins T., Spiteri C., Laukkonen E., Heiskanen A.-S., Austner K., Semeradova S. &
390 Künitzer A., 2012. *Hydromorphological alterations and pressures in European rivers,*
391 *lakes, transitional and coastal waters: thematic assessment for EEA Water 2012 report.*
392 VITUKI, NIVA, IWRS, Ecologic, Ecologic Institute, Stichting Deltares, SYKE,
393 CENIA, Prague.
- 394 Figarski T. & Kajtoch Ł., 2015. Alterations of riverine ecosystems adversely affect
395 bird assemblages. *Hydrobiologia* 744: 287–296.
- 396 Friend P.F. & Sinha R., 1993. Braiding and meandering parameters. *Geol. Soc.*
397 *London Spec. Publ.* 75: 105–111.
- 398 Garrison B.A., 1999. Bank Swallow (*Riparia riparia*). In A. Poole, F. Gill (Eds.),
399 *The birds of North America*, 414, Inc., Philadelphia.
- 400 Girvetz E.H., 2010. Removing erosion control projects increases bank swallow

- 401 (*Riparia riparia*) population viability modeled along the Sacramento River, California,
402 USA. *Biol. Conserv.* 143: 828–838.
- 403 Gulickx M.M.C., Beecroft R. & Green A., 2007. Creation of artificial Sand Martin
404 *Riparia riparia* burrows at Kingfishers Bridge. *Conservation Evidence*. 4: 51–53.
- 405 Gunn J. & Bailey D., 1993. Limestone quarrying and quarry reclamation in Britain.
406 *Environ. Geol.* 21: 167–172.
- 407 Hastie T. & Tibshirani R., 1990. *Generalized Additive Models*. Chapman and Hall,
408 New York.
- 409 Haywood S.M., 1979. Mineral landscapes: The next ten years? *Environ. Geochem.*
410 *Health* 1: 25–30.
- 411 Heneberg P., 2007. Sand martin (*Riparia riparia*) in the Czech Republic at the turn
412 of the millenium. *Linzer biol. Beitr.* 39: 293–312.
- 413 del Hoyo J., Elliott A. & Christie D.A., 2004. *Handbook of the birds of the world*,
414 Volume 9. Cotingas to pipits and wagtails. Lynx Editions, Barcelona.
- 415 Jankowiak Ł & Ławicki Ł, 2014. Marginal habitats as important refugia for riparian
416 birds during flood years. *Bird Study*. 61: 125–129.
- 417 Jones G., 1987a. Selection against large size in the Sand Martin. *Ibis*. 129: 274–280.
- 418 Jones G., 1987b. Selection against large size in the Sand Martin *Riparia riparia*
419 during a dramatic population crash. *Ibis*. 129: 274–280.
- 420 Kajtoch Ł & Figarski T, 2013. Short-term restoration of riverine bird assemblages
421 after a severe flood. *Bird Study*. 60: 327–334.
- 422 Kitowski I., Komosa A. & Chodorowski J., 2015. Absorbed radiation dose from
423 radon to the Sand Martin *Riparia riparia* during breeding at the sand mines in Eastern
424 Poland. *Int. J. Environ. Res.* 9: 1097–1106.

- 425 Krištofik J., Šustek Z. & Gajdoš P., 1994. Arthropods in nests of the Sand Martin
426 (*Riparia riparia* Linnaeus, 1758) in south Slovakia. Biol. Bratislava 49: 683–690.
- 427 Legwaila I.A., Lange E. & Cripps J., 2015. Quarry Reclamation in England: a
428 Review of techniques. Jasmr 4: 55–79.
- 429 Masoero G., Tamietti A., Boano G. & Caprio E., 2016. Apparent constant adult
430 survival of a Sand Martin *Riparia riparia* population in relation to climatic variables.
431 Ardea 104: 253–262.
- 432 Milgrom T., 2008. Environmental aspects of rehabilitating abandoned quarries: Israel
433 as a case study. Landsc. Urban Plan. 87: 172–179.
- 434 Moffatt K.C., Crone E.E., Holl K.D., Schlorff R.W. & Garrison B.A., 2005.
435 Importance of hydrologic and landscape heterogeneity for restoring bank swallow
436 (*Riparia riparia*) colonies along the Sacramento River, California. Restor. Ecol. 13:
437 391–402.
- 438 Morgan R.A., 1979. Sand Martin nest record cards. Bird Study 26: 129–132.
- 439 Muzzi E. & Mongardi G., 2016. Comparison of revegetation techniques on mineral
440 clay soil: analysis of quantitative response of vegetation cover. Ital. J. Agron. 11: 164–
441 170.
- 442 Newton I., 1998. Population Limitation in Birds. Academic Press, London.
- 443 Norman D. & Peach W.J., 2013. Density-dependent survival and recruitment in a
444 long-distance Palaearctic migrant, the Sand Martin *Riparia riparia*. Ibis. 155: 284–296.
- 445 Pannekoek J. & Van Strien, A., 2006. TRIM version 3.54 (Trends & Indices for
446 Monitoring data). Voorburg: Statistics Netherlands.
- 447 Persson C., 1987a. Population processes in south-west Scanian Sand martins
448 (*Riparia riparia*). J. Zool. 1: 671–691.

- 449 Persson C., 1987b. Age structure, sex ratios and survival rates in a south Swedish
450 Sand martin (*Riparia riparia*) population, 1964 to 1984. *J. Zool.* 1: 639–670.
- 451 R Core Team, 2017. R: A language and environment for statistical computing. R
452 Foundation for Statistical Computing. Available at: <https://www.r-project.org/>
- 453 Sanogo S., Fink A.H., Omotosho J.A., Ba A., Redl R. & Ermert V., 2015. Spatio-
454 temporal characteristics of the recent rainfall recovery in West Africa. *Int. J. Climatol.*
455 35: 4589–4605.
- 456 Svensson S., 1986. Number of pairs, timing of egg-laying and clutch size in a
457 subalpine Sand Martin *Riparia riparia* colony, 1968-1985. *Ornis Scand.* 17: 221–229.
- 458 Szép T., 1995. Survival rates of Hungarian Sand Martins and their relationship with
459 Sahel rainfall. *J. Appl. Stat.* 22: 891–904.
- 460 Szép T. & Møller A.P., 1999. Cost of parasitism and host immune defence in the
461 Sand Martin *Riparia riparia*: A role for parent-offspring conflict? *Oecologia* 119: 9–15.
- 462 Szép T., Szabó D.Z. & Vallner J., 2003. Integrated population monitoring of Sand
463 Martin *Riparia riparia* - an opportunity to monitor the effects of environmental
464 disasters along the river Tisza. *Ornis Hungarica* 12: 169–182.
- 465 Wood S.N., 2011. Fast stable restricted maximum likelihood and marginal likelihood
466 estimation of semiparametric generalized linear models. *J. R. Stat. Soc. Ser. B* 73: 3–36.
- 467 Yu G.-A., Disse M., Tong L. & Yu Y., 2015. Evolution of channel networks and
468 morphology of a dryland River under human impacts – A case from Tarim River in
469 Northwest China. In E-proceedings of the 36th IAHR World Congress, 28 June - 3 July
470 p 11. The Hague, the Netherlands.
- 471 Zuquette L.V., Colares O.J. & Pejon O.J., 2002. Environmental degradation related
472 to human activities, Fortaleza metropolitan region, State of Ceará, Brazil. *Bull. Eng.*

473 Geol. Environ. 61: 241–251.

474

475

476

477 **Figure captions**

478

479 Fig. 1

480 Location of the study area, showing the River Po, and the active (circles) and
481 abandoned (squares) quarries.

482

483 Fig. 2

484 Breeding sites of Sand Martins in the quarries.

485

486 Fig. 3

487 Changes course of the River Po from 1922 to 2008. Size of lines is for presentational
488 purposes only (i.e. it is not related to channel width).

489

490 Fig. 4

491 Number of nests per year from 1970 to 2016 counted during the censuses of the
492 breeding colonies in riverbanks (light grey) and in quarries (dark grey) in relation to the
493 amount of sand and gravel extracted from quarries from 1980 and 2016 (dashed) and to
494 the trend of the sinuosity index of the River Po course from 1922 to 2008 (thick solid
495 line). The sinuosity index was calculated for the River Po courses in the years 1922, 1948,
496 1955, 1963, 1978, 1991 and 2008. The solid lines (light grey for riverbanks and dark for
497 quarries) are smooths (edf = 7.06 and 8.11 respectively) fitted from a GAM specifying a
498 Poisson data distribution, and broken lines are the 95% confidence interval around the
499 smoothed trend. Note that the census of the colonies with an exhaustive count of all active
500 nests started in 2000.

501 Fig. 5 (a) Millions of m^3 (Mm³) of extracted construction aggregates in Piedmont from
502 1960 to 1985 (source: Regione Piemonte - Documento di programmazione delle attività
503 estrattive I Stralcio – Relazione -
504 http://www.regione.piemonte.it/industria/cave/dpae_1.htm accessed on 27 October
505 2016).

506 (b) Thousands of m^3 of extracted construction aggregates in the 15 quarries of the
507 study area from 1982 to 2016 (data from the Piemonte Region's database).