

Block-Separation of Variables: a Form of Partial Separation for Natural Hamiltonians

Claudia Maria CHANU and Giovanni RASTELLI

Dipartimento di Matematica, Università di Torino, Torino, via Carlo Alberto 10, Italy

E-mail: claudiamaria.chanu@unito.it, giovanni.rastelli@unito.it

Received August 07, 2018, in final form February 14, 2019; Published online February 23, 2019

<https://doi.org/10.3842/SIGMA.2019.013>

Abstract. We study twisted products $H = \alpha^r H_r$ of natural autonomous Hamiltonians H_r , each one depending on a separate set, called here separate r -block, of variables. We show that, when the twist functions α^r are a row of the inverse of a block-Stäckel matrix, the dynamics of H reduces to the dynamics of the H_r , modified by a scalar potential depending only on variables of the corresponding r -block. It is a kind of partial separation of variables. We characterize this block-separation in an invariant way by writing in block-form classical results of Stäckel separation of variables. We classify the block-separable coordinates of \mathbb{E}^3 .

Key words: Stäckel systems; partial separation of variables; position-dependent time parametrisation

2010 Mathematics Subject Classification: 70H05; 37J15; 70H06

1 Introduction

In [28] Paul Stäckel started the study of (complete) separation of variables in orthogonal coordinates for the Hamilton–Jacobi equation of natural Hamiltonians with N degrees of freedom. The characterization given by Stäckel is both coordinate dependent – involving $N \times N$ Stäckel matrices (see Theorem 2.1 below) – and invariant – involving N quadratic first integrals of the Hamiltonian. In the following years, the theory was widely developed by Levi-Civita [21], Eisenhart [13, 14] and many others (see [18, 19, 24] for more complete references). We point out that complete separation implies the completeness of the separated integral of the Hamilton–Jacobi equation, because of the existence of N independent constants of motion in involution and, consequently, the Liouville integrability of the Hamiltonian system.

Stäckel himself considered in [29] the case of partial separation of variables and obtained a sufficient characterization of it in terms of Stäckel matrices of reduced dimension and of a corresponding number of quadratic first integrals of the Hamiltonian. Partial separation of variables gained much less interest than complete separation. This fact is certainly related with the little use of partial separation in the search of solutions of the Hamilton–Jacobi equation. Moreover, partial separation does not guarantee the existence of complete integrals of the partially separated Hamilton–Jacobi equation, neither Liouville integrability. Nevertheless, as pointed out by [20], in this case the Jacobi method of inversion can, sometimes, produce additional first integrals of the Hamiltonian. Recent papers develop partial separation theory for Hamilton–Jacobi and Schrödinger equations, improving somehow the results of Stäckel, by giving a more detailed characterization of the metric coefficients in partially separable coordinates and by providing further conditions for the separation of the quantum systems [17, 22]. Partial separation of Hamilton–Jacobi and Helmholtz equations on four-dimensional manifolds is briefly considered in [4]. A different approach to non-complete additive separation is represented by non-regular separation which relies on the existence of an additively separated solution on proper submanifolds only [5, 6, 19].

In our study, we shift for the first time the interest from the Hamilton–Jacobi equation to the dynamics of the system. We observe that the partial separation introduced by Stäckel, as well as the complete separation, establishes a dynamical relationship between H and the (partially) separated equations when these are considered as Hamiltonians on submanifolds of the original phase space. Namely, we find that the projection of the orbits of H on these submanifolds, spanned by the separated blocks of coordinates, coincides with the orbits of the separated Hamiltonians on the same submanifolds. The only difference is a position-dependent rescaling of the corresponding Hamiltonian parameters. As a consequence, the dynamics of H can be decomposed into a number of lower-dimensional Hamiltonian systems, allowing in some case a simpler analysis of the original system. The separated blocks of coordinates, considered together, form a N -dimensional coordinate system on the base manifold of H and in these coordinates the N -dimensional metric tensor takes a block-diagonal form. This fact motivates the name we choose for this kind of separation. We prefer to not use the expression *partial separation* since it is already associated with Hamilton–Jacobi theory, which we do not consider here. By shifting the focus from Hamilton–Jacobi theory to the dynamics, we remove the obstruction represented by the completeness of the integral of the Hamilton–Jacobi equation, which is strictly connected with Stäckel theory of complete separation of variables. Indeed, partial separation does not imply the existence of a complete integral, so that the Jacobi method (the construction of a canonical transformation to a trivially integrable Hamiltonian) generally fails. On the contrary, our dynamical interpretation of block separation is basically insensitive to complete or partial separation. We find useful and natural to relate block-separation with the structure of *twisted product* that the Hamiltonian assumes when the separation is possible. This allows us to state our results in a form very close to analogous results in classical Stäckel separation, analogy missing in all the other studies about partial separation. Namely, we can characterize block-separation by introducing “block” versions of celebrated Levi-Civita and Eisenhart equations, and of more recent theorems about complete separation.

The main result of block-separation is the splitting of a N -dimensional Hamiltonian system into lower-dimensional systems (the blocks). Thus, methods of analysis disposable only for low-dimensional systems become available, such as, for example, the topological classification of integrable Hamiltonian systems [3].

In Section 2 we recall the basic theorems about Stäckel complete separation of variables which we rewrite in block-separable form. In Section 3 we define twisted products of Hamiltonians and state some relevant properties of them. In Section 4 we give a dynamical interpretation of Stäckel separation, providing examples of the related properties of time-scaling. In Section 5 we introduce block-separation and our main results about its characterization, with the block-like formulations of Levi-Civita, Eisenhart and other theorems, and we provide an invariant characterization of block separation. The explicit example we deal with is the four-body Calogero system. In Section 6, we characterize, at least with necessary conditions, all the possible block-separable coordinates of \mathbb{E}^3 . Section 7 contains our final considerations and comments.

2 Outline of Stäckel separation

We briefly recall the principal theorems regarding complete separation of the Hamilton–Jacobi equation, see [18] and [1] for further details. The theory of complete additive separation of the Hamilton–Jacobi equation begins with the work of Stäckel [28, 29] about separability of the Hamilton–Jacobi equation in orthogonal coordinates. The Einstein summation convention on equal indices is understood, unless otherwise stated.

Theorem 2.1. *In a given orthogonal coordinate system (q^i) , the Hamilton–Jacobi equation*

$$H = \frac{1}{2} \left(g^{ii} \left(\frac{\partial W}{\partial q^i} \right)^2 + V(q) \right) = c_1, \quad (2.1)$$

admits a complete integral in the separated form,

$$W = \sum_{i=1}^N W_i(q^i, c_1, c_a), \quad a = 2, \dots, N, \quad \det \left(\frac{\partial^2 W}{\partial q^i \partial c_j} \right) \neq 0, \quad c_j = (c_1, c_a),$$

if and only if

- 1) *there exists a $N \times N$ matrix S which is invertible and such that each element of its j -th row depends on q^j only, such that the (g^{ii}) are a row of S^{-1} . The matrix S is called Stäckel matrix.*
- 2) *V is a Stäckel multiplier, i.e., there exist N functions $v_i(q^i)$ such that*

$$V = v_i(q^i)g^{ii}.$$

As a consequence of (1) and (2), there exist $N - 1$ independent quadratic first integrals (K_a) of H ($a = 2, \dots, N$) such that $(c_i) = (c_1, c_a)$ are the constant values of (H, K_a) , where $(K_a) = K_2, \dots, K_N$.

A complete integral of (2.1) is then determined by the N separated equations

$$\left(\frac{dW_r}{dq^r} \right)^2 + v_r = 2S_r^i c_i,$$

where all S_r^i depend ($i = 1, \dots, N$) on the coordinate q^r only.

We remark that completeness for the integral W of the Hamilton–Jacobi equation means that it depends on N parameters (c_j) , constants of motion, such that

$$\det \left(\frac{\partial W}{\partial q^i \partial c_j} \right) \neq 0.$$

Therefore, the (c_j) can be part of a new set of canonical coordinates in which the Hamiltonian flow becomes trivially integrable.

Later, Levi-Civita [21] obtained necessary and sufficient conditions for the complete separability of a generic Hamiltonian in a general coordinate system. For natural Hamiltonians and orthogonal coordinates, $H = \frac{1}{2}g^{ii}p_i^2 + V(q)$, the Levi-Civita equations split into

$$g^{ii}g^{jj}\partial_{ij}g^{kk} - g^{ii}\partial_i g^{jj}\partial_j g^{kk} - g^{jj}\partial_j g^{ii}\partial_i g^{kk} = 0, \quad (2.2)$$

with $i \neq j$ not summed, $i, j, k = 1, \dots, N$, for the components of the metric tensor, and

$$g^{ii}g^{jj}\partial_{ij}V - g^{ii}\partial_i g^{jj}\partial_j V - g^{jj}\partial_j g^{ii}\partial_i V = 0,$$

with $i \neq j$ not summed, $i, j = 1, \dots, N$, called *Bertrand–Darboux equations* and whose solution (for g^{ii} satisfying (2.2)) is V in the form of a Stäckel multiplier.

In [13], Eisenhart provided a geometrical characterization of complete separation of variables in orthogonal coordinates introducing Killing tensors and, later, determined all the possible

orthogonal separable coordinate systems of \mathbb{E}^3 . He determined eleven types of orthogonal separable coordinate systems, which are described, for example, in [24]. The fundamental Eisenhart equations

$$\partial_i \lambda_j = (\lambda_i - \lambda_j) \partial_i \ln |g^{jj}|, \quad i, j = 1, \dots, N$$

characterize the eigenvalues of a Killing tensor: let K be a 2-tensor with eigenvalues (λ_i) and eigenvectors (∂_i) , then K is a Killing tensor if and only if the Eisenhart equations are satisfied. The integrability conditions of the Eisenhart equations for a Killing tensor with simple eigenvalues coincide with the Levi-Civita equations (2.2).

The coordinate systems can be geometrically understood as foliations of hypersurfaces called *coordinate webs*. The separability of a coordinate web can be characterized by a single *characteristic Killing tensor*, i.e., a symmetric Killing 2-tensor with pointwise simple eigenvalues and normally integrable eigenvectors, which determine in each point of the space (up to possible singular sets of zero measure) the basis of coordinate vectors [1]. We recall that a symmetric *Killing 2-tensor* K is defined by the equivalent equations

$$[g, K] = 0, \quad \nabla_{(i} K_{j)k} = 0,$$

where $[\cdot, \cdot]$ is the Schouten bracket and ∇ is the covariant derivative with respect to the metric g .

Theorem 2.2. *The Hamilton–Jacobi equation of a natural Hamiltonian with scalar potential V is completely separable in an orthogonal coordinate web if and only if there exists a characteristic Killing 2-tensor K whose eigenvectors are normal to the foliations of the web and such that $d(KdV) = 0$.*

The last condition is equivalent to say that V is a Stäckel multiplier in the orthogonal coordinates associated with K . In the formula of Theorem 2.2 K is considered as a linear operator mapping one-forms into one-forms.

Necessary and sufficient conditions for a Killing tensor to be characteristic are given in [10].

Theorem 2.3 (Tonolo–Schouten–Nijenhuis [26, 27, 30]). *A 2-tensor K with real distinct eigenvalues has normal eigenvectors if and only if the following conditions are satisfied*

$$N_{[ij}^l g_{k]l} = 0, \quad N_{[ij}^l K_{k]l} = 0, \quad N_{[ij}^l K_{k]m} K_l^m = 0,$$

where N_{jk}^i are the components of the Nijenhuis tensor of K_j^i defined by

$$N_{jk}^i = K_l^i K_{[j,k]}^l + K_{[j}^l K_{k],l}^i.$$

An equivalent formulation of Theorem 2.2 involves N independent quadratic first integrals, therefore, N Killing 2-tensors, instead of a single characteristic Killing tensor [1].

Theorem 2.4. *The natural Hamiltonian H is separable in some orthogonal coordinates (q^i) , if and only if*

- 1) *there exist other $N - 1$ independent quadratic in the momenta functions K_a such that*

$$\{H, K_a\} = 0,$$

- 2) *the Killing two-tensors (k_a) associated with (K_a) are simultaneously diagonalized with pointwise independent eigenvalues and have common normally integrable eigenvectors.*

It follows that $\{K_a, K_b\} = 0$.

The original formulation of the theorem requires the reality of the eigenvalues, however, this request is unnecessary if one accepts also complex separable coordinates [11].

3 Twisted products of Hamiltonians

Let

$$M = \times_{r=1}^n M_r,$$

be the product of n Riemannian or pseudo-Riemannian manifolds (M_r, g_r) of dimension n_r , so that $\dim(M) = n_1 + \dots + n_n = N$, and let α^r be n non zero functions on M . The manifold M with metric tensor

$$G = \alpha^1 g_1 + \dots + \alpha^n g_n,$$

is a Riemannian or pseudo-Riemannian manifold called *twisted product manifold* of the (M_r) with twist functions (α^r) [23]. In the case when $\alpha^1 = 1$ and $\alpha^2, \dots, \alpha^n$ are functions on M_1 , the manifold M is called *warped product*. We extend to functions on T^*M and T^*M_r the concept of twisted and warped products in a natural way. In particular, for each r we consider natural Hamiltonians

$$H_r = \frac{1}{2} (g_r^{r_i r_j} p_{r_i} p_{r_j} + V_r(q^{r_i})),$$

where (q^{r_i}, p_{r_i}) , $i = 1, \dots, n_r$, are canonical coordinates on T^*M_r , we construct *twisted product*

$$H = \alpha^r H_r = \alpha^1 H_1 + \dots + \alpha^n H_n,$$

of the H_r with twist functions $\alpha^r \in \mathcal{F}(M)$.

Then, H is a natural Hamiltonian on T^*M with metric G and potential $V = \alpha^r V_r$.

The manifold M is naturally endowed with block-diagonal coordinates (q^{r_i}) such that the components of G are in the form

$$G^{r_i r_j} = \alpha^r g_r^{r_i r_j}, \quad G^{r_i s_j} = 0, \quad s \neq r,$$

we call these coordinates *twisted coordinates*.

We have now $n + 1$ Hamiltonians, each one with its own Hamiltonian parameter. We call t the Hamiltonian parameter of H and τ_r the Hamiltonian parameter of H_r . From Hamilton's equations we get

$$\frac{dq^{r_i}}{dt} = \frac{\partial H}{\partial p_{r_i}} = \alpha^r \frac{\partial H_r}{\partial p_{r_i}} = \alpha^r \frac{dq^{r_i}}{d\tau_r},$$

and

$$\frac{dp_{r_i}}{dt} = -\frac{\partial H}{\partial q^{r_i}} = -\alpha^r \frac{\partial H_r}{\partial q^{r_i}} - H_s \frac{\partial \alpha^s}{\partial q^{r_i}} = \alpha^r \frac{dp_{r_i}}{d\tau_r} - H_s \frac{\partial \alpha^s}{\partial q^{r_i}}.$$

Therefore, the relation between the Hamiltonian vector fields X_H of H and X_r of H_r is

$$X_H = \bar{X}_1 + \dots + \bar{X}_N - H_s \frac{\partial \alpha^s}{\partial q^{r_i}} \frac{\partial}{\partial p_{r_i}},$$

where

$$\bar{X}_r = \alpha^r X_r, \quad r \text{ not summed},$$

is the rescaled Hamiltonian vector field of H_r .

4 Stäckel systems as twisted Hamiltonians

In this section we study Stäckel systems in their nature of twisted Hamiltonians. Our aim is to enlighten the relations among the dynamics of the N -dimensional Hamiltonian system determined by H and the dynamics of the N one-dimensional Hamiltonians H_r , so that $n_1 = \dots = n_N = 1$, determined by the separated equations of H . Separation of variables for the Hamilton–Jacobi equation of H will not be of primary interest in what follows. See Section 2 for definitions of Stäckel matrix, Stäckel multiplier and separated equations.

Let be $H_r = \frac{1}{2}(p_r^2 + V_r)$ and assume that α^r are a row (say the first one) of the inverse of a Stäckel matrix S for given coordinates (q^r) . Then, the twisted product $H = \alpha^r H_r$ admits separation of variables and we have the separated equations

$$H_r = S_r^i c_i, \quad (4.1)$$

where c_i are N constants, corresponding to the N constants of motion K_i of $H = K_1$ and

$$K_a = (S^{-1})_a^r H_r, \quad a = 2, \dots, N.$$

The Hamilton–Jacobi complete separated integral $W = W_1(q^1, c_i) + \dots + W_N(q^N, c_i)$ is given by integration of

$$\left(\frac{dW_r}{dq^r} \right)^2 + V_r = 2S_r^i c_i.$$

The Hamilton’s equations of H , in time t , are

$$\begin{aligned} \dot{q}^r &= \alpha^r p_r, \\ \dot{p}_r &= -\partial_r \alpha^i H_i - \alpha^i \partial_r H_i = -\partial_r \alpha^i S_i^j c_j - \frac{1}{2} \alpha^r \frac{d}{dq^r} V_r, \end{aligned} \quad (4.2)$$

where we use the separated equations (4.1) to replace H_r along the integral curves. Since $\alpha^r = (S^{-1})_1^r$ is a row of the inverse of S , we have

$$\partial_r \alpha^r S_i^j = \partial_r (\alpha^i S_i^j) - \alpha^i \partial_r S_i^j = \partial_r (\delta_1^j) - \alpha^r \frac{d}{dq^r} S_r^j = -\alpha^r \frac{dS_r^j}{dq^r}, \quad (4.3)$$

and the same for all other elements of the rows of S^{-1} . Then, we can write

$$\dot{p}_r = \alpha^r \frac{d}{dq^r} \left(c_j S_r^j - \frac{1}{2} V_r \right). \quad (4.4)$$

Let γ_P be the integral curve of X_H containing a point $P \in T^*M$. We consider the values $c_i = K_i(P)$ and we introduce the Hamiltonians

$$\tilde{H}_r = H_r - c_j S_r^j,$$

with Hamiltonian parameters $\tilde{\tau}_r$. We can write the equations of Hamilton for \tilde{H}_r as

$$\frac{d}{d\tilde{\tau}_r} q^r = p_r = \partial^r \tilde{H}_r, \quad (4.5)$$

$$\frac{d}{d\tilde{\tau}_r} p_r = \frac{d}{dq^r} \left(c_j S_r^j - \frac{1}{2} V_r \right) = -\partial_r \tilde{H}_r. \quad (4.6)$$

Therefore,

Proposition 4.1. *For each orbit of H , the N Hamiltonian vector fields $X_{\tilde{H}_r}$ of the \tilde{H}_r are proportional to the components with respect to (∂_r, ∂^r) of the Hamiltonian vector field X_H of H , where α^r are the proportionality functions.*

Proof. Let be

$$X_{\tilde{H}_r} = \frac{\partial \tilde{H}_r}{\partial p_r} \partial_r - \frac{\partial \tilde{H}_r}{\partial q^r} \partial^r,$$

and

$$(X_H)_r = \dot{q}^r \partial_r + \dot{p}_r \partial^r.$$

Due to (4.5) and (4.6) we can write (4.2) and (4.4) as

$$\dot{q}^r = \alpha^r \partial^r \tilde{H}_r, \quad \dot{p}_r = -\alpha^r \partial_r \tilde{H}_r,$$

and it follows immediately

$$(X_H)_r = \alpha^r X_{\tilde{H}_r}. \quad \blacksquare$$

Remark 4.2. After Proposition 4.1 we can put

$$\alpha^r = \frac{d\tilde{\tau}_r}{dt},$$

and consider the twist functions as determining position-dependent time-scalings between the Hamiltonian parameters t and $\tilde{\tau}_r$.

From Proposition 4.1 follows the important result

Proposition 4.3. *The projection of each orbit of H on each coordinate manifold (q^r, p_r) coincides with the orbit of $\tilde{H}_r = H_r - c_j S_r^j$.*

Remark 4.4. The Lagrange equations of the dynamics of \tilde{H}_r , expressed in times $\tilde{\tau}_r$ are

$$\frac{d^2 q^r}{d\tilde{\tau}_r^2} = \frac{d}{dq^r} \left(c_j S_r^j - \frac{1}{2} V_r \right).$$

Remark 4.5. Observe that

$$\tilde{H} = \alpha^r \tilde{H}_r = H - c_1, \quad \tilde{K}_j = (S^{-1})_j^r \tilde{H}_r = K_j - c_j, \quad j \neq 1,$$

i.e., the Stäckel systems associated with H and \tilde{H} coincide up to additive constants. To the constants (c_i) for (H, K_a) correspond the constants $(\tilde{c}_i = 0)$ for (\tilde{H}, \tilde{K}_a) .

Example 4.6. *Twisted product of pendula.* In order to show the effect of the time-scaling described above, we consider the twisted product of the following three one-dimensional Hamiltonians

$$H_i = \frac{1}{2} (p_i^2 - \cos q^i), \quad i = 1, 2, \quad H_3 = \frac{1}{2} p_3^2,$$

corresponding to two pendula and a purely inertial term, coupled together by the first row of the inverse of the 3×3 matrix

$$S = \begin{pmatrix} 2 & 1 + q^1 & 2(q^1)^2 + 2 \\ 3 & q^2 & (q^2)^3 + 2 \\ 4 & q^3 & (q^3)^2 + 1 \end{pmatrix},$$




Figure 1. Projections of the orbits of H and \tilde{H}_1 on (p_1, q^1) .




Figure 2. Projections of the orbits of H and \tilde{H}_1 on (t, p_1, q^1) and (τ_1, p_1, q^1) .

which is a Stäckel matrix in a neighborhood of the origin, since the Taylor expansion up to the second order terms of its determinant Δ around $(0, 0, 0)$ is $\Delta = 5 + 5q^1 - 6q^2 + 2q^3$. The elements of the matrix S^{-1} are therefore quite complicated rational functions that we do not need to compute explicitly but make the coupling of the H_r suitable to enhance the effect of the time-scaling. We take as (α^r) the first row of S^{-1} and consider

$$H = \alpha^r H_r = c_1.$$

The quadratic first integrals of H are determined by the remaining rows of (S^{-1})

$$K_a = (S^{-1})_a^r H_r = c_a, \quad a = 2, 3.$$

We already know from the previous section that, despite the complicated expression of the coupling terms (α^r) , the relation among the dynamics of H and of the separated Hamiltonians $\tilde{H}_r = H_r - c_a S_r^a$ reduces to a simple position-dependent time scaling.

We plot the numerical evaluation of the systems of Hamiltonian H and

$$\tilde{H}_1 = H_1 - 2c_1 - (1 + q^1)c_2 - 2((q^1)^2 + 1)c_3$$

respectively, and project the orbits on (q^1, p_1) , obtaining, for the initial conditions $p_1 = 0$, $p_2 = 0$, $p_3 = 0$, $q^1 = 0.2$, $q^2 = -0.2$, $q^3 = 0$ and consequently $c_1 = H = 0.09494666248$, $c_2 = 0.0916913483$, $c_3 = -0.3797866499$, the graphs in Fig. 1, where we see that the orbits on (q^1, p_1) of the two systems coincide. But, if we include the dependence on the different Hamiltonian parameters (denoted in both the graphs as t), we get Fig. 2 and we can see how the dependence on the Hamiltonian parameters can be extremely different in the two cases.

Example 4.7. *Twisted product with constant coefficients of harmonic oscillators.* Taking twisted products of Hamiltonians seems an interesting way to establish an interaction among Hamiltonian systems. An example, even if somehow trivial, is provided by the twisted product with constant coefficients of harmonic oscillators. Let

$$H_i = \frac{1}{2}(p_i^2 + \omega_i^2(q^i)^2), \quad i = 1, \dots, n$$

be a finite set of harmonic oscillators. Let

$$H = \alpha^i H_i, \quad \alpha^i \in \mathbb{R}^+,$$

be their twisted product with constant twist functions. The H_i are all constants of motion of H and there is actually no interaction among them. However, some effect of the twisted product is nevertheless evident. The Hamilton equations of H are

$$\begin{aligned} \frac{dq^i}{dt} &= \alpha^i \frac{dq^i}{d\tau_i} = \alpha^i p_i, \\ \frac{dp_i}{dt} &= -\alpha^i \frac{dp_i}{d\tau_i} = -\alpha^i \omega_i^2 q^i. \end{aligned}$$

The general solution of these equations is

$$q^i(t) = c_1^i \sin(\alpha^i \omega_i t) + c_2^i \cos(\alpha^i \omega_i t), \quad c_j^i \in \mathbb{R}.$$

We see that, for example, the choice $\alpha^i = k/\omega_i$, for any real positive k , determines a time-scaling that gives to all the oscillators the same frequency k with respect to t (as well as any other real positive frequency for different choices of k for each i). Namely, the rescaling is in this case

$$t = \sum_{i=1}^n \frac{1}{\alpha^i} \tau_i + t_0, \quad \text{or} \quad \tau_i = \alpha^i t + \tau_i^0.$$

The frequency of each oscillator H_i with respect to its own Hamiltonian parameter τ_i remains clearly ω_i .

5 Block-separation

The results of the previous section can be generalized as follows, leading to a kind of partial separation of variables that we call block-separation.

Let M be a N -dimensional manifold. Let us consider a partition of a coordinate system on M organized as follows. For $n \leq N$, consider for each integer $r = 1, \dots, n$ the integers n_r such that

$$N = n_1 + \dots + n_n.$$

The coordinate system is therefore composed of n blocks, and for each $r \leq n$ we have an r -block of coordinates that we denote as $(q^{r_1}, \dots, q^{r_{n_r}})$. We call M_r the manifold spanned by the r -block of coordinates. Consider T^*M with the conjugate r -block momenta $(p_{r_1}, \dots, p_{r_{n_r}})$.

Let us consider

$$H_r = \frac{1}{2} g_r^{r_i r_j} p_{r_i} p_{r_j} + V_r(q^{r_k}),$$

and the n block-separated equations

$$H_r = S_r^a c_a, \quad a = 1, \dots, n,$$

where we assume that $g_r^{r_i r_j}$, S_r^a and V_r are functions of coordinates of the r -block only and c_a are constants. If we assume that the $n \times n$ matrix (S_r^a) is invertible, and we call it *block-Stäckel matrix*, then we can write the n equations

$$(S^{-1})_a^r H_r = c_a. \quad (5.1)$$

We denote $\alpha^r = (S^{-1})_1^r$ and call

$$H = \alpha^r H_r, \quad K_a = (S^{-1})_a^r H_r, \quad a = 2, \dots, n.$$

Hence, H is in the form of twisted product and it is a natural Hamiltonian whose metric tensor G is block-diagonalized, with components

$$G^{r_i r_j} = \alpha^r g_r^{r_i r_j}, \quad G^{r_i s_j} = 0, \quad s \neq r,$$

and whose scalar potential has the form

$$V = \alpha^r V_r,$$

while the scalar potentials in K_a are

$$W_a = (S^{-1})_a^r V_r. \quad (5.2)$$

A necessary condition for the procedure of above is that the (5.1) are indeed constants of motion of H .

Proposition 5.1. *The n functions (H, K_a) are all independent, quadratic in the momenta and pairwise in Poisson involution.*

Proof. Since S is a block-Stäckel matrix, in analogy to (4.3) we have

$$\partial_{r_i} \alpha_a^s S_s^j = \partial_{r_i} (\alpha_a^s S_s^j) - \alpha_a^s \partial_{r_i} S_s^j = \partial_{r_i} (\delta_1^j) - \alpha_a^r \partial_{r_i} S_r^j = -\alpha_a^r \partial_{r_i} S_r^j,$$

where r is not summed and $\alpha_a^r = (S^{-1})_a^r$. Then, from the definition of Poisson bracket and of block-separated coordinates, we get the statement. ■

For the dynamics of H , an analogue of Proposition 4.3 holds.

Proposition 5.2. *The dynamics of H coincides in each r -block with the dynamics of*

$$\tilde{H}_r = H_r - c_a S_r^a,$$

*up to a reparametrization of the Hamiltonian parameter given by $\alpha^r = d\tilde{\tau}_r/dt$. So, the projections of the orbits of H on each T^*M_r coincide with the orbits of \tilde{H}_r .*

Proof. The proof follows the same reasoning of the proof of Proposition 4.3. It follows that, denoting with $(X_H)_r$ the r -block component of the Hamiltonian vector field X_H of H ,

$$(X_H)_r = \dot{q}^{r_i} \partial_{r_i} + \dot{p}_{r_i} \partial^{r_i},$$

we have

$$(X_H)_r = \alpha^r X_{\tilde{H}_r},$$

where $X_{\tilde{H}_r}$ is the Hamiltonian vector field of \tilde{H}_r . Therefore, if $(X_H)_r$ is tangent to any submanifold $f \subseteq T^*M$, then also $X_{\tilde{H}_r}$ is, and vice-versa. Hence, just as for the Stäckel systems, the dynamics of H is determined in each r -block, up to reparametrizations of the Hamiltonian parameter, by the dynamics of the \tilde{H}_r , with the difference that the \tilde{H}_r are no longer one-dimensional. ■

In this way, the time-independent dynamics of H can be exactly decomposed into the n lower-dimensional separated dynamics of Hamiltonians \tilde{H}_r . The \tilde{H}_r share with the H_r , factors of the twisted product H , the same inertial terms, while the scalar potential is modified by the addition of the term $-c_a S_r^a$.

Partial separation of Hamilton–Jacobi equation was introduced by di Pirro in [12] and generalized by Stäckel in [29]. He introduced the $n \times n$ matrix S and his results are analogue to our Proposition 5.1. Stäckel obtained sufficient conditions for partial separation of the Hamilton–Jacobi equation of natural Hamiltonians. His work has been extended more recently in [17] and [22], including the study of partial separation of the Schrödinger equation, obtaining again sufficient conditions for partial separation, and a more detailed form of the components of the metric tensor in partially separable coordinates. We remark that, by introducing twisted products, our characterization of block-separation provides necessary and sufficient conditions for it, in analogy with Stäckel theory of complete separation. We do not make here a strict comparison between our results and those of [29] and [17], since these last results are strictly related to Hamilton–Jacobi theory and there is no consideration of the dynamical relations among the N -dimensional Hamiltonian and the separated Hamiltonians, which is our main interest. The detailed characterization of the partially separable metric’s components in [17] should eventually coincide with a similar characterization of block-separable metrics. We do not consider here the distinction between linear and quadratic in the momenta first integrals (from linear first integrals one can always obtain quadratic ones). It is remarkable that in the last century very few works have been devoted to partial separation of Hamilton–Jacobi equation. This is understandable when one considers that Hamilton–Jacobi theory is of not easy application, apart the simplest cases, even when completely separated integrals of the Hamilton–Jacobi equations do exist. Some applications of partially separated integrals of the Hamilton–Jacobi equation, in order to generate new *possible* first integrals of the Hamiltonian, are presented in [20] and [25]. Our approach based upon the block-separated dynamics, instead of the partially separated Hamilton–Jacobi equation, appears to be completely new and could be more suitable for applications of the theory, particularly in the analysis of systems with many degrees of freedom.

5.1 Block-Eisenhart and block-Levi-Civita equations

We can see, with some surprise, that the characterisation of block-separation includes tools developed for Stäckel complete separation. Indeed, we can formulate classical results by Eisenhart and Levi-Civita in block form. If we assume that (q^i) are twisted coordinates (q^{r_i}) , then for G we have

$$G_{r_i r_j} = \frac{1}{\alpha^r} g_{r_i r_j}^r, \quad r \text{ not summed,}$$

so that

$$G_{r_k a} G^{a s_j} = \delta_{r_k}^{s_j}.$$

Moreover, if we assume that the coordinates (q^{r_i}) are block-separated

$$k_a^{r_i r_j} = (S^{-1})_a^r g_r^{r_i r_j},$$

where k_a^{bc} are the components of the 2-tensor associated with K_a . Then,

$$(k_a)_{r_j}^{r_i} = \frac{1}{\alpha^r} (S^{-1})_a^r \delta_{r_j}^{r_i},$$

and we can consider the functions

$$\lambda_a^r = \frac{1}{\alpha^r} (S^{-1})_a^r, \quad r \text{ not summed},$$

as the analogue of eigenvalues of Killing tensors k_a in Stäckel theory. They are indeed the eigenvalues with respect to G of the Killing tensors (k_a) associated with the first integrals (K_a) .

Proposition 5.3. *In block-separated coordinates, we have that*

$$\{H, K_a\} = 0,$$

if and only if the block-Eisenhart equations

$$\partial_{r_k} \lambda_a^s = (\lambda_a^r - \lambda_a^s) \partial_{r_k} \ln |\alpha^s|, \quad (5.3)$$

hold, with $r, s = 1, \dots, n$, for all r_i, s_j in the respective separated blocks, and (5.2) hold.

Proof. By expanding

$$\{H, K_a\} = 0,$$

in block-separable coordinates, collecting homogeneous terms in the momenta and dividing by $\alpha^r \alpha^s$, from the higher order terms in the momenta we have,

$$(\lambda_a^s - \lambda_a^r) \partial_{r_k} g^{s_i s_j} + g^{s_i s_j} [\partial_{r_k} \lambda_a^s - (\lambda_a^r - \lambda_a^s) \partial_{r_k} \ln |\alpha^s|] = 0, \quad (5.4)$$

for all r, s, r_k, s_i, s_j in the respective blocks. If $r = s$ the equations become

$$g^{s_i s_j} \partial_{s_k} \lambda_a^s = 0,$$

and, if $r \neq s$, then $\partial_{r_k} g^{s_i s_j} = 0$. Hence, (5.4) is equivalent to

$$g^{s_i s_j} [\partial_{r_k} \lambda_a^s - (\lambda_a^r - \lambda_a^s) \partial_{r_k} \ln |\alpha^s|] = 0.$$

If $g^{s_i s_j} = 0$, the equations are identically satisfied, otherwise, we have (5.3). Since not all the $g^{s_i s_j}$ are zero, we have the statement. The first-order terms in the momenta vanish if and only if (5.2) hold. \blacksquare

By definition of the λ_a^s and of the α^s , we have the equations

$$S_r^s \alpha^r = \delta_1^s, \quad S_r^s \lambda_a^r \alpha^r = \delta_a^s.$$

We observe that, after putting $\alpha^r = g^{rr}$, the previous equations are identical to the relations typical of Stäckel systems. In the same way, the block-Eisenhart equations become the standard Eisenhart equations.

Proposition 5.4. *The block-Eisenhart equations (5.3) hold if and only if (S_a^r) is a block-Stäckel matrix.*

As for the Stäckel systems, the block-Levi-Civita equations can be considered as the integrability conditions of the block-Eisenhart equations. The derivation is essentially the same as in [1].

It is therefore straightforward to see that the block-diagonalized coordinates (q^1, \dots, q^N) are block-separated for the Hamiltonian H if and only if the block-Levi-Civita equations

$$\alpha^r \alpha^s \partial_{r_i s_j} \alpha^m - \alpha^r \partial_{r_i} \alpha^s \partial_{s_j} \alpha^m - \alpha^s \partial_{s_j} \alpha^r \partial_{r_i} \alpha^m = 0,$$

are satisfied, where the coordinates r_i, s_j are in different blocks and $m = 1, \dots, n$. The scalar potential V is already in the form of a block-Stäckel multiplier, thanks to the form of H , and satisfies

$$\alpha^r \alpha^s \partial_{r_i s_j} V - \alpha^r \partial_{r_i} \alpha^s \partial_{s_j} V - \alpha^s \partial_{s_j} \alpha^r \partial_{r_i} V = 0.$$

See (2.2) for a comparison.

5.2 Invariant characterization

As in Stäckel theory, we can use the previous results for an invariant characterization of block-separation. Therefore, we have the analogue of the Eisenhart–Kalnins–Miller–Benenti theorem [1].

Proposition 5.5. *The twisted Hamiltonian $H = \alpha^r H_r$, $r = 1, \dots, n$, is block-separated in twisted coordinates (q^{r_i}) if and only if*

- 1) *there exist other $n - 1$ independent quadratic in the momenta functions $K_a = \alpha_a^r H_r$ such that*

$$\{H, K_a\} = 0,$$

- 2) *the Killing two-tensors (k_a) are simultaneously block-diagonalized and have common normally integrable eigenspaces*

Moreover, it follows that $\{K_a, K_b\} = 0$.

Remark 5.6. The proof of the previous statement relies on the assumption that each H_r depends only on coordinates in T^*M_r . Otherwise, if the metric tensor of H is only block-diagonal in (q^{r_i}) , conditions 1) and 2) are only necessary.

Consequently, the strategy for finding block-separated coordinates of a given N -dimensional natural Hamiltonian H is the following

- Find a number $n \leq N$ of independent quadratic first integrals (K_a) of H in involution among themselves, whose associated Killing tensors (k_a) admit common block-diagonalized normally integrable eigenspaces. The number n corresponds to the number of blocks. The dimension of the common eigenspaces equals the dimension of each block.
- At this point, we have block-diagonalized coordinates and we can write $H = \alpha^r H_r$ for some functions H_r , the α^r being determined by the block-Stäckel matrix determined by (k_a) .
- The last step consists in checking that each H_r depends only on coordinates in T^*M_r .

This is indeed the procedure applied in Example 5.9.

We recall (see [15]) that in a Riemannian manifold any symmetric 2-tensor is pointwise diagonalizable, that is there exist n vector fields E_i , pointwise orthogonal eigenvectors of k , such that $k = \sum_i \lambda^i E_i \otimes E_i$. The λ^i are the roots of the characteristic equation $\det(k - \lambda g) = 0$ and the geometric multiplicity of each eigenvalue λ^i coincides with its algebraic multiplicity.

In our case, the algebraic multiplicity of each λ_a^r is n_r at least (for some a , we can have $\lambda_a^r = \lambda_a^s$, and the algebraic multiplicity is in this case $n_r + n_s$).

Proposition 5.5 provides an invariant characterization of block-separable coordinates in terms of what we can call *block-Killing–Stäckel algebras* generated by the (k_a) . See [1, 2] for a definition of Killing–Stäckel algebras.

If we assume that for N -dimensional 2-tensors T_λ^κ to each eigenvalue of algebraic multiplicity n_r it corresponds a space of n_r linearly independent covariant eigenvectors $\{X_a\}$, we can consider the $(N - n_r)$ -dimensional space of vectors E_{N-n_r} such that $\langle X_a, E^b \rangle = 0, \forall E^b \in E_{N-n_r}$. We assume that E_{N-n_r} is a regular distribution of constant rank $N - n_r$.

The necessary and sufficient condition for the integrability of the distributions E_{N-n_r} is given by the Haantjes theorem [16].

Theorem 5.7. *Let T_κ^λ be a tensor such that to each root with multiplicity n_r of the characteristic equation belongs a set of n_r linearly independent covariant eigenvectors. Then the E_{N-n_r} determined by these vectors are integrable if and only if*

$$H_{\nu\sigma}^\kappa T_\mu^\nu T_\lambda^\sigma - 2H_{\nu[\lambda}^\sigma T_{\mu]}^\nu T_\sigma^\kappa + H_{\mu\lambda}^\nu T_\sigma^\kappa T_\nu^\sigma = 0,$$

where

$$H_{\mu\lambda}^\kappa = 2T_{[\mu}^\nu \partial_{|\nu|} T_{\lambda]}^\kappa - 2T_\nu^\kappa \partial_{[\mu} T_{\lambda]}^\nu,$$

is the so-called Haantjes tensor of T .

Therefore, in analogy with the characterization of the Stäckel separable coordinate systems, we have

Proposition 5.8. *A natural Hamiltonian admits block-separable coordinates only if its metric tensor admits a symmetric Killing 2-tensor T satisfying the Haantjes theorem and its scalar potential V satisfies $d(TdV) = 0$. We call T the characteristic tensor of the block-separable coordinates.*

The coordinates are therefore divided into n blocks, where n is the number of the pointwise different eigenvalues of T , the dimension of each block equals the multiplicity of the corresponding eigenspace.

An analogue result about characteristic Killing tensors of Stäckel systems is given in [10] making use of theorems due to Tonolo, Schouten and Nijenhuis (see Section 2). The main difference is due to the fact that, in that case, the eigenvalues of the tensors are simple.

The characterization of block-separation via Killing 2-tensors is extremely powerful in view of applications. For example, in any Riemannian manifold of constant curvature, all Killing tensors, of any order, are linear combinations with constant coefficients of symmetric products of Killing vectors, i.e., isometries [18]. Many common computer-algebra softwares include specific commands for the determination of Killing tensors of Riemannian manifolds.

Example 5.9. *The four-body Calogero system.* The N -body Calogero system is the Hamiltonian system of N points of unitary mass on a line, subject to the interaction

$$V = \sum_{i=1}^{N-1} \sum_{j=i+1}^N (x_i - x_j)^{-2}.$$

The Hamiltonian is therefore

$$H_{(N)} = \frac{1}{2} \sum_{i=1}^N p_i^2 + V,$$

in Cartesian coordinates (x^i) and it is known to be maximally superintegrable for any N and multiseparable for $N < 4$ [2, 31].

In constant curvature manifolds, quadratic first integrals $K_a = \frac{1}{2} k_a^{ij} p_i p_j + W_a$ of natural Hamiltonians can be determined in a systematic way. Indeed, see for example [2], K_a is a first integral of H if and only if the functions k_a^{ij} are the components of a symmetric Killing 2-tensor k_a and $dW_a = k_a dV$. It follows that a necessary condition on k_a for K_a to be a first integral of H is $d(k_a dV) = 0$. As in any constant curvature manifold, the generic Killing 2-tensor of \mathbb{E}^4 is a linear combination (depending on 50 real parameters) of symmetric product of pairs of Killing vectors. By imposing the condition $d(k_a dV) = 0$ to the elements of this space, one finds that $H_{(4)}$ admits, other than the Hamiltonian, only two quadratic independent first integrals in

involution, and not the three necessary for standard Stäckel separation. The two quadratic first integrals of $H_{(4)}$ can be chosen as follows

$$K_a = \frac{1}{2} k_a^{ij} p_i p_j + W_a,$$

where W_a are suitable functions that we will make explicit later on, and

$$k_1^{ii} = \sum_{j,k} x^j x^k, \quad j, k = 1, \dots, 4, \quad j < k, \quad j, k \neq i,$$

$$k_1^{rs} = \frac{1}{2} \left((x^l)^2 + (x^m)^2 + x^r x^s - \sum_{j < k} x^j x^k \right),$$

where $j, k = 1, \dots, 4$, l, m, r, s all different,

$$k_2^{ii} = \sum_{j \neq i} (x^j)^2, \quad k_2^{ij} = -x^i x^j, \quad i \neq j.$$

The eigenvalues of k_1 are

$$\{0, (x^1)^2 + (x^2)^2 + (x^3)^2 + (x^4)^2\}$$

with multiplicity 1, and

$$\sum_{i,j} x^i x^j - \frac{1}{2} ((x^1)^2 + (x^2)^2 + (x^3)^2 + (x^4)^2), \quad i, j = 1, \dots, 4, \quad i < j,$$

with multiplicity 2.

The eigenvalues of k_2 are 0, of multiplicity 1, and

$$(x^1)^2 + (x^2)^2 + (x^3)^2 + (x^4)^2,$$

of multiplicity 3. Since the tensors of components k_1^{ij} and k_2^{ij} commute as linear operators and the metric is positive definite, they can be diagonalized simultaneously in some coordinate system.

By using some properties of the eigenvalues of Killing tensors [9], one finds that these coordinates $(r, \phi_1, \phi_2, \phi_3)$ are spherical and determined by the consecutive transformations [7]

$$z^1 = 2^{-1/2} (x^1 - x^2),$$

$$z^2 = 6^{-1/2} (x^1 + x^2 - 2x^3),$$

$$z^3 = 12^{-1/2} (x^1 + x^2 + x^3 - 3x^4),$$

$$z^4 = 2^{-1} (x^1 + x^2 + x^3 + x^4),$$

and

$$z^4 = r \cos \phi_1,$$

$$z^3 = r \sin \phi_1 \cos \phi_2,$$

$$z^2 = r \sin \phi_1 \sin \phi_2 \cos \phi_3,$$

$$z^1 = r \sin \phi_1 \sin \phi_2 \sin \phi_3.$$

The scalar potential V of $H_{(4)}$ becomes in these coordinates

$$V = \frac{1}{r^2 \sin^2 \phi_1} f(\phi_2, \phi_3),$$

where $f(\phi_2, \phi_3)$ is a rather complicated rational function of trigonometric functions of ϕ_1, ϕ_2 .
Therefore, the Hamiltonian becomes

$$H_{(4)} = \alpha^1 H_1 + \alpha^2 H_2 + \alpha^3 H_3,$$

with

$$\alpha^1 = 1, \quad \alpha^2 = \frac{1}{r^2}, \quad \alpha^3 = \frac{1}{r^2 \sin^2 \phi_1},$$

$$H_1 = \frac{1}{2} p_r^2 + V_1, \quad H_2 = \frac{1}{2} p_{\phi_1}^2 + V_2, \quad H_3 = \frac{1}{2} \left(p_{\phi_2}^2 + \frac{1}{\sin^2 \phi_2} p_{\phi_3}^2 \right) + V_3,$$

where

$$V = \alpha^i V_i,$$

with $V_1 = 0, V_2 = 0, V_3 = f(\phi_2, \phi_3)$. Moreover,

$$K_1 = H_2 + \frac{1 - 2 \sin^2 \phi_1}{\sin^2 \phi_1} H_3, \quad K_2 = H_2 + \frac{1}{\sin^2 \phi_1} H_3.$$

So that the inverse of the block-Stäckel matrix is

$$S^{-1} = \begin{pmatrix} 1 & \frac{1}{r^2} & \frac{1}{r^2 \sin^2 \phi_1} \\ 0 & 1 & \frac{1 - 2 \sin^2 \phi_1}{\sin^2 \phi_1} \\ 0 & 1 & \frac{1}{\sin^2 \phi_1} \end{pmatrix},$$

and the block-Stäckel matrix

$$S = \begin{pmatrix} 1 & 0 & -\frac{1}{r^2} \\ 0 & \frac{1}{2 \sin^2 \phi_1} & \frac{2 \sin^2 \phi_1 - 1}{2 \sin^2 \phi_1} \\ 0 & -\frac{1}{2} & \frac{1}{2} \end{pmatrix}.$$

Therefore, the block-separated Hamiltonians are

$$\tilde{H}_1 = H_1 - c_1 + \frac{1}{r^2} c_3,$$

$$\tilde{H}_2 = H_2 - \frac{1}{2 \sin^2 \phi_1} c_2 - \frac{2 \sin^2 \phi_1 - 1}{2 \sin^2 \phi_1} c_3,$$

$$\tilde{H}_3 = H_3 + \frac{1}{2} c_2 - \frac{1}{2} c_3.$$

The dynamics of the original Hamiltonian H is therefore decomposed into three separated blocks, corresponding to the two dynamics of Hamiltonians \tilde{H}_1, \tilde{H}_2 , with one degree of freedom, and the two-degrees of freedom dynamics generated by \tilde{H}_3 .

Example 5.10. *Killing tensor with an eigenvalue of multiplicity $N - 1$.* If H admits a single quadratic first integral, this one determines block-separable coordinates if it has exactly one eigenvalue of multiplicity one and another one of multiplicity $N - 1$, so that we have a 2×2 Stäckel matrix. Indeed, from block-Eisenhart equations we have

$$X_1 \lambda_1 = 0, \quad X_{2_i} \lambda_2 = 0, \quad i = 1, \dots, N - 1,$$

where X_1 is the eigenvector corresponding to the eigenvalue λ_1 of multiplicity one and X_{2_i} the eigenvectors of λ_2 of multiplicity $N - 1$. Hence, provided λ_1 is not a constant, we have that the submanifolds $\lambda_1 = \text{const}$ are orthogonal to the eigenvector X_1 , which is therefore normally integrable. We can put in this case $X_1 = \partial_1$ and the block separation is essentially determined by the equations

$$\lambda_1(q^2, \dots, q^N) = \text{const},$$

moreover $\lambda_2(q^1)$.

We find in this way another (partial) analogy with Stäckel separation, since in that case, the existence of a single Killing 2-tensor with distinct eigenvalues in dimension two is enough to determine Stäckel separable coordinates and the eigenvalues themselves, if not constants, generate the separable coordinates.

In [9] we show that Stäckel coordinates can be completely determined by the eigenvalues of the associated Killing two-tensors. Part of those results can be easily extended to block-separable systems. However, we leave the analysis of these questions for future researches.

6 Block-separable coordinates of \mathbb{E}^3

In dimension three, only two types of block-separable coordinates can exist, plus the trivial case of a single three-dimensional block. So, or each block is one-dimensional, and the coordinates are standard separable orthogonal coordinates, or one block is one-dimensional and the other one is two-dimensional. In this last case, by denoting the separable coordinates as (u, v, w) , the geodesic Hamiltonian is

$$H = \alpha^1 H_1 + \alpha^2 H_2,$$

with

$$H_1 = g_1(u) p_u^2,$$

and, since any 2-dimensional Riemannian manifold is locally conformally flat,

$$H_2 = g_2(v, w) (p_v^2 + p_w^2),$$

where the choice of local Cartesian coordinates on the manifolds $u = \text{const}$ is not restrictive. Since g_1 can always be set equal to 1 by a rescaling of u , we can assume $g_1 = 1$ and call g_2 simply g .

The corresponding general block-Stäckel matrix has the form

$$S = \begin{pmatrix} S_1^1(u) & S_1^2(u) \\ S_2^1(v, w) & S_2^2(v, w) \end{pmatrix}.$$

Since

$$\alpha^1 = (S^{-1})_1^1, \quad \alpha^2 = (S^{-1})_1^2,$$

the contravariant metric G associated with H has components

$$G^{uu} = \frac{S_2^2(v, w)}{\Delta}, \quad G^{vv} = G^{ww} = \frac{-S_1^2(u)g}{\Delta},$$

with $\Delta = S_1^1(u)S_2^2(v, w) - S_1^2(u)S_2^1(v, w)$.

We restrict ourselves to the space \mathbb{E}^3 by imposing that the Riemann tensor of the metric G is identically zero. We have

$$R_{2132} = -\frac{3(\partial_u S_1^2 S_1^1 - \partial_u S_1^1 S_1^2)(\partial_w S_2^2 S_2^1 - \partial_w S_2^1 S_2^2)}{4\Delta S_1^2 g},$$

and

$$R_{3123} = -\frac{3(\partial_u S_1^2 S_1^1 - \partial_u S_1^1 S_1^2)(\partial_v S_2^2 S_2^1 - \partial_v S_2^1 S_2^2)}{4\Delta S_1^2 g}.$$

Hence, two cases are possible

- i) $(\partial_u S_1^2 S_1^1 - \partial_u S_1^1 S_1^2) = 0$,
- ii) $(\partial_w S_2^2 S_2^1 - \partial_w S_2^1 S_2^2) = 0$ and $(\partial_v S_2^2 S_2^1 - \partial_v S_2^1 S_2^2) = 0$.

Case i

We have

$$S_1^1 = aS_1^2,$$

where a is a constant. We observe that the components of G become

$$G^{uu} = \frac{S_2^2}{S_1^2(aS_2^2 - S_2^1)}, \quad G^{vv} = G^{ww} = -\frac{g}{aS_2^2 - S_2^1},$$

that is, we can write without restrictions

$$G^{uu} = h(u)^{-2}f(v, w)^{-2}, \quad G^{vv} = G^{ww} = l(v, w)^{-2},$$

with the obvious definitions of h , f and l .

A further rescaling of u allows to set $h = 1$ and u can always be considered as an ignorable coordinate, therefore, associated with a Killing vector ∂_u .

The unknown functions are now reduced to two, and we can consider the remaining components of the Riemann tensor. The resulting equations are

$$\begin{aligned} l(\partial_{vv}l + \partial_{ww}l) - (\partial_v l)^2 - (\partial_w l)^2 &= 0, \\ l\partial_{vv}f - \partial_v l\partial_v f + \partial_w l\partial_w f &= 0, \\ l\partial_{ww}f + \partial_v l\partial_v f - \partial_w l\partial_w f &= 0, \\ l\partial_{vw}f - \partial_v l\partial_w f - \partial_w l\partial_v f &= 0. \end{aligned} \tag{6.1}$$

Proposition 6.1. *The coordinate leaves $u = \text{const}$ are planes and ∂_u is proportional to a Killing vector.*

Proof. Equation (6.1) means that the Riemann tensor of the coordinate leaves $u = \text{const}$ is zero. ■

The block-Stäckel matrix is in this case

$$S = \begin{pmatrix} a & 1 \\ \left(1 - \frac{a}{f^2}\right)gl^2 & -\frac{gl^2}{f^2} \end{pmatrix},$$

and the Hamiltonian $H = \alpha_1^1 H_1 + \alpha_1^2 H_2$ and the first integral $K = \alpha_2^1 H_1 + \alpha_2^2 H_2$ are

$$H = \frac{1}{f^2} p_u^2 + \frac{1}{l^2} (p_v^2 + p_w^2), \quad K = \left(1 - \frac{a}{f^2}\right) p_u^2 - \frac{a}{l^2} (p_v^2 + p_w^2).$$

Since $K = p_u^2 - aH$, we have that the first integral associated with this kind of block-separation is simply p_u^2 . An expected result, because the variable u is ignorable.

We observe that the solution of the equation (6.1) is

$$l = c_2 e^{\frac{c_0}{2}(w^2 - v^2) - c_1 v - c_3 w}, \quad (6.2)$$

where (c_i) are real constants. The substitution of (6.2) in the remaining equations yields

$$\begin{aligned} \partial_{vv} f + (c_1 - c_0 v) \partial_v f - (c_3 + c_0 w) \partial_w f &= 0, \\ \partial_{ww} f - (c_1 - c_0 v) \partial_v f + (c_3 + c_0 w) \partial_w f &= 0, \\ \partial_{vw} f + (c_3 + c_0 w) \partial_v f + (c_1 - c_0 v) \partial_w f &= 0. \end{aligned} \quad (6.3)$$

The sum of the first two equations implies that f is a solution of the Laplace equation. Then,

$$f = f_1(v + iw) + f_2(v - iw).$$

The substitution of this equation into (6.3) gives, after few manipulations, the equivalent system

$$\begin{aligned} f_1'' + (c_1 - ic_3 - c_0(v + iw)) f_1' &= 0, \\ f_2'' + (c_1 + ic_3 - c_0(v - iw)) f_2' &= 0. \end{aligned}$$

Example 6.2. *Rotational and cylindrical coordinates.* Given in the Euclidean plane any coordinate system with a symmetry axis, the coordinates of \mathbb{E}^3 obtained by rotating the plane around the symmetry axis and by taking as third coordinate the angle of rotation, are of this form. If f is constant and l^{-2} represents a metric in the Euclidean plane (v, w) , then the cylindrical coordinate system with the plane (v, w) as base is of this form.

Case ii

A similar analysis for Case ii gives the equivalent condition

$$S_2^1 = a S_2^2.$$

We have

$$G^{uu} = \frac{1}{S_1^1 - a S_1^2}, \quad G^{vv} = G^{ww} = -\frac{S_1^2 g}{S_2^2 (S_1^1 - a S_1^2)},$$

and we can write without restrictions

$$G^{uu} = h(u)^2, \quad G^{vv} = G^{ww} = l(u)^2 f(v, w)^2,$$

with the obvious definitions of h , f and l .

Again, a rescaling of u allows to set $h = 1$, in this case, u is not necessarily ignorable, but the metric has the structure of a warped metric. Therefore, ∂_u must be parallel to a conformal Killing vector [8].

The equations arising from the imposition that the Riemann tensor is equal to zero are now

$$\begin{aligned} l''l - 2(l')^2 &= 0, \\ l^4(f(\partial_{vv}f + \partial_{ww}f) - (\partial_v f)^2 - (\partial_w f)^2) - (l')^2 &= 0. \end{aligned} \quad (6.4)$$

The block-Stäckel matrix is in this case

$$S = \begin{pmatrix} 1 - al^2 & -l^2 \\ a\frac{g}{f^2} & \frac{g}{f^2} \end{pmatrix},$$

and the Hamiltonian $H = \alpha_1^1 H_1 + \alpha_1^2 H_2$ and first integral $K = \alpha_2^1 H_1 + \alpha_2^2 H_2$ are

$$H = p_u^2 + l^2 f^2 (p_v^2 + p_w^2), \quad K = -ap_u^2 + (1 - al^2) f^2 (p_v^2 + p_w^2).$$

Since $K = f^2 (p_v^2 + p_w^2) - aH$, we have that the first integral associated with this kind of block-separation is

$$f^2 (p_v^2 + p_w^2),$$

as expected, due to the warped form of the metric.

The solutions of (6.4) are

$$l = -(c_1 u + c_2)^{-1},$$

that, substituted in the second equation, give

$$f(\partial_{uu}f + \partial_{ww}f) - (\partial_v f)^2 - (\partial_w f)^2 - c_1^2 = 0. \quad (6.5)$$

We remark that (6.5) means that the Ricci scalar of the submanifolds $u = \text{const}$ is

$$R = 2l^2(u)c_1^2,$$

and its Riemann tensor has non null components

$$R_{yzyz} = \frac{c_1^2}{f^2}.$$

Therefore,

Proposition 6.3. *In Case ii, all the coordinate leaves orthogonal to ∂_u are planes ($c_1 = 0$) or spheres. The vector ∂_u is proportional to a conformal Killing vector.*

Example 6.4. *Spherical and cylindrical coordinates.* An example of Type ii of block separation is given by spherical-type coordinates, where u is the radius of the spheres and (v, w) are any coordinates on the sphere. If $c_1 = 0$, $c_2 \neq 0$, then the coordinates are cylindrical with the plane (v, w) as base.

Remark 6.5. From the previous remarks it follows that, if a coordinate system is block-separable in $n > m$ blocks, then, not necessarily it is block separable in m blocks too. Indeed, if we consider the ellipsoidal coordinates in \mathbb{E}^3 , they are Stäckel-separable, then block-separable in three blocks, but they cannot be block-separable in 2 blocks, since these coordinates do not include planes or spheres. This fact is not surprising, since it is known that Stäckel separation in ellipsoidal coordinates cannot be achieved by successive separation of the single variables, and block-separation into two blocks in dimension three means exactly that one of the variables can be separated from the others.

The classification of 4D block-separable coordinates, even in Euclidean spaces, appears to be much less simple than in 3D spaces. We must leave the classification of block-separable coordinates in 4D Euclidean spaces for other works. Partial separation of Hamilton–Jacobi and Helmholtz equations on 4D Riemannian manifolds is briefly considered in [4].

7 Conclusions and future directions

By introducing the idea of twisted product of natural Hamiltonians and the analysis of the consequent relations among the Hamiltonian flows of the product and of its factors, we provide a new, dynamical, interpretation of classical partial separation of variables of Hamilton–Jacobi equation, as well as of complete separation. We find that our block-separation, when possible, allows the reconstruction of the orbits of the product Hamiltonian from the orbits of the several lower-dimensional block-separated Hamiltonians. We characterize block-separation in an invariant way, by adapting classical results of complete separation theory for the Hamilton–Jacobi equation. A deeper characterisation of the twisted form of the Hamiltonian is in progress. Furthermore, we extend Eisenhart’s classification of completely separable coordinate systems, the Stäckel systems, in \mathbb{E}^3 to block-separable coordinate systems, finding essentially coordinate-blocks of rotational, cylindrical, and spherical type. We are confident that the possibility of reducing the analysis of the dynamics of Hamiltonians with many degrees of freedom to the dynamics of its lower-dimensional, block-separated Hamiltonians can find many applications, even in the field of numerical computations. We do not consider here the block-separation of Schrödinger’s and other related equations of mathematical physics. Studies on partial separation of these equations are somehow more developed than those on partial separation of Hamilton–Jacobi equation, probably because, for these equations, it is less relevant the absence of completeness in partial separation and the consequent impossibility of application of the Jacobi’s canonical transformation. We will study block-separation of Schrödinger and related equations in next papers.

Acknowledgements

We are grateful to Raymond McLenaghan, Krishan Rajaratnam and Piergiulio Tempesta for conversations useful to clarify some points of our research and to the anonymous referees which gave a relevant contribution to the improvement of this article.

References

- [1] Benenti S., Intrinsic characterization of the variable separation in the Hamilton–Jacobi equation, *J. Math. Phys.* **38** (1997), 6578–6602.
- [2] Benenti S., Chanu C., Rastelli G., The super-separability of the three-body inverse-square Calogero system, *J. Math. Phys.* **41** (2000), 4654–4678.
- [3] Bolsinov A.V., Fomenko A.T., Integrable Hamiltonian systems. Geometry, topology, classification, *Chapman & Hall/CRC*, Boca Raton, FL, 2004.
- [4] Boyer C.P., Kalnins E.G., Miller Jr. W., Separable coordinates for four-dimensional Riemannian spaces, *Comm. Math. Phys.* **59** (1978), 285–302.
- [5] Broadbridge P., Chanu C.M., Miller Jr. W., Solutions of Helmholtz and Schrödinger equations with side condition and nonregular separation of variables, *SIGMA* **8** (2012), 089, 31 pages, [arXiv:1209.2019](https://arxiv.org/abs/1209.2019).
- [6] Chanu C., Geometry of non-regular separation, in *Symmetries and Overdetermined Systems of Partial Differential Equations*, *IMA Vol. Math. Appl.*, Vol. 144, *Springer*, New York, 2008, 305–317.
- [7] Chanu C., Degiovanni L., Rastelli G., Three and four-body systems in one dimension: integrability, super-integrability and discrete symmetries, *Regul. Chaotic Dyn.* **16** (2011), 496–503, [arXiv:1309.0089](https://arxiv.org/abs/1309.0089).

- [8] Chanu C., Degiovanni L., Rastelli G., Generalizations of a method for constructing first integrals of a class of natural Hamiltonians and some remarks about quantization, *J. Phys. Conf. Ser.* **343** (2012), 012101, 15 pages, [arXiv:1111.0030](https://arxiv.org/abs/1111.0030).
- [9] Chanu C., Rastelli G., Eigenvalues of Killing tensors and separable webs on Riemannian and pseudo-Riemannian manifolds, *SIGMA* **3** (2007), 021, 21 pages, [arXiv:nlin.SI/0612042](https://arxiv.org/abs/nlin.SI/0612042).
- [10] Cochran C.M., McLenaghan R.G., Smirnov R.G., Equivalence problem for the orthogonal webs on the 3-sphere, *J. Math. Phys.* **52** (2011), 053509, 22 pages, [arXiv:1009.4244](https://arxiv.org/abs/1009.4244).
- [11] Degiovanni L., Rastelli G., Complex variables for separation of the Hamilton–Jacobi equation on real pseudo-Riemannian manifolds, *J. Math. Phys.* **48** (2007), 073519, 23 pages, [arXiv:nlin.SI/0612051](https://arxiv.org/abs/nlin.SI/0612051).
- [12] Di Pirro G.A., Sugli integrali primi quadratici delle equazioni della meccanica, *Ann. Mat. Pura Appl.* **24** (1896), 315–334.
- [13] Eisenhart L.P., Separable systems of Stäckel, *Ann. of Math.* **35** (1934), 284–305.
- [14] Eisenhart L.P., Stäckel systems in conformal Euclidean space, *Ann. of Math.* **36** (1935), 57–70.
- [15] Eisenhart L.P., Riemannian geometry, 8th ed., *Princeton Landmarks in Mathematics*, Princeton University Press, Princeton, NJ, 1997.
- [16] Haantjes J., On X_m -forming sets of eigenvectors, *Indag. Math.* **58** (1955), 158–162.
- [17] Havas P., Separation of variables in the Hamilton–Jacobi, Schrödinger, and related equations. II. Partial separation, *J. Math. Phys.* **16** (1975), 2476–2489.
- [18] Kalnins E.G., Separation of variables for Riemannian spaces of constant curvature, *Pitman Monographs and Surveys in Pure and Applied Mathematics*, Vol. 28, Longman Scientific & Technical, Harlow, John Wiley & Sons, Inc., New York, 1986.
- [19] Kalnins E.G., Kress J.M., Miller Jr. W., Separation of variables and superintegrability: the symmetry of solvable systems, IOP Publishing, Bristol, 2018.
- [20] Lehmann-Filhés H.R., Über die Verwendung unvollständiger Integrale der Hamilton–Jacobischer partiellen Differentialgleichung, *Astron. Nachr.* **165** (1904), 209–216.
- [21] Levi-Civita T., Sulla integrazione della equazione di Hamilton–Jacobi per separazione di variabili, *Math. Ann.* **59** (1904), 383–397.
- [22] Marcus R.A., Separation of sets of variables in quantum mechanics, *J. Chem. Phys.* **41** (1964), 603–609.
- [23] Meumertzhaim M., Reckziegel H., Schaaf M., Decomposition of twisted and warped product nets, *Results Math.* **36** (1999), 297–312.
- [24] Moon P., Spencer D.E., Field theory handbook. Including coordinate systems, differential equations and their solutions, 2nd ed., *Springer-Verlag*, Berlin, 1988.
- [25] Ni A.V., Pak C.W., Finding analytic solutions on active arcs of the optimal trajectory in a gravitational field and their applications, *Autom. Remote Control* **78** (2017), 313–331.
- [26] Nijenhuis A., X_{n-1} -forming sets of eigenvectors, *Indag. Math.* **54** (1951), 200–212.
- [27] Schouten J.A., Über Differentialkomitanten zweier kontravarianter Grössen, *Proc. K. Ned. Acad. Amsterdam* **43** (1940), 449–452.
- [28] Stäckel P., Sur une classe de problèmes de dynamique, *C. R. Acad. Sci. Paris* **116** (1893), 485–487.
- [29] Stäckel P., Ueber quadratische Integrale der Differentialgleichungen der Dynamik, *Ann. Mat. Pura Appl.* **25** (1897), 55–60.
- [30] Tonolo A., Sulle varietà Riemanniane normali a tre dimensioni, *Pont. Acad. Sci. Acta* **13** (1949), 29–53.
- [31] Wojciechowski S., Superintegrability of the Calogero–Moser system, *Phys. Lett. A* **95** (1983), 279–281.