

Bryn Mawr College
Scholarship, Research, and Creative Work at Bryn Mawr
College

Psychology Faculty Research and Scholarship

Psychology

2017

Orexin 2 receptor regulation of the hypothalamic–pituitary–adrenal (HPA) response to acute and repeated stress

Laura A. Grafe

Bryn Mawr College, lgrafe@brynmawr.edu

Darrell Eacret

Sandra Luz

Anthony L. Gotter

John J. Renger

See next page for additional authors

[Let us know how access to this document benefits you.](#)

Follow this and additional works at: https://repository.brynmawr.edu/psych_pubs

 Part of the [Psychology Commons](#)

Custom Citation

Grafe, Laura A., Darrell Eacret, Sandra Luz, Anthony L. Gotter, John J. Renger, Chris J. Winrow, Seema Bhatnagara. 2017. "Orexin 2 receptor regulation of the hypothalamic–pituitary–adrenal (HPA) response to acute and repeated stress." *Neuroscience* 348.21: 313–323.

This paper is posted at Scholarship, Research, and Creative Work at Bryn Mawr College. https://repository.brynmawr.edu/psych_pubs/69

For more information, please contact repository@brynmawr.edu.

Authors

Laura A. Grafe, Darrell Eacret, Sandra Luz, Anthony L. Gotter, John J. Renger, Chris J. Winrow, and Seema Bhatnagar

**Orexin 2 receptor regulation of the hypothalamic-pituitary-adrenal (HPA)
response to acute and repeated stress**

Laura A. Grafe¹, Darrell Eacret¹, Sandra Luz¹, Anthony L. Gotter², John J. Renger²,
Chris J. Winrow², Seema Bhatnagar^{1,3}

Department of Anesthesiology and Critical Care, Children's Hospital of Philadelphia,
Philadelphia, PA, USA 19104¹; Department of Neuroscience, Merck Research
Laboratories, West Point, PA, USA 19486²; University of Pennsylvania Perelman
School of Medicine, Philadelphia, Pennsylvania, USA 19104³

Corresponding Author:

Seema Bhatnagar

Abramson Research Center, Suite 402B

Children's Hospital of Philadelphia

3615 Civic Center Boulevard

Philadelphia, PA 19104

Tel: 267-426-0951

Email: hatnagars@email.chop.edu

Abstract

Orexins are hypothalamic neuropeptides that have a documented role in mediating the acute stress response. However, their role in habituation to repeated stress, and the role of orexin receptors (OX₁R and OX₂R) in the stress response, has yet to be defined. Orexin neuronal activation and levels in the cerebrospinal fluid were found to be stimulated with acute restraint, but were significantly reduced by day five of repeated restraint. As certain disease states such as panic disorder are associated with increased central orexin levels and failure to habituate to repeated stress, the effect of activating orexin signaling via Designer Receptors Exclusively Activated by Designer Drugs (DREADDs) on the hypothalamic-pituitary-adrenal (HPA) response was evaluated after repeated restraint. While vehicle-treated rats displayed habituation of Adrenocorticotropic Hormone (ACTH) from day 1 to day 5 of restraint, stimulating orexins did not further increase ACTH beyond vehicle levels for either acute or repeated restraint. We delineated the roles of orexin receptors in acute and repeated stress by using a selective OX₂R antagonist (MK-1064). Pretreatment with MK-1064 reduced day 1 ACTH levels, but did not allow further habituation on day 5 compared with vehicle-treated rats, indicating that endogenous OX₂R activity plays a role in acute stress, but not in habituation to repeated stress. However, in restrained rats with further stimulated orexins by DREADDs, MK-1064 decreased ACTH levels on day 5. Collectively, these results indicate that the OX₂R plays a role in acute stress, and can prevent habituation to repeated stress under conditions of high orexin release.

Key words: MK-1064, orexin receptor, Hypocretin, Stress, Habituation, DREADDs

Orexins (also called hypocretins) are peptides generated from the prepro-orexin precursor that is exclusively localized in cells of the lateral and posterior hypothalamic region (de Lecea et al., 1998; Sakurai et al., 1998). Prepro-orexin is cleaved into two highly structurally related and highly conserved peptides, orexin A and orexin B, which bind to two G-protein coupled orexin receptors, orexin 1 and orexin 2 receptors (OX₁R and OX₂R, respectively) (de Lecea et al., 1998; Sakurai et al., 1998). Orexin A has nearly equal affinities for both receptors, while Orexin B has higher affinity for OX₂R (Gotter et al., 2012b). While many brain regions express both receptors, some regions exhibit differential expression (Marcus et al., 2001). For example, the cingulate cortex and locus coeruleus selectively express OX₁R, while the paraventricular nuclei of the hypothalamus (PVN) and shell neurons of nucleus accumbens preferentially express OX₂R (Marcus et al., 2001; Trivedi et al., 1998).

Orexins are implicated in a wide variety of neuroendocrine and behavioral responses including arousal, food intake, cognitive function, autonomic responses, emotional memory, and the stress response (Sakurai, 2014). The most salient phenotype of orexin knockouts is reduced arousal resulting in narcolepsy-like behavior, which has led to the identification and development of OXR antagonists for the pharmacological treatment of insomnia (Gotter et al., 2012a). Conversely, i.c.v. administration of exogenous orexin A to rats or orexin receptor agonists to mice promote arousal and wakefulness (Hagan et al., 1999; Nagahara et al., 2015). Orexins are also important in regulating the neurobiological systems that respond to stressful stimuli. For example, orexins promote the hypothalamic-pituitary-adrenal (HPA) axis response (Winsky-Sommerer et al., 2004, 2005; Berridge et al., 2010; Johnson et al., 2012). Specifically, central orexin administration (either Orexin A or Orexin B) increases HPA hormones (Jászberényi et al., 2000; Kuru et al., 2000; Spinazzi et al., 2006). Conversely, orexin neurons are activated by corticotropin releasing hormone and respond to stressors such as forced swim stress (Winsky-Sommerer et al., 2005; Chang et al., 2007; Furlong et al., 2009; Chen et al., 2013). Both orexin 1 and 2 receptors have been implicated in acute stress; the role of each receptor appears dependent on the paradigm of stress used (Chang et al., 2007; Gozzi et al., 2013; Bonaventure et al., 2015).

While this evidence indicates a role for orexins in activating the HPA activity under conditions of acute stress, the role of orexin signaling in response to repeated stress is not known. With repeated exposure to moderately intense stressors, individuals typically habituate to that stress as indicated by decreasing responsivity in behavioral, HPA and autonomic measures (Grissom et al., 2008; Grissom and Bhatnagar, 2009). Failure to habituate to a stressor is a hallmark of stress-related illnesses such as post-traumatic stress disorder (PTSD) and panic disorder (Johnson et al., 2012). Moreover, patients with panic anxiety symptoms have higher levels of orexin in their cerebrospinal fluid (CSF) (Johnson et al., 2010). If involved in habituation to a stressor, orexins may be a good target for treatment for some of these stress-related psychiatric illnesses.

To determine the role of orexins in acute and repeated restraint, we first characterized orexin neuronal activation and levels in the CSF in both conditions. We found these measures to be increased with acute restraint, but they significantly decreased with repeated restraint. In order to increase orexin activation prior to each of five daily 30-min restraints (to model higher orexin levels as observed in panic disorder), we used Designers Receptors Exclusively Activated by Designer Drugs (DREADDs) driven by the orexin promoter. We found that even with additional orexin stimulation, HPA activation habituated by day 5 of restraint. However, we were able to delineate the roles of orexin receptors in acute and repeated stress using a selective OX₂R specific antagonist (MK-1064) (Roecker et al., 2014). Prior to this, the roles of orexin receptors in the stress response, whether acute or repeated, had not yet been fully characterized. As OX₂R is abundant in the PVN and is known to play a role in arousal, it may mediate excitatory effects at this site where the HPA response is initiated (Shirasaka et al., 2001; Gotter et al., 2012b). In support of this, we found that OX₂R receptor promotes the acute stress response, and it partially prevents habituation to repeated stress only in conditions of high orexin release.

EXPERIMENTAL PROCEDURES

Animals

Adult male Sprague Dawley rats (225-250g) were obtained from Charles River Laboratories (Wilmington, MA, USA). Rats were singly housed in polycarbonate enclosures with standard bedding and with food and water available ad libitum. Animals were acclimated to a 12-h light-dark cycle with lights on at 06:15 and lights off at 18:15 in a temperature controlled vivarium for at least 5 days prior to administration of any stress protocols. All experiments took place during the inactive phase between 0800-1200 (Zeitgeber Time 01:45-05:45). Experiments followed the NIH Guide for the Care and Use of Laboratory Animals and were approved by the Children's Hospital of Philadelphia Research Institute's Animal Care and Use Committee.

Stereotaxic delivery of DREADDs-containing viruses

Designer Receptors Exclusively Activated by Designer Drugs (DREADDs) are constructs packaged into viruses that contain synthetic G protein coupled receptors (GPCRs) and can be activated by the otherwise pharmacologically inert ligand Clozapine-N-Oxide (CNO). We obtained the CMV-hM3Dq-mCitrine plasmid from Dr. Bryan Roth (University of North Carolina, Chapel Hill, NC). Slice electrophysiology has demonstrated that CNO application to cells expressing this Gq coupled designer receptor causes depolarization and increases firing rate (Alexander et al., 2009). We next obtained a 1295 bp promoter for human preproorexin gene (Ple112) from Addgene plasmid No. 29004 (gift of Dr. Elizabeth Simpson, Univ. of British Columbia). This promoter was subcloned upstream of the hM3Dq-mCitrine region to replace the construct's CMV promoter to drive transgene expression specifically in orexin neurons. The fragment Ple112-hm3Dq-mCitrine was then subcloned between the Inverted Terminal Repeats (ITRs) of the AAV2 genome. In a separate study, we found AAV1 serotype displayed optimal tropism for Sprague Dawley rat hypothalamic neurons when we delivered in vivo and compared to AAV5,8,9 expression of GFP reporters driven by common constitutively active promoters Synapsin and CB7. Based on this finding, The University of Pennsylvania Vector Core produced a recombinant adenovirus rAAV2/1-Ple112-hM3Dq-mCitrine (using AAV1 serotype capsid for optimal transduction in orexin

neurons) for our use.

Surgeries were performed in aseptic conditions. Animals were anesthetized using a cocktail of ketamine, xylazine, and acepromazine. Using stereotaxic technique, DREADDs-containing virus (10^9 titer, 1ul bilaterally) was injected into the lateral hypothalamus (coordinates: 2.5 mm caudal to bregma, 1.8 mm from mid-line and 8 mm ventral to dura mater). After surgery, animals were injected with meloxicam (2mg/kg). Animals recovered for 4 weeks while virus expressed before any further experiments were performed.

Immunofluorescence staining for visualizing the virus tag was conducted as follows. Rats were rapidly decapitated and brains were flash frozen in 2-methyl butane. Coronal sections for the lateral hypothalamus were cut on a cryostat into six serial sets of 20-um-thick sections directly onto the slide. Rostral-caudal coordinates (relative to bregma) for the lateral hypothalamic area analyzed were -2.12 mm to -3.60 mm. Tissue was fixed for 30 minutes in 4% paraformaldehyde (Electron Microscopy Sciences; Hatfield, PA). Sections were washed several times with Phosphate Buffered Saline (PBS; pH 7.4) with 0.3% Triton-X (Sigma-Aldrich; St Louis, MO), and then incubated with primary antibodies for both Orexin A (1:250, sc-8070; Santa Cruz Biotechnology, Santa Cruz CA) and GFP (1:500, ab290; AbCam, Cambridge, UK) in a solution of PBS with 5% Normal Donkey Serum (EMD Millipore; Billerica, MA) and 0.3% Triton-X overnight. As the mCitrine tag on the DREADDs-containing virus originates from *Aequorea victoria* jellyfish, GFP antibodies are known to react with these proteins (Le et al., 2006). Sections were rinsed several times in PBS, followed by the addition of AlexaFluor488 Donkey anti-goat and AlexaFluor647 Donkey anti-rabbit (1:200, A-11055 and A-31573; Life Technologies, Carlsbad, CA). Sections were given a final set of washes and coverslipped with fluoromount. Images were acquired with a Leitz DMR microscope with a digital camera (Leica Microsystems; Buffalo Grove, IL). The NIH Image J colocalization plugin was used to determine percent orexin cells transduced by the virus. Quantification revealed $71 \pm 5.4\%$ of neurons within the lateral hypothalamus exhibited the DREADDs tag. Roughly 1% of these cells were not dual labeled with orexin A antibody.

Drugs

Clozapine N-Oxide (CNO; Sigma-Aldrich; St Louis, MO) was dissolved in saline and 8% DMSO and injected intraperitoneally at 2 mg/kg in a 5 mg/mL solution. This dose is in accordance with doses used in previous DREADDs studies in rats (Farrell and Roth, 2013). MK-1064, a selective OX₂R antagonist, (Roecker et al., 2014) was dissolved overnight in 20% Vitamin E d- α -tocopherol polyethylene glycol 1000 succinate (Vit E-TPGS) (Sigma-Aldrich; St Louis, MO). Rats were orally administered either 20% Vit E-TPGS as vehicle or 30 mg/kg MK-1064, both at 1 mL/kg for consistent volume. All animals were given a vehicle dose 1 day prior to the start of the restraint paradigm to habituate to the oral gavage procedure. Animals received injections of vehicle (20% Vit E TPGS, p.o.) or MK-1064 and vehicle (saline and 8% DMSO, i.p.) or CNO 90 minutes prior to the start of the 30-minute restraint. This timing was chosen based on the fact that both MK-1064 and CNO have been shown to promote behavioral effects in the rat within 30 minutes of administration and effects last up to 4 hours after administration (Alexander et al., 2009; Farrell and Roth, 2013; Hasegawa et al., 2014; Roecker et al., 2014). Additionally, CNO was given at the time of MK-1064 injection to minimize the effects of repeated handling prior to restraint.

Assaying orexin levels in the cerebrospinal fluid (CSF)

DREADDs-expressing rats were injected *IP* with either vehicle or CNO 90 minutes prior to either CSF collection. In a separate experiment, DREADDs-expressing rats were injected *IP* with vehicle or CNO 90 minutes prior to stress exposure. Specifically, rats were either exposed to 1 or 5 days of 30-min restraint (*IP* injections were given daily prior to each restraint for the 5-day group). Animals were restrained in Plexiglas restrainers (9 x 3.5 inches) made from the University of Pennsylvania Machine Shop. Plexiglas inserts (7.5 x 3.25 inches) were placed inside the restrainers appropriate for the size of each male rat so that they could not turn around in the restrainer. As orexins are known to exhibit a circadian rhythm (Taheri et al., 1999; Fujiki et al., 2001; Zeitzer, 2013), animals were restrained within 2 hours after lights on, a time at which orexin levels reach a nadir. At 15 min into the 1st or 5th restraint, CSF was collected (rough 105 minutes post *IP* injection). Briefly, rats were anesthetized with a

cocktail of ketamine and xylazine, placed in the ear bars of a stereotaxic apparatus with their nose pointing downward, and a 25-gauge needle was used to extract 100 μ L of CSF from the cisterna magna. CSF was ejected into eppendorf tubes and frozen at -80°C until radioimmunoassay. CSF levels of orexin were assayed with a Radioimmunoassay kit from Phoenix Pharmaceuticals (Burlingame, CA). The minimum levels of detection for orexin was 80 pg/ml. Intra-and interassay variability was 5-7% and 12-15%, respectively.

Assaying orexin neuronal activation

Two additional groups of rats were injected with the DREADDs construct and exposed to either 1 or 5 days of restraint with injections of vehicle or CNO administered as previously described. Brains were collected for one group of rats on Day 1 of restraint (30 min after restraint ended), while brains were not collected in the other cohort until Day 5 of restraint. The lateral hypothalamus of these brains were further analyzed for orexin neural activation. Specifically, sections of the lateral hypothalamus were cut and fixed as described above in DREADDs immunofluorescent staining. Sections were incubated with c-Fos antibody (1:1250, sc-52; Santa Cruz Biotechnology, Santa Cruz, CA) followed by Biotin-SP-conjugated AffiniPure Donkey-Antirabbit IgG (1:200, Jackson ImmunoResearch), which was visualized with a 3'3'-diaminobenzidine reaction (Sigma-Aldrich, St Louis, MO). Sections were then stained for orexin A (1:50, sc-8070; Santa Cruz Biotechnology, Santa Cruz CA) followed by biotinylated Horse Anti-Goat antibody (1:500, BA-9500; Vector Laboratories, Burlingame, CA.), which was visualized with a Nova Red reaction (SK-4800, Vector Laboratories, Burlingame, CA). Sections were mounted with permount (Electron Microscopy Sciences, Fort Washington, PA). Images were acquired with a Leitz DMR microscope with a digital camera (Leica Microsystems; Buffalo Grove, IL). 4 sections per animal (with rostral-caudal coordinates (relative to bregma) ranging from -2.04 to -3.24) were analyzed using NIH Image J. Specifically, a standardized box encompassed the lateral hypothalamus (calculated using Paxinos & Watson Rat Brain Atlas) and dual labeled cells were counted by hand. All image analysis was performed with the experimenter blind to the treatment groups.

Assessing HPA reactivity to acute or repeated restraint with orexin manipulations

A naïve cohort of rats injected with DREADDs-containing virus was exposed to either 1 or 5 consecutive days of restraint with injections of vehicle or CNO *IP* and either vehicle or MK-1064 administered *PO* 90 min prior to each restraint. See **Fig. 3A** for depiction of experimental paradigm. Animals were weighed on Day 1 and Day 5 of restraint. Video cameras were set up above the restrainers in order to record struggle behavior on day 2 of restraint. While rats are in the restrainer for 30 minutes, most of the struggle behavior occurs within the first 10 minutes. Therefore, we analyzed struggle behavior during this time period. A trained investigator blind to experimental groups hand scored struggle behavior – defined as attempts to escape, or intense movement of the animal while in the restrainer. Blood was collected on Day 1 and Day 5 of restraint to assess the HPA response to acute and repeated restraint, respectively. Briefly, on day 1, tail blood was taken at 0 minutes (prior to being placed in restraint), again at 15 minutes and 30 minutes (during restraint), and at 60 minutes (recovery time point in the home cage). Plasma corticosterone and ACTH were assayed with a Radioimmunoassay kit from MP Biomedical (Orangeburg, NY). The minimum levels of detection for ACTH and corticosterone were 5.7pg/ml and 0.6µg/dl, respectively. Intra- and interassay variability was less than 10%.

Statistical Analysis

Data are presented as the mean \pm the standard error of the mean. For habituation of vehicle-treated animals, a 2-way ANOVA assessed Day and Time variables (Day 1 and 5 were compared, as well as the time point throughout restraint). A t-test was used to evaluate orexin neural activation and orexin levels in the CSF in control versus CNO treated animals on Day 1 and Day 5 of restraint. A 2-way ANOVA assessed Day and Drug variables for orexin neural activation and orexin A in the CSF throughout repeated restraint. For struggle behavior and body weight, a 2-way ANOVA assessed CNO and 2-SORA variables. For all other ACTH data, a 2-way ANOVA assessed Drug and Time variables on Day 1 and Day 5 separately. When warranted, Tukey's-corrected t-tests were performed. All hypothesis tests used $\alpha=0.05$ as the

criterion level of significance. Statistical analysis was conducted with GraphPad Prism (GraphPad Software, La Jolla, CA, USA) in order to identify statistical differences.

RESULTS

Orexin levels habituate with repeated stress

Habituation of the HPA response to 5 days of restraint stress was assessed by evaluating plasma ACTH levels prior to, during and after 30 minutes of restraint stress on day 1 and day 5 of restraint (paradigm and data in **Fig. 1A**). Consistent with previous results (Grissom and Bhatnagar, 2009), male rats in this paradigm habituated to 5 days of repeated restraint with regard to plasma ACTH levels. This effect was significant at 30 minutes of restraint. However, we did not observe habituation or group differences in our corticosterone assays, consistent with previous lab results that it takes 8-10 days of restraint to see corticosterone effects ((Grissom et al., 2008); data not shown).

After demonstrating ACTH habituation to repeated restraint, we examined changes in orexins from one day to five days of daily 30-min restraint. To assess this, orexin neuronal activation and CSF levels of orexin were evaluated on both day 1 and day 5 of restraint. Representative images of the cFos/orexin dual stain on day 1 and day 5 of restraint are displayed in **Fig. 1B**. An average of 40 orexin cells per section were dual labeled with cFos after a single 30-min restraint, and this was significantly reduced to nearly half that amount by day 5 of restraint stress. In terms of percentages of total orexin neurons present, a single 30-min restraint activated 75% of orexin neurons, whereas only 40% were active on day 5 of restraint stress (data not shown). Orexin A levels in CSF showed a similar pattern to orexin neuronal activation, since orexin A levels in CSF habituated after 5 days of restraint stress (**Fig. 1C**). Specifically, orexin A levels were significantly reduced to approximately 66% of the day 1 value. In summary, both HPA and orexin activity decreased with repeated restraint stress.

Stimulation of orexins prior to each of five daily 30-min restraints

As certain disease states such as panic disorder are associated with increased central orexin levels and failure to habituate to repeated stress, we further stimulated orexins prior to each of five daily 30-min restraints. To do this, DREADDs were injected and expressed selectively in orexin neurons to activate these neurons. Representative fluorescent images of DREADDs expression at four weeks are displayed in **Fig. 2A**, along with a depiction of the spread of viral expression along four levels of Bregma (anterior to posterior) in **Fig. 2B**. Quantification revealed $71 \pm 5.4\%$ of neurons within the lateral hypothalamus exhibited the DREADDs tag (data not shown). Orexin levels in the CSF are quantified for vehicle- or CNO-treated DREADDs-expressing rats in non-stress conditions in **Fig. 2C**. Relative to vehicle, CNO significantly increased the level of orexins in the CSF compared with vehicle-treated rats when evaluated 90 minutes after treatment. These data validated the functionality of the DREADDs construct.

Moreover, orexin neuronal activation with DREADDs stimulation prior to each of five daily 30-min restraints were quantified (**Fig 2D**). Compared with vehicle treatment, CNO significantly increased the number of orexin neurons dual labeled with cFos (and hence neural activation) on day 1 of restraint. The level of CNO-induced orexin neuron recruitment on day 5, however, remained significantly below that seen on day 1 of restraint stress after CNO (just as was seen with vehicle-treated groups on day 1 compared with day 5).

Lastly, orexin levels in the CSF with DREADDs stimulation prior to each of five daily 30-min restraints were quantified (**Fig 2E**). Relative to vehicle-treated animals, CNO induced orexin A in CSF to significantly greater levels on day 1, verifying that DREADDs-mediated CNO activation of orexin neurons resulted in augmented release of orexin A neuropeptide. While CNO treatment also increased orexin A levels on day 5 of restraint compared with vehicle treatment, CNO-treated animals exhibited significantly reduced orexin A levels on day 5 compared with day 1 of restraint, just as it did in vehicle-treated animals. Together, these data indicate that the DREADDs constructs increased orexin activation and levels in the CSF, but repeated restraint stress partially counteracts these effects by lowering orexin activity and levels in the CSF compared to day 1 of restraint.

OX₂R underlies struggle behavior and body weight change during restraint stress

The contribution of orexin receptor signaling in both struggle behavior and body weight change in response to restraint stress was determined through use of a selective OX₂R antagonist, MK-1064. Specifically, after DREADDs expression, rats underwent 5 consecutive days of 30-minute restraint, with either vehicle or CNO to stimulate DREADDs prior to each restraint, and with either vehicle or MK-1064 to antagonize the orexin 2 receptor (paradigm depicted in **Fig. 3A**). Struggle behavior indicates the number of attempts to escape during restraint, which is an important component of the stress response and habituates with repeated restraint (Grissom et al., 2008). As demonstrated in **Fig. 3B**, the CNO/vehicle group spent significantly more time struggling than the other three treatment groups. Essentially, stimulating orexins with CNO increases struggle behavior, and blocking OX₂R with MK-1064 reverses this effect. Coincident with changes observed in struggle behavior, body weight decreased from the first to last day of restraint in levels of high orexin, but this was eliminated with MK-1064 pretreatment (**Fig. 3C**). Thus, while all rats lost weight over the 5 days of restraint stress, further stimulating orexins prior to each 30-min restraint exacerbated this effect and these results appear mediated through OX₂R.

OX₂R is involved in the ACTH response to acute restraint stress

The role of orexin signaling in the HPA response to acute restraint stress was determined by measuring plasma ACTH levels in response to 30 minutes of restraint in DREADDs-expressing rats pretreated with either vehicle or CNO in combination with either vehicle or the OX₂R antagonist, MK-1064. CNO pretreatment did not increase ACTH levels compared with that from vehicle-treated rats (**Fig. 4A**). Pretreatment with MK-1064, however, significantly attenuated the ACTH response in combination with either vehicle (**Fig. 4B**), or CNO pre-treatment (**Fig. 4C**), indicating that OX₂R is responsible for mediating at least part of the ACTH response to acute restraint. Importantly, the ACTH response was similarly attenuated by MK-1064 in either the

presence or absence of CNO (**Fig. 4D**), indicating that activation of orexin signaling through mechanisms not involving OX₂R contributes little to the ACTH response on this first day of restraint stress.

Orexin signaling through OX₂R and non-OX₂R mechanisms have opposing influences on ACTH responses to repeated restraint stress

The role of orexin signaling in mediating the HPA response to repeated stress was evaluated in **Fig. 5**. Specifically, rats were evaluated for plasma ACTH levels on the 5th day of restraint stress following treatment with CNO and/or MK-1064. CNO treatment did not change ACTH levels on day 5 of restraint compared with vehicle-only controls (**Fig. 5A**). In the absence of CNO, blockade of endogenous OX₂R activity by MK-1064 also did not further affect plasma levels of ACTH compared with vehicle (**Fig. 5B**), unlike that seen on the first day of restraint stress. However, in rats treated with CNO to induce orexin neuron activation in combination with MK-1064, the ACTH response was attenuated further (**Fig. 5C**). Because ACTH levels are elevated following stimulation of orexin neuron activation (CNO/Vehicle) relative to that following stimulation of orexin neuron activation plus antagonism of the OX₂R (CNO/MK-1064), these results indicate that OX₂R activity has the capacity to increase ACTH levels after 5 days of restraint stress. In other words, exogenous OX₂R activity prevents the full habituation to repeated restraint. Indeed, ACTH responses are reduced by the CNO/MK-1064 manipulation relative to the Vehicle/MK-1064 treatment (**Fig. 5D**). This comparison evaluates the influence of total orexin neuron activity minus that of OX₂R, indicating that possibly OX₁R, or other mechanisms induced by orexin neuron activity, has the capacity to further reduce ACTH responses to repeated restraint stress.

DISCUSSION

While orexins are known to play a role in the acute stress response, it was unclear how they mediated the response to repeated stressors, and the contribution of orexin receptors to both acute and repeated stress was undetermined. With repeated

exposure to modest stressors, individuals typically habituate to that stress by decreasing responsivity, particularly in HPA measures (Grissom et al., 2008; Grissom and Bhatnagar, 2009). Failure to habituate is a hallmark of stress-related illnesses such as post-traumatic stress disorder (PTSD) and panic disorder (Johnson et al., 2012). The present studies demonstrated HPA habituation to repeated restraint in male rodents, and determined that both orexin neuronal activation and levels in the CSF decrease after repeated restraint. As central orexins are augmented in some disease states such as panic disorder (where patients fail to habituate to repeated stress), high levels of orexin were stimulated via DREADDs prior to each of five daily 30-min restraints. Our data confirm that our DREADDs construct was efficacious in stimulating orexin neuronal activation and increasing orexin levels in the CSF in non-stress conditions, as well as in acute and repeated restraint. We were also able to discern the contribution of orexin receptor signaling in acute and repeated stress through use of a selective OX₂R antagonist, MK-1064. We found that the OX₂R contributed to struggle behavior and body weight decrease during repeated restraint. Moreover, the OX₂R played a role in the plasma ACTH response to acute stress and repeated stress, although the latter was only in conditions of high orexin release.

Orexins and the HPA axis reciprocally interact. Specifically, central orexin increases corticosterone and ACTH, whereas CRH depolarizes orexin neurons (Russell et al., 2001; Samson et al., 2002; Winsky-Sommerer et al., 2004). The activation of orexin neurons during acute stress has been examined by a limited number of studies using a wide variety of stress paradigms (Chang et al., 2007; Furlong et al., 2009; Chen et al., 2013). As much of the previous literature has focused on the role of orexin A in stress (Ida et al., 2000; Russell et al., 2001; Sakamoto et al., 2004), our studies specifically examined orexin A activity both through dual labeling of orexin A neurons with cFos (a measure of neural activation) and orexin A levels in the CSF in acute and repeated stress. Importantly, our results indicate that acute restraint stress induces very high levels of orexin neuron activation (75%), accompanied by increased levels of orexin A in the CSF. We also show, for the first time, that orexin activity decreases with habituation to repeated stress. Thus, with repeated exposure to a moderately intense stressor, both the HPA and orexin response diminish. When inducing additional orexin

activation via DREADDs prior to each of five daily 30-min restraints, we were unable to increase orexin neuron activation to the levels seen with acute stress. Based on the literature, CNO does not appear to become less effective after repeated administration, so this is likely not the cause for these results (Kozorovitskiy et al., 2012). Moreover, this reduced responsiveness of orexin neurons does not appear due to depletion of available orexin neuropeptide stores, since DREADDs-induced CSF levels of orexin A and orexin neuron activation as measured by c-fos expression were both attenuated during habituation. It is important to note that Orexin B also plays an important role in the stress response (Kuru et al., 2000; Shirasaka et al., 2001; Spinazzi et al., 2006), so assessing orexin B levels in the future may shed more light on how the orexins are involved in repeated stress.

Struggle behavior and body weight changes are important components of the response to stressful stimuli. Our data show that exogenous orexin stimulation increased struggle behavior and decreased body weight. This was reversed by the OX₂R antagonist, indicating that OX₂R plays a role in these components of the stress response in conditions of high orexin release. This effect on struggle behavior is consistent with literature support for the OX₂R in promoting arousal (Gotter et al., 2012b). Moreover, arousal is an important component of the stress response (Giardino and de Lecea, 2014), so it is likely that DREADDs-induced orexin activation is increasing struggle behavior by increasing arousal. However, our data indicate that DREADDs differentially modulate the HPA response and struggling behavior, thus, mechanisms above and beyond those regulated by orexins may be involved. [Although orexin antagonists are known to promote sleep \(Gotter et al., 2016\), they still allow arousability to salient stimuli \(Tannenbaum et al., 2014, 2016\). Thus, it is unlikely that reduction in struggle behavior is due to the sedative effects of the OX₂R antagonist. However, as we did not directly measure sleep using telemetry devices, this possibility cannot be ruled out.](#)

It is unclear where OX₂R_s are acting to regulate struggle behavior and body weight gain. Previous literature has indicated that rats with high amygdala excitability show increased struggling behavior (Anisman et al., 1997; McIntyre et al., 1999). Given

that OX₂R_s are expressed in medial but not central amygdala, this subnuclei of the amygdala could be an important substrate for regulating struggle behavior (Marcus et al., 2001). Alternatively, both central and peripheral regions may play a role in OX₂R mediation of body weight gain. For example, central orexins acting on hypothalamic and dopaminergic brain regions have been implicated in increasing spontaneous physical activity and nonexercise thermogenesis (Kotz et al., 2012). Moreover, orexins acting on white adipose tissue have been shown to promote lipolysis (Perez-Leighton et al., 2014). Thus, it is possible that the reduction in body weight observed with CNO treatment is a result of increased energy expenditure and lipolysis by animals in this treatment group.

There have been few studies delineating the contribution of each orexin receptor to the acute HPA response. One study reports that pretreatment with an OX₂R antagonist prevents forced swim-induced increases in plasma ACTH and corticosterone (Chang et al., 2007). This is consistent with our findings that OX₂R contributes to the HPA response induced by acute restraint stress. However, as MK-1064 pretreatment only reduced the HPA response by half, it is still possible that the other components of the stress response may play a role in regulating the acute HPA response to restraint. Possible components would not include the contribution of OX₁R, however, as the ACTH response was similarly attenuated by MK-1064 in either the presence or absence of CNO. As OX₂R is preferentially expressed (over OX₁R) in the paraventricular nucleus of the hypothalamus, antagonizing OX₂R (but not necessarily OX₁R) would likely reduce the acute HPA response (Shirasaka et al., 2001; Gotter et al., 2012b). However, other sites expressing OX₂R such as the amygdala, paraventricular nucleus of the thalamus (PVT), and hippocampus may also be involved (Marcus et al., 2001).

We found that under endogenous orexin levels elicited by restraint alone, blocking OX₂R had no effect on ACTH levels on day 5 of restraint. Thus, in these conditions, OX₂R does not affect habituation. However, our data measuring orexin levels in the CSF revealed that compared with one day of 30-min restraint, orexin levels decreased by five days of daily 30-min restraint. Hence, blocking OX₂R in low orexin conditions may not have an effect. However, when orexins are stimulated with

DREADDs, blocking OX₂R further attenuates ACTH responses. This indicates OX₂R plays a role in repeated restraint stress only in conditions of high orexin release. This also indicates that OX₂R desensitization does not appear to be involved in typical habituation. However, the ultimate effects of orexins throughout stress might be dependent on the balance of signaling from both orexin receptors (Scott et al., 2011). Though, not much is known about orexin receptor regulation after stress. While the OX₂R activity promotes ACTH release and prevents habituation in high levels of orexin such as that seen during acute stress conditions, one possibility is that OX₁R activated by orexin release could promote habituation, as CNO/MK-1064 treated animals have even lower plasma ACTH levels than Vehicle/MK-1064 treated animals after repeated restraint. Thus, one interpretation of the results is that OX₁R underlies habituation. Direct tests of OX₁R function are required to confirm this interpretation. Interestingly, previous data from our lab indicates that the OX₁R in the posterior paraventricular thalamus is necessary in adapting to repeated swim stress, allowing facilitation to a novel restraint stressor (Heydendael et al., 2011). However, it is not clear whether orexins are involved in facilitation after habituating to repeated restraint. Overall, we believe the evidence suggests that orexin neuron has the capacity to influence habituation through opposing OX₂R and non- OX₂R mechanisms.

The results from this study improve our understanding of the role of orexin receptors in regulating the HPA response to both acute and repeated stress. We have confirmed the involvement of OX₂R in acute stress, and have shown that it has the capacity to promote ACTH levels opposing habituation to repeated stress in conditions of exogenously elevated orexin signaling. Future studies should identify not only the neuronal pathways regulating the reduced responsiveness of orexin neurons during habituation, but also investigate the specific brain regions where each receptor may be acting to influence these changes in HPA activity, struggle behavior, and body weight gain. While there are high concentrations of OX₂R in the PVN, the locus coeruleus preferentially expresses OX₁R (Peyron et al., 1998; Marcus et al., 2001). Additionally, other stress regulatory regions with high concentrations of orexin input include the PVT and the amygdala. These experiments would allow for a greater understanding of

mechanisms associated with hyperactivation of orexin signaling observed in psychiatric illness such as panic disorder.

ACKNOWLEDGEMENTS

This work was supported by Merck & Co. Inc. LG was supported by award number T32NS007413 from the National Institute of Neurological Disorders and Stroke (NINDS). The content is the sole responsibility of the authors and does not necessarily represent the official views of the NINDS of the National Institutes of Health. We gratefully acknowledge Lauren Wilson and Jane Dobkin for their help in coding struggle behavior, as well as Amanda Cornfeld for validating DREADDs expression by fluorescent labeling.

DISCLOSURES

ALG, CJW and JJR are employees of Merck & Co., Inc., Kenilworth, NJ (known as Merck Sharp & Dohme Corp. (MSD) outside the United States and Canada), have received salary and research support from the company, and may own stock/stock options in the company.

REFERENCES

- Alexander GM, Rogan SC, Abbas AI, Armbruster BN, Pei Y, Allen JA, Nonneman RJ, Hartmann J, Moy SS, Nicoletis MA, McNamara JO, Roth BL (2009) Remote control of neuronal activity in transgenic mice expressing evolved G protein-coupled receptors. *Neuron* 63:27–39 Available at: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2751885&tool=pmcentrez&rendertype=abstract> [Accessed January 21, 2014].
- Anisman H, Lu ZW, Song C, Kent P, McIntyre DC, Merali Z (1997) Influence of psychogenic and neurogenic stressors on endocrine and immune activity: differential effects in fast and slow seizing rat strains. *Brain Behav Immun* 11:63–74 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9193768> [Accessed January 7, 2016].
- Berridge CW, España RA, Vittoz NM (2010) Hypocretin/orexin in arousal and stress. *Brain Res* 1314:91–102 Available at: <http://www.sciencedirect.com/science/article/pii/S0006899309018939> [Accessed August 30, 2014].
- Bonaventure P, Yun S, Johnson PL, Shekhar A, Fitz SD, Shireman BT, Lebold TP, Nepomuceno D, Lord B, Wennerholm M, Shelton J, Carruthers N, Lovenberg T, Dugovic C (2015) A Selective Orexin-1 Receptor Antagonist Attenuates Stress-Induced Hyperarousal without Hypnotic Effects. *J Pharmacol Exp Ther* 352:590–601 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25583879> [Accessed January 6, 2016].
- Chang H, Saito T, Ohiwa N, Tateoka M, Deocaris CC, Fujikawa T, Soya H (2007) Inhibitory effects of an orexin-2 receptor antagonist on orexin A- and stress-induced ACTH responses in conscious rats. *Neurosci Res* 57:462–466 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17188385> [Accessed April 20, 2015].
- Chen X, Wang H, Lin Z, Li S, Li Y, Bergen HT, Vrontakis ME, Kirouac GJ (2013) Orexins (hypocretins) contribute to fear and avoidance in rats exposed to a single episode of footshocks. *Brain Struct Funct* 219:2103–2118 Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23955372> [Accessed January 28, 2014].

de Lecea L, Kilduff TS, Peyron C, Gao X, Foye PE, Danielson PE, Fukuhara C, Battenberg EL, Gautvik VT, Bartlett FS, Frankel WN, van den Pol AN, Bloom FE, Gautvik KM, Sutcliffe JG (1998) The hypocretins: hypothalamus-specific peptides with neuroexcitatory activity. *Proc Natl Acad Sci U S A* 95:322–327 Available at: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=18213&tool=pmcentrez&rendertype=abstract> [Accessed January 23, 2015].

Farrell MS, Roth BL (2013) Pharmacosynthetics: Reimagining the pharmacogenetic approach. *Brain Res* 1511:6–20 Available at: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3562395&tool=pmcentrez&rendertype=abstract> [Accessed March 18, 2015].

Fujiki N, Yoshida Y, Ripley B, Honda K, Mignot E, Nishino S (2001) Changes in CSF hypocretin-1 (orexin A) levels in rats across 24 hours and in response to food deprivation. *Neuroreport* 12:993–997 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11303775> [Accessed February 25, 2015].

Furlong TM, Vianna DML, Liu L, Carrive P (2009) Hypocretin/orexin contributes to the expression of some but not all forms of stress and arousal. *Eur J Neurosci* 30:1603–1614 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19811530> [Accessed May 16, 2014].

Giardino WJ, de Lecea L (2014) Hypocretin (orexin) neuromodulation of stress and reward pathways. *Curr Opin Neurobiol* 29:103–108 Available at: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4267967&tool=pmcentrez&rendertype=abstract> [Accessed March 21, 2015].

Gotter AL, Forman MS, Harrell CM, Stevens J, Svetnik V, Yee KL, Li X, Roecker AJ, Fox S V., Tannenbaum PL, Garson SL, Lepelcure I De, Calder N, Rosen L, Struyk A, Coleman PJ, Herring WJ, Renger JJ, Winrow CJ (2016) Orexin 2 Receptor Antagonism is Sufficient to Promote NREM and REM Sleep from Mouse to Man. *Sci Rep* 6:27147 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/27256922> [Accessed February 13, 2017].

- Gotter AL, Roecker AJ, Hargreaves R, Coleman PJ, Winrow CJ, Renger JJ (2012a) Orexin receptors as therapeutic drug targets. *Prog Brain Res* 198:163–188 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22813974> [Accessed March 18, 2014].
- Gotter AL, Webber AL, Coleman PJ, Renger JJ, Winrow CJ (2012b) International Union of Basic and Clinical Pharmacology. LXXXVI. Orexin receptor function, nomenclature and pharmacology. *Pharmacol Rev* 64:389–420 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22759794> [Accessed March 18, 2014].
- Gozzi A, Lepore S, Vicentini E, Merlo-Pich E, Bifone A (2013) Differential effect of orexin-1 and CRF-1 antagonism on stress circuits: a fMRI study in the rat with the pharmacological stressor Yohimbine. *Neuropsychopharmacology* 38:2120–2130 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23736277> [Accessed January 28, 2014].
- Grissom N, Bhatnagar S (2009) Habituation to repeated stress: get used to it. *Neurobiol Learn Mem* 92:215–224 Available at: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2773683&tool=pmcentrez&rendertype=abstract> [Accessed March 20, 2014].
- Grissom N, Kerr W, Bhatnagar S (2008) Struggling behavior during restraint is regulated by stress experience. *Behav Brain Res* 191:219–226 Available at: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2477735&tool=pmcentrez&rendertype=abstract> [Accessed March 20, 2014].
- Hagan JJ et al. (1999) Orexin A activates locus coeruleus cell firing and increases arousal in the rat. *Proc Natl Acad Sci U S A* 96:10911–10916 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10485925> [Accessed September 18, 2017].
- Hasegawa E, Yanagisawa M, Sakurai T, Mieda M (2014) Orexin neurons suppress narcolepsy via 2 distinct efferent pathways. *J Clin Invest* 124:604–616 Available at: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3904620&tool=pmcentrez&rendertype=abstract> [Accessed March 31, 2014].
- Heydendael W, Sharma K, Iyer V, Luz S, Piel D, Beck S, Bhatnagar S (2011)

Orexins/hypocretins act in the posterior paraventricular thalamic nucleus during repeated stress to regulate facilitation to novel stress. *Endocrinology* 152:4738–4752 Available at:

<http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3230061&tool=pmcentrez&rendertype=abstract> [Accessed January 25, 2014].

Ida T, Nakahara K, Murakami T, Hanada R, Nakazato M, Murakami N (2000) Possible involvement of orexin in the stress reaction in rats. *Biochem Biophys Res Commun* 270:318–323 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10733946> [Accessed August 11, 2014].

Jászberényi M, Bujdosó E, Pataki I, Telegdy G (2000) Effects of orexins on the hypothalamic-pituitary-adrenal system. *J Neuroendocrinol* 12:1174–1178 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11106974> [Accessed August 11, 2014].

Johnson PL, Molosh A, Fitz SD, Truitt WA, Shekhar A (2012) Orexin, stress, and anxiety/panic states. *Prog Brain Res* 198:133–161 Available at: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3665356&tool=pmcentrez&rendertype=abstract> [Accessed January 27, 2014].

Johnson PL, Truitt W, Fitz SD, Minick PE, Dietrich A, Sanghani S, Träskman-Bendz L, Goddard AW, Brundin L, Shekhar A (2010) A key role for orexin in panic anxiety. *Nat Med* 16:111–115 Available at: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2832844&tool=pmcentrez&rendertype=abstract> [Accessed August 3, 2014].

Kotz C, Nixon J, Butterick T, Perez-Leighton C, Teske J, Billington C (2012) Brain orexin promotes obesity resistance. *Ann N Y Acad Sci* 1264:72–86 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22803681> [Accessed August 30, 2016].

Kozorovitskiy Y, Saunders A, Johnson CA, Lowell BB, Sabatini BL (2012) Recurrent network activity drives striatal synaptogenesis. *Nature* 485:646–650 Available at: <http://www.nature.com/doi/10.1038/nature11052> [Accessed January 5, 2017].

Kuru M, Ueta Y, Serino R, Nakazato M, Yamamoto Y, Shibuya I, Yamashita H (2000) Centrally administered orexin/hypocretin activates HPA axis in rats. *Neuroreport*

11:1977–1980 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10884055>
[Accessed January 6, 2016].

Le T, Liang Z, Patel H, Yu MH, Sivasubramaniam G, Slovitt M, Tanentzapf G, Mohanty N, Paul SM, Wu VM, Beitel GJ (2006) A new family of *Drosophila* balancer chromosomes with a w- dfd-GMR yellow fluorescent protein marker. *Genetics* 174:2255–2257 Available at:
<http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1698648&tool=pmcentrez&rendertype=abstract> [Accessed November 12, 2015].

Marcus JN, Aschkenasi CJ, Lee CE, Chemelli RM, Saper CB, Yanagisawa M, Elmquist JK (2001) Differential expression of orexin receptors 1 and 2 in the rat brain. *J Comp Neurol* 435:6–25 Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/11370008> [Accessed March 19, 2014].

McIntyre DC, Kent P, Hayley S, Merali Z, Anisman H (1999) Influence of psychogenic and neurogenic stressors on neuroendocrine and central monoamine activity in fast and slow kindling rats. *Brain Res* 840:65–74 Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/10517953> [Accessed January 7, 2016].

Nagahara T, Saitoh T, Kutsumura N, Irukayama-Tomobe Y, Ogawa Y, Kuroda D, Gouda H, Kumagai H, Fujii H, Yanagisawa M, Nagase H (2015) Design and Synthesis of Non-Peptide, Selective Orexin Receptor 2 Agonists. *J Med Chem* 58:7931–7937 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26267383>
[Accessed September 8, 2016].

Perez-Leighton CE, Billington CJ, Kotz CM (2014) Orexin modulation of adipose tissue. *Biochim Biophys Acta* 1842:440–445 Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/23791983> [Accessed August 30, 2016].

Peyron C, Tighe DK, van den Pol AN, de Lecea L, Heller HC, Sutcliffe JG, Kilduff TS (1998) Neurons containing hypocretin (orexin) project to multiple neuronal systems. *J Neurosci* 18:9996–10015 Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/9822755> [Accessed August 11, 2014].

Roecker AJ et al. (2014) Discovery of 5''-chloro-N-[(5,6-dimethoxypyridin-2-yl)methyl]-

2,2':5',3''-terpyridine-3'-carboxamide (MK-1064): a selective orexin 2 receptor antagonist (2-SORA) for the treatment of insomnia. *ChemMedChem* 9:311–322 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24376006> [Accessed January 15, 2015].

Russell SH, Small CJ, Dakin CL, Abbott CR, Morgan DG, Ghatei MA, Bloom SR (2001) The central effects of orexin-A in the hypothalamic-pituitary-adrenal axis in vivo and in vitro in male rats. *J Neuroendocrinol* 13:561–566 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11412343> [Accessed August 11, 2014].

Sakamoto F, Yamada S, Ueta Y (2004) Centrally administered orexin-A activates corticotropin-releasing factor-containing neurons in the hypothalamic paraventricular nucleus and central amygdaloid nucleus of rats: possible involvement of central orexins on stress-activated central CRF neurons. *Regul Pept* 118:183–191 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15003835> [Accessed February 15, 2015].

Sakurai T et al. (1998) Orexins and orexin receptors: a family of hypothalamic neuropeptides and G protein-coupled receptors that regulate feeding behavior. *Cell* 92:573–585 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9491897> [Accessed February 15, 2015].

Sakurai T (2014) The role of orexin in motivated behaviours. *Nat Rev Neurosci* 15:719–731 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25301357> [Accessed October 10, 2014].

Samson WK, Taylor MM, Follwell M, Ferguson A V (2002) Orexin actions in hypothalamic paraventricular nucleus: physiological consequences and cellular correlates. *Regul Pept* 104:97–103 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11830283> [Accessed August 11, 2014].

Scott MM, Marcus JN, Pettersen A, Birnbaum SG, Mochizuki T, Scammell TE, Nestler EJ, Elmquist JK, Lutter M (2011) Hcrtr1 and 2 signaling differentially regulates depression-like behaviors. *Behav Brain Res* 222:289–294 Available at: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3474296&tool=pmcentre>

z&rendertype=abstract [Accessed October 30, 2015].

Shirasaka T, Miyahara S, Kunitake T, Jin QH, Kato K, Takasaki M, Kannan H (2001) Orexin depolarizes rat hypothalamic paraventricular nucleus neurons. *Am J Physiol Regul Integr Comp Physiol* 281:R1114-8 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11557617> [Accessed June 5, 2015].

Spinazzi R, Andreis PG, Rossi GP, Nussdorfer GG (2006) Orexins in the regulation of the hypothalamic-pituitary-adrenal axis. *Pharmacol Rev* 58:46–57 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16507882> [Accessed August 5, 2014].

Taheri S, Mahmoodi M, Opacka-Juffry J, Ghatei MA, Bloom SR (1999) Distribution and quantification of immunoreactive orexin A in rat tissues. *FEBS Lett* 457:157–161 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10486585> [Accessed October 30, 2015].

Tannenbaum PL, Stevens J, Binns J, Savitz AT, Garson SL, Fox S V., Coleman P, Kuduk SD, Gotter AL, Marino M, Tye SJ, Uslaner JM, Winrow CJ, Renger JJ (2014) Orexin receptor antagonist-induced sleep does not impair the ability to wake in response to emotionally salient acoustic stimuli in dogs. *Front Behav Neurosci* 8:182 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24904334> [Accessed February 15, 2017].

Tannenbaum PL, Tye SJ, Stevens J, Gotter AL, Fox S V., Savitz AT, Coleman PJ, Uslaner JM, Kuduk SD, Hargreaves R, Winrow CJ, Renger JJ (2016) Inhibition of Orexin Signaling Promotes Sleep Yet Preserves Salient Arousability in Monkeys. *Sleep* 39:603–612 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26943466> [Accessed February 15, 2017].

Winsky-Sommerer R, Boutrel B, de Lecea L (2005) Stress and arousal: the corticotrophin-releasing factor/hypocretin circuitry. *Mol Neurobiol* 32:285–294 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16385142> [Accessed February 15, 2015].

Winsky-Sommerer R, Yamanaka A, Diano S, Borok E, Roberts AJ, Sakurai T, Kilduff TS, Horvath TL, de Lecea L (2004) Interaction between the corticotropin-releasing

factor system and hypocretins (orexins): a novel circuit mediating stress response. *J Neurosci* 24:11439–11448 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15601950> [Accessed August 11, 2014].

Zeitler JM (2013) Control of sleep and wakefulness in health and disease. *Prog Mol Biol Transl Sci* 119:137–154 Available at: <http://www.sciencedirect.com/science/article/pii/B9780123969712000063> [Accessed August 11, 2015].

Figure Legends

Fig. 1. Orexin levels habituate with repeated restraint stress (n = 10/group). (A) Left: Depiction of Restraint Paradigm and Blood collection. Right: Plasma ACTH data for Day 1 and Day 5 of restraint. Vehicle/Vehicle-treated animals display habituation over the 5 days of restraint stress, with decreased plasma ACTH values on day 5 compared to day 1 (2-Way ANOVA, Day: $F(1, 67) = 8.2, p = 0.006$; Time: $F(1,67) = 12.7, p < 0.0001$). (B) Total orexin neurons dual labeled with cFos on day 1 versus day 5 of restraint stress (T test, $p < 0.05$). Right, representative 10x magnification images of the lateral hypothalamus with cFos (black) and orexin (red) dual stain on Day 1 and 5 of restraint. (C) Orexin levels in the CSF on day 1 and 5 of restraint stress (T test, $p < 0.05$). * $p < 0.05$, ** $p < 0.01$, *** $p < 0.0001$

Fig. 2. Stimulation of orexins prior to each of five daily 30-min restraints (n = 10/group). (A) Representative images of DREADDs expression in the lateral hypothalamus at 20x

magnification. Orexin Neurons = red, DREADDs tag = green, orexin neurons transfected with virus = yellow. (B) Viral expression along four levels of Bregma (anterior to posterior) in a cartoon using rat brain atlas images. Each red dot represents 10 cells expressing the viral tag. (C) Orexin A levels in CSF following vehicle or CNO treatment (non-restraint stress conditions; T test, $p < 0.001$). (D) Total orexin neurons dual labeled with cFos in acute and repeated restraint stress following either vehicle or CNO pretreatment (2-Way ANOVA, Day: ($F(1,62) = 9.6$, $p < 0.001$; CNO: ($F(1,62) = 4.1$, $p < 0.05$, Interaction: ($F(1,62) = 0.01$, $p = 0.95$). (E) Orexin A levels in the CSF following acute and repeated restraint stress with either vehicle or CNO pretreatment (2-Way ANOVA, Day: ($F(1,37) = 26.47$, CNO: $p < 0.0001$; Interaction: ($F(1,37) = 15.05$, $p = 0.0004$; ($F(1,37) = 2.6$, $p = 0.11$). * $p < 0.05$, *** $p < 0.001$

Fig. 3. OX₂R underlies struggle behavior and body weight change during restraint stress ($n=7$ /group). (A) Timeline of experimental procedures. DREADDs-containing viruses were injected into the lateral hypothalamus 4 weeks prior to the start of the restraint paradigm. MK-1064 and CNO were administered orally or subcutaneously, respectively, 90 minutes prior to each 30-minute restraint for 5 consecutive days. Tail blood was collected at 0, 15, 30, and 60 minutes on day 1 of restraint for hormone assays. Struggle Behavior was quantified on Day 2 of restraint. Tail blood was collected at 0, 15, and 30 minutes on day 5 of restraint. Rats were euthanized at 60 minutes, with both blood and brain collected at this time. (B) Orexin activation by CNO increases struggle behavior (2-Way ANOVA, CNO: ($F(1,22) = 9.8$, $p = 0.005$; MK-1064: ($F(1,22) = 7.4$, $p = 0.013$; Interaction: ($F(1,22) = 4.7$, $p = 0.042$). (C) Body weight change after 5 days of restraint stress in rats differentially treated with CNO and/or MK-1064 (2-Way ANOVA, MK-1064: ($F(1,23) = 4.3$, $p = 0.049$; Interaction: ($F(1,23) = 3.1$, $p = 0.09$). ** $p < 0.01$

Fig. 4. OX₂R is involved in the ACTH response to acute restraint stress ($n = 12$ /group). ACTH levels were evaluated on the first day of restraint stress following vehicle/vehicle versus CNO/vehicle pretreatment to evaluate orexin neuron activation (A; no significance by ANOVA), vehicle/vehicle versus vehicle/MK-1064 pretreatment to

evaluate the role of endogenous OX₂R (B; significant: 2-Way ANOVA, MK-1064: (F(1,63) = 9.3, p = 0.003), CNO/MK-1064 versus CNO/vehicle pretreatment to assess the effect of exogenous OX₂R activity induction (C; significant: 2-Way ANOVA, Drug: F(1,71) = 4.6, p = 0.035), or Vehicle/MK-1064 versus CNO/MK-1064 pretreatment to understand the role of orexin-induced mechanisms other than OX₂R (D; no significance by ANOVA). *p<0.05, **p<0.01

Fig. 5. OX₂R and non-OX₂R mediated orexin mechanisms have opposing influences on ACTH responses to repeated restraint stress. (n=12/group) ACTH levels were evaluated on the fifth day of restraint stress following vehicle/vehicle versus CNO/vehicle pretreatment to evaluate orexin neuron activation (A; no significance by ANOVA), vehicle/vehicle versus vehicle/MK-1064 pretreatment to evaluate the role of endogenous OX₂R (B; no significance by ANOVA), CNO/MK-1064 versus CNO/vehicle pretreatment to assess the effect of exogenous OX₂R activity induction (C; significant: 2-Way ANOVA, Drug: F(1,45) = 13.1, p = 0.001), or Vehicle/MK-1064 versus CNO/MK-1064 pretreatment to understand the role of orexin-induced mechanisms other than OX₂R (D; significant: 2-Way ANOVA, Drug: F(1,40) = 12.02, p<0.01). *p<0.05, **p<0.01