

CRYSTALLOGRAPHIC
COMMUNICATIONS

ISSN 2056-9890

Received 14 March 2015

Accepted 18 March 2015

Edited by W. T. A. Harrison, University of
Aberdeen, Scotland

Keywords: crystal structure; clozapinium;
crystal structure; molecular configuration;
hydrogen bonding; supramolecular assembly

CCDC references: 1054724; 1054723;
1054722

Supporting information: this article has
supporting information at journals.iucr.org/e

The crystal structures of three clozapinium salts: different molecular configurations, and supramolecular assembly in one, two and three dimensions


Manpreet Kaur,^a Jerry P. Jasinski,^{b*} Hemmige S. Yathirajan,^a Channappa N. Kavitha^a and Christopher Glidewell^c

^aDepartment of Studies in Chemistry, University of Mysore, Manasagangotri, Mysore 570 006, India, ^bDepartment of Chemistry, Keene State College, 229 Main Street, Keene, NH 03435-2001, USA, and ^cSchool of Chemistry, University of St Andrews, St Andrews, Fife KY16 9ST, Scotland. *Correspondence e-mail: jjasinski@keene.edu

The structures of three salts derived from clozapine, 8-chloro-11-(4-methylpiperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine, are reported, namely, clozapinium 3,5-dinitrobenzoate dimethyl sulfoxide monosolvate, $C_{18}H_{20}ClN_4^+ \cdot C_7H_3N_2O_6^- \cdot C_2H_6OS$, (I), where the dimethyl sulfoxide component is disordered over two sets of atomic sites having occupancies 0.627 (2) and 0.373 (2); clozapinium hydrogen maleate 0.21-hydrate, $C_{18}H_{20}ClN_4^+ \cdot C_4H_3O_4^- \cdot 0.21H_2O$, (II), and clozapinium 2-hydroxybenzoate, $C_{18}H_{20}ClN_4^+ \cdot C_7H_5O_3^-$, (III). In all three salts, the protonation site is the methylated N atom of the piperazine ring, and the dimensions and conformations of the fused tricyclic system are very similar. However, differences are apparent in the piperazine component: in both compounds (II) and (III), the unprotonated N atom of this ring has a pyramidal geometry, but in compound (I) this atom has a planar geometry. In compound (III), both N-substituents in this ring occupy equatorial sites, but in compound (II) the fused tricyclic system occupies an axial site of the piperazine ring. The independent components of compound (I) are linked within the selected asymmetric unit by a combination of $N-H \cdots O$ and $C-H \cdots O$ hydrogen bonds, and these hydrogen-bonded aggregates are linked into chains by an aromatic $\pi-\pi$ stacking interaction. In compound (II), the components are linked into sheets by a combination of $O-H \cdots O$, $N-H \cdots O$ and $C-H \cdots O$ hydrogen bonds, and in compound (III), a combination of $N-H \cdots O$, $C-H \cdots O$ and $C-H \cdots N$ hydrogen bonds links the components into a three-dimensional framework structure. Comparisons are made with some similar compounds.

1. Chemical context

Clozapine, 8-chloro-11-(4-methylpiperazin-1-yl)-5*H*-dibenzo[*b,e*][1,4]diazepine, is a well established medication used in the treatment of schizophrenia which in general leads to a lower incidence of adverse side effects such as Parkinsonian-type symptoms than some other treatments (Breier *et al.*, 1994). Although structures have been reported for both the free base itself (Petcher & Weber, 1976) and for its doubly-protonated di-cation, as the dibromide salt (Fillers & Hawkinson, 1982), there appear to be no reports of the structures of any mono-protonated clozapine derivatives. Accordingly, we have now determined the structures of three such salts with a variety of counter-ions. Of these salts, the 3,5-dinitrobenzoate crystallizes from dimethyl sulfoxide as a stoichiometric monosolvate (I) (Fig. 1, Scheme 1); however, the hydrogen maleate crystallizes from the same solvent as a partial hydrate (II) (Fig. 2);



and the 2-hydroxybenzoate crystallizes from a 1:1 mixture of acetonitrile and methanol in a solvent-free form (III) (Fig. 3). A number of other such salts were prepared, but no crystals suitable for single-crystal X-ray diffraction have so far been obtained from these, despite attempts to prepare crystals using a range of solvents. The aims of the present study are firstly to establish the site of protonation in the mono-protonated cations; secondly, to compare the conformations of the clozapinium cations; and thirdly, to explore the supramolecular assembly in these three salts.





Figure 1
The independent components of compound (I), showing the atom-labelling scheme and the N—H···O hydrogen bonds within the selected asymmetric unit. Displacement ellipsoids are drawn at the 30% probability level and the major and minor orientations of the disordered dimethyl sulfoxide component, containing atoms S31 and S41, respectively, have occupancies 0.627 (2) and 0.373 (2).

selected asymmetric unit is a charge-assisted hydrogen bond (Gilli *et al.*, 1994) and it is nearly linear with fairly short H···O and N···O distances (Table 1). There are also some short C—H···O contacts between the cation and the disordered DMSO components, but those to the major component, in particular, have long H···O distances, and C—H···O angles which are less than 140° (*cf.* Wood *et al.*, 2009) so these may be better regarded as adventitious contacts rather than structurally significant interactions.

Compound (II) consists of a clozapinium cation, a hydrogen maleate anion which contains a short intra-anion O—H···O hydrogen bond (Table 1), and a partial occupancy water molecule whose refined site occupancy is 0.210 (7): despite the short O···O distance in the intra-anion hydrogen bond, the H atom is significantly displaced from a position equidistant from the two O atoms involved (Table 1). The two ionic components are linked by a nearly linear charge-assisted hydrogen bond (Fig. 2), while the partially occupied water site forms hydrogen bonds to two anions (Table 1).

2. Structural commentary

Compound (I) consists of a clozapinium cation, in which protonation has occurred at the protonated N atom of the piperazine ring, as is also observed in both of compounds (II) and (III), a 3,5-dinitrobenzoate anion, and a molecule of dimethyl sulfoxide (DMSO), which is disordered over two orientations having site occupancies of 0.627 (2) and 0.273 (2), respectively (Fig. 1). It was possible to select an asymmetric unit for (I) in which the three components are linked by N—H···O hydrogen bonds (Fig. 1, Table 1). The N—H···O hydrogen bond between the two ionic components within the

Table 1

Hydrogen bonds and short intramolecular contacts (Å, °) for compounds (I)–(III).

Cg1 and Cg2 represent the centroids of rings C5A/C6–C9/C9A and C1–C4/C4A/C11A, respectively

Compound	<i>D</i> –H... <i>A</i>	<i>D</i> –H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> –H... <i>A</i>
(I)	N5–H5...O31	0.83 (3)	2.10 (3)	2.921 (4)	170 (3)
	N14–H14...O22	1.00 (3)	1.58 (3)	2.575 (3)	176 (2)
	C4–H4...O31	0.95	2.58	3.342 (4)	137
	C4–H4...O41	0.95	2.38	3.208 (7)	146
	C6–H6...O31	0.95	2.54	3.313 (5)	139
(II)	N5–H5...O22 ⁱ	0.86 (3)	2.24 (3)	3.084 (2)	170 (3)
	N14–H14...O23	0.87 (3)	1.83 (2)	2.688 (2)	173 (2)
	O21–H21...O24	1.00 (5)	1.47 (5)	2.420 (3)	157 (5)
	O31–H31A...O24	0.90	2.05	2.913 (11)	160
	O31–H31B...O21 ⁱⁱ	0.90	2.37	3.232 (11)	161
	C12–H12A...O22 ⁱⁱⁱ	0.99	2.48	3.308 (2)	141
	C15–H15A...Cg1 ^{iv}	0.99	2.95	3.642 (2)	128
C15–H15B...Cg2 ^{iv}	0.99	2.95	3.759 (2)	139	
(III)	N14–H14...O22	0.89 (6)	1.75 (6)	2.612 (4)	164 (5)
	O23–H23...O21	1.01 (11)	1.52 (10)	2.507 (5)	161 (9)
	C4–H4...O22 ^v	0.95	2.23	3.261 (4)	166
	C9–H9...O22 ^v	0.95	2.25	3.202 (4)	176
	C12–H12A...O21	0.99	2.44	3.306 (5)	146
	C12–H12B...N5 ^{vi}	0.99	2.56	3.539 (4)	170
	C24–H24...Cg1 ^{vii}	0.95	2.83	3.637 (5)	144

Symmetry codes: (i) $-1 + x, \frac{3}{2} - y, -\frac{1}{2} + z$; (ii) $-x, 2 - y, 1 - z$; (iii) $2 - x, 1 - y, 1 - z$; (iv) $1 - x, \frac{1}{2} + y, \frac{1}{2} - z$; (v) $\frac{1}{2} + x, -\frac{1}{2} + y, z$; (vi) $x, 1 - y, \frac{1}{2} + z$; (vii) $-\frac{1}{2} + x, \frac{1}{2} + y, -1 + z$.

There is again an intra-anion O–H...O hydrogen bond in the 2-hydroxybenzoate component of compound (III) and, again, the two ionic components are linked by a fairly short, charge-assisted hydrogen bond (Fig. 3). It is of interest to note both the general similarity in the dimensions of the two intra-anion hydrogen bonds in compounds (II) and (III), and also that in each of (I)–(III), the site of the protonation of the




Figure 3
The independent components of compound (III), showing the atom-labelling scheme, and the O–H...O and N–H...O hydrogen bonds within the selected asymmetric unit. Displacement ellipsoids are drawn at the 30% probability level.

clozapine is the methylated atom N14 of the piperazine ring. In each case, this N–H bond participates in a short charge-assisted hydrogen bond between the ionic components. As discussed below, this is the only N–H...O hydrogen bond involving the ionic components in both compound (I) and compound (III).

In the clozapinium cations of compounds (I)–(III), the fused tricyclic units exhibit very similar conformations, as




Figure 2
The independent components of compound (II), showing the atom-labelling scheme, and the O–H...O and N–H...O hydrogen bonds within the selected asymmetric unit. Displacement ellipsoids are drawn at the 30% probability level and the water molecule, containing atom O31, has occupancy 0.210 (7).

Table 2
Selected geometric parameters (°) for compounds (I)–(III).

'Dihedral' denotes the dihedral angles between the mean planes of rings C1–C4/C4A/C11A and C5A/C6–C9/C9A.

Parameter	(I)	(II)	(III)
C11–N11–C12	120.80 (19)	118.45 (15)	118.3 (3)
C11–N11–C16	126.01 (19)	122.04 (15)	121.7 (3)
C12–N11–C16	112.91 (18)	111.09 (14)	111.0 (3)
C4A–N5–C5A–C6	–117.8 (2)	–116.17 (19)	–117.1 (3)
C5A–N5–C4A–C4	115.1 (3)	120.07 (19)	115.3 (3)
C1–C11A–C11–N10	–129.2 (3)	–137.5 (2)	–139.6 (3)
C9–C9A–N10–C11	146.3 (2)	141.11 (19)	141.9 (3)
C9A–N10–C11–C11A	–7.9 (4)	0.9 (3)	1.0 (5)
N10–C11–N11–C12	6.3 (3)	2.7 (3)	5.5 (4)
C11–N11–C12–C13	–120.8 (2)	–90.0 (2)	151.8 (3)
Dihedral	62.21 (11)	60.97 (9)	59.07 (16)




Figure 4

A stereoview of part of the crystal structure of compound (I), showing the formation of a π -stacked chain of hydrogen-bonded ion pairs. For the sake of clarity, H atoms not involved in the hydrogen bonds (shown as dashed lines) have been omitted, as have the disordered DMSO molecules.

shown by the relevant torsional and dihedral angles (Table 2) which define the relative orientations of the three rings. It is interesting to note that corresponding pairs of torsional angles involving either one or the other of the two aryl rings consistently have similar magnitudes but opposite signs, indicative of near mirror symmetry, provided the difference in the atomic types N10 and C11 is ignored, with the pseudo mirror containing the bond N5–H5 and passing through the midpoint of the bond N10–C11. However, there are some interesting differences between (I)–(III) in respect of the piperazine rings, which in each compound adopt a chair conformation, with protonation at the methylated atom N14, where the methyl atom C17 always occupies the equatorial site. While the geometry at N11 is planar within experimental uncertainty in compound (I), it is pyramidal in each of (II) and (III). In addition, the atom C11 (and hence the bulky tricyclic system) occupies the equatorial site at N11 in compound (III), but in compound (II) the tricyclic unit unexpectedly occupies the axial site at atom N11, as indicated by the values of the torsional angles C11–N11–C12–C13 for these two compounds (Table 2).

3. Supramolecular features

The supramolecular assembly in compounds (I)–(III) provides structures in one, two and three dimensions respectively. There are no hydrogen bonds in the structure of compound (I) other than those within the selected asymmetric unit (Table 1, Fig. 1). However, the hydrogen-bonded ionic components are linked into a chain by an aromatic π – π stacking interaction. The C5A,C6–C9,C9A ring in the cation at (x, y, z) makes a dihedral angle of only $1.34(12)^\circ$ with the C21–C26 ring in the anion at $(x, \frac{1}{2} - y, \frac{1}{2} + z)$. The distance between the centroids of these two rings is $3.4583(14) \text{ \AA}$ and the shortest perpendicular




Figure 5

Part of the crystal structure of compound (II) showing the formation of a centrosymmetric four-ion aggregate. For the sake of clarity, the unit-cell outline and the H atoms bonded to C atoms not involved in the motif shown have been omitted. Atoms marked with an asterisk (*) are at the symmetry position $(2 - x, 1 - y, 1 - z)$.

distance between the centroid of one ring and the plane of the other is $3.2761(1) \text{ \AA}$, corresponding to a ring-centroid offset of *ca* 1.11 \AA . This stacking interaction links the hydrogen-bonded ionic components into a chain running parallel to the [001] direction (Fig. 4). Two chains of this type, related to one another by inversion, pass through each unit cell, but there are no direction-specific interactions between adjacent chains. The DMSO molecules are pendent from the chains but otherwise play no part in the supramolecular assembly, so that their role may be largely that of filling otherwise empty cavities within the structure formed by the ionic components.

In compound (II) a combination of O–H...O, N–H...O and C–H...O hydrogen bonds (Table 1) links the independent components into complex sheets, but the sheet formation can readily be analysed in terms of a small number of fairly simple sub-structures (Ferguson *et al.*, 1998*a,b*; Gregson *et al.*, 2000). Ion pairs (Fig. 2) which are related by inversion are linked by C–H...O hydrogen bonds, forming a cyclic centrosymmetric aggregate characterized by an $R_4^4(22)$ (Bernstein *et al.*, 1995) motif, with the reference aggregate centred at $(1, \frac{1}{2}, \frac{1}{2})$ (Fig. 5). In a second sub-structure, ion pairs which are related by a glide plane are linked by N–H...O




Figure 6
A stereoview of part of the crystal structure of compound (II) showing the formation of hydrogen-bonded $C_2^2(16)$ chain parallel to [201]. For the sake of clarity, H atoms bonded to C atoms and the partial-occupancy water molecules have been omitted.

hydrogen bonds to form a $C_2^2(16)$ chain running parallel to the [201] direction (Fig. 6). This chain motif directly links the reference four-ion aggregate (Fig. 5) centred at $(1, \frac{1}{2}, \frac{1}{2})$ to the four symmetry-related aggregates centred at $(0, 0, 0)$, $(0, 1, 0)$, $(2, 0, 1)$ and $(2, 1, 1)$, so forming a sheet lying parallel to $(1\ 0\ \bar{2})$ (Fig. 7). Embedded within this sheet is a further cyclic motif, which is formally centrosymmetric, containing two anions and two water molecules. However, since the occupancy of the water sites is only 0.210 (7), if either of the two water sites in this motif is occupied there is a high probability that the other such site will be unoccupied: indeed, in the majority of cases, neither site will be occupied.




Figure 7
A stereoview of part of the crystal structure of compound (II) showing the formation of a hydrogen-bonded sheet lying parallel to $(10\bar{2})$. For the sake of clarity, H atoms bonded to C atoms not involved in the motifs shown have been omitted.




Figure 8
A stereoview of part of the crystal structure of compound (III) showing the formation of a hydrogen-bonded $C_2^2(11)$ chain running parallel to $[1\bar{1}0]$. For the sake of clarity, H atoms not involved in the motif shown have been omitted.

The independent components of compound (III) are linked into a three-dimensional framework structure by a combination of $N-H\cdots O$, $O-H\cdots O$ and $C-H\cdots N$ hydrogen bonds (Table 1), and again the formation of the framework is most readily analysed in terms of three one-dimensional substructures. In the simplest of these sub-structures, the $C-H\cdots O$ hydrogen bond involving atom C4 links ion pairs related by translation into a $C_2^2(11)$ chain running parallel to the $[1\bar{1}0]$ direction (Fig. 8). The second sub-structure involves both $C-H\cdots N$ and $C-H\cdots O$ hydrogen bonds: cations related by the c -glide plane at $y = 0.5$ are linked by $C-H\cdots N$ hydrogen bonds into $C(7)$ chains running parallel to the [001] direction, and similarly related ion pairs are linked by the $C-H\cdots O$ hydrogen bond involving atom C9 to form a $C_2^2(11)$ chain also running parallel to [001], such that the combined effect of these two hydrogen bonds generates a $C(7)\ C_2^2(11)[R_3^2(19)]$





Figure 9
A stereoview of part of the crystal structure of compound (III) showing the formation of a hydrogen-bonded $C(7)\ C_2^2(11)[R_3^2(19)]$ chain of rings running parallel to [001]. For the sake of clarity, H atoms bonded to the C atoms not involved in the motif shown have been omitted.



Figure 10

A stereoview of part of the crystal structure of compound (III) showing the formation of a hydrogen-bonded chain running parallel to [101]. For the sake of clarity, H atoms bonded to the C atoms not involved in the motif shown have been omitted.

chain of rings running parallel to [001] (Fig. 9). Finally, the alternating action of the hydrogen bonds involving, atom C4 on the one hand, and atoms C9 and C15 on the other (Table 1) generates a complex chain running parallel to the [101] direction (Fig. 10).

4. Database survey

It is of interest briefly to compare the structures reported here for the salts (I)–(III) with those of some closely related analogues, in particular clozapine itself, compound (IV) (see Scheme 2) and the dihydrobromide salt (V).



In the free base clozapine, which crystallizes in the space group $P2_12_12_1$ (Petcher & Weber, 1976), the geometry at the

piperazinyl N atom corresponding to atom N11 in compounds (I)–(III) is very nearly planar, with a sum of interbond angles of $357(2)^\circ$, and there are no direction-specific interactions between the molecules: in particular, the N–H bond does not participate in any kind of hydrogen bond. In the salt (V), which crystallizes in space group $P2_1/c$ (Fillers & Hawkinson, 1982), the protonation sites are the N atoms corresponding to atoms N10 and N14 in compounds (I)–(III), so that the doubly bonded N atom of the diazepine ring is protonated in preference to the second N atom of the piperazine ring where, as in (IV), the geometry is nearly planar, with a sum of interbond angles of $357(2)^\circ$: the individual ionic components in (V) are linked by charge-assisted N–H \cdots Br hydrogen bonds, such that cations related by a 2_1 screw axis are bridged by one of the two independent anions to form a $C_2^1(7)$ chain, from which the anions of the second type are pendent. Loxapine, compound (VI), is similar to clozapine but differs from it in two respects: the nature of the hetero-atoms in the seven-membered ring, and the location of the Cl substituent. Here again there are no direction-specific interactions between the molecules (Petcher & Weber, 1976). The overall molecular shapes of the molecules of compounds (IV) and (VI) are extremely similar, and it was suggested (Petcher & Weber, 1976) that the structures observed in the solid state represented a preferred form which persists in aqueous solutions and at the site of neuroleptic receptors. However, in the presence of charge-assisted hydrogen bonds in compounds (I)–(III), reported here, which are probably slightly stronger than those between the molecules of (IV) and (VI) and adjacent water molecules in solution, the molecular configurations show some significant differences, as noted above, so that no preferred configuration is apparent from the structures of (I)–(III).

5. Synthesis and crystallization

Clozapine was a gift from R L Fine Chem, Bengaluru, Karnataka, India. Equimolar quantities of clozapine and the appropriate acid (1.53 mmol of each component) were dissolved in methanol at 333 K. The solutions were permitted to cool to ambient temperature, and the resulting crystals were then collected by filtration, and dried over phosphorus(V) oxide. Crystals suitable for single-crystal X-ray diffraction were obtained by slow evaporation, at ambient temperature and in the presence of air, of solutions in dimethyl sulfoxide, for compounds (I) and (II), and a mixture (1:1 v/v) of acetonitrile and methanol for compound (III).

6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 3. The H atoms bonded to C or N atoms in the ionic components of compounds (I)–(III) were all located in difference maps. The H atoms bonded to C atoms were then treated as riding atoms in geometrically idealized positions with C–H distances of 0.95 Å (alkenyl and aromatic), 0.98 Å (CH₃) or 0.99 Å (CH₂) and with $U_{iso}(H) =$

Table 3
Experimental details.

	(I)	(II)	(III)
Crystal data			
Chemical formula	$C_{18}H_{20}ClN_4^+ \cdot C_7H_3N_2O_6^- \cdot C_2H_6OS$	$C_{18}H_{20}ClN_4^+ \cdot C_4H_3O_4^- \cdot 0.21H_2O$	$C_{18}H_{20}ClN_4^+ \cdot C_7H_5O_3^-$
M_r	617.07	446.68	464.94
Crystal system, space group	Monoclinic, $P2_1/c$	Monoclinic, $P2_1/c$	Monoclinic, Cc
Temperature (K)	173	173	173
a, b, c (Å)	15.3593 (2), 15.7685 (2), 11.8679 (2)	9.7166 (3), 9.9699 (2), 23.1059 (6)	17.4296 (5), 15.3728 (5), 8.6359 (3)
β (°)	91.8097 (14)	96.800 (3)	90.325 (3)
V (Å ³)	2872.89 (7)	2222.60 (10)	2313.88 (13)
Z	4	4	4
Radiation type	Cu $K\alpha$	Cu $K\alpha$	Cu $K\alpha$
μ (mm ⁻¹)	2.34	1.84	1.75
Crystal size (mm)	0.26 × 0.22 × 0.18	0.46 × 0.32 × 0.22	0.42 × 0.36 × 0.20
Data collection			
Diffractionmeter	Agilent Eos Gemini	Agilent Eos Gemini	Agilent Eos Gemini
Absorption correction	Multi-scan (<i>CrysAlis RED</i> ; Agilent, 2012)	Multi-scan (<i>CrysAlis RED</i> ; Agilent, 2012)	Multi-scan (<i>CrysAlis RED</i> ; Agilent, 2012)
T_{min}, T_{max}	0.424, 0.656	0.440, 0.668	0.399, 0.705
No. of measured, independent and observed [$I > 2\sigma(I)$] reflections	19784, 5527, 4623	8634, 4228, 3552	7244, 4069, 3962
R_{int}	0.032	0.026	0.048
$(\sin \theta/\lambda)_{max}$ (Å ⁻¹)	0.614	0.614	0.619
Refinement			
$R[F^2 > 2\sigma(F^2)], wR(F^2), S$	0.056, 0.152, 1.04	0.048, 0.133, 1.03	0.056, 0.141, 1.08
No. of reflections	5527	4228	4069
No. of parameters	409	295	308
No. of restraints	6	0	2
H-atom treatment	H atoms treated by a mixture of independent and constrained refinement	H atoms treated by a mixture of independent and constrained refinement	H atoms treated by a mixture of independent and constrained refinement
$\Delta\rho_{max}, \Delta\rho_{min}$ (e Å ⁻³)	0.89, -0.97	0.46, -0.30	0.53, -0.36
Absolute structure	–	–	Flack x determined using 1674 quotients $[(I^+) - (I^-)] / [(I^+) + (I^-)]$ (Parsons <i>et al.</i> , 2013)
Absolute structure parameter	–	–	-0.022 (17)

Computer programs: *CrysAlis PRO* and *CrysAlis RED* (Agilent, 2012), *SHELXS97* (Sheldrick, 2008), *SHELXL2014* (Sheldrick, 2015) and *PLATON* (Spek, 2009).

$kU_{eq}(C)$, where $k = 1.5$ for the methyl groups, which were permitted to rotate but not to tilt, and 1.2 for all other H atoms bonded to C atoms. For the H atoms bonded to N or O atoms, the atomic coordinates were refined with $U_{iso}(H) = 1.2U_{eq}(N)$ or $1.5U_{eq}(O)$, giving the N–H and O–H distances shown in Table 1.

In compound (I), the dimethyl sulfoxide component is disordered over two sets of atomic sites having unequal occupancy (*cf.* Fig. 1). For the minor disorder component, the bonded distances and the one-angle non-bonded distances were all restrained to be the same as the corresponding distances in the major component subject to uncertainties of 0.005 Å and 0.01 Å respectively. The anisotropic displacement parameters for those pairs of partial-occupancy C and O atoms occupying essentially the same physical space were constrained to be identical, and the H atoms of the dimethyl sulfoxide components were included as riding atoms with C–H distances 0.95 Å and $U_{iso}(H) = 1.5U_{eq}(C)$. Subject to these conditions, independent refinement of the site occupancies for the two disorder components gave values of 0.613 (3) and 0.359 (3); thereafter the occupancies were constrained to sum to unity, giving final values of 0.627 (2) and 0.373 (2). At this

stage of the refinements there were no significant features in the difference maps for compounds (I) and (III), but for (II) there was a single significant peak, $1.51 e \text{ \AA}^{-3}$, which was within plausible hydrogen-bonding distance of two O atoms. Examination of the structures of compounds (I) and (III) using *PLATON* (Spek, 2009) showed that there were no solvent-accessible voids in these structures. However, in compound (II), there was a total void volume of *ca* 88 Å³ per unit cell, and examination of the structure of (II) using the *SQUEEZE* tool (Spek, 2015) within *PLATON* disclosed the presence of an addition 8.8 electrons per unit cell, equivalent to 0.22 molecules of water per ion pair. Accordingly, the large residual was modeled as the O atom, denoted O31, of a partial occupancy water molecule, which was refined isotropically: it was not possible to locate the H atoms associated with atom O31 in difference maps, but they were included in calculated positions with O–H 0.90 Å and $U_{iso}(H) = 1.5U_{iso}(O)$. Subject to these conditions, the occupancy of the water molecule refined to a value of 0.210 (7), very close to that indicated by *SQUEEZE*. It should be emphasized here that the application of the *SQUEEZE* procedure referred to above was intended only to estimate the number of electrons not yet accounted for

at that stage of the refinement, and that the refinements at every stage were undertaken with the original data, independent of SQUEEZE. For compound (III), the correct orientation of the structure with respect to the polar axis directions was established by means of the Flack x parameter (Flack, 1983), $x = -0.022$ (17), calculated (Parsons *et al.*, 2013) using 1674 quotients of the type $[(I^+) - (I^-)] / [(I^+) + (I^-)]$.

Acknowledgements

MK thanks the UOM for research facilities and is also grateful to CPEPA, UGC, for the award of a JRF. JPJ acknowledges the NSF-MRI program (grant No. 1039027) for funds to purchase the X-ray diffractometer.

References

- Agilent (2012). *CrysAlis PRO* and *CrysAlis RED*. Agilent Technologies Ltd, Yarnton, England.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Breier, A., Buchanan, R. W., Kirkpatrick, B., David, O. R., Irish, D., Summerfelt, A. & Carpenter, J. W. Jr (1994). *Am. J. Psychiatry*, **151**, 20–26.
- Ferguson, G., Glidewell, C., Gregson, R. M. & Meehan, P. R. (1998a). *Acta Cryst.* **B54**, 129–138.
- Ferguson, G., Glidewell, C., Gregson, R. M. & Meehan, P. R. (1998b). *Acta Cryst.* **B54**, 139–150.
- Fillers, J. P. & Hawkinson, S. W. (1982). *Acta Cryst.* **B38**, 1750–1753.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Gilli, P., Bertolasi, V., Ferretti, V. & Gilli, G. (1994). *J. Am. Chem. Soc.* **116**, 909–915.
- Gregson, R. M., Glidewell, C., Ferguson, G. & Lough, A. J. (2000). *Acta Cryst.* **B56**, 39–57.
- Parsons, S., Flack, H. D. & Wagner, T. (2013). *Acta Cryst.* **B69**, 249–259.
- Petcher, T. J. & Weber, H.-P. (1976). *J. Chem. Soc. Perkin Trans. 2*, pp. 1415–1420.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Spek, A. L. (2015). *Acta Cryst.* **C71**, 9–18.
- Wood, P. A., Allen, F. H. & Pidcock, E. (2009). *CrystEngComm*, **11**, 1563–1571.

supporting information

Acta Cryst. (2015). E71, 406–413 [doi:10.1107/S205698901500554X]

The crystal structures of three clozapinium salts: different molecular configurations, and supramolecular assembly in one, two and three dimensions

Manpreet Kaur, Jerry P. Jasinski, Hemmige S. Yathirajan, Channappa N. Kavitha and Christopher Glidewell

Computing details

For all compounds, data collection: *CrysAlis PRO* (Agilent, 2012); cell refinement: *CrysAlis PRO* (Agilent, 2012); data reduction: *CrysAlis RED* (Agilent, 2012); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL2014* (Sheldrick, 2015) and *PLATON* (Spek, 2009).

(I) Clozapinium 3,5-dinitrobenzoate dimethyl sulfoxide monosolvate

Crystal data

$C_{18}H_{20}ClN_4^+ \cdot C_7H_3N_2O_6^- \cdot C_2H_6OS$

$M_r = 617.07$

Monoclinic, $P2_1/c$

$a = 15.3593$ (2) Å

$b = 15.7685$ (2) Å

$c = 11.8679$ (2) Å

$\beta = 91.8097$ (14)°

$V = 2872.89$ (7) Å³

$Z = 4$

$F(000) = 1288$

$D_x = 1.427$ Mg m⁻³

Cu $K\alpha$ radiation, $\lambda = 1.54184$ Å

Cell parameters from 5527 reflections

$\theta = 4.0$ – 71.3°

$\mu = 2.34$ mm⁻¹

$T = 173$ K

Block, colourless

$0.26 \times 0.22 \times 0.18$ mm

Data collection

Agilent Eos Gemini
diffractometer

Radiation source: Enhance (Cu) X-ray Source

ω scans

Absorption correction: multi-scan
(*CrysAlis RED*; Agilent, 2012)

$T_{\min} = 0.424$, $T_{\max} = 0.656$

19784 measured reflections

5527 independent reflections

4623 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.032$

$\theta_{\max} = 71.3^\circ$, $\theta_{\min} = 4.0^\circ$

$h = -17 \rightarrow 18$

$k = -19 \rightarrow 15$

$l = -14 \rightarrow 14$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.056$

$wR(F^2) = 0.152$

$S = 1.04$

5527 reflections

409 parameters

6 restraints

Hydrogen site location: difference Fourier map

H atoms treated by a mixture of independent
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0639P)^2 + 2.7986P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.89$ e Å⁻³

$\Delta\rho_{\min} = -0.97$ e Å⁻³

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C1	0.35763 (15)	0.19065 (15)	0.64680 (19)	0.0349 (5)	
H1	0.3526	0.1546	0.5828	0.042*	
C2	0.40839 (15)	0.16561 (16)	0.7394 (2)	0.0396 (5)	
H2	0.4388	0.1132	0.7385	0.048*	
C3	0.41440 (16)	0.21776 (18)	0.8335 (2)	0.0428 (6)	
H3	0.4452	0.1990	0.8996	0.051*	
C4	0.37586 (16)	0.29647 (17)	0.8315 (2)	0.0406 (5)	
H4	0.3830	0.3332	0.8945	0.049*	
C4A	0.32657 (14)	0.32265 (15)	0.7379 (2)	0.0344 (5)	
N5	0.28834 (14)	0.40442 (13)	0.73413 (19)	0.0398 (5)	
H5	0.296 (2)	0.4270 (19)	0.797 (3)	0.048*	
C5A	0.32343 (14)	0.45611 (15)	0.6491 (2)	0.0374 (5)	
C6	0.36736 (17)	0.53078 (17)	0.6785 (3)	0.0479 (6)	
H6	0.3744	0.5458	0.7557	0.057*	
C7	0.40067 (17)	0.58291 (18)	0.5976 (3)	0.0534 (7)	
H7	0.4290	0.6345	0.6180	0.064*	
C8	0.39217 (16)	0.55887 (17)	0.4864 (3)	0.0491 (7)	
C18	0.43801 (6)	0.62363 (5)	0.38475 (8)	0.0715 (3)	
C9	0.34849 (16)	0.48559 (16)	0.4543 (2)	0.0422 (6)	
H9	0.3430	0.4709	0.3767	0.051*	
C9A	0.31242 (14)	0.43313 (14)	0.5354 (2)	0.0355 (5)	
N10	0.25884 (12)	0.36645 (12)	0.49775 (17)	0.0361 (4)	
C11	0.25516 (14)	0.29455 (14)	0.54945 (19)	0.0333 (5)	
C11A	0.31376 (14)	0.26801 (14)	0.64620 (19)	0.0328 (5)	
N11	0.19690 (14)	0.23520 (13)	0.50890 (17)	0.0393 (5)	
C12	0.14714 (16)	0.24954 (16)	0.4036 (2)	0.0399 (5)	
H12A	0.1621	0.2052	0.3484	0.048*	
H12B	0.1634	0.3052	0.3720	0.048*	
C13	0.05023 (16)	0.24786 (15)	0.4213 (2)	0.0396 (5)	
H13A	0.0340	0.2952	0.4712	0.047*	
H13B	0.0184	0.2551	0.3480	0.047*	
N14	0.02552 (12)	0.16556 (12)	0.47312 (16)	0.0344 (4)	
H14	0.0420 (18)	0.1196 (18)	0.420 (2)	0.041*	
C15	0.07363 (16)	0.15561 (16)	0.5833 (2)	0.0387 (5)	
H15A	0.0576	0.1010	0.6181	0.046*	
H15B	0.0570	0.2018	0.6348	0.046*	
C16	0.17085 (16)	0.15805 (15)	0.5671 (2)	0.0375 (5)	
H16A	0.2016	0.1551	0.6415	0.045*	
H16B	0.1882	0.1080	0.5227	0.045*	

C17	-0.06991 (17)	0.1565 (2)	0.4843 (3)	0.0531 (7)	
H17A	-0.0826	0.1017	0.5190	0.080*	
H17B	-0.0987	0.1595	0.4095	0.080*	
H17C	-0.0915	0.2024	0.5316	0.080*	
C21	0.10211 (14)	-0.00223 (14)	0.1606 (2)	0.0328 (5)	
C22	0.14232 (15)	-0.07486 (15)	0.2025 (2)	0.0392 (5)	
H22	0.1445	-0.0862	0.2811	0.047*	
C23	0.17913 (16)	-0.13049 (15)	0.1277 (3)	0.0457 (6)	
C24	0.17770 (16)	-0.11803 (18)	0.0129 (3)	0.0483 (7)	
H24	0.2032	-0.1573	-0.0370	0.058*	
C25	0.13710 (15)	-0.04533 (18)	-0.0255 (2)	0.0431 (6)	
C26	0.09975 (14)	0.01300 (15)	0.0456 (2)	0.0362 (5)	
H26	0.0729	0.0628	0.0158	0.043*	
C27	0.06407 (15)	0.06224 (15)	0.2402 (2)	0.0359 (5)	
O21	0.03166 (12)	0.12736 (11)	0.19957 (16)	0.0466 (4)	
O22	0.07009 (12)	0.04261 (11)	0.34434 (15)	0.0442 (4)	
N23	0.22603 (19)	-0.20507 (16)	0.1753 (3)	0.0687 (8)	
O23	0.2293 (3)	-0.21397 (19)	0.2776 (3)	0.1090 (11)	
O24	0.26036 (17)	-0.25228 (15)	0.1091 (3)	0.0895 (9)	
N25	0.13695 (16)	-0.0268 (2)	-0.1471 (2)	0.0619 (7)	
O25	0.17166 (18)	-0.0786 (2)	-0.2086 (2)	0.1021 (11)	
O26	0.10244 (16)	0.03914 (18)	-0.18044 (18)	0.0696 (7)	
S31	0.31808 (7)	0.56967 (6)	1.00103 (8)	0.0448 (3)	0.627 (2)
O31	0.3389 (3)	0.4857 (2)	0.9475 (3)	0.0655 (11)	0.627 (2)
C31	0.4020 (7)	0.6409 (4)	0.9689 (7)	0.083 (3)	0.627 (2)
H31A	0.3963	0.6574	0.8894	0.125*	0.627 (2)
H31B	0.4586	0.6135	0.9830	0.125*	0.627 (2)
H31C	0.3979	0.6914	1.0165	0.125*	0.627 (2)
C32	0.3429 (6)	0.5547 (4)	1.1451 (4)	0.081 (2)	0.627 (2)
H32A	0.4050	0.5416	1.1557	0.122*	0.627 (2)
H32B	0.3082	0.5076	1.1735	0.122*	0.627 (2)
H32C	0.3293	0.6066	1.1865	0.122*	0.627 (2)
S41	0.39889 (18)	0.53363 (13)	1.05795 (17)	0.0678 (8)	0.373 (2)
O41	0.3879 (5)	0.4760 (4)	0.9582 (6)	0.0655 (11)	0.373 (2)
C41	0.4247 (12)	0.6356 (6)	1.0098 (12)	0.083 (3)	0.373 (2)
H41A	0.3746	0.6590	0.9670	0.125*	0.373 (2)
H41B	0.4749	0.6323	0.9610	0.125*	0.373 (2)
H41C	0.4388	0.6723	1.0744	0.125*	0.373 (2)
C42	0.2944 (5)	0.5526 (6)	1.1067 (8)	0.081 (2)	0.373 (2)
H42A	0.2725	0.5012	1.1426	0.122*	0.373 (2)
H42B	0.2554	0.5683	1.0432	0.122*	0.373 (2)
H42C	0.2967	0.5989	1.1618	0.122*	0.373 (2)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0350 (11)	0.0347 (11)	0.0348 (11)	0.0028 (9)	-0.0013 (9)	-0.0032 (9)
C2	0.0350 (12)	0.0400 (13)	0.0437 (13)	0.0097 (10)	-0.0005 (10)	0.0000 (10)

C3	0.0363 (12)	0.0554 (15)	0.0363 (12)	0.0065 (11)	-0.0053 (10)	0.0002 (11)
C4	0.0372 (12)	0.0489 (14)	0.0357 (12)	0.0012 (11)	-0.0011 (10)	-0.0089 (10)
C4A	0.0283 (11)	0.0352 (12)	0.0398 (12)	0.0012 (9)	0.0030 (9)	-0.0039 (9)
N5	0.0391 (11)	0.0353 (11)	0.0452 (12)	0.0046 (9)	0.0042 (9)	-0.0088 (9)
C5A	0.0257 (10)	0.0322 (11)	0.0543 (14)	0.0039 (9)	0.0006 (10)	-0.0038 (10)
C6	0.0348 (13)	0.0416 (14)	0.0670 (17)	-0.0027 (11)	-0.0037 (12)	-0.0089 (12)
C7	0.0365 (13)	0.0398 (14)	0.084 (2)	-0.0080 (11)	-0.0040 (13)	-0.0051 (14)
C8	0.0301 (12)	0.0383 (13)	0.079 (2)	-0.0014 (10)	0.0071 (12)	0.0105 (13)
C18	0.0632 (5)	0.0533 (4)	0.0991 (6)	-0.0144 (4)	0.0176 (4)	0.0167 (4)
C9	0.0332 (12)	0.0361 (12)	0.0573 (15)	0.0052 (10)	0.0037 (11)	0.0031 (11)
C9A	0.0237 (10)	0.0291 (11)	0.0534 (14)	0.0042 (9)	-0.0015 (9)	0.0004 (10)
N10	0.0305 (9)	0.0315 (10)	0.0459 (11)	0.0015 (8)	-0.0031 (8)	0.0014 (8)
C11	0.0293 (11)	0.0331 (11)	0.0373 (12)	0.0029 (9)	-0.0015 (9)	-0.0012 (9)
C11A	0.0288 (10)	0.0328 (11)	0.0365 (11)	0.0005 (9)	-0.0016 (9)	-0.0009 (9)
N11	0.0423 (11)	0.0353 (10)	0.0394 (10)	-0.0076 (9)	-0.0112 (9)	0.0057 (8)
C12	0.0437 (13)	0.0390 (13)	0.0362 (12)	-0.0066 (10)	-0.0098 (10)	0.0055 (10)
C13	0.0435 (13)	0.0321 (12)	0.0426 (13)	0.0041 (10)	-0.0073 (10)	0.0016 (10)
N14	0.0316 (10)	0.0342 (10)	0.0373 (10)	0.0035 (8)	-0.0008 (8)	-0.0004 (8)
C15	0.0434 (13)	0.0372 (12)	0.0354 (12)	0.0004 (10)	-0.0015 (10)	0.0017 (10)
C16	0.0402 (12)	0.0327 (12)	0.0389 (12)	-0.0025 (10)	-0.0094 (10)	0.0053 (9)
C17	0.0336 (13)	0.0621 (18)	0.0636 (18)	0.0058 (12)	0.0029 (12)	0.0058 (14)
C21	0.0254 (10)	0.0292 (11)	0.0436 (12)	-0.0045 (9)	-0.0015 (9)	-0.0033 (9)
C22	0.0352 (12)	0.0313 (11)	0.0510 (14)	-0.0026 (10)	-0.0029 (10)	-0.0001 (10)
C23	0.0332 (12)	0.0296 (12)	0.0742 (19)	-0.0001 (10)	-0.0026 (12)	-0.0076 (12)
C24	0.0305 (12)	0.0480 (15)	0.0665 (18)	-0.0061 (11)	0.0048 (11)	-0.0243 (13)
C25	0.0290 (11)	0.0542 (15)	0.0459 (14)	-0.0113 (11)	0.0010 (10)	-0.0122 (11)
C26	0.0262 (11)	0.0364 (12)	0.0458 (13)	-0.0076 (9)	-0.0025 (9)	-0.0015 (10)
C27	0.0308 (11)	0.0323 (12)	0.0445 (13)	-0.0032 (9)	-0.0008 (9)	-0.0059 (10)
O21	0.0480 (10)	0.0359 (9)	0.0554 (11)	0.0104 (8)	-0.0054 (8)	-0.0052 (8)
O22	0.0519 (10)	0.0388 (9)	0.0420 (9)	0.0027 (8)	0.0013 (8)	-0.0051 (7)
N23	0.0586 (16)	0.0345 (13)	0.113 (3)	0.0082 (12)	-0.0038 (16)	-0.0085 (15)
O23	0.142 (3)	0.0668 (18)	0.117 (3)	0.0490 (19)	-0.017 (2)	0.0159 (17)
O24	0.0714 (16)	0.0442 (12)	0.153 (3)	0.0188 (12)	0.0088 (16)	-0.0227 (15)
N25	0.0389 (12)	0.100 (2)	0.0472 (13)	-0.0174 (14)	0.0037 (10)	-0.0139 (15)
O25	0.0687 (16)	0.179 (3)	0.0587 (15)	0.0199 (19)	0.0102 (12)	-0.0409 (18)
O26	0.0705 (15)	0.0915 (18)	0.0462 (12)	-0.0312 (14)	-0.0086 (10)	0.0062 (12)
S31	0.0570 (7)	0.0362 (5)	0.0410 (5)	-0.0028 (4)	-0.0024 (4)	-0.0033 (4)
O31	0.094 (3)	0.0484 (16)	0.0543 (16)	-0.009 (2)	0.009 (2)	-0.0199 (13)
C31	0.081 (6)	0.060 (2)	0.111 (7)	-0.030 (3)	0.040 (6)	-0.037 (3)
C32	0.145 (7)	0.054 (2)	0.044 (3)	0.021 (4)	-0.006 (3)	-0.002 (2)
S41	0.109 (2)	0.0467 (11)	0.0475 (11)	0.0257 (11)	0.0001 (11)	-0.0104 (8)
O41	0.094 (3)	0.0484 (16)	0.0543 (16)	-0.009 (2)	0.009 (2)	-0.0199 (13)
C41	0.081 (6)	0.060 (2)	0.111 (7)	-0.030 (3)	0.040 (6)	-0.037 (3)
C42	0.145 (7)	0.054 (2)	0.044 (3)	0.021 (4)	-0.006 (3)	-0.002 (2)

Geometric parameters (Å, °)

C1—C2	1.385 (3)	C16—H16A	0.9900
C1—C11A	1.394 (3)	C16—H16B	0.9900
C1—H1	0.9500	C17—H17A	0.9800
C2—C3	1.388 (4)	C17—H17B	0.9800
C2—H2	0.9500	C17—H17C	0.9800
C3—C4	1.375 (4)	C21—C22	1.386 (3)
C3—H3	0.9500	C21—C26	1.386 (3)
C4—C4A	1.387 (3)	C21—C27	1.517 (3)
C4—H4	0.9500	C22—C23	1.381 (4)
C4A—C11A	1.397 (3)	C22—H22	0.9500
C4A—N5	1.417 (3)	C23—C24	1.376 (4)
N5—C5A	1.417 (3)	C23—N23	1.482 (4)
N5—H5	0.83 (3)	C24—C25	1.376 (4)
C5A—C6	1.396 (3)	C24—H24	0.9500
C5A—C9A	1.402 (4)	C25—C26	1.384 (4)
C6—C7	1.375 (4)	C25—N25	1.473 (4)
C6—H6	0.9500	C26—H26	0.9500
C7—C8	1.375 (4)	C27—O21	1.233 (3)
C7—H7	0.9500	C27—O22	1.274 (3)
C8—C9	1.384 (4)	N23—O24	1.215 (4)
C8—C18	1.746 (3)	N23—O23	1.222 (4)
C9—C9A	1.397 (4)	N25—O26	1.227 (4)
C9—H9	0.9500	N25—O25	1.228 (4)
C9A—N10	1.400 (3)	S31—O31	1.507 (3)
N10—C11	1.291 (3)	S31—C32	1.756 (5)
C11—N11	1.372 (3)	S31—C31	1.760 (5)
C11—C11A	1.496 (3)	C31—H31A	0.9800
N11—C16	1.461 (3)	C31—H31B	0.9800
N11—C12	1.461 (3)	C31—H31C	0.9800
C12—C13	1.510 (4)	C32—H32A	0.9800
C12—H12A	0.9900	C32—H32B	0.9800
C12—H12B	0.9900	C32—H32C	0.9800
C13—N14	1.491 (3)	S41—O41	1.497 (4)
C13—H13A	0.9900	S41—C42	1.749 (6)
C13—H13B	0.9900	S41—C41	1.756 (6)
N14—C17	1.483 (3)	C41—H41A	0.9800
N14—C15	1.490 (3)	C41—H41B	0.9800
N14—H14	1.00 (3)	C41—H41C	0.9800
C15—C16	1.512 (3)	C42—H42A	0.9800
C15—H15A	0.9900	C42—H42B	0.9800
C15—H15B	0.9900	C42—H42C	0.9800
C2—C1—C11A	120.9 (2)	N11—C16—C15	111.57 (19)
C2—C1—H1	119.5	N11—C16—H16A	109.3
C11A—C1—H1	119.5	C15—C16—H16A	109.3
C1—C2—C3	119.4 (2)	N11—C16—H16B	109.3

C1—C2—H2	120.3	C15—C16—H16B	109.3
C3—C2—H2	120.3	H16A—C16—H16B	108.0
C4—C3—C2	120.2 (2)	N14—C17—H17A	109.5
C4—C3—H3	119.9	N14—C17—H17B	109.5
C2—C3—H3	119.9	H17A—C17—H17B	109.5
C3—C4—C4A	120.5 (2)	N14—C17—H17C	109.5
C3—C4—H4	119.8	H17A—C17—H17C	109.5
C4A—C4—H4	119.8	H17B—C17—H17C	109.5
C4—C4A—C11A	120.0 (2)	C22—C21—C26	119.6 (2)
C4—C4A—N5	120.8 (2)	C22—C21—C27	120.5 (2)
C11A—C4A—N5	119.2 (2)	C26—C21—C27	119.9 (2)
C4A—N5—C5A	112.30 (19)	C23—C22—C21	118.8 (2)
C4A—N5—H5	109 (2)	C23—C22—H22	120.6
C5A—N5—H5	110 (2)	C21—C22—H22	120.6
C6—C5A—C9A	120.1 (2)	C24—C23—C22	123.5 (2)
C6—C5A—N5	119.9 (2)	C24—C23—N23	118.9 (3)
C9A—C5A—N5	120.0 (2)	C22—C23—N23	117.6 (3)
C7—C6—C5A	121.1 (3)	C25—C24—C23	116.1 (2)
C7—C6—H6	119.4	C25—C24—H24	122.0
C5A—C6—H6	119.4	C23—C24—H24	122.0
C8—C7—C6	118.6 (3)	C24—C25—C26	123.0 (3)
C8—C7—H7	120.7	C24—C25—N25	118.4 (3)
C6—C7—H7	120.7	C26—C25—N25	118.5 (3)
C7—C8—C9	121.7 (3)	C25—C26—C21	119.1 (2)
C7—C8—C18	118.3 (2)	C25—C26—H26	120.4
C9—C8—C18	120.0 (2)	C21—C26—H26	120.4
C8—C9—C9A	120.3 (3)	O21—C27—O22	126.7 (2)
C8—C9—H9	119.9	O21—C27—C21	118.2 (2)
C9A—C9—H9	119.9	O22—C27—C21	115.0 (2)
C9—C9A—N10	117.8 (2)	O24—N23—O23	124.6 (3)
C9—C9A—C5A	118.1 (2)	O24—N23—C23	117.1 (3)
N10—C9A—C5A	123.6 (2)	O23—N23—C23	118.3 (3)
C11—N10—C9A	122.8 (2)	O26—N25—O25	124.3 (3)
N10—C11—N11	118.1 (2)	O26—N25—C25	118.2 (3)
N10—C11—C11A	125.2 (2)	O25—N25—C25	117.6 (3)
N11—C11—C11A	116.6 (2)	O31—S31—C32	104.5 (2)
C1—C11A—C4A	118.7 (2)	O31—S31—C31	107.6 (3)
C1—C11A—C11	121.9 (2)	C32—S31—C31	99.1 (4)
C4A—C11A—C11	119.5 (2)	S31—C31—H31A	109.5
C11—N11—C16	126.01 (19)	S31—C31—H31B	109.5
C11—N11—C12	120.80 (19)	H31A—C31—H31B	109.5
C16—N11—C12	112.91 (18)	S31—C31—H31C	109.5
N11—C12—C13	111.7 (2)	H31A—C31—H31C	109.5
N11—C12—H12A	109.3	H31B—C31—H31C	109.5
C13—C12—H12A	109.3	S31—C32—H32A	109.5
N11—C12—H12B	109.3	S31—C32—H32B	109.5
C13—C12—H12B	109.3	H32A—C32—H32B	109.5
H12A—C12—H12B	107.9	S31—C32—H32C	109.5

N14—C13—C12	109.72 (19)	H32A—C32—H32C	109.5
N14—C13—H13A	109.7	H32B—C32—H32C	109.5
C12—C13—H13A	109.7	O41—S41—C42	106.5 (4)
N14—C13—H13B	109.7	O41—S41—C41	108.6 (5)
C12—C13—H13B	109.7	C42—S41—C41	99.8 (5)
H13A—C13—H13B	108.2	S41—C41—H41A	109.5
C17—N14—C15	112.1 (2)	S41—C41—H41B	109.5
C17—N14—C13	112.7 (2)	H41A—C41—H41B	109.5
C15—N14—C13	109.17 (18)	S41—C41—H41C	109.5
C17—N14—H14	104.9 (16)	H41A—C41—H41C	109.5
C15—N14—H14	110.5 (16)	H41B—C41—H41C	109.5
C13—N14—H14	107.3 (16)	S41—C42—H42A	109.5
N14—C15—C16	110.49 (19)	S41—C42—H42B	109.5
N14—C15—H15A	109.6	H42A—C42—H42B	109.5
C16—C15—H15A	109.6	S41—C42—H42C	109.5
N14—C15—H15B	109.6	H42A—C42—H42C	109.5
C16—C15—H15B	109.6	H42B—C42—H42C	109.5
H15A—C15—H15B	108.1		
C11A—C1—C2—C3	1.1 (4)	C11A—C11—N11—C16	16.8 (3)
C1—C2—C3—C4	-5.1 (4)	N10—C11—N11—C12	6.3 (3)
C2—C3—C4—C4A	3.6 (4)	C11A—C11—N11—C12	-169.7 (2)
C3—C4—C4A—C11A	1.8 (4)	C11—N11—C12—C13	-120.8 (2)
C3—C4—C4A—N5	-178.5 (2)	C16—N11—C12—C13	53.4 (3)
C4—C4A—N5—C5A	115.1 (3)	N11—C12—C13—N14	-56.7 (3)
C11A—C4A—N5—C5A	-65.2 (3)	C12—C13—N14—C17	-175.3 (2)
C4A—N5—C5A—C6	-117.8 (2)	C12—C13—N14—C15	59.5 (2)
C4A—N5—C5A—C9A	63.5 (3)	C17—N14—C15—C16	175.4 (2)
C9A—C5A—C6—C7	-0.1 (4)	C13—N14—C15—C16	-59.1 (2)
N5—C5A—C6—C7	-178.8 (2)	C11—N11—C16—C15	121.5 (2)
C5A—C6—C7—C8	-1.9 (4)	C12—N11—C16—C15	-52.4 (3)
C6—C7—C8—C9	2.3 (4)	N14—C15—C16—N11	55.3 (3)
C6—C7—C8—C18	-177.8 (2)	C26—C21—C22—C23	-0.1 (3)
C7—C8—C9—C9A	-0.8 (4)	C27—C21—C22—C23	-177.5 (2)
C18—C8—C9—C9A	179.38 (18)	C21—C22—C23—C24	-0.4 (4)
C8—C9—C9A—N10	171.3 (2)	C21—C22—C23—N23	176.6 (2)
C8—C9—C9A—C5A	-1.3 (3)	C22—C23—C24—C25	0.3 (4)
C6—C5A—C9A—C9	1.7 (3)	N23—C23—C24—C25	-176.7 (2)
N5—C5A—C9A—C9	-179.6 (2)	C23—C24—C25—C26	0.3 (4)
C6—C5A—C9A—N10	-170.4 (2)	C23—C24—C25—N25	177.4 (2)
N5—C5A—C9A—N10	8.2 (3)	C24—C25—C26—C21	-0.7 (3)
C9—C9A—N10—C11	146.3 (2)	N25—C25—C26—C21	-177.8 (2)
C5A—C9A—N10—C11	-41.5 (3)	C22—C21—C26—C25	0.6 (3)
C9A—N10—C11—N11	176.4 (2)	C27—C21—C26—C25	178.1 (2)
C9A—N10—C11—C11A	-7.9 (4)	C22—C21—C27—O21	177.5 (2)
C2—C1—C11A—C4A	4.3 (3)	C26—C21—C27—O21	0.0 (3)
C2—C1—C11A—C11	-177.0 (2)	C22—C21—C27—O22	-2.0 (3)
C4—C4A—C11A—C1	-5.7 (3)	C26—C21—C27—O22	-179.5 (2)

N5—C4A—C11A—C1	174.6 (2)	C24—C23—N23—O24	-0.8 (4)
C4—C4A—C11A—C11	175.5 (2)	C22—C23—N23—O24	-177.9 (3)
N5—C4A—C11A—C11	-4.1 (3)	C24—C23—N23—O23	177.6 (3)
N10—C11—C11A—C1	-129.2 (3)	C22—C23—N23—O23	0.5 (4)
N11—C11—C11A—C1	46.5 (3)	C24—C25—N25—O26	-178.3 (2)
N10—C11—C11A—C4A	49.5 (3)	C26—C25—N25—O26	-1.1 (4)
N11—C11—C11A—C4A	-134.7 (2)	C24—C25—N25—O25	1.6 (4)
N10—C11—N11—C16	-167.1 (2)	C26—C25—N25—O25	178.8 (3)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N5—H5...O31	0.83 (3)	2.10 (3)	2.921 (4)	170 (3)
N14—H14...O22	1.00 (3)	1.58 (3)	2.575 (3)	176 (2)
C4—H4...O31	0.95	2.58	3.342 (4)	137
C4—H4...O41	0.95	2.38	3.208 (7)	146
C6—H6...O31	0.95	2.54	3.313 (5)	139

(II) Clozapinium hydrogen maleate 0.21-hydrate*Crystal data*C₁₈H₂₀CIN₄⁺·C₄H₃O₄⁻·0.21H₂O*M_r* = 446.68Monoclinic, *P*2₁/*c**a* = 9.7166 (3) Å*b* = 9.9699 (2) Å*c* = 23.1059 (6) Å β = 96.800 (3)°*V* = 2222.60 (10) Å³*Z* = 4*F*(000) = 936.4*D_x* = 1.335 Mg m⁻³Cu *K*α radiation, λ = 1.54184 Å

Cell parameters from 4228 reflections

 θ = 3.9–71.2° μ = 1.84 mm⁻¹*T* = 173 K

Block, colourless

0.46 × 0.32 × 0.22 mm

*Data collection*Agilent Eos Gemini
diffractometer

Radiation source: Enhance (Cu) X-ray Source

 ω scansAbsorption correction: multi-scan
(*CrysAlis RED*; Agilent, 2012)*T_{min}* = 0.440, *T_{max}* = 0.668

8634 measured reflections

4228 independent reflections

3552 reflections with *I* > 2σ(*I*)*R_{int}* = 0.026 θ_{\max} = 71.2°, θ_{\min} = 3.9°*h* = -11→11*k* = -12→7*l* = -28→27*Refinement*Refinement on *F*²

Least-squares matrix: full

R[*F*² > 2σ(*F*²)] = 0.048*wR*(*F*²) = 0.133*S* = 1.03

4228 reflections

295 parameters

0 restraints

Hydrogen site location: difference Fourier map

H atoms treated by a mixture of independent
and constrained refinement $w = 1/[\sigma^2(F_o^2) + (0.069P)^2 + 0.8133P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.46 \text{ e } \text{Å}^{-3}$ $\Delta\rho_{\min} = -0.30 \text{ e } \text{Å}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C1	0.6656 (2)	0.4997 (2)	0.23723 (9)	0.0361 (4)	
H1	0.7048	0.4466	0.2691	0.043*	
C2	0.7438 (2)	0.5303 (2)	0.19279 (9)	0.0417 (5)	
H2	0.8362	0.4981	0.1940	0.050*	
C3	0.6866 (2)	0.6083 (2)	0.14640 (9)	0.0409 (5)	
H3	0.7417	0.6340	0.1169	0.049*	
C4	0.5503 (2)	0.6488 (2)	0.14272 (8)	0.0356 (4)	
H4	0.5110	0.6994	0.1100	0.043*	
C4A	0.46942 (19)	0.61605 (19)	0.18684 (8)	0.0291 (4)	
N5	0.32745 (17)	0.65240 (17)	0.18232 (7)	0.0326 (3)	
H5	0.307 (2)	0.703 (3)	0.1527 (11)	0.039*	
C5A	0.23552 (18)	0.54101 (19)	0.18333 (8)	0.0296 (4)	
C6	0.1401 (2)	0.5111 (2)	0.13518 (8)	0.0357 (4)	
H6	0.1364	0.5660	0.1014	0.043*	
C7	0.0501 (2)	0.4028 (2)	0.13556 (9)	0.0385 (5)	
H7	-0.0150	0.3835	0.1026	0.046*	
C8	0.05780 (19)	0.3239 (2)	0.18497 (9)	0.0360 (4)	
C18	-0.05340 (6)	0.18656 (6)	0.18694 (3)	0.0560 (2)	
C9	0.14961 (19)	0.3526 (2)	0.23364 (9)	0.0334 (4)	
H9	0.1507	0.2984	0.2675	0.040*	
C9A	0.24090 (18)	0.46068 (19)	0.23353 (8)	0.0290 (4)	
N10	0.32420 (16)	0.48876 (16)	0.28580 (6)	0.0301 (3)	
C11	0.45129 (18)	0.52545 (18)	0.28708 (8)	0.0283 (4)	
C11A	0.52924 (19)	0.54587 (19)	0.23581 (8)	0.0293 (4)	
N11	0.52983 (16)	0.53872 (17)	0.34067 (6)	0.0315 (4)	
C12	0.46621 (19)	0.50463 (19)	0.39301 (7)	0.0300 (4)	
H12A	0.5381	0.4687	0.4229	0.036*	
H12B	0.3960	0.4336	0.3834	0.036*	
C13	0.39818 (18)	0.62443 (19)	0.41749 (8)	0.0302 (4)	
H13A	0.3214	0.6569	0.3889	0.036*	
H13B	0.3592	0.5986	0.4536	0.036*	
N14	0.50291 (17)	0.73320 (17)	0.43053 (7)	0.0324 (4)	
H14	0.565 (3)	0.702 (2)	0.4574 (11)	0.039*	
C15	0.5702 (2)	0.7683 (2)	0.37746 (8)	0.0382 (5)	
H15A	0.6437	0.8358	0.3878	0.046*	
H15B	0.5004	0.8082	0.3477	0.046*	
C16	0.63243 (19)	0.6451 (2)	0.35254 (8)	0.0361 (4)	
H16A	0.6701	0.6694	0.3160	0.043*	
H16B	0.7101	0.6120	0.3805	0.043*	

C17	0.4417 (3)	0.8518 (2)	0.45611 (11)	0.0561 (6)	
H17A	0.3690	0.8896	0.4278	0.084*	
H17B	0.5140	0.9193	0.4661	0.084*	
H17C	0.4017	0.8251	0.4914	0.084*	
C21	1.1468 (2)	0.6834 (2)	0.54450 (9)	0.0415 (5)	
O21	1.1086 (2)	0.7228 (3)	0.49282 (8)	0.0910 (9)	
H21	1.006 (5)	0.739 (5)	0.488 (2)	0.137*	
O22	1.26642 (15)	0.69322 (19)	0.56647 (7)	0.0533 (4)	
C22	1.0433 (2)	0.6232 (3)	0.57919 (10)	0.0490 (6)	
H22	1.0817	0.5844	0.6151	0.059*	
C23	0.9056 (2)	0.6141 (3)	0.56863 (10)	0.0494 (6)	
H23	0.8618	0.5719	0.5984	0.059*	
C24	0.8103 (2)	0.6601 (2)	0.51734 (8)	0.0356 (4)	
O23	0.68470 (15)	0.6486 (2)	0.51985 (7)	0.0526 (4)	
O24	0.85884 (18)	0.7068 (3)	0.47375 (8)	0.0810 (8)	
O31	0.7719 (11)	0.9745 (11)	0.5046 (5)	0.074 (4)*	0.210 (7)
H31A	0.8183	0.9016	0.4951	0.111*	0.210 (7)
H31B	0.8247	1.0490	0.5081	0.111*	0.210 (7)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0326 (9)	0.0429 (11)	0.0324 (9)	0.0035 (8)	0.0016 (7)	-0.0027 (8)
C2	0.0333 (10)	0.0513 (13)	0.0421 (11)	0.0024 (9)	0.0106 (8)	-0.0098 (10)
C3	0.0439 (11)	0.0483 (12)	0.0328 (10)	-0.0098 (9)	0.0148 (8)	-0.0075 (9)
C4	0.0435 (11)	0.0355 (10)	0.0273 (9)	-0.0063 (8)	0.0031 (8)	-0.0016 (8)
C4A	0.0319 (9)	0.0278 (9)	0.0271 (8)	-0.0028 (7)	0.0018 (7)	-0.0056 (7)
N5	0.0350 (8)	0.0308 (8)	0.0308 (8)	0.0020 (7)	-0.0011 (6)	0.0024 (7)
C5A	0.0259 (8)	0.0304 (9)	0.0321 (9)	0.0043 (7)	0.0013 (7)	-0.0053 (7)
C6	0.0331 (9)	0.0416 (11)	0.0310 (9)	0.0058 (8)	-0.0016 (7)	-0.0024 (8)
C7	0.0277 (9)	0.0469 (12)	0.0388 (10)	0.0021 (8)	-0.0047 (8)	-0.0118 (9)
C8	0.0250 (9)	0.0357 (10)	0.0467 (11)	-0.0019 (8)	0.0017 (8)	-0.0088 (9)
C18	0.0448 (3)	0.0505 (3)	0.0690 (4)	-0.0191 (2)	-0.0089 (3)	-0.0033 (3)
C9	0.0279 (9)	0.0355 (10)	0.0365 (9)	0.0002 (8)	0.0022 (7)	-0.0017 (8)
C9A	0.0245 (8)	0.0322 (9)	0.0301 (9)	0.0035 (7)	0.0020 (7)	-0.0059 (7)
N10	0.0299 (8)	0.0341 (8)	0.0261 (7)	-0.0022 (6)	0.0020 (6)	-0.0023 (6)
C11	0.0294 (9)	0.0293 (9)	0.0258 (8)	0.0004 (7)	0.0016 (7)	-0.0011 (7)
C11A	0.0297 (9)	0.0325 (9)	0.0254 (8)	-0.0012 (7)	0.0028 (7)	-0.0038 (7)
N11	0.0308 (8)	0.0407 (9)	0.0227 (7)	-0.0074 (7)	0.0025 (6)	0.0006 (6)
C12	0.0341 (9)	0.0316 (9)	0.0237 (8)	-0.0051 (7)	0.0012 (7)	0.0029 (7)
C13	0.0274 (8)	0.0358 (10)	0.0268 (8)	-0.0034 (7)	0.0001 (7)	0.0030 (7)
N14	0.0388 (9)	0.0302 (8)	0.0265 (7)	-0.0031 (7)	-0.0026 (6)	0.0028 (6)
C15	0.0458 (11)	0.0390 (11)	0.0281 (9)	-0.0157 (9)	-0.0022 (8)	0.0080 (8)
C16	0.0297 (9)	0.0537 (12)	0.0241 (8)	-0.0129 (9)	-0.0004 (7)	0.0012 (8)
C17	0.0829 (18)	0.0339 (11)	0.0526 (14)	0.0015 (12)	0.0122 (13)	-0.0047 (10)
C21	0.0347 (10)	0.0498 (12)	0.0383 (10)	-0.0006 (9)	-0.0032 (8)	-0.0123 (9)
O21	0.0376 (9)	0.183 (3)	0.0495 (10)	-0.0273 (13)	-0.0053 (8)	0.0325 (14)
O22	0.0315 (8)	0.0719 (12)	0.0541 (9)	0.0031 (7)	-0.0052 (7)	-0.0144 (8)

C22	0.0418 (11)	0.0594 (15)	0.0421 (11)	0.0017 (10)	-0.0102 (9)	0.0144 (10)
C23	0.0419 (11)	0.0644 (15)	0.0401 (11)	-0.0090 (11)	-0.0029 (9)	0.0215 (11)
C24	0.0354 (10)	0.0418 (11)	0.0278 (9)	-0.0107 (9)	-0.0041 (7)	0.0051 (8)
O23	0.0344 (8)	0.0847 (13)	0.0367 (8)	-0.0113 (8)	-0.0040 (6)	0.0215 (8)
O24	0.0400 (9)	0.158 (2)	0.0418 (9)	-0.0252 (11)	-0.0089 (7)	0.0419 (12)

Geometric parameters (Å, °)

C1—C2	1.382 (3)	C12—C13	1.507 (3)
C1—C11A	1.399 (3)	C12—H12A	0.9900
C1—H1	0.9500	C12—H12B	0.9900
C2—C3	1.387 (3)	C13—N14	1.493 (2)
C2—H2	0.9500	C13—H13A	0.9900
C3—C4	1.377 (3)	C13—H13B	0.9900
C3—H3	0.9500	N14—C17	1.478 (3)
C4—C4A	1.397 (3)	N14—C15	1.498 (2)
C4—H4	0.9500	N14—H14	0.87 (3)
C4A—C11A	1.397 (3)	C15—C16	1.513 (3)
C4A—N5	1.418 (2)	C15—H15A	0.9900
N5—C5A	1.427 (3)	C15—H15B	0.9900
N5—H5	0.86 (3)	C16—H16A	0.9900
C5A—C6	1.394 (3)	C16—H16B	0.9900
C5A—C9A	1.405 (3)	C17—H17A	0.9800
C6—C7	1.390 (3)	C17—H17B	0.9800
C6—H6	0.9500	C17—H17C	0.9800
C7—C8	1.381 (3)	C21—O22	1.216 (3)
C7—H7	0.9500	C21—O21	1.270 (3)
C8—C9	1.380 (3)	C21—C22	1.485 (3)
C8—C18	1.748 (2)	O21—H21	1.00 (5)
C9—C9A	1.396 (3)	C22—C23	1.334 (3)
C9—H9	0.9500	C22—H22	0.9500
C9A—N10	1.400 (2)	C23—C24	1.487 (3)
N10—C11	1.285 (2)	C23—H23	0.9500
C11—N11	1.382 (2)	C24—O23	1.234 (2)
C11—C11A	1.494 (2)	C24—O24	1.251 (3)
N11—C16	1.459 (2)	O31—H31A	0.8959
N11—C12	1.462 (2)	O31—H31B	0.9007
C2—C1—C11A	120.84 (19)	N11—C12—H12B	109.2
C2—C1—H1	119.6	C13—C12—H12B	109.2
C11A—C1—H1	119.6	H12A—C12—H12B	107.9
C1—C2—C3	119.49 (19)	N14—C13—C12	109.42 (15)
C1—C2—H2	120.3	N14—C13—H13A	109.8
C3—C2—H2	120.3	C12—C13—H13A	109.8
C4—C3—C2	120.47 (18)	N14—C13—H13B	109.8
C4—C3—H3	119.8	C12—C13—H13B	109.8
C2—C3—H3	119.8	H13A—C13—H13B	108.2
C3—C4—C4A	120.39 (19)	C17—N14—C13	111.41 (17)

C3—C4—H4	119.8	C17—N14—C15	112.05 (17)
C4A—C4—H4	119.8	C13—N14—C15	110.95 (14)
C11A—C4A—C4	119.47 (17)	C17—N14—H14	106.2 (16)
C11A—C4A—N5	119.70 (16)	C13—N14—H14	106.3 (16)
C4—C4A—N5	120.83 (17)	C15—N14—H14	109.6 (16)
C4A—N5—C5A	113.90 (15)	N14—C15—C16	110.77 (16)
C4A—N5—H5	110.1 (16)	N14—C15—H15A	109.5
C5A—N5—H5	112.8 (17)	C16—C15—H15A	109.5
C6—C5A—C9A	119.55 (18)	N14—C15—H15B	109.5
C6—C5A—N5	120.79 (18)	C16—C15—H15B	109.5
C9A—C5A—N5	119.66 (16)	H15A—C15—H15B	108.1
C7—C6—C5A	121.36 (19)	N11—C16—C15	111.51 (16)
C7—C6—H6	119.3	N11—C16—H16A	109.3
C5A—C6—H6	119.3	C15—C16—H16A	109.3
C8—C7—C6	118.39 (18)	N11—C16—H16B	109.3
C8—C7—H7	120.8	C15—C16—H16B	109.3
C6—C7—H7	120.8	H16A—C16—H16B	108.0
C9—C8—C7	121.44 (19)	N14—C17—H17A	109.5
C9—C8—C18	118.90 (17)	N14—C17—H17B	109.5
C7—C8—C18	119.65 (15)	H17A—C17—H17B	109.5
C8—C9—C9A	120.55 (19)	N14—C17—H17C	109.5
C8—C9—H9	119.7	H17A—C17—H17C	109.5
C9A—C9—H9	119.7	H17B—C17—H17C	109.5
C9—C9A—N10	117.09 (17)	O22—C21—O21	121.8 (2)
C9—C9A—C5A	118.67 (17)	O22—C21—C22	118.8 (2)
N10—C9A—C5A	124.00 (17)	O21—C21—C22	119.42 (19)
C11—N10—C9A	122.17 (15)	C21—O21—H21	109 (3)
N10—C11—N11	118.36 (16)	C23—C22—C21	131.1 (2)
N10—C11—C11A	126.65 (16)	C23—C22—H22	114.5
N11—C11—C11A	114.80 (15)	C21—C22—H22	114.5
C4A—C11A—C1	119.06 (17)	C22—C23—C24	129.9 (2)
C4A—C11A—C11	120.61 (16)	C22—C23—H23	115.1
C1—C11A—C11	120.28 (17)	C24—C23—H23	115.1
C11—N11—C16	122.04 (15)	O23—C24—O24	122.78 (19)
C11—N11—C12	118.45 (15)	O23—C24—C23	117.38 (18)
C16—N11—C12	111.09 (14)	O24—C24—C23	119.84 (19)
N11—C12—C13	111.93 (15)	C24—O24—H21	110.7 (19)
N11—C12—H12A	109.2	H31A—O31—H31B	113.2
C13—C12—H12A	109.2		
C11A—C1—C2—C3	0.3 (3)	C4—C4A—C11A—C11	171.79 (17)
C1—C2—C3—C4	-3.7 (3)	N5—C4A—C11A—C11	-8.6 (3)
C2—C3—C4—C4A	2.4 (3)	C2—C1—C11A—C4A	4.3 (3)
C3—C4—C4A—C11A	2.3 (3)	C2—C1—C11A—C11	-173.05 (19)
C3—C4—C4A—N5	-177.34 (18)	N10—C11—C11A—C4A	45.2 (3)
C11A—C4A—N5—C5A	-59.5 (2)	N11—C11—C11A—C4A	-139.81 (18)
C4—C4A—N5—C5A	120.07 (19)	N10—C11—C11A—C1	-137.5 (2)
C4A—N5—C5A—C6	-116.17 (19)	N11—C11—C11A—C1	37.5 (3)

C4A—N5—C5A—C9A	64.0 (2)	N10—C11—N11—C16	-142.60 (19)
C9A—C5A—C6—C7	-0.3 (3)	C11A—C11—N11—C16	42.0 (2)
N5—C5A—C6—C7	179.87 (17)	N10—C11—N11—C12	2.7 (3)
C5A—C6—C7—C8	-0.4 (3)	C11A—C11—N11—C12	-172.74 (16)
C6—C7—C8—C9	1.6 (3)	C11—N11—C12—C13	-91.0 (2)
C6—C7—C8—C18	-179.85 (15)	C16—N11—C12—C13	57.8 (2)
C7—C8—C9—C9A	-2.0 (3)	N11—C12—C13—N14	-57.45 (19)
C18—C8—C9—C9A	179.39 (14)	C12—C13—N14—C17	-178.53 (16)
C8—C9—C9A—N10	175.80 (17)	C12—C13—N14—C15	55.9 (2)
C8—C9—C9A—C5A	1.2 (3)	C17—N14—C15—C16	179.67 (18)
C6—C5A—C9A—C9	-0.1 (3)	C13—N14—C15—C16	-55.1 (2)
N5—C5A—C9A—C9	179.71 (16)	C11—N11—C16—C15	91.5 (2)
C6—C5A—C9A—N10	-174.23 (16)	C12—N11—C16—C15	-56.0 (2)
N5—C5A—C9A—N10	5.6 (3)	N14—C15—C16—N11	54.9 (2)
C9—C9A—N10—C11	141.11 (19)	O22—C21—C22—C23	171.5 (3)
C5A—C9A—N10—C11	-44.6 (3)	O21—C21—C22—C23	-8.7 (5)
C9A—N10—C11—N11	-173.90 (17)	C21—C22—C23—C24	1.3 (5)
C9A—N10—C11—C11A	0.9 (3)	C22—C23—C24—O23	-174.8 (3)
C4—C4A—C11A—C1	-5.6 (3)	C22—C23—C24—O24	6.0 (5)
N5—C4A—C11A—C1	174.05 (17)		

Hydrogen-bond geometry (Å, °)

Cg1 and Cg2 represent the centroids of the rings (C5A,C6–C9,C9A) and (C1–C4,C4A,C11A), respectively.

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N5—H5...O22 ⁱ	0.86 (3)	2.24 (3)	3.084 (2)	170 (3)
N14—H14...O23	0.87 (3)	1.82 (3)	2.688 (2)	173 (2)
O21—H21...O24	1.00 (5)	1.46 (5)	2.420 (3)	157 (5)
O31—H31A...O24	0.90	2.05	2.913 (11)	160
O31—H31B...O21 ⁱⁱ	0.90	2.37	3.232 (11)	161
C12—H12A...O22 ⁱⁱⁱ	0.99	2.48	3.308 (2)	141
C15—H15A...Cg1 ^{iv}	0.99	2.95	3.642 (2)	128
C15—H15B...Cg2 ^{iv}	0.99	2.95	3.759 (2)	139

Symmetry codes: (i) $x-1, -y+3/2, z-1/2$; (ii) $-x+2, -y+2, -z+1$; (iii) $-x+2, -y+1, -z+1$; (iv) $-x+1, y+1/2, -z+1/2$.

(III) Clozapinium 2-hydroxybenzoate

Crystal data

$C_{18}H_{20}ClN_4^+ \cdot C_7H_5O_3^-$

$M_r = 464.94$

Monoclinic, *Cc*

$a = 17.4296 (5) \text{ \AA}$

$b = 15.3728 (5) \text{ \AA}$

$c = 8.6359 (3) \text{ \AA}$

$\beta = 90.325 (3)^\circ$

$V = 2313.88 (13) \text{ \AA}^3$

$Z = 4$

$F(000) = 976$

$D_x = 1.335 \text{ Mg m}^{-3}$

Cu $K\alpha$ radiation, $\lambda = 1.54184 \text{ \AA}$

Cell parameters from 4069 reflections

$\theta = 3.8\text{--}72.6^\circ$

$\mu = 1.75 \text{ mm}^{-1}$

$T = 173 \text{ K}$

Block, colourless

$0.42 \times 0.36 \times 0.20 \text{ mm}$

Data collection

Agilent Eos Gemini
diffractometer
Radiation source: Enhance (Cu) X-ray Source
 ω scans
Absorption correction: multi-scan
(*CrysAlis RED*; Agilent, 2012)
 $T_{\min} = 0.399$, $T_{\max} = 0.705$
7244 measured reflections

4069 independent reflections
3962 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.048$
 $\theta_{\max} = 72.6^\circ$, $\theta_{\min} = 3.8^\circ$
 $h = -21 \rightarrow 21$
 $k = -10 \rightarrow 18$
 $l = -10 \rightarrow 10$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.056$
 $wR(F^2) = 0.141$
 $S = 1.08$
4069 reflections
308 parameters
2 restraints
Hydrogen site location: difference Fourier map

H atoms treated by a mixture of independent
and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.1069P)^2]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.53 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.36 \text{ e } \text{\AA}^{-3}$
Absolute structure: Flack x determined using
1674 quotients $[(I^+)-(I^-)]/[(I^+)+(I^-)]$ (Parsons *et al.*, 2013)
Absolute structure parameter: -0.022 (17)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.68573 (19)	0.4925 (2)	0.8347 (4)	0.0296 (6)
H1	0.6566	0.5324	0.8945	0.036*
C2	0.7650 (2)	0.4966 (2)	0.8402 (4)	0.0346 (7)
H2	0.7899	0.5394	0.9018	0.042*
C3	0.80783 (19)	0.4372 (2)	0.7545 (4)	0.0338 (7)
H3	0.8623	0.4405	0.7552	0.041*
C4	0.77094 (19)	0.3730 (2)	0.6681 (4)	0.0322 (7)
H4	0.8003	0.3310	0.6136	0.039*
C4A	0.69150 (18)	0.3700 (2)	0.6611 (3)	0.0273 (6)
N5	0.65422 (16)	0.30589 (18)	0.5690 (3)	0.0306 (6)
H5	0.688 (3)	0.276 (3)	0.511 (6)	0.037*
C5A	0.60999 (17)	0.2452 (2)	0.6565 (3)	0.0290 (6)
C6	0.6279 (2)	0.1568 (2)	0.6588 (4)	0.0368 (7)
H6	0.6702	0.1363	0.6004	0.044*
C7	0.5850 (2)	0.0983 (2)	0.7448 (5)	0.0403 (8)
H7	0.5969	0.0380	0.7442	0.048*
C8	0.5249 (2)	0.1300 (2)	0.8312 (5)	0.0369 (7)
Cl8	0.47247 (6)	0.05843 (6)	0.94789 (14)	0.0561 (3)
C9	0.50474 (19)	0.2173 (2)	0.8300 (4)	0.0332 (7)

H9	0.4622	0.2369	0.8886	0.040*
C9A	0.54720 (18)	0.2761 (2)	0.7424 (4)	0.0287 (6)
N10	0.51934 (16)	0.36170 (18)	0.7350 (3)	0.0294 (5)
C11	0.56224 (18)	0.4295 (2)	0.7362 (3)	0.0267 (6)
C11A	0.64754 (19)	0.4310 (2)	0.7428 (3)	0.0272 (6)
N11	0.52619 (16)	0.50963 (18)	0.7496 (3)	0.0303 (6)
C12	0.44338 (19)	0.5111 (2)	0.7757 (4)	0.0334 (7)
H12A	0.4281	0.4591	0.8356	0.040*
H12B	0.4159	0.5098	0.6751	0.040*
C13	0.4218 (2)	0.5929 (2)	0.8641 (4)	0.0357 (7)
H13A	0.3654	0.5949	0.8774	0.043*
H13B	0.4458	0.5914	0.9682	0.043*
N14	0.44753 (17)	0.6726 (2)	0.7804 (3)	0.0331 (6)
H14	0.423 (3)	0.675 (3)	0.690 (7)	0.040*
C15	0.5313 (2)	0.6687 (2)	0.7446 (4)	0.0338 (7)
H15A	0.5614	0.6712	0.8422	0.041*
H15B	0.5457	0.7195	0.6805	0.041*
C16	0.55040 (19)	0.5851 (2)	0.6580 (4)	0.0306 (6)
H16A	0.5237	0.5847	0.5565	0.037*
H16B	0.6063	0.5821	0.6393	0.037*
C17	0.4297 (3)	0.7522 (3)	0.8713 (5)	0.0482 (9)
H17A	0.4588	0.7513	0.9687	0.072*
H17B	0.3747	0.7539	0.8937	0.072*
H17C	0.4439	0.8038	0.8114	0.072*
C21	0.28195 (18)	0.6783 (2)	0.3237 (4)	0.0306 (6)
C22	0.2327 (2)	0.6192 (3)	0.2489 (5)	0.0462 (9)
C23	0.1947 (3)	0.6426 (4)	0.1142 (7)	0.0657 (14)
H23A	0.1623	0.6020	0.0627	0.079*
C24	0.2041 (3)	0.7258 (5)	0.0548 (6)	0.0673 (15)
H24	0.1774	0.7422	-0.0369	0.081*
C25	0.2518 (3)	0.7851 (3)	0.1272 (5)	0.0509 (10)
H25	0.2576	0.8419	0.0856	0.061*
C26	0.2911 (2)	0.7617 (2)	0.2602 (4)	0.0354 (7)
H26	0.3245	0.8022	0.3089	0.042*
C27	0.32403 (17)	0.6517 (2)	0.4680 (4)	0.0286 (6)
O21	0.31483 (16)	0.57583 (18)	0.5174 (4)	0.0445 (6)
O22	0.36650 (14)	0.70751 (16)	0.5327 (3)	0.0332 (5)
O23	0.2212 (2)	0.5383 (2)	0.3064 (5)	0.0661 (10)
H23	0.259 (6)	0.544 (6)	0.394 (13)	0.099*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0314 (15)	0.0289 (13)	0.0285 (14)	0.0049 (11)	0.0008 (12)	-0.0026 (11)
C2	0.0345 (17)	0.0331 (15)	0.0363 (16)	-0.0012 (12)	-0.0036 (13)	-0.0015 (12)
C3	0.0239 (13)	0.0341 (15)	0.0433 (18)	0.0016 (11)	-0.0013 (12)	0.0033 (12)
C4	0.0311 (16)	0.0340 (15)	0.0317 (16)	0.0088 (12)	0.0028 (12)	0.0018 (12)
C4A	0.0305 (16)	0.0285 (14)	0.0229 (13)	0.0044 (11)	0.0013 (11)	0.0035 (10)

N5	0.0311 (13)	0.0348 (13)	0.0259 (12)	0.0058 (11)	0.0038 (10)	-0.0053 (10)
C5A	0.0255 (15)	0.0335 (15)	0.0279 (15)	0.0030 (11)	-0.0050 (12)	-0.0046 (11)
C6	0.0329 (15)	0.0346 (17)	0.0429 (19)	0.0062 (12)	-0.0030 (13)	-0.0084 (13)
C7	0.0354 (17)	0.0276 (15)	0.058 (2)	0.0028 (12)	-0.0092 (15)	-0.0047 (13)
C8	0.0300 (15)	0.0340 (16)	0.0466 (19)	-0.0048 (12)	-0.0060 (14)	0.0011 (13)
C18	0.0461 (5)	0.0410 (5)	0.0813 (8)	-0.0089 (4)	0.0056 (5)	0.0139 (4)
C9	0.0243 (14)	0.0387 (16)	0.0364 (17)	-0.0001 (11)	-0.0050 (12)	-0.0021 (12)
C9A	0.0241 (13)	0.0336 (15)	0.0281 (14)	0.0035 (11)	-0.0074 (11)	-0.0035 (11)
N10	0.0253 (11)	0.0345 (13)	0.0284 (13)	0.0068 (10)	-0.0013 (10)	0.0003 (10)
C11	0.0272 (15)	0.0338 (15)	0.0190 (13)	0.0063 (11)	0.0000 (10)	0.0004 (10)
C11A	0.0277 (15)	0.0296 (13)	0.0242 (14)	0.0034 (11)	0.0004 (11)	0.0022 (10)
N11	0.0272 (13)	0.0328 (13)	0.0310 (13)	0.0088 (10)	0.0028 (10)	0.0038 (10)
C12	0.0269 (15)	0.0355 (16)	0.0378 (17)	0.0087 (12)	0.0056 (12)	0.0048 (12)
C13	0.0338 (16)	0.0443 (18)	0.0290 (15)	0.0149 (14)	0.0048 (12)	0.0022 (13)
N14	0.0352 (14)	0.0367 (14)	0.0271 (14)	0.0137 (11)	-0.0073 (11)	-0.0033 (10)
C15	0.0329 (15)	0.0329 (15)	0.0357 (16)	0.0062 (12)	-0.0082 (13)	0.0000 (12)
C16	0.0315 (15)	0.0311 (15)	0.0292 (14)	0.0069 (11)	0.0017 (12)	0.0038 (12)
C17	0.053 (2)	0.049 (2)	0.0425 (19)	0.0227 (18)	-0.0116 (17)	-0.0150 (16)
C21	0.0250 (13)	0.0386 (16)	0.0282 (15)	0.0046 (12)	-0.0008 (11)	-0.0034 (12)
C22	0.0366 (18)	0.053 (2)	0.049 (2)	-0.0049 (16)	-0.0060 (16)	-0.0092 (17)
C23	0.053 (3)	0.093 (4)	0.051 (3)	-0.011 (3)	-0.021 (2)	-0.014 (3)
C24	0.052 (3)	0.114 (5)	0.036 (2)	0.012 (3)	-0.0204 (19)	0.002 (2)
C25	0.047 (2)	0.068 (3)	0.0374 (19)	0.0157 (19)	-0.0009 (17)	0.0149 (18)
C26	0.0311 (16)	0.0442 (18)	0.0307 (16)	0.0052 (13)	-0.0020 (13)	0.0025 (13)
C27	0.0212 (13)	0.0339 (15)	0.0308 (15)	0.0054 (11)	0.0030 (11)	-0.0010 (11)
O21	0.0373 (14)	0.0400 (13)	0.0562 (16)	-0.0004 (11)	-0.0038 (11)	0.0128 (12)
O22	0.0315 (11)	0.0375 (11)	0.0304 (11)	0.0034 (9)	-0.0075 (9)	-0.0002 (8)
O23	0.060 (2)	0.0520 (17)	0.086 (3)	-0.0213 (17)	-0.0142 (18)	-0.0066 (18)

Geometric parameters (Å, °)

C1—C2	1.383 (5)	C12—H12B	0.9900
C1—C11A	1.400 (5)	C13—N14	1.493 (5)
C1—H1	0.9500	C13—H13A	0.9900
C2—C3	1.395 (5)	C13—H13B	0.9900
C2—H2	0.9500	N14—C17	1.487 (4)
C3—C4	1.393 (5)	N14—C15	1.496 (4)
C3—H3	0.9500	N14—H14	0.88 (6)
C4—C4A	1.386 (4)	C15—C16	1.524 (4)
C4—H4	0.9500	C15—H15A	0.9900
C4A—C11A	1.404 (4)	C15—H15B	0.9900
C4A—N5	1.422 (4)	C16—H16A	0.9900
N5—C5A	1.429 (4)	C16—H16B	0.9900
N5—H5	0.90 (5)	C17—H17A	0.9800
C5A—C6	1.394 (5)	C17—H17B	0.9800
C5A—C9A	1.408 (4)	C17—H17C	0.9800
C6—C7	1.387 (6)	C21—C26	1.403 (5)
C6—H6	0.9500	C21—C22	1.406 (5)

C7—C8	1.379 (6)	C21—C27	1.499 (4)
C7—H7	0.9500	C22—O23	1.355 (6)
C8—C9	1.387 (5)	C22—C23	1.383 (7)
C8—C18	1.752 (4)	C23—C24	1.389 (9)
C9—C9A	1.394 (5)	C23—H23A	0.9500
C9—H9	0.9500	C24—C25	1.381 (9)
C9A—N10	1.405 (4)	C24—H24	0.9500
N10—C11	1.283 (5)	C25—C26	1.381 (5)
C11—N11	1.388 (4)	C25—H25	0.9500
C11—C11A	1.488 (4)	C26—H26	0.9500
N11—C12	1.462 (4)	C27—O21	1.253 (4)
N11—C16	1.467 (4)	C27—O22	1.261 (4)
C12—C13	1.520 (4)	O23—H23	1.00 (11)
C12—H12A	0.9900		
C2—C1—C11A	121.4 (3)	N14—C13—C12	111.1 (3)
C2—C1—H1	119.3	N14—C13—H13A	109.4
C11A—C1—H1	119.3	C12—C13—H13A	109.4
C1—C2—C3	119.3 (3)	N14—C13—H13B	109.4
C1—C2—H2	120.3	C12—C13—H13B	109.4
C3—C2—H2	120.3	H13A—C13—H13B	108.0
C4—C3—C2	120.1 (3)	C17—N14—C13	110.8 (3)
C4—C3—H3	120.0	C17—N14—C15	110.4 (3)
C2—C3—H3	120.0	C13—N14—C15	111.3 (3)
C4A—C4—C3	120.3 (3)	C17—N14—H14	110 (3)
C4A—C4—H4	119.9	C13—N14—H14	108 (3)
C3—C4—H4	119.9	C15—N14—H14	106 (3)
C4—C4A—C11A	120.3 (3)	N14—C15—C16	110.5 (3)
C4—C4A—N5	120.0 (3)	N14—C15—H15A	109.5
C11A—C4A—N5	119.7 (3)	C16—C15—H15A	109.5
C4A—N5—C5A	113.8 (2)	N14—C15—H15B	109.5
C4A—N5—H5	111 (3)	C16—C15—H15B	109.5
C5A—N5—H5	108 (3)	H15A—C15—H15B	108.1
C6—C5A—C9A	119.7 (3)	N11—C16—C15	109.8 (3)
C6—C5A—N5	121.5 (3)	N11—C16—H16A	109.7
C9A—C5A—N5	118.8 (3)	C15—C16—H16A	109.7
C7—C6—C5A	121.3 (3)	N11—C16—H16B	109.7
C7—C6—H6	119.4	C15—C16—H16B	109.7
C5A—C6—H6	119.4	H16A—C16—H16B	108.2
C8—C7—C6	118.3 (3)	N14—C17—H17A	109.5
C8—C7—H7	120.9	N14—C17—H17B	109.5
C6—C7—H7	120.9	H17A—C17—H17B	109.5
C7—C8—C9	122.1 (3)	N14—C17—H17C	109.5
C7—C8—C18	119.3 (3)	H17A—C17—H17C	109.5
C9—C8—C18	118.6 (3)	H17B—C17—H17C	109.5
C8—C9—C9A	119.7 (3)	C26—C21—C22	118.7 (3)
C8—C9—H9	120.1	C26—C21—C27	121.2 (3)
C9A—C9—H9	120.1	C22—C21—C27	120.0 (3)

C9—C9A—N10	116.6 (3)	O23—C22—C23	118.4 (4)
C9—C9A—C5A	119.0 (3)	O23—C22—C21	121.1 (4)
N10—C9A—C5A	124.1 (3)	C23—C22—C21	120.5 (4)
C11—N10—C9A	124.0 (3)	C22—C23—C24	119.6 (5)
N10—C11—N11	117.2 (3)	C22—C23—H23A	120.2
N10—C11—C11A	126.6 (3)	C24—C23—H23A	120.2
N11—C11—C11A	115.8 (3)	C25—C24—C23	120.9 (4)
C1—C11A—C4A	118.5 (3)	C25—C24—H24	119.6
C1—C11A—C11	120.3 (3)	C23—C24—H24	119.6
C4A—C11A—C11	121.2 (3)	C24—C25—C26	119.9 (5)
C11—N11—C12	118.3 (3)	C24—C25—H25	120.0
C11—N11—C16	121.7 (3)	C26—C25—H25	120.0
C12—N11—C16	111.0 (2)	C25—C26—C21	120.5 (4)
N11—C12—C13	109.7 (3)	C25—C26—H26	119.8
N11—C12—H12A	109.7	C21—C26—H26	119.8
C13—C12—H12A	109.7	O21—C27—O22	124.0 (3)
N11—C12—H12B	109.7	O21—C27—C21	118.3 (3)
C13—C12—H12B	109.7	O22—C27—C21	117.8 (3)
H12A—C12—H12B	108.2	C22—O23—H23	96 (5)
C11A—C1—C2—C3	-1.1 (5)	N11—C11—C11A—C1	32.9 (4)
C1—C2—C3—C4	-1.9 (5)	N10—C11—C11A—C4A	38.8 (4)
C2—C3—C4—C4A	2.8 (5)	N11—C11—C11A—C4A	-148.7 (3)
C3—C4—C4A—C11A	-0.8 (5)	N10—C11—N11—C12	5.5 (4)
C3—C4—C4A—N5	178.1 (3)	C11A—C11—N11—C12	-167.7 (3)
C4—C4A—N5—C5A	115.3 (3)	N10—C11—N11—C16	-138.6 (3)
C11A—C4A—N5—C5A	-65.7 (3)	C11A—C11—N11—C16	48.2 (4)
C4A—N5—C5A—C6	-117.1 (3)	C11—N11—C12—C13	151.8 (3)
C4A—N5—C5A—C9A	62.6 (4)	C16—N11—C12—C13	-60.6 (3)
C9A—C5A—C6—C7	-0.2 (5)	N11—C12—C13—N14	56.6 (4)
N5—C5A—C6—C7	179.6 (3)	C12—C13—N14—C17	-177.1 (3)
C5A—C6—C7—C8	-1.3 (5)	C12—C13—N14—C15	-53.8 (4)
C6—C7—C8—C9	2.3 (6)	C17—N14—C15—C16	177.3 (3)
C6—C7—C8—C18	-177.1 (3)	C13—N14—C15—C16	53.8 (4)
C7—C8—C9—C9A	-1.6 (5)	C11—N11—C16—C15	-152.7 (3)
C18—C8—C9—C9A	177.7 (3)	C12—N11—C16—C15	60.9 (4)
C8—C9—C9A—N10	174.2 (3)	N14—C15—C16—N11	-56.8 (4)
C8—C9—C9A—C5A	0.0 (5)	C26—C21—C22—O23	-179.2 (4)
C6—C5A—C9A—C9	0.8 (5)	C27—C21—C22—O23	1.2 (6)
N5—C5A—C9A—C9	-178.9 (3)	C26—C21—C22—C23	0.7 (6)
C6—C5A—C9A—N10	-172.9 (3)	C27—C21—C22—C23	-178.9 (4)
N5—C5A—C9A—N10	7.4 (5)	O23—C22—C23—C24	178.5 (5)
C9—C9A—N10—C11	141.9 (3)	C21—C22—C23—C24	-1.4 (8)
C5A—C9A—N10—C11	-44.2 (5)	C22—C23—C24—C25	0.9 (9)
C9A—N10—C11—N11	-171.4 (3)	C23—C24—C25—C26	0.3 (8)
C9A—N10—C11—C11A	1.0 (5)	C24—C25—C26—C21	-1.0 (6)
C2—C1—C11A—C4A	3.1 (5)	C22—C21—C26—C25	0.5 (5)
C2—C1—C11A—C11	-178.6 (3)	C27—C21—C26—C25	-180.0 (3)

C4—C4A—C11A—C1	-2.1 (4)	C26—C21—C27—O21	-178.5 (3)
N5—C4A—C11A—C1	179.0 (3)	C22—C21—C27—O21	1.1 (5)
C4—C4A—C11A—C11	179.6 (3)	C26—C21—C27—O22	1.7 (4)
N5—C4A—C11A—C11	0.6 (4)	C22—C21—C27—O22	-178.7 (3)
N10—C11—C11A—C1	-139.6 (3)		

Hydrogen-bond geometry (Å, °)

Cg1 represents the centroid of the ring (C5A,C6–C9,C9A).

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
N14—H14...O22	0.89 (6)	1.75 (6)	2.612 (4)	164 (5)
O23—H23...O21	1.01 (11)	1.52 (10)	2.507 (5)	166 (9)
C4—H4...O22 ⁱ	0.95	2.33	3.261 (4)	166
C9—H9...O22 ⁱⁱ	0.95	2.25	3.202 (4)	176
C12—H12B...O21	0.99	2.44	3.306 (5)	146
C15—H15A...N5 ⁱⁱ	0.99	2.56	3.539 (4)	170
C24—H24...Cg1 ⁱⁱⁱ	0.95	2.83	3.637 (5)	144

Symmetry codes: (i) $x+1/2, y-1/2, z$; (ii) $x, -y+1, z+1/2$; (iii) $x-1/2, y+1/2, z-1$.