

1 Title:
2 Isothermic and fixed intensity heat acclimation methods elicit equal increases in Hsp72 mRNA.
3
4 2. Submission Type:
5 Original Investigation
6
7 3. Names of Authors:
8 ¹ Oliver R. Gibson, University of Brighton
9 ¹ Jessica A. Mee, University of Brighton
10 ² Lee Taylor, University of Bedfordshire
11 ² James A. Tuttle, University of Bedfordshire
12 ¹ Peter W. Watt, University of Brighton
13 ¹ Neil S. Maxwell, University of Brighton
14
15 4. Contact Details:
16 ¹ Oliver Gibson, o.r.gibson@brighton.ac.uk Centre for Sport and Exercise Science and Medicine (SESAME),
17 University of Brighton, Welkin Human Performance Laboratories, Denton Road, Eastbourne, UK
18
19 ² Muscle Cellular and Molecular Physiology (MCMP) and Applied Sport and Exercise Science (ASEP)
20 Research Groups, Department of Sport Science and Physical Activity, Institute of Sport and Physical Activity
21 Research (ISPAR), University of Bedfordshire, Bedford Campus, Polhill Avenue, Bedfordshire, UK
22
23 5. Preferred Running Head
24 Hsp72 responses to Heat Acclimation methods.
25
26 6. Abstract Word Count
27 214
28
29 7. Text Word Count
30 4,777
31
32 8. Number of Figures and Table
33 Three figures. Two tables.
34
35 9. Keywords:
36 Thermoregulation, Heat Stress, Cellular Stress Response, Hyperthermia, Thermotolerance, Heat Shock Protein
37 72, Heat Illness.

38 **Abstract**

39 Thermotolerance, to which Heat shock protein-72 (Hsp72) contributes, is an acquired state achieved following
40 heat acclimation (HA), eliciting cellular adaption and protection against thermal stress. Optimal HA methods
41 achieving the greatest heat shock response (HSR) are equivocal; therefore investigation of methods provoking
42 the greatest sustained HSR is required to optimise cellular adaptation.

43

44 Twenty four males performed short term HA (STHA; five sessions) and long term HA (LTHA; STHA plus
45 further five sessions) utilising fixed intensity (FIXED; workload = 50% $\dot{V}O_{2peak}$), continuous isothermic HA
46 (ISO_{CONT}; target rectal temperature (T_{rec}) = 38.5°C) or progressive isothermic HA (ISO_{PROG}; target T_{rec} = 38.5°C
47 for STHA then target T_{rec} = 39.0°C for LTHA). Leukocyte Hsp72 mRNA was measured pre and post day 1, day
48 5 and day 10 of HA via qRT-PCR to determine the HSR.

49

50 Hsp72 mRNA increased ($p < 0.05$) pre to post day 1, pre to post day 5, and pre to post day 10 in FIXED, ISO-
51 _{CONT} and ISO_{PROG}, but no differences were observed between methods ($p > 0.05$). The equal Hsp72 mRNA
52 increases occurring from consistent, reduced or increased endogenous strain following STHA and LTHA
53 suggest that transcription occurs following attainment of sufficient endogenous criteria. These data give
54 confidence that all reported HA methods increase Hsp72 mRNA and are capable of eliciting adaptations towards
55 thermotolerance.

56

57 **Introduction**

58 Repeated exposure to stressful thermal environments initiates a phenotypic heat adaptation known as heat
59 acclimation (HA) (Garrett et al. 2011), an element of which has been identified as thermotolerance (Moseley
60 1997). Thermotolerance (Moseley 1997), or acquired cellular thermotolerance (McClung et al. 2008), describes
61 the cellular adaptation accompanying systemic changes (Magalhães et al. 2010b; Sawka et al. 2011; Hom et al.
62 2012) induced by successful HA. Acquired cellular thermotolerance confers cytoprotection against subsequent
63 thermal exposure, translating to complimentary reductions in endogenous physiological and systemic strain
64 (Yamada et al. 2007; McClung et al. 2008). An established element of acquired cellular thermotolerance
65 involves changes in heat shock proteins (HSP) (Moseley 1997); in particular increases in the inducible, and
66 thermosensitive protein heat shock protein HSPA1A (HSP72) (McClung et al. 2008; Beckham et al. 2008;
67 Kampinga et al. 2009) following transcription of its gene (Hsp72 mRNA) as part of the heat shock response
68 (HSR).

69
70 Increased basal HSP72 is commonly reported following repeated exercise-heat stress, as is the inducibility of
71 the protein (Maloyan et al. 1999; McClung et al. 2008; Selkirk et al. 2009; Magalhães et al. 2010b; Amorim et
72 al. 2011). Previously, extracellular HSP72 (eHSP72) has been used as a marker of the stress response. In spite of
73 an established eHSP72 response to sufficient exercise-heat stress (Marshall et al. 2006; Yamada et al. 2007;
74 Ogura et al. 2008; Magalhães et al. 2010b; Périard et al. 2012; Gibson et al. 2014), the mechanisms leading to an
75 increase in circulating concentration are equivocal (Fleshner and Johnson 2005; Lancaster and Febbraio 2005b;
76 Lancaster and Febbraio 2005a). Additionally, the biological role of eHSP72 appears more closely linked to an
77 immunological response, rather than a process favourably augmenting thermotolerance, and the associated
78 cytoprotective adaptations (Asea 2006). The measurement of intracellular HSP72 is optimal for determining
79 cellular responses to HA (Magalhães et al. 2010b). HA increases basal HSP72, improving the cellular defence of
80 heat stress, and also leading to augmented translation during heat stress (Maloyan et al. 1999). The measurement
81 of HSP72 gene expression (Hsp72 mRNA) therefore offers an alternative marker of the magnitude of the
82 cellular stress response, and subsequent initiation of protein transcription required for increased thermotolerance
83 (Maloyan and Horowitz 2002). Based upon previous data (Maloyan et al. 1999) HA should increase the
84 measured Hsp72 mRNA transcription, a process primarily regulated by Heat shock factor protein 1 (HSF-1) as
85 part of the HSR (Kregel 2002).

86

87 HSF1 activation involves a complex series of regulatory events, including nuclear localization, oligomerisation
88 and acquisition of HSE–DNA binding, ultimately resulting in the transcription of Hsp72 mRNA (Sarge et al.
89 1993), this in response to the magnitude of thermal and physiological challenge (Maloyan et al. 1999; McClung
90 et al. 2008).

91

92 Fixed intensity HA methods (Houmard et al. 1990; Nielsen et al. 1993; Nielsen et al. 1997; Cheung and
93 McLellan 1998; Kresfelder et al. 2006; Marshall et al. 2007; Yamada et al. 2007; Watkins et al. 2008;
94 Sandström et al. 2008; Lorenzo et al. 2010; Lorenzo and Minson 2010; Amorim et al. 2011; Castle et al. 2011)
95 derive exercise intensity from pre acclimation baseline testing with the workload and exogenous environment
96 consistent day to day. Whilst thermal stress may be sufficient for the initial sessions of HA, with ongoing
97 adaptation, the relative potentiating stimuli may diminish along with the rate of adaptation, even to the extent
98 that the latter stage of HA are analogous to a reduction in stress (Taylor and Cotter 2006; Taylor 2014).
99 Isothermic HA, also known as controlled hyperthermia, (Patterson et al. 2004; Magalhães et al. 2006; Garrett et
100 al. 2009; Magalhães et al. 2010a; Magalhães et al. 2010b; Hom et al. 2012; Garrett et al. 2012; Castle et al.
101 2012; Garrett et al. 2014; Patterson et al. 2014) imposes session-by-session workloads based upon targeted
102 endogenous criteria (core temperature $\geq 38.5^{\circ}\text{C}$), thus sustaining potentiating stimuli throughout the intervention
103 via a combination of active then passive heat exposure (Fox et al. 1963).

104

105 The aim of the present study was to identify differences in Hsp72 mRNA response to exogenously controlled,
106 fixed intensity HA, an endogenously controlled isothermic HA method, and a progressive endogenous
107 isothermic HA method. We hypothesised that Hsp72 mRNA would increase following completion of an acute
108 HA session, irrespective of the method used; however isothermic methods would sustain the magnitude of
109 increase throughout acclimation due to sustained elevations in core temperature, with an increase in target core
110 temperature progressively increasing transcription.

111

112 **Methods**

113 *Participants*

114 Twenty-four healthy males were assigned into fixed intensity HA (FIXED) (n = 8) continuous isothermic HA
115 (ISO_{CONT}) (n = 8) or progressive isothermic HA (ISO_{PROG}) (n = 8), see Table 1 for descriptive characteristics.
116 Confounding variables of smoking, caffeine, glutamine, alcohol, generic supplementation, prior thermal,
117 hypoxic, and hyperbaric exposures were all controlled in line with previous work in the field (Taylor et al. 2011;

118 Gibson et al. 2014). Following full description of experimental procedures, the methods were approved by the
119 institutional ethics committee. All participants completed medical questionnaires and provided written informed
120 consent following the principles outlined by the Declaration of Helsinki of 1975, as revised in 2013.

121

122 *Preliminary Testing*

123 Participants consumed 500 mL of water 2 h before all preliminary and experimental exercise sessions (Sawka et
124 al. 2007). A urine osmometer (Alago Vitech Scientific, Pocket PAL-OSMO, UK) ensured consistent hydration
125 prior to each experimental session (Garrett et al. 2014) in accordance with established urine osmolality (<700
126 mOsm·Kg⁻¹ H₂O (Sawka et al. 2007)), if this criterion was not met participants consumed 500 mL of water and
127 rested until hydration criteria was achieved. Prior to the $\dot{V}O_{2peak}$ determination, height (cm) was measured using
128 a fixed stadiometer (Detecto Physicians Scales; Cranlea & Co., Birmingham, UK) and NBM recorded to 0.01 kg
129 from digital scales (ADAM GFK 150, USA). Body fat (%) was calculated (Siri 1956) from body density,
130 derived from a four site skin fold calculation (Durnin and Womersley, 1974) using skin fold calipers
131 (Harpenden, Burgess Hill, UK) with body surface area also calculated later (Du Bois and Du Bois 1916).

132

133 $\dot{V}O_{2peak}$ (L·min⁻¹) was determined from an incremental test on a cycle ergometer (Monark e724, Vansbro,
134 Sweden) in temperate conditions (20°C, 40% relative humidity (RH)). Saddle position was adjusted by the
135 participant to their preferred cycling position and remained unchanged for all experimental trials. Starting
136 intensity was set at 80 W with resistance applied to the flywheel eliciting 24 W·min⁻¹ increases at the constant
137 cadence of 80 rpm. Heart rate (HR; b·min⁻¹) was monitored continually during all exercise tests by telemetry
138 (Polar Electro Oyo, Kempele, Finland). Expired metabolic gas was measured using an online system (Metamax
139 3X, Cortex, Germany). $\dot{V}O_{2peak}$ was considered the highest $\dot{V}O_2$ obtained in any 10 s period.

140

141 *Heat Acclimation Protocol*

142 Each HA testing session was conducted in the morning (08:00 ± 01:00 h) to minimise daily variation in
143 performance (Drust et al. 2005). Following provision of a urine sample and measurement of NBM, each
144 participant was equipped with a rectal thermistor (Henleys Medical, UK, Meter logger Model 401, Yellow
145 Springs Instruments, Yellow Springs, Missouri, USA) and a HR monitor. Resting measures, including pre- and
146 post-session venous blood samples, were taken whilst participants were seated in temperate laboratory
147 conditions. Following resting measures, participants mounted a cycle ergometer (Monark, e724, Vansbro,

148 Sweden) located inside an environmental chamber and commenced exercising ($40.2 \pm 0.4^\circ\text{C}$, $39.0 \pm 7.8\%$ RH;
149 WatFlow control system; TISS, Hampshire, UK). FIXED participants performed all ten 90 min sessions cycling
150 continuously at a workload corresponding to 50% $\dot{V}O_{2\text{peak}}$ (80 rpm; 50% $\dot{V}O_{2\text{peak}} = 1.90 \pm 0.30 \text{ L}\cdot\text{min}^{-1}$, power at
151 50% $\dot{V}O_{2\text{peak}} = 125 \pm 30 \text{ W}$). ISO_{CONT} (65% $\dot{V}O_{2\text{peak}} = 2.19 \pm 0.34 \text{ L}\cdot\text{min}^{-1}$, $175 \pm 27 \text{ W}$) and ISO_{PROG} (65%
152 $\dot{V}O_{2\text{peak}} = 2.46 \pm 0.46 \text{ L}\cdot\text{min}^{-1}$, $197 \pm 36 \text{ W}$) participants began exercising at a workload corresponding to 65% of
153 $\dot{V}O_{2\text{peak}}$ until a target T_{rec} of 38.5°C or 39.0°C was achieved, respectively. ISO_{CONT} targeted a T_{rec} of 38.5°C for
154 all ten sessions, whereas ISO_{PROG} targeted a T_{rec} of 38.5°C for the first five sessions progressing to a T_{rec} of
155 39.0°C for the final five sessions. In both ISO_{CONT} and ISO_{PROG}, once target T_{rec} had been reached, power was
156 adjusted every 5 min, first by a 25% $\dot{V}O_{2\text{peak}}$ reduction, and then adjusted ($\pm 5\% \dot{V}O_{2\text{peak}}$, or seated rest) to
157 maintain the experimental T_{rec} for a total session duration of 90 min, exercising duration was calculated based
158 upon the duration of cycling required to reach, and then maintain the target T_{rec} in ISO_{CONT} and ISO_{PROG}. All
159 participants in ISO_{CONT} and ISO_{PROG} were required to rest during both STHA (ISO_{CONT} = $23 \pm 9 \text{ min}\cdot\text{session}^{-1}$;
160 ISO_{PROG} = $37 \pm 9 \text{ min}\cdot\text{session}^{-1}$) and LTHA (ISO_{CONT} = $19 \pm 10 \text{ min}\cdot\text{session}^{-1}$; ISO_{PROG} = $30 \pm 9 \text{ min}\cdot\text{session}^{-1}$),
161 exercise was resumed once core temperature reduced below 38.5°C . During each testing session HR, T_{rec} and
162 power output, were recorded every 5 min, a visual representation of the exercise intensities and T_{rec} responses to
163 STHA and LTHA are presented in Figure 1.

164

165 *Blood Sampling, RNA extraction and One step reverse transcription quantitative polymerase chain reaction*
166 *(RT-QPCR)*

167 Venous blood samples were taken immediately pre- and post- exercise-heat exposure on the first, fifth and tenth
168 experimental sessions for FIXED, ISO_{CONT} and ISO_{PROG}. All blood samples were drawn from the antecubital
169 vein into 6 mL EDTA Vacuette tubes (Grenier BIO-one, UK). 1 mL of venous blood was pipetted into 10 mL of
170 1 in 10 red blood cell lysis solution (10X red blood Cell Lysis Solution, Miltenyi Biotech, UK). Samples were
171 incubated for 15 min at room temperature then isolated via centrifugation at 400G for 5 min and washed twice
172 in 2 mL PBS at 400G for 5 min to isolate all leukocytes. RNA was then extracted via the previously validated
173 acid guanidium thiocyanate–phenol–chloroform extraction method (Chomczynski and Sacchi 1987). Quantity
174 was determined at an optical density of 260 nm while quality was determined via the 260/ 280 and 260/ 230
175 ratios using a nanodrop spectrophotometer (Nanodrop 2000c Thermo Scientific).

176

177 Hsp72 relative mRNA expression (Hsp72 mRNA) was quantified using RT-QPCR. Primers β 2-Microglobulin
178 (β 2-M; NCBI Accession number: NM_004048; Forward CCGTGTGAACCATGTGACT, Reverse,
179 TGCGGCATCTTCAAACCT) and Hsp72 (NCBI Accession number: NM_005345; Forward
180 CGCAACGTGCTCATCTTTGA, Reverse TCGCTTGTCTGGCTGATGT) were designed using primer
181 design software (Primer Quest and Oligoanalyzer - Integrated DNA technologies). 20 μ L reactions containing
182 10 μ L SYBR-Green RT-PCR Mastermix (Quantifast SYBRgreen Kit, Qiagen), 0.15 μ L forward primer, 0.15
183 μ L reverse primer, 0.2 μ L reverse transcription mix (Quantifast RT Mix, Qiagen) and 9.5 μ L sample (70 ng
184 RNA/ μ L) were prepared in separate tubes. Each PCR reaction (Rotorgene Q, Qiagen, Manchester, UK) was
185 then performed as follows: 10 min, 50°C (reverse transcription), 5 min 95°C (transcriptase inactivation and
186 initial denaturation); followed by: 10 s, 95°C (denaturation), 30 s, 60°C (annealing and extension) for 40 cycles.
187 Fluorescence was measured following each cycle as a result of the incorporation of SYBR green dye into the
188 amplified PCR product. Melt curves (50 to 95°C; Ramp protocol 5s stages) were analysed for each reaction to
189 ensure only the single gene of interest was amplified. A comparative critical threshold (CT) method was used to
190 quantify Hsp72 mRNA in comparison with β 2-M (Schmittgen and Livak 2008).

191

192 *Statistical Analysis*

193 All outcome variables were first checked for normality using Kolmogorov-Smirnov and sphericity using the
194 Greenhouse Geisser method prior to further analysis. Two way mixed design ANOVA were performed to
195 determine differences in dependent variables between HA methods for STHA and LTHA timescales (between
196 HA methods and Day 1, Day 5 and Day 10). A three way mixed design ANOVA was performed on the Hsp72
197 mRNA data to determine differences between pre and post value (repeated measures – within subjects) on
198 different days (repeated measures – within subjects) from independent HA methods (between subjects).
199 Adjusted Bonferroni comparisons were used as post hoc analyses, determining where differences existed within
200 ANOVA when a time or interaction was found. Data are reported as mean \pm SD, with two-tailed significance
201 was accepted at $p < 0.05$.

202

203 **Results**

204 *Participant Characteristics*

205 No differences ($p > 0.05$) existed between groups for descriptive variables height, NBM, BSA, body fat % or
206 $\dot{V}O_{2peak}$. A difference ($p < 0.05$) was observed for age whereby ISO_{PROG} was older than FIXED (+6.5 years).

207

208 *Evidence of Heat Acclimation*

209 Resting T_{rec} was reduced ($p = 0.002$), and sweat loss increased ($p = 0.002$) overall, with a significant reduction
210 between Day 1 and Day 10 ($p = 0.003$ and $p = 0.002$ respectively), no interaction effects were observed for
211 resting T_{rec} ($p = 0.592$) or sweat loss ($p = 0.281$), figure 2. Resting HR demonstrated a significant overall effect
212 ($p < 0.001$) and interaction effect ($p = 0.009$), with significant differences observed between Day 1 and Day 5 (p
213 < 0.001) and Day 1 and Day 10 ($p = 0.001$) in ISO_{CONT}, and a difference between ISO_{PROG} and FIXED ($p =$
214 0.043), and ISO_{PROG} and ISO_{CONT} ($p = 0.015$) on Day 1, and between FIXED and ISO_{CONT} ($p = 0.038$), and
215 FIXED and ISO_{PROG} ($p = 0.023$) on Day 10, figure 2.

216

217 *Session Specific Data*

218 Exercising duration ($p = 0.001$), mean session intensity ($p = 0.002$), total work done ($p < 0.001$), mean T_{rec} ($p =$
219 0.002), duration $T_{rec} \geq 38.5^{\circ}\text{C}$ ($p = 0.011$), mean HR ($p = 0.019$), and peak HR ($p < 0.001$) all demonstrated
220 overall differences between days, no between day difference was observed for peak T_{rec} ($p = 0.226$) or duration
221 $T_{rec} \geq 39.0^{\circ}\text{C}$ ($p = 0.245$).

222

223 Exercising duration ($p = 0.004$), mean session intensity ($p = 0.000$), total work done ($p = 0.004$), mean T_{rec} ($p =$
224 0.010), peak T_{rec} ($p = 0.004$), duration $T_{rec} \geq 38.5^{\circ}\text{C}$ ($p = 0.008$), duration $T_{rec} \geq 39.0^{\circ}\text{C}$ ($p = 0.005$) all
225 demonstrated interaction effects, no interaction effect was observed for mean HR ($p = 0.077$) or peak HR ($p =$
226 0.588). See Table 2 for full post hoc analysis.

227

228 No differences between days or the interaction effect were observed for mean exercising intensity ($p = 0.124$; p
229 $= 0.061$), change T_{rec} ($p = 0.227$; $p = 0.109$).

230

231 *Hsp72 mRNA responses*

232 No differences in Hsp72 mRNA were observed between days ($p = 0.236$) or across HA methods between days
233 ($p = 0.167$). Hsp72 mRNA did increase Pre to Post overall ($p < 0.001$), and Pre to Post over time ($p = 0.034$);
234 Day 1 ($p < 0.001$), Day 5 ($p < 0.001$) and Day 10 ($p < 0.001$). No Pre to Post difference occurred between HA
235 methods ($p = 0.069$) or for the Pre to Post, between day, between HA methods interaction ($p = 0.217$); on Day 1
236 (FIXED; 2.3 ± 1.0 to 6.4 ± 2.8 , ISO_{CONT}; 1.9 ± 0.6 to 4.4 ± 1.1 and ISO_{PROG}; 1.9 ± 0.8 to 7.1 ± 2.9), Day 5
237 (FIXED; 2.3 ± 0.8 to 4.2 ± 2.2 , ISO_{CONT}; 2.3 ± 0.8 to 5.3 ± 2.5 and ISO_{PROG}; 2.2 ± 0.5 to 6.3 ± 2.2) and Day 10
238 (FIXED; 2.3 ± 0.7 to 4.3 ± 2.0 , ISO_{CONT}; 2.1 ± 0.7 to 4.3 ± 1.3 and ISO_{PROG}; 2.0 ± 0.5 to 6.1 ± 1.7).

239

240 **Discussion**

241 The aim of this experiment was to determine whether there was a difference in the change in leukocyte Hsp72
242 mRNA expression between fixed intensity, continuous isothermic, and progressive isothermic methods during
243 STHA and LTHA. Participants were successfully matched for anthropometric descriptive data and $\dot{V}O_{2peak}$.
244 ISO_{PROG} participants were observed as older than FIXED although the magnitude of difference is not
245 physiologically relevant with regards to heat stress responses (Kenny et al. 2010). An anticipated increase in
246 Hsp72 mRNA expression was observed pre to post each session of exercise-heat stress across all groups overall.
247 No statistical difference in Hsp72 mRNA existed between HA methods, either pre or post acclimation on day 1,
248 day 5 or day 10.

249

250 In spite of diminished endogenous stress in FIXED due to the ongoing HA adaptations the reduction was not to
251 the extent that mRNA was statistically reduced on day 5 or day 10. Consequently equal signals for the
252 attainment of thermotolerance are present in FIXED (active heat acclimation) as ISO_{CONT} and ISO_{PROG} methods
253 (active and passive acclimation). This is an important observation which suggests that exercise per se is not as
254 significant as hyperthermia. No significant pre to post increase in Hsp72 mRNA was observed by implementing
255 a progressive increase in core temperature/hyperthermia (38.5°C to 39.0°C) suggesting targeting a T_{rec} of 38.5°C
256 is sufficient. The reduced endogenous thermal strain (mean T_{rec} , peak T_{rec} , and duration $T_{rec} \geq 38.5^\circ\text{C}$) did not
257 attenuate Hsp72 mRNA responses observed following FIXED between day 1 and day 5 (following STHA) and
258 day 10 (following LTHA) (Table 2). Previous data from our laboratory has shown FIXED day 1 presents
259 equivalent endogenous strain to that elicited at 50% $\dot{V}O_{2peak}$ in 40°C, whereas day 10 presents strain equivalent
260 to working at the same intensity in just 30°C (Gibson et al. 2014). This reduction in strain due to the ongoing
261 adaptive process of HA. The attenuated endogenous criteria were not apparent within isothermic methods
262 demonstrating the effectiveness of these methods at targeting core temperatures. Correspondingly Hsp72
263 increases were also maintained each day as previously within the field (Magalhães et al. 2010b). Our data
264 further implicates these endogenous thermoregulatory markers as the most relevant signals for manipulating
265 Hsp72 mRNA (Magalhães et al. 2010b) with all the methods tested providing sufficient endogenous stimuli for
266 Hsp72 mRNA transcription. Different duration exercising and workload intensity across day 1 and day 5 and
267 day 10 do not appear relevant contributors to the Hsp72 mRNA response within our experimental design, and
268 are in accordance with previous suggestions (Hom et al. 2012). These observations, that hyperthermia rather

269 than exercise is an important signal for Hsp72 transcription is supported by the equal post exercise expression
270 using active then passive acclimation in ISO_{CONT} and ISO_{PROG}, as active only in FIXED. This is in agreement
271 with other passive heating data (Maloyan et al. 1999). It is not known if this is true of the mean exercise
272 intensity required of each method which, despite not being significantly different between methods, may
273 influence the magnitude of the mRNA response during heat acclimation (e.g. if the FIXED intensity group
274 exercised at an intensity >50% $\dot{V}O_{2peak}$). Increased relative exercise intensity proportionally increases metabolic
275 heat production, thus increasing core temperature (Mora-Rodriguez et al. 2008) which is associated with
276 increased HSP72 (Mestre-Alfaro et al. 2012). This exogenous parameter of exercise-heat stress therefore cannot
277 be disassociated from changes in Hsp72 mRNA in spite of a secondary rather than causal role (Liu et al. 2000;
278 Milne and Noble 2002; Liu et al. 2004).

279

280 Reduced thermal endogenous strain, particularly the attenuated magnitude and rate of core temperature increase,
281 may be most pertinent to the observed reductions in Hsp72 mRNA transcription in this study. These endogenous
282 criteria have been considered as important in other measures of HSP responses to acclimation (Magalhães et al.
283 2010b). Post acclimation day increases in Hsp72 mRNA indicated that the stress presented at the start of HA,
284 and after STHA and LTHA all surpassed the minimum required endogenous strain to elicit increased
285 transcription of Hsp72 mRNA in leukocytes across HA methods. The Hsp72 mRNA response provides further
286 evidence of the importance of providing a consistent stressor for adaptation, via the facilitation of consistent or
287 elevations in core temperature throughout STHA and LTHA. Sustained Hsp72 mRNA increases demonstrate the
288 continued stimulation of the pathway responsible for thermotolerance - the cellular stress response to heat. As
289 Hsp72 mRNA continued to elevate throughout the HA period, complete HSP72 protein mediated acclimation
290 benefits had not been achieved in any method, despite adaptive phenotypic HA responses following both STHA
291 and LTHA (Horowitz and Kodesh 2010). It is currently unknown whether an upper adaptive limit to HA or
292 thermotolerance exists at a cellular level. HA increases baseline HSP72 and blunts inducibility of HSP72 *ex vivo*
293 heat shock (McClung et al. 2008). Theoretically, once stress is presented to a cell, thermotolerance through
294 optimised HSP72 affords sufficient cytoprotection and therefore, normal cell function and homeostasis is
295 maintained without further transcription (Kregel 2002). Implementation of isothermic methods give the greatest
296 efficacy towards continual and consistent magnitudes of Hsp72 mRNA transcription and concurrent increases in
297 HSP72 which are associated with thermotolerance *in vitro* (Kregel 2002), *in vivo* (Maloyan et al. 1999), and HA
298 improvements in heat tolerance (Patterson et al. 2004). Augmented HSP72, enhances cell tolerance to
299 subsequent heat insults translating to enhanced organ, systemic and whole body tolerance (Beckham et al. 2008)

300 and when considering the heat shock response (HSR) to the stress stimuli, a repressed HSF-1 activity. HA and
301 thermotolerance are associated, with greater physiological HA adaptation blunting HSP72 induction to heat
302 shock *ex vivo*, with HA accompanied by elevated baseline and improved regulation of HSP72 (Yamada et al.
303 2007; McClung et al. 2008). It is known that HSR inhibition impairs cellular and systemic adaptations
304 associated with thermotolerance and HA in exercising humans via reductions in circulating cytokines and
305 cellular and systemic markers of heat strain (Kuennen et al. 2011). Phenotypic adaptations occurring throughout
306 STHA and LTHA do not delay or mitigate the HSR requirement of the tested HA methods, with sufficient if not
307 consistent core temperature increases (Hom et al. 2012) augmenting synergistic cellular thermotolerance
308 (Maloyan et al. 1999; Horowitz et al. 2004) alongside systemic HA phenotype adaptations (Moseley 1997).

309

310 Both final/peak, and absolute change in T_{rec} appear to have an effect on HSP72 changes during HA (Magalhães
311 et al. 2010b), this has been previously shown by extracellular HSP72 release (Périard et al. 2012; Gibson et al.
312 2014), and now Hsp72 mRNA, indicating elevated thermal stress. Mechanistically, failure for ISO_{PROG} to elicit
313 significant differences in Hsp72 mRNA in spite of differential mean, peak, and change in T_{rec} in comparison
314 with ISO_{CONT} suggest progressively increasing the endogenous thermal strain through isothermic HA may not
315 augment additional phenotypic HA or acquired cellular thermotolerance. A required “threshold” for the
316 transcription of Hsp72 mRNA appears to be surpassed by ISO_{CONT} over both STHA and LTHA time scales
317 irrespective of a 0.5°C increase in the target temperature suggesting the rate of transcription may be maximal
318 following attainment of an internal temperature of 38.5°C. Maximal mean $T_{rec} \geq 38.5^\circ\text{C}$ were higher in this study
319 and others showing increased HSP72 (McClung et al. 2008; Magalhães et al. 2010b) compared with others
320 where mean $T_{rec} < 38.5^\circ\text{C}$ (Yamada et al. 2007; Hom et al. 2012), no data is available for the duration spent at
321 this T_{rec} . A “threshold” for HA appears to be surpassed by ISO_{CONT} and ISO_{PROG} over LTHA with no further
322 benefit of a 38.5°C to 39.0°C progression in the “threshold”. We observed no difference in Hsp72 mRNA
323 transcription between 38.5°C and 39.0°C T_{rec} , suggesting mean temperature alone may not be the most
324 important signal for increase or that an optimal Hsp72 mRNA transcription rate may occur once a suggested
325 threshold of 38.5°C (T_{rec}) has been surpassed (Morton et al. 2009; Amorim et al. 2011).

326

327 It appears that despite achieving consistent core temperatures, isothermic methods contain some degree of
328 variability in the acute sessional, and adaptive responses. This variability in the response to the isothermic
329 should be acknowledged as a potential limitation of the method. Figure 1 demonstrates that the resting
330 temperature of ISO_{PROG} was lower than the other groups, most notably when compared with ISO_{CONT} during

331 STHA. Additionally ISO_{PROG} required a lower final exercise intensity in than ISO_{CONT}, this despite similar
332 temperature during STHA and higher temperature during LTHA. The variability in isothermic methods is most
333 identifiable from exercise/rest durations between ISO_{CONT} and ISO_{PROG}, and following the progression from
334 STHA to LTHA. Additional duration at rest in LTHA is counter intuitive with heat gain decreasing with
335 adaptation thus greater work is required to achieve the target temperature. This appears true of the initial bout
336 of exercise where attainment of the target temperature is delayed in LTHA compared to STHA (figure 1).
337 Mechanistically, the additional duration at rest in LTHA, compared to STHA is facilitated by the requirement
338 for exercise to be maintained longer during the initial bout of exercise to achieve the target temperature. The
339 result of this is a reduced requirement for participants to resume exercise following rest as the 90 minute session
340 ends before temperature reduces below the target threshold. During STHA, the time to target core temperature is
341 achieved earlier in the session than in LTHA. A greater duration then remains for heat dissipation and
342 temperature reduction, consequently initiating a resumption of exercise in accordance of the requirements of the
343 protocol. The extended first exercise bout in LTHA reduces the time remaining in the session for resuming
344 exercise and thus participants demonstrate less work/lower average intensity of work later in the session. The
345 greater duration of the initial bout of exercise prior to cessation also rationalises some of the differences between
346 ISO_{CONT} and ISO_{PROG} during LTHA. The requirement for a greater change in core temperature in ISO_{PROG},
347 requires participants to exercise for longer initially to attain the higher temperature as such they again perform
348 less work later in the session. These limitations demonstrate the importance of future research optimising
349 isothermic methods so that a greater consistency of protocol administration, and potentially consistency of
350 Hsp72 mRNA transcription is achieved. A larger sample size may reduce the variability in the protocol
351 administration, and may strengthen the observations of the Hsp72 mRNA particularly trends towards reductions
352 in FIXED which may become statistically different given prolonged acclimation (i.e. +10 days) or a greater
353 sample size. It was observed that Hsp72 mRNA Post day 5 ($p = 0.100$) and post day 10 ($p = 0.082$) reduced non
354 significantly in comparison to day 1, an observation not true of ISOCONT (Post day 1 vs. Post day 5 $p = 0.998$;
355 Post day 1 vs. Post day 10 $p = 1.000$) or ISOPROG (Post day 1 vs. Post day 5 $p = 1.000$; Post day 1 vs. Post day
356 10 $p = 0.677$). An explanation for this may relate to the variability in the change in FIXED, physiologically this
357 might be rationalised by individual differences in acclimation rate, and thus endogenous criteria using this
358 protocol; an element that might be further clarified by a larger sample size.

359

360 Future work could involve tissue viability/*ex vivo* experiments to quantify the increased thermotolerance
361 induced between HA methods alongside the measurement of the HSP72 protein, the absence of which is a

362 limitation of the present experiment. Analysis of the acute Hsp72 mRNA response to the first session of
363 progressive isothermic HA would allow analysis of increased hyperthermia from 38.5°C to 39°C to be
364 quantified, although the measurement of mRNA presents a limitation in itself as no data is available to confirm
365 intracellular HSP72 increases, with differential HA methods eliciting different gains in total protein which may
366 in itself augment a changing mRNA/protein ratio. Cellular thermotolerance is unlikely to be explicit to HSP72
367 alone, with a number of genes associated with the cellular stress response to hyperthermia. Therefore a wider
368 genomic and molecular analysis would facilitate further insight into the adaptive mechanisms (Sonna et al.
369 2002). Data suggests an endogenous threshold/minimum criteria may exist for Hsp72 mRNA or HSP72 protein
370 increases as proposed by others (Amorim et al. 2008; Morton et al. 2009; Magalhães et al. 2010b; Périard et al.
371 2012; Gibson et al. 2014). Further investigation of precise endogenous signals leading to greatest intracellular
372 Hsp72 mRNA and HSP72 increases in leukocytes and muscle is warranted to enable links between HA and
373 thermotolerance, to be further examined. This could be facilitated by extended HA durations beyond ten
374 sessions to determine whether in FIXED further diminished endogenous strain would see a continued
375 attenuation of the post session mRNA transcription, or via an experiment where either lower isothermic
376 temperatures are targeted, or changes from baseline implemented to elicit graded minimum thresholds.
377 Individual variability associated with metabolic heat production and retention and the respective effects they
378 may have on Hsp72 mRNA expression could be eliminated by modifying the isothermic method to administer
379 the exercise based upon a fixed relative rate of heat production (Cramer and Jay 2014), further optimising
380 acquired cellular thermotolerance through repeated exercise-heat stress at an optimised asymptote of core
381 temperature increase.

382

383 **Perspectives**

384 Continuous and progressive isothermic HA elicit and sustain similar endogenous systemic strain. This is in
385 contrast to fixed intensity HA which elicits less varied, but diminishing thermoregulatory strain following the
386 procurement of STHA and LTHA adaptations. Hsp72 mRNA transcription, a marker of the cellular stress
387 response to hyperthermia and an important component of thermotolerance, demonstrated equal sessional
388 increases utilising all HA methods. The equal Hsp72 mRNA increases occurring after equal, reduced or
389 increased core temperature following STHA and LTHA suggest that as long as a minimum endogenous criteria
390 is surpassed, additional endogenous thermoregulatory strain is not of further benefit, nor is continual exercise
391 load crucial so long as hyperthermia is present. These data give confidence that all reported HA methods
392 increase Hsp72 mRNA and are capable of eliciting adaptations towards thermotolerance.

393

394 **Acknowledgements**

395 The authors would like to thank the volunteers for their participation in this investigation. Funding and support
396 was obtained from the University of Brighton and the University of Bedfordshire.

397

398 **Conflict of Interest**

399 The authors declare that they have no competing interests such as funding or personal financial interest.

400 **References**

- 401 Amorim F, Yamada P, Robergs R, Schneider S, Moseley P. Effects of whole-body heat acclimation on cell
402 injury and cytokine responses in peripheral blood mononuclear cells. *Eur J Appl Physiol.* 2011;111:1609–18.
- 403 Amorim FT, Yamada PM, Robergs R a, Schneider SM, Moseley PL. The effect of the rate of heat storage on
404 serum heat shock protein 72 in humans. *Eur J Appl Physiol.* 2008;104:965–72.
- 405 Asea A. Initiation of the Immune Response by Extracellular Hsp72: Chaperokine Activity of Hsp72. *Curr*
406 *Immunol Rev.* 2006;2:209–215.
- 407 Beckham JT, Wilmink GJ, Mackanos MA, Takahashi K, Contag CH, Takahashi T, Jansen ED. Role of HSP70
408 in cellular thermotolerance. *Lasers Surg Med.* 2008;40:704–15.
- 409 Castle P, Mackenzie RW, Maxwell N, Webborn ADJ, Watt PW. Heat acclimation improves intermittent
410 sprinting in the heat but additional pre-cooling offers no further ergogenic effect. *J Sports Sci.* 2011;29:1125–
411 34.
- 412 Castle PC, Kularatne BP, Brewer J, Mauger AR, Austen RA, Tuttle JA, Sculthorpe N, Mackenzie RW, Maxwell
413 NS, Webborn ADJ. Partial heat acclimation of athletes with spinal cord lesion. *Eur J Appl Physiol.* 2012;;109–
414 115.
- 415 Cheung SS, McLellan TM. Heat acclimation, aerobic fitness, and hydration effects on tolerance during
416 uncompensable heat stress. *J Appl Physiol.* 1998;84:1731–1739.
- 417 Chomczynski P, Sacchi N. Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-
418 chloroform extraction. *Anal Biochem.* 1987;162:156–9.
- 419 Cramer MN, Jay O. Selecting the correct exercise intensity for unbiased comparisons of thermoregulatory
420 responses between groups of different mass and surface area. *J Appl Physiol.* 2014;116:1123–1132.
- 421 Drust B, Waterhouse J, Atkinson G, Edwards B, Reilly T. Circadian rhythms in sports performance--an update.
422 *Chronobiol Int.* 2005;22:21–44.
- 423 Du Bois D, Du Bois EF. A formula to estimate the approximate surface area if height and weight be known.
424 *Arch Intern Med.* 1916;17:863–871.
- 425 Durnin J V, Womersley J Body fat assessed from total body density and its estimation from skinfold thickness:
426 measurements on 481 men and women aged from 16 to 72 years. *British Journal of Nutrition.* 1974; 32:77–97.
- 427 Fleshner M, Johnson JD. Endogenous extra-cellular heat shock protein 72: releasing signal(s) and function. *Int J*
428 *Hyperth.* 2005;21:457–471.
- 429 Fox RH, Goldsmith R, Kidd DJ, Lewis HE. Acclimatization to heat in man by controlled elevation of body
430 temperature. *J Physiol.* 1963;166:530–47.

431 Garrett AT, Creasy R, Rehrer NJ, Patterson MJ, Cotter JD. Effectiveness of short-term heat acclimation for
432 highly trained athletes. *Eur J Appl Physiol.* 2012;112:1827–37.

433 Garrett AT, Goosens NG, Rehrer NJ, Patterson MJ, Harrison J, Sammut I, Cotter JD. Short-term heat
434 acclimation is effective and may be enhanced rather than impaired by dehydration. *Am J Hum Biol.*
435 2014;26:311–320.

436 Garrett AT, Goosens NG, Rehrer NJ, Rehrer NG, Patterson MJ, Cotter JD. Induction and decay of short-term
437 heat acclimation. *Eur J Appl Physiol.* 2009;107:659–70.

438 Garrett AT, Rehrer NJ, Patterson MJ. Induction and decay of short-term heat acclimation in moderately and
439 highly trained athletes. *Sport Med.* 2011;41:757–71.

440 Gibson OR, Dennis A, Parfitt T, Taylor L, Watt PW, Maxwell NS. Extracellular Hsp72 concentration relates to
441 a minimum endogenous criteria during acute exercise-heat exposure. *Cell Stress Chaperones.* 2014;19:389–400.

442 Hom LL, Lee EC-H, Apicella JM, Wallace SD, Emmanuel H, Klau JF, Poh PYS, Marzano S, Armstrong LE,
443 Casa DJ, Maresh CM. Eleven days of moderate exercise and heat exposure induces acclimation without
444 significant HSP70 and apoptosis responses of lymphocytes in college-aged males. *Cell Stress Chaperones.*
445 2012;17:29–39.

446 Horowitz M, Eli-Berchoer L, Wapinski I, Friedman N, Kodesh E. Stress-related genomic responses during the
447 course of heat acclimation and its association with ischemic-reperfusion cross-tolerance. *J Appl Physiol.*
448 2004;97:1496–507.

449 Horowitz M, Kodesh E. Molecular signals that shape the integrative responses of the heat-acclimated
450 phenotype. *Med Sci Sports Exerc.* 2010;42:2164–72.

451 Houmard JA, Costill DL, Davis JA, Mitchell JB, Pascoe DD, Robergs RA. The influence of exercise intensity
452 on heat acclimation in trained subjects. *Med Sci Sport Exerc.* 1990;22:615–620.

453 Kampinga HH, Hageman J, Vos MJ, Kubota H, Tanguay RM, Bruford E a, Cheetham ME, Chen B, Hightower
454 LE. Guidelines for the nomenclature of the human heat shock proteins. *Cell Stress Chaperones.* 2009;14:105–
455 11.

456 Kenny GP, Gagnon D, Dorman LE, Hardcastle SG, Jay O. Heat balance and cumulative heat storage during
457 exercise performed in the heat in physically active younger and middle-aged men. *Eur J Appl Physiol.*
458 2010;109:81–92.

459 Kregel KC. Heat shock proteins: modifying factors in physiological stress responses and acquired
460 thermotolerance. *J Appl Physiol.* 2002;92:2177–86.

461 Kresfelder TL, Claassen N, Cronjé MJ. Hsp70 Induction and hsp70 Gene polymorphisms as Indicators of
462 acclimatization under hyperthermic conditions. *J Therm Biol.* 2006;31:406–415.

463 Kuennen M, Gillum T, Dokladny K, Bedrick E, Schneider S, Moseley P. Thermotolerance and heat acclimation
464 may share a common mechanism in humans. *Am J Physiol Regul Integr Comp Physiol.* 2011;301:R524–33.

465 Lancaster GI, Febbraio M a. Mechanisms of stress-induced cellular HSP72 release: implications for exercise-
466 induced increases in extracellular HSP72. *Exerc Immunol Rev.* 2005;11:46–52.

467 Lancaster GI, Febbraio MA. Exosome-dependent trafficking of HSP70: a novel secretory pathway for cellular
468 stress proteins. *J Biol Chem.* 2005;280:23349–23355.

469 Liu Y, Lormes W, Baur C, Opitz-Gress A, Altenburg D, Lehmann M, Steinacker JM. Human skeletal muscle
470 HSP70 response to physical training depends on exercise intensity. *Int J Sports Med.* 2000;21:351–5.

471 Liu Y, Lormes W, Wang L, Reissnecker S, Steinacker JM. Different skeletal muscle HSP70 responses to high-
472 intensity strength training and low-intensity endurance training. *Eur J Appl Physiol.* 2004;91:330–5.

473 Lorenzo S, Halliwill JR, Sawka MN, Minson CT. Heat acclimation improves exercise performance. *J Appl*
474 *Physiol.* 2010;109:1140–7.

475 Lorenzo S, Minson CT. Heat acclimation improves cutaneous vascular function and sweating in trained cyclists.
476 *J Appl Physiol.* 2010;109:1736–43.

477 Magalhães FC, Passos RLF, Fonseca MA, Oliveira KPM, Ferreira-Júnior JB, Martini ARP, Lima MRM,
478 Guimarães JB, Baraúna VG, Silami-Garcia E, Rodrigues LOC. Thermoregulatory efficiency is increased after
479 heat acclimation in tropical natives. *J Physiol Anthropol.* 2010;29:1–12.

480 Magalhães FDC, Amorim FT, Passos RLF, Fonseca MA, Oliveira KPM, Lima MRM, Guimarães JB, Ferreira-
481 Júnior JB, Martini ARP, Lima NR V, Soares DD, Oliveira EM, Rodrigues LOC. Heat and exercise acclimation
482 increases intracellular levels of Hsp72 and inhibits exercise-induced increase in intracellular and plasma Hsp72
483 in humans. *Cell Stress Chaperones.* 2010;15:885–95.

484 Magalhães FDC, Machado-Moreira CA, Vimieiro-Gomes AC, Silami-Garcia E, Lima NRV, Rodrigues LOC.
485 Possible Biphasic Sweating Response during Short-term Heat Acclimation Protocol for Tropical Natives. *J*
486 *Physiol Anthropol.* 2006;25:215–219.

487 Maloyan A, Horowitz M. beta-Adrenergic signaling and thyroid hormones affect HSP72 expression during heat
488 acclimation. *J Appl Physiol.* 2002;93:107–15.

489 Maloyan A, Palmon A, Horowitz M. Heat acclimation increases the basal HSP72 level and alters its production
490 dynamics during heat stress. *Am J Physiol.* 1999;276:R1506–15.

491 Marshall HC, Campbell SA, Roberts CW, Nimmo MA. Human physiological and heat shock protein 72
492 adaptations during the initial phase of humid-heat acclimation. *J Therm Biol.* 2007;32:341–348.

493 Marshall HC, Ferguson RA, Nimmo MA. Human resting extracellular heat shock protein 72 concentration
494 decreases during the initial adaptation to exercise in a hot, humid environment. *Cell Stress Chaperones.*
495 2006;11:129–134.

496 McClung JP, Hasday JD, He JR, Montain SJ, Chevront SN, Sawka MN, Singh IS. Exercise-heat acclimation in
497 humans alters baseline levels and ex vivo heat inducibility of HSP72 and HSP90 in peripheral blood
498 mononuclear cells. *Am J Physiol Regul Integr Comp Physiol.* 2008;294:R185–91.

499 Mestre-Alfaro A, Ferrer MD, Banquells M, Riera J, Drobic F, Sureda A, Tur JA, Pons A. Body temperature
500 modulates the antioxidant and acute immune responses to exercise. *Free Radic Res.* 2012;46:799–808.

501 Milne KJ, Noble EG. Exercise-induced elevation of HSP70 is intensity dependent. *J Appl Physiol.*
502 2002;93:561–8.

503 Mora-Rodriguez R, Del Coso J, Estevez E. Thermoregulatory responses to constant versus variable-intensity
504 exercise in the heat. *Med Sci Sports Exerc.* 2008;40:1945–52.

505 Morton JP, Kayani AC, McArdle A, Drust B. The Exercise-Induced Stress Response of Skeletal Muscle, with
506 Specific Emphasis on Humans. *Sport Med.* 2009;39:643–662.

507 Moseley PL. Heat shock proteins and heat adaptation of the whole organism. *J Appl Physiol.* 1997;83:1413–
508 1417.

509 Nielsen B, Hales JR, Strange S, Christensen NJ, Warberg J, Saltin B. Human circulatory and thermoregulatory
510 adaptations with heat acclimation and exercise in a hot, dry environment. *J Physiol.* 1993;460:467–485.

511 Nielsen B, Strange S, Christensen NJ, Warberg J, Saltin B. Acute and adaptive responses in humans to exercise
512 in a warm, humid environment. *Pflugers Arch.* 1997;434:49–56.

513 Ogura Y, Naito H, Akin S, Ichinoseki-Sekine N, Kurosaka M, Kakigi R, Sugiura T, Powers SK, Katamoto S,
514 Demirel H a. Elevation of body temperature is an essential factor for exercise-increased extracellular heat shock
515 protein 72 level in rat plasma. *Am J Physiol Regul Integr Comp Physiol.* 2008;294:R1600–7.

516 Patterson M. Sustained and generalized extracellular fluid expansion following heat acclimation. *J Physiol.*
517 2004;559:327–34.

518 Patterson MJ, Stocks JM, Taylor N a S. Whole-body fluid distribution in humans during dehydration and
519 recovery, before and after humid-heat acclimation induced using controlled hyperthermia. *Acta Physiol (Oxf).*
520 2014;210:899–912.

521 Patterson MJ, Stocks JM, Taylor NAS. Humid heat acclimation does not elicit a preferential sweat redistribution
522 toward the limbs. *Am J Physiol Regul Integr Comp Physiol.* 2004;286:R512–8.

523 Périard JD, Ruell P, Caillaud C, Thompson MW. Plasma Hsp72 (HSPA1A) and Hsp27 (HSPB1) expression
524 under heat stress: influence of exercise intensity. *Cell Stress Chaperones.* 2012;17:375–83.

525 Sandström ME, Siegler JC, Lovell RJ, Madden L a, McNaughton L. The effect of 15 consecutive days of heat-
526 exercise acclimation on heat shock protein 70. *Cell Stress Chaperones.* 2008;13:169–75.

527 Sarge KD, Murphy SP, Morimoto RI. Activation of heat shock gene transcription by heat shock factor 1
528 involves oligomerization, acquisition of DNA-binding activity, and nuclear localization and can occur in the
529 absence of stress. *Mol Cell Biol.* 1993;13:1392–407.

530 Sawka MN, Burke LM, Eichner ER, Maughan RJ, Montain SJ, Stachenfeld NS. American College of Sports
531 Medicine position stand. Exercise and fluid replacement. *Med Sci Sports Exerc.* 2007;39:377–90.

532 Sawka MN, Leon LR, Montain SJ, Sonna LA. Integrated physiological mechanisms of exercise performance,
533 adaptation, and maladaptation to heat stress. *Compr Physiol.* 2011;1:1883–928.

534 Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative CT method. *Nat Protoc.*
535 2008;3:1101–1108.

536 Selkirk GA, McLellan TM, Wright HE, Rhind SG. Expression of intracellular cytokines, HSP72, and apoptosis
537 in monocyte subsets during exertional heat stress in trained and untrained individuals. *Am J Physiol Regul
538 Integr Comp Physiol.* 2009;296:R575–86.

539 Siri WE. The gross composition of the body. *Adv Biol Med Phys.* 1956;4:239–280.

540 Sonna LA, Fujita J, Gaffin SL, Lilly CM. Invited review: Effects of heat and cold stress on mammalian gene
541 expression. *J Appl Physiol.* 2002;92:1725–42.

542 Taylor L, Midgley AW, Christmas B, Hilman AR, Madden L a, Vince R V, McNaughton LR. Daily hypoxia
543 increases basal monocyte HSP72 expression in healthy human subjects. *Amino Acids.* 2011;40:393–401.

544 Taylor N, Cotter J. Heat adaptation: guidelines for the optimisation of human performance. *Int Sport Med J.*
545 2006;7:33–57.





546 Taylor NAS. Human Heat Adaptation. *Compr Physiol.* 2014;4:325–365.

547 Watkins AM, Cheek DJ, Harvey AE, Blair KE, Mitchell JB. Heat Acclimation and HSP-72 Expression in
548 Exercising Humans. *Int J Sports Med.* 2008;29:269–276.

549 Yamada PM, Amorim FT, Moseley P, Robergs R, Schneider SM. Effect of heat acclimation on heat shock
550 protein 72 and interleukin-10 in humans. *J Appl Physiol.* 2007;103:1196–204.

551

552



53

54

55 Figure 1. Mean \pm SD T_{rec} (top; $^{\circ}\text{C}$) and exercise intensity (bottom; $\% \dot{V}O_{2peak}$) for the first five sessions (STHA: left) and all ten sessions (LTHA: right) of fixed intensity (FIXED,
56 $n = 8$), continuous isothermic (ISO_{CONT} , $n = 8$), and progressive isothermic (ISO_{PROG} , $n = 8$) heat acclimation methods. Error bars have been removed from T_{rec} data for clarity.
57





Figure 2 Mean \pm SD Changes in resting T_{rec} , resting heart rate and sweat rate following STHA (Day 1 to 5) utilising fixed intensity (FIXED), continuous isothermic (ISO_{CONT}), and progressive isothermic (ISO_{PROG}) methods. * denotes significant difference overall from Day 1 ($p < 0.05$). # denotes significant difference within group and Day ($p < 0.05$). ^ denotes significant difference from ISO_{PROG} within group and Day ($p < 0.05$). † denotes significant difference from FIXED within group and Day 1 ($p < 0.05$).

566
567
568
569



570
 571
 572 Figure 3 Mean \pm SD Hsp72 mRNA pre and post sessions on Day 1, Day 5 and Day 10 of fixed intensity (FIXED)
 573 continuous isothermic (ISO_{CONT}), and progressive isothermic (ISO_{PROG}) methods. * denotes significant Pre to Post
 574 difference within session ($p < 0.05$).
 575
 576
 577

Table 1. Mean \pm SD Participant characteristics for fixed intensity (FIXED), continuous isothermic (ISO_{CONT}), and progressive isothermic (ISO_{PROG}) heat acclimation methods.

	FIXED	ISO _{CONT}	ISO _{PROG}
Age (years)	19.9 \pm 1.0	22.6 \pm 5.5	26.1 \pm 4.9*
Height (cm)	179.3 \pm 5.8	177.9 \pm 5.8	179.5 \pm 6.6
Body Mass (kg)	79.2 \pm 18.3	74.2 \pm 6.9	75.1 \pm 8.8
BSA (m ²)	1.97 \pm 0.21	1.92 \pm 0.11	1.94 \pm 0.11
Body fat (%)	14.9 \pm 7.7	14.8 \pm 2.2	14.1 \pm 3.5
$\dot{V}O_{2peak}$ (L.min ⁻¹)	3.61 \pm 0.90	3.62 \pm 0.69	3.79 \pm 0.55

*denotes significantly difference from FIXED ($p < 0.05$)

593 Table 2. Mean \pm SD Protocol, thermoregulatory and physiological data characterising exercise – heat stress on day one, day five and day ten of fixed intensity
 594 (FIXED), continuous isothermic (ISO_{CONT}), and progressive isothermic (ISO_{PROG}) methods.

	Day 1			Day 5			Day 10		
	FIXED	ISO _{CONT}	ISO _{PROG}	FIXED	ISO _{CONT}	ISO _{PROG}	FIXED	ISO _{CONT}	ISO _{PROG}
Exercising Duration (min)	90.0 \pm 0.0	61.9 \pm 10.7†	56.3 \pm 16.6†	90.0 \pm 0.0	76.3 \pm 15.5*	53.1 \pm 10.3†^	90.0 \pm 0.0	78.8 \pm 15.8*	70.0 \pm 9.3* #
Mean Session Intensity (% $\dot{V}O_{2peak}$)	49.7 \pm 0.6	36.6 \pm 5.3†	36.7 \pm 11.2†	50.0 \pm 0.0	47.0 \pm 8.3*	32.3 \pm 8.6†^	50.0 \pm 0.0	50.5 \pm 9.5*	45.8 \pm 8.0*#
Mean Exercising Intensity (% $\dot{V}O_{2peak}$)	49.7 \pm 0.6	52.6 \pm 8.2	58.8 \pm 5.1	50.0 \pm 0.0	57.4 \pm 4.9	56.8 \pm 5.9	50.0 \pm 0.0	58.7 \pm 7.0	58.9 \pm 6.2
Total Work Done (kJ)	656 \pm 166	498 \pm 81	554 \pm 102	673 \pm 165	657 \pm 100*	500 \pm 152	684 \pm 164	719 \pm 126*	708 \pm 176*#
Mean T _{rec} (°C)	38.17 \pm 0.17	38.15 \pm 0.23	38.21 \pm 0.25	37.85 \pm 0.22*	38.10 \pm 0.19	38.27 \pm 0.24†	37.74 \pm 0.19*	38.04 \pm 0.23†	38.18 \pm 0.21†
Peak T _{rec} (°C)	38.92 \pm 0.26	38.65 \pm 0.32	38.87 \pm 0.18	38.52 \pm 0.43*	38.66 \pm 0.25	38.91 \pm 0.24	38.40 \pm 0.33*	38.67 \pm 0.23	39.06 \pm 0.37†
Δ T _{rec} (°C)	1.81 \pm 0.60	1.53 \pm 0.37	2.01 \pm 0.33	1.47 \pm 0.74	1.74 \pm 0.20	2.10 \pm 0.42	1.56 \pm 0.72	1.95 \pm 0.32	2.32 \pm 0.61†
Duration T _{rec} \geq 38.5°C (min)	32.5 \pm 8.5	28.8 \pm 15.1	44.4 \pm 21.3	13.1 \pm 16.0*	22.5 \pm 20.7	51.3 \pm 18.5†^	5.0 \pm 8.0*	29.4 \pm 23.5†	35.6 \pm 18.6†
Duration T _{rec} \geq 39.0°C (min)	5.6 \pm 12.1	0.0 \pm 0.0	1.9 \pm 3.7	1.3 \pm 3.5	2.5 \pm 7.1	6.9 \pm 14.4	0.0 \pm 0.0	0.0 \pm 0.0	20.0 \pm 16.0#†^
Mean HR (b.min ⁻¹)	159 \pm 12	151 \pm 13	144 \pm 9	149 \pm 21	148 \pm 9	140 \pm 8	146 \pm 14	151 \pm 8	144 \pm 14
Peak HR (b.min ⁻¹)	176 \pm 12	183 \pm 9	182 \pm 11	171 \pm 26	172 \pm 12	174 \pm 8	164 \pm 13	174 \pm 11	171 \pm 13

595 Notes: Exercising duration is cumulative time spent exercising during each of the 90 min sessions. Mean session intensity is calculated from each participant's relative exercise intensity during
596 each five min period including rest periods during the given session. Mean exercise intensity is calculated from each participant's relative exercise intensity during each five min period
597 excluding rest periods during the given session.
598

599 * denotes difference from Day 1 within respective method ($p < 0.05$). # denotes difference from Day 5 within respective method ($p < 0.05$).

600 † denotes difference from FIXED within respective day ($p < 0.05$). ^ denotes difference from ISO_{CONT} within respective day ($p < 0.05$).

601

602