

VU Research Portal

Severe role impairment associated with mental disorders

WHO WMH-ICS Collaborators

published in

Depression and Anxiety
2018

DOI (link to publisher)

[10.1002/da.22778](https://doi.org/10.1002/da.22778)

document version

Publisher's PDF, also known as Version of record

document license

Article 25fa Dutch Copyright Act

[Link to publication in VU Research Portal](#)

citation for published version (APA)

WHO WMH-ICS Collaborators (2018). Severe role impairment associated with mental disorders: Results of the WHO World Mental Health Surveys International College Student Project. *Depression and Anxiety*, 35(9), 802-814. <https://doi.org/10.1002/da.22778>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.


- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl



RESEARCH ARTICLE



Severe role impairment associated with mental disorders: Results of the WHO World Mental Health Surveys International College Student Project

Jordi Alonso MD, PhD^{1,2,3} | Philippe Mortier PhD⁴ | Randy P. Auerbach PhD^{5,6} | Ronny Bruffaerts PhD⁴ | Gemma Vilagut PhD^{1,3} | Pim Cuijpers PhD⁷ | Koen Demyttenaere PhD⁴ | David D. Ebert PhD⁸ | Edel Ennis PhD⁹ | Raul A. Gutiérrez-García PhD¹⁰ | Jennifer Greif Green PhD¹¹ | Penelope Hasking PhD¹² | Christine Lochner MA, PhD¹³ | Matthew K. Nock PhD¹⁴ | Stephanie Pinder-Amaker PhD^{5,6} | Nancy A. Sampson BA¹⁵ | Alan M. Zaslavsky PhD¹⁵ | Ronald C. Kessler PhD¹⁵ | WHO WMH-ICS Collaborators¹

¹Health Services Research Group, IMIM (Hospital del Mar Medical Research Institute), Barcelona, Spain

²Pompeu Fabra University (UPF), Barcelona, Spain

³CIBER en Epidemiología y Salud Pública (CIBERESP), Madrid, Spain

⁴Universitair Psychiatrisch Centrum–Katholieke Universiteit Leuven (UPC–KUL), Campus Gasthuisberg, Leuven, Belgium

⁵Department of Psychiatry, Harvard Medical School, Boston, MA, USA

⁶Center for Depression, Anxiety and Stress Research, McLean Hospital, Belmont, MA, USA

⁷Department of Clinical, Neuro and Developmental Psychology, Amsterdam Public Health Research Institute, Vrije Universiteit Amsterdam, Amsterdam, the Netherlands

⁸Department for Psychology, Clinical Psychology and Psychotherapy Friedrich-Alexander University Erlangen Nuremberg, Erlangen, Germany

⁹School of Psychology, Ulster University, Londonderry, United Kingdom

¹⁰Campus Salamanca, De La Salle Bajío University, León, Mexico

¹¹School of Education, Boston University, Boston, MA, USA

¹²School of Psychology & Speech Pathology, Curtin University, Perth, Australia

¹³SAMRC/UCT/SU Unit on Risk & Resilience in Mental Disorders, Department of Psychiatry, Stellenbosch University, Stellenbosch, South Africa

¹⁴Department of Psychology, Harvard University, Cambridge, MA, USA

¹⁵Department of Health Care Policy, Harvard Medical School, Boston, MA, USA

Correspondence

Jordi Alonso, Health Services Research Group, IMIM (Hospital del Mar Medical Research Institute), Barcelona, Spain.
Email: jalonso@imim.es

Background: College entrance is a stressful period with a high prevalence of mental disorders.

Aims: To assess the role impairment associated with 12-month mental disorders among incoming first-year college students within a large cross-national sample.

Methods: Web-based self-report surveys assessing the prevalence of DSM-IV mental disorders and health-related role impairment (Sheehan Disability Scale) were obtained and analyzed from 13,984 incoming first-year college students (Response = 45.5%), across 19 universities in eight countries. Impairment was assessed in the following domains: home management, work (e.g., college-related problems), close personal relationships, and social life.

Results: Mean age of the sample was 19.3 ($SD = 0.59$) and 54.4% were female. Findings showed that 20.4% of students reported any severe role impairment (10% of those without a mental disorder vs. 42.9% of those with at least one disorder, $P < 0.01$). In bivariate analyses, panic disorder, and mania were associated most frequently with severe impairment (60.6% and 57.5%, respectively). Students reporting three or more mental disorders had almost fivefold more

frequently severe impairment relative to those without mental disorders. Multiple logistic regression showed that major depression (OR = 4.0; 95%CI = 3.3, 4.8), generalized anxiety (OR = 3.9; 95%CI = 3.1, 4.8), and panic disorder (OR = 2.9; 95%CI 2.4, 4.2) were associated with the highest odds of severe impairment. Only minimal deviations from these overall associations were found across countries.

Conclusion: Mental disorders among first-year college students are associated with substantial role impairment. Providing preventative interventions targeting mental disorders and associated impairments is a critical need for institutions to address.

KEYWORDS

anxiety disorders, depression, disability, drug use disorders, intimate relationships, role impairment, social function, suicide/self-harm, university students

1 | INTRODUCTION

College students are a key group in society in terms of human capital (Abel & Deitz, 2012) as they will drive future economic growth and innovation. They are an increasingly higher proportion of the population younger than 25 in developed countries (OECD, 2016). College is a peak period for mental disorders (Auerbach et al., 2016; Blanco et al., 2008; Cho et al., 2015; Eisenberg, Golberstein, & Gollust, 2007; Ibrahim, Kelly, Adams, & Glazebrook, 2013; Kendler, Myers, & Dick, 2015; Mojtabai et al., 2015; Vazquez, Torres, Otero, & Diaz, 2011), particularly mood, anxiety, and substance use disorders. A recent meta-analysis reported that an average of 30.6% of the students suffered from depression (Ibrahim et al., 2013). This high prevalence is significant for both the distress caused and its associated impairment in academic performance (Auerbach et al., 2016, 2018; Bruffaerts et al., 2012, 2018; Skidmore, Kaufman, & Crowell, 2016) as well as suicidal thoughts and behaviors (Mortier, Auerbach, et al., 2018; Mortier, Cuijpers, et al., 2018). However, research is needed to clarify how mental disorders may affect important functioning across key life domains.

Mental disorders are a significant contributor to disability in the general population (Ormel et al., 2008; Verger, Guagliardo, Gilbert, Rouillon, & Kovess-Masfetty, 2010). Among students, mental disorders have a substantial impact on academic performance (Auerbach et al., 2016; Kessler, Foster, Saunders, & Stang, 1995), and prematriculation onset disorders are strong predictors of college attrition (Auerbach et al., 2016). More broadly, mental disorders are associated with lower employment in adulthood (Mojtabai et al., 2015). Nevertheless, impairment in other domains, including home management/chores, work, close personal relationships, and social life remains poorly understood. An enhanced understanding of the mental disorder-related role impairment is critical, as this may inform specific care needs of students. Addressing those needs may potentially reduce individual suffering of patients and their families as well as increase human capital of our societies. A French study reported that the presence of any 12-month mental disorder was associated with a 10-fold likelihood of "marked" or "extreme" impairment in the Sheehan Disability Scale among university students (Verger et al., 2010). Nevertheless, the amount and type of disability associated with mental disorders among university students has been

assessed only in a few, single country studies (Klemenc-Ketis, Kersnik, Eder, & Colaric, 2011; Verger et al., 2010). The WHO World Mental Health Surveys International College Student Project (WMH-ICS) was developed to obtain accurate longitudinal information about the prevalence, correlates and impact of mental, substance, and behavioral disorders among college students throughout the world [https://www.hcp.med.harvard.edu/wmh/college_student_survey.php]. The aims in doing this were to assess unmet need for treatment, to develop a practical method of targeting students in need of outreach, and to lay the groundwork for the implementation and evaluation of preventive and clinical interventions. The initial phase of the initiative, which is the focus of the current report, involves carrying out surveys with representative samples in a total of 19 colleges of eight countries of incoming college students to estimate prevalence of mental disorders, associated impairments, adverse social and academic consequences, and patterns of help-seeking.

The objectives of this paper are to estimate among incoming first-year college students: (a) the prevalence of role impairment (home management/chores, college-related work, close relationships, and social life) associated with mental disorders; and (b) the modification of this association by comorbid mental disorders. A secondary aim was to test whether there was between-country variation in role impairment associated with mental disorders.

2 | METHOD

2.1 | Study design

Cross-sectional baseline data from surveys of participants at the WMH-ICS project in a total of 19 colleges of eight countries were used for this study. Most participating sites are following first-year students every 12 months (e.g., from the first to final year of studies). However, data collection remains ongoing, and thus, follow-up data are beyond the scope of the current manuscript.

2.2 | Participants

The initial round of WMH-ICS surveys were carried out in eight countries (Australia, Belgium, Germany, Mexico, Northern Ireland,

South Africa, Spain, and the United States). Details on the participating countries and college surveys are provided in Table 1. Web-based self-report questionnaires were administered to all incoming first-year students from 19 participating universities across these countries (7 private, 12 public) between October 2014 and February 2017. Inclusion criteria for participating in the survey were the same in all countries: (a) university students 18+ years-old; (b) enrolled in the first year and for the first time in the participating universities; and (c) fluent in the official language of the country. We excluded all those participants not complying with all the inclusion criteria and those who refused to provide their informed consent. A total of 14,371 questionnaires were completed, with sample sizes ranging from a low of 633 in Australia to a high of 4,580 in Belgium. The weighted by achieved sample size mean response rate across all surveys was 45.5%. An overview of the participating sample and invitation procedures carried out in each country is provided in Online Data Supplement 1. The sample for the analyses reported here was restricted to students identifying themselves as male or female who were full-time students ($n = 13,984$).

Sociodemographic differences between respondents and nonrespondents were assessed when constructing the nonresponse survey weights. Females were over-represented in the survey. Data were weighted to adjust for differences between survey respondents and nonrespondents on whatever sociodemographic information was made available about the student body by University officials using poststratification weights (Groves & Couper, 1998).

2.3 | Procedures

All incoming first-year students in the participating universities were invited to participate in a web-based self-report health survey. The initial mode of contact varied across universities, with the survey being either a part of a health evaluation in some universities, a part of the registration process in others, or implemented as a stand-alone survey administered to students via their student email addresses in still others. In all cases, other than in Mexico, potential respondents were invited to participate, and initial nonrespondents were recontacted through a series of personalized reminder emails containing unique electronic links to the survey. Ten universities implemented conditional incentives in the final stages of refusal conversion (e.g., a raffle for store credit coupons, movie passes). In addition, one site (Spain) used an “end-game strategy” consisting of a random sample of nonrespondents at the end of the normal recruitment period that was offered incentives for participation. Respondents to these end-game interviews were given a weight equal to $1/p$, where p represented the proportion of nonrespondents at the end of the normal recruitment period that was included in the end-game, to adjust for the under-sampling of these hard-to-recruit respondents. The situation was different in Mexico, where students were invited to participate in conjunction with mandatory activities, which varied from school to school (e.g., student health evaluations; tutoring sessions), with time set aside for completing the survey during the sessions. Informed consent was obtained before administering the questionnaires in all countries. Procedures for obtaining informed consent and protecting human participants were approved and monitored for compliance by the institutional review

boards of the organizations coordinating the surveys in each country. At the end of the survey, all respondents received a general notification which provides information on how to access specialized mental health services if they considered it necessary for them. Students endorsing recent and/or severe suicide thoughts and behaviors received information about available resources within their school and/or community.

2.4 | Measures

2.4.1 | Role impairment

Severity of health-related role impairment during the past 12 months was assessed using an adapted version of the Sheehan Disability Scale (Leon, Olsson, Portera, Farber, & Sheehan, 1997; Ormel et al., 2008). This was completed for four role domains home management/chores, college-related and other work, close personal relationships, and social life. Home management specified the activities “cleaning, shopping, and working around the house, apartment or yard.” College-related and other work specified “ability to work as well as most of other people.” Close personal relationships were specified as “the ability to initiate and maintain close personal relationships,” whereas social life was asked without further specification. A 0–10 visual analogue scale was used to rate the degree of impairment for each domain. The scale was labeled as no interference (0), mild (1–3), moderate (4–6), severe (7–9), and very severe (10) interference. Severe self-reported role impairment was defined as having a 7–10 rating (Kessler & Ustun, 2004; Wittchen, Nelson, & Lachner, 1998). In our study Cronbach's alpha (α) for the overall SDS scale was 0.876.

2.4.2 | Mental disorders

Due to the size and logistical complexities of the surveys, it was impossible to administer an in-depth psychiatric diagnostic interview to each student. Instead, the survey instrument consisted of a broad range of short validated self-report screening scales. These included the Composite International Diagnostic Interview Screening Scales (CIDI-SC) (Kessler & Ustun, 2004; Kessler et al., 2013) for major depressive episode, mania/hypomania (broad mania), generalized anxiety disorder (GAD), panic disorder (PD), and drug use disorder (abuse or dependence either on cannabis, cocaine, or any other street drug, or on a prescription drug either used without a prescription or with used more than prescribed to get high, buzzed, or numbed out); and the AUDIT screening scale for alcohol use disorder (Saunders, Aasland, Babor, de la Fuente, & Grant, 1993). The CIDI-SC scales have concordance with blinded clinical diagnoses in the range $AUC = 0.70$ – 0.78 (16). In line with recommendation (Babor, Higgins-Biddle, Saunders, & Monteiro, 2001), we defined alcohol use disorder (abuse or dependence) in the AUDIT as having a total score of 8+ and a score of 4+ on the AUDIT dependence score. This version of AUDIT scoring has concordance with clinical diagnosis in the range $AUC = 0.78$ – 0.91 (Reinert & Allen, 2002). Additional items taken from the CIDI (Kessler & Ustun, 2004) were used to assess age of onset of each disorder and number of months with symptoms in the past year. Participants with a positive response for lifetime disorder were assessed for the number of months in the past year that they experienced symptoms. All

TABLE 1 WMH-ICS sample characteristics

Country	Number of participating universities	Total size of the universities	Number of first-year students eligible	Number of first-year students participated	Response rate	Survey field dates	Sampling and procedures
Australia	1 Public	~45,000	9,042	633	7.0%	2016	All first-year students were invited to participate through e-mail. Five reminder emails were sent with personalized links to the survey. Conditional incentives were applied (movie passes).
Belgium	1 Public	~40,000	8,530	4,580	53.7%	2014–2016	All first-year students were invited for a psycho-medical check-up in the student mental health center. Surveys were completed in the waiting room. Students who did not show up for the psycho-medical check-up received up to eight reminder emails. Conditional incentives were applied (store credit coupons).
Germany	1 Public	~40,000	5,064	677	13.4%	2016–2017	All first-year students were invited to participate through e-mail. Six reminder emails were sent with personalized links to the survey. Conditional incentives were applied (store credit coupons).
Mexico	4 Private/2 public	~28,000	5,293	4,199	79.3%	2016	All first-year students were eligible for the survey. Initial contact differed by university; survey included in an obligatory health evaluation (one university), as part of obligatory group tutoring sessions (one university), or as part of required classes (two universities) or teacher evaluations (two universities). Two universities sent reminder emails (tutors sent out emails to their tutees; in a required class of personal development, reminders were sent out by faculty). No incentives were applied.
Northern Ireland	1 Public	~25,000	4,359	739	17.0%	2015	All first-year students due to register were invited to participate. Following registration, ID numbers and links to the survey were provided. Five reminder emails/text messages were sent with personalized links to the survey. A sixth reminder involved a researcher telephoning nonresponders. All responders were entered into a number of draws to win an iPad.
South Africa	1 Public	~30,000	5,338	686	12.9%	2015	All first-year students were invited to participate through e-mail. Eight reminder emails and one text message were sent with personalized links to the survey. Conditional incentives were applied (5x R1000 draw).
Spain	5 Public	~96,000	16,332	2,118	13.0%	2014–2015	All first-year students were eligible for the survey. Initial contact differed by university (information stands, information sessions in classrooms, through the university's website). Four reminder emails were sent with personalized links to the survey. Conditional monetary incentives were applied. Additionally, an end-game strategy was implemented by selecting a random proportion of nonrespondents and offering all of them a monetary incentive.
United States	3 Private	~21,800	4,382	739	16.9%	2015–2016	All first-year students were invited to participate through e-mail. Three reminder emails were sent with personalized links to the survey. Conditional incentives were applied (gift cards).
Total	12 Public/7 private	~326,000	58,340	14,371	45.5% ^a	2014–2017	

^aWeighted by achieved sample size.

respondents indicating that they experienced symptoms for 1 month or more in the past year were coded positive for past year disorder. A detailed report of the prevalence of mental disorders in this sample is under review (Auerbach et al., 2018). Students endorsing recent and/or severe suicide thoughts and behaviors received information about available resources within their school and/or community.

2.4.3 | Sociodemographic factors

Questions on sociodemographic characteristics were administered at the beginning of the survey in order to promote the warming up of the respondent. Sociodemographic factors examined in the survey included: gender, age, parental education, parental marital status, urbanicity, religious background, and sexual orientation. Gender was assessed by asking respondents whether they identified themselves as male, female, transgender (male-to-female/female-to-male), or "other."

Respondent age was categorized into three categories (18 years/19 years/20 or more years old). Parental educational level was assessed for father and mother separately (none/elementary school/secondary school/some postsecondary education/university graduate/doctoral degree), and was categorized into high (university graduate or more), medium (some postsecondary education), and low (secondary school or less) based on the highest-of-both parents' educational level. Parental marital status was dichotomized into "parents not married or parent(s) deceased" versus "parents married and both alive." Respondents were asked about the level of urbanicity of the place they were raised (small city/large city/town or village/suburbs/rural area), and their religious background (categorized into Christian/Other religion/No religion). Students were asked whether they identified themselves as heterosexual, gay or lesbian, bisexual, asexual, not sure, and other. *Those indicating "other" were presented an open text item "How would you describe your sexual orientation?" All free text responses were inspected in the data cleaning process, and, if applicable, recoded to one of the five other response options.* Additional questions were asked about the extent to which respondents reported being attracted to men and women and the gender(s) of people they had sex with (if any) in the past 5 years. To create the final sexual orientation variables, respondents were categorized into the following categories: heterosexual with no same-sex attraction, heterosexual with same-sex attraction, nonheterosexual without same-sex sexual intercourse, and nonheterosexual with same-sex sexual intercourse.

2.4.4 | College-related factors

The college-related factors examined in the survey included: high school performance, the most important reason to go to university, and residence during first semester. Respondents were asked where they ranked academically compared to other students at the time of their high school graduation (from top 5% to bottom 10%; categorized into quartiles). They were also asked, out of nine possible reasons, what their most important reason was to go to university. Based on the results of a tetrachoric factor analysis (details available on request), the most important reason to go to university was categorized into extrinsic reasons (i.e., family wanted me to/my friends were going/teachers advised me to/did not want to get a job right away)

versus intrinsic reasons (to achieve a degree/I enjoy learning and studying/to study a subject that really interests me/to improve job prospects generally/to train for specific type of job). Finally, respondents were also asked where they were living during the first semester of the academic year (parents', other relative's, or own home/university or college hall of residence/shared house, apartment, or flat/private hall of residence/other), and if they already worked or expected to work on a student job.

2.5 | Analysis

All analyses were conducted with SAS version 9.4 (Inc., 2010). In addition, multiple imputation (MI) by chained equations (Van Buuren, 2012) was used to adjust for within-survey item nonresponse, random internal subsampling of survey sections, and missing data due to skip logic errors that occurred in a few surveys. Cross-tabulations were used to estimate the prevalence of severe role impairment among those with 12-month mental disorder, and are reported as weighted within-country proportions, with associated MI-adjusted standard errors obtained through the Taylor series linearization method. To obtain pooled estimates of prevalence across countries, each country was given an equal sum of weights.

Logistic regression analyses were used to identify 12-month mental disorders as predictors for any severe role impairment, and in each separate role domain of severe role impairment. Regression coefficients and their MI-based standard errors were exponentiated to generate odds ratios (OR) and associated 95% confidence intervals (CI). A series of five models was evaluated: (M1) type only, i.e., model including a separate dummy variables for each of the six types of mental disorder; (M2) continuous number only, i.e., one variable indicating number of disorders ranging from zero to six; (M3) categorical number of disorders only, i.e., a series of dummy variables indicating exactly one, exactly two, and three or more disorders; (M4) both type and continuous number of comorbid disorders and (M5) both type and categorical number, excluding the dummy indicating exactly one mental disorder. This series of models was conducted to evaluate whether it is the joint effect of specific types of disorder and/or the interactions between disorders that explains the data best (Kessler et al., 2010; Nock, Borges, & Ono, 2012). Model 1 implicitly assumed that the joint effect of the multiple disorders is the product of the odds ratios. The assumptions in Models 2 and 3 were that the disorder is unimportant once number of disorders is known, and the main effect coefficients for the various component disorders are assumed to be the same. Additionally, Model 3, with dummy variables for number of predictor disorders, implicitly allows for interactions in the sense that the coefficients associated with having exactly two or exactly three or exactly n disorders can be significantly different from the product of the coefficient associated with having exactly one disorder. The model that includes terms for both type and number of predictor disorders, in comparison, allows both for differences in effects of the different disorders and for interactions. Specifically, the coefficients associated with types of disorders can be interpreted as the coefficients of the outcome among individuals who had one and only one specific mental disorder compared to respondents with none of the disorders, while the

coefficients associated with number of disorders are multipliers of the combination of coefficients associated with the component disorders. All models were adjusted for sociodemographic and college-related variables, and country membership. Akaike's information criterion was used to select the best model. Based on this model, population attributable risk proportions (PARP) were calculated to estimate the potential reduction in severe role impairment if we were able to eliminate the six 12-month mental disorders under study from the population, assuming a full causal relationship between disorder and role impairment (Krynska & Martin, 2009). We then examined between-country variation in associations by including predictor-by-country interactions and an adjusted interaction dummy coding scheme that kept the product of all country-specific ORs equal to one. The latter method allowed us to detect significant between-country variation by evaluating the statistical significance of deviation of within-country coefficients from the median 1.0 value. Statistical significance in all analyses was evaluated using two-sided MI-based tests with significance level α set at 0.05.

3 | RESULTS

3.1 | Sample description

The final sample included in this study is of 13,984 students, representing a weighted by achieved sample size mean response rate of 45.5% (Table 1). Mean age of the participants was 19.33 ($SD = 0.59$), and 54.4% were female. The characteristics of the sample are presented in the Online Data Supplement Table 1.

3.2 | Prevalence of mental disorders and severe role impairment

As shown in Table 2 (first column), 12-month prevalence of any mental disorder was 31.4%, and ranged from 3% for drug use disorder to 18.5% for major depression. A detail account of the prevalence of mental disorders in this sample has been reported elsewhere (Auerbach et al., 2018). A 20.4% of the sample reported any severe role impairment (Table 2, 3rd column), 10% for those with no mental disorder and 42.9% of those with any mental disorder ($P < 0.001$). Students with panic disorder and broad mania showed the highest proportion with severe impairment (60.6 and 57.5%, respectively). Results suggest a dose-dependent effect whereby the higher the number of disorders, the greater the impairment (almost 69% among students reporting three or more mental disorders). The highest level of impairment was observed for social life and close relationships.

The frequency of any severe role impairment across the different participating sites is summarized in Table 3. As expected, severe role impairment is much higher among students with than without mental disorders across all sites. However, there is considerable variation in the prevalence of any severe impairment, from the lowest in Mexico and Belgium (8.9 and 10.6%, respectively) to the highest in Australia and Germany (38.7 and 27.4%, respectively). These site differences in prevalence of severe impairment persist among students with any and those with no mental disorder.

TABLE 2 Severe role impairment by 12-month mental disorder in the WMH-ICS surveys

	% (SE)	Any severe role impairment % (SE)	Home management % (SE)	College-related work % (SE)	Close relationships % (SE)	Social life % (SE)	Exactly 1 severe role impairment % (SE)	Exactly 2 severe role impairments % (SE)	3 or more severe role impairments % (SE)
Full sample (n = 13,984)	100 (0.0)	20.4 (0.6)	6.6 (0.4)	9.6 (0.4)	11.1 (0.4)	12.1 (0.5)	8.9 (0.4)	5.8 (0.3)	5.7 (0.3)
No mental disorder	68.6 (0.6)	10.0 (0.5)	3.3 (0.3)	4.1 (0.3)	4.8 (0.4)	5.4 (0.4)	5.4 (0.4)	2.4 (0.2)	2.2 (0.3)
Any mental disorder	31.4 (0.6)	42.9 (1.2)	13.7 (0.9)	21.7 (1.0)	24.8 (1.0)	26.8 (1.1)	16.4 (0.9)	13.0 (0.8)	13.5 (0.9)
Type of disorder									
Major depressive episode	18.5 (0.5)	51.2 (1.5)	16.7 (1.3)	27.4 (1.4)	30.9 (1.5)	33.3 (1.5)	17.7 (1.2)	15.6 (1.1)	17.9 (1.3)
Generalized anxiety disorder	16.7 (0.5)	53.1 (1.6)	18.1 (1.4)	27.8 (1.5)	30.6 (1.5)	34.8 (1.6)	19.1 (1.4)	15.5 (1.2)	18.5 (1.3)
Panic disorder	4.5 (0.3)	60.6 (3.4)	24.4 (3.0)	35.2 (3.3)	32.4 (3.3)	37.6 (3.4)	21.6 (3.1)	16.9 (2.6)	22.1 (3.0)
Broad mania	3.1 (0.2)	57.5 (3.8)	24.5 (3.7)	32.9 (3.6)	35.7 (3.7)	35.3 (3.7)	19.9 (3.4)	14.3 (2.7)	23.3 (3.4)
Alcohol abuse or dependence	6.3 (0.3)	34.2 (2.8)	13.0 (2.1)	15.7 (2.1)	19.4 (2.3)	21.4 (2.4)	13.8 (2.1)	9.5 (1.7)	10.9 (1.9)
Drug abuse or dependence	3.0 (0.2)	48.4 (4.2)	22.3 (3.8)	25.8 (3.7)	31.5 (3.9)	23.9 (3.6)	20.8 (3.7)	8.7 (2.5)	18.9 (3.3)
Number of disorders									
Exactly one mental disorder	17.2 (0.5)	30.2 (1.5)	7.6 (0.9)	13.7 (1.1)	16.4 (1.2)	17.5 (1.3)	13.1 (1.2)	10.6 (1.0)	6.5 (0.8)
Exactly two mental disorders	9.4 (0.4)	52.9 (2.2)	16.7 (1.8)	26.9 (2.0)	32.0 (2.1)	35.0 (2.1)	18.5 (1.8)	16.2 (1.7)	18.2 (1.8)
Three or more mental disorders	4.8 (0.3)	68.8 (3.0)	29.4 (3.2)	40.1 (3.3)	40.7 (3.2)	43.8 (3.3)	24.3 (3.1)	15.5 (2.4)	29.0 (3.0)

Note. To obtain pooled estimates of prevalence, each country was given an equal sum of weights. SE, standard error.

TABLE 3 Any severe role impairment by 12-month mental disorder in the WMH-ICS surveys, by country

	Australia	Belgium	Germany	Mexico	Northern Ireland	South Africa	Spain	United States	F(ndf, ddf) [P-value] ^a
	% (SE)	% (SE)	% (SE)	% (SE)	% (SE)	% (SE)	% (SE)	% (SE)	
Full sample	38.7 (2.3)	10.6 (0.5)	27.4 (1.9)	8.9 (0.5)	22.2 (1.6)	20.9 (1.8)	17.3 (1.3)	17.0 (1.5)	65.24(7,88432) [<0.01]*
No mental disorder	16.9 (2.4)	5.6 (0.4)	15.1 (2.1)	5.4 (0.4)	9.8 (1.4)	11.2 (1.8)	9.7 (1.3)	9.3 (1.4)	14.35(7,5920) [<0.01]*
Any mental disorder	67.2 (3.4)	31.9 (1.6)	48.9 (3.3)	20.0 (1.5)	43.4 (3.1)	41.3 (3.8)	32.6 (2.7)	37.8 (3.5)	28.09(7,179424) [<0.01]*
Type of disorder									
Major depressive episode	75.4 (4.0)	41.6 (2.2)	51.6 (3.9)	27.5 (2.5)	52.0 (4.0)	48.6 (5.0)	38.8 (3.5)	50.6 (5.2)	14.51(7,971377) [<0.01]*
Generalized anxiety disorder	78.1 (3.7)	43.8 (2.7)	58.7 (4.9)	29.0 (2.5)	52.5 (4.0)	47.6 (5.1)	39.9 (3.9)	43.7 (4.9)	15.25(7,1274302) [<0.01]*
Panic disorder	75.1 (6.2)	37.9 (5.0)	65.5 (8.9)	28.6 (5.8)	62.5 (10.1)	57.7 (12.9)	71.4 (7.6)	46.0 (9.6)	6.25(7,281) [<0.01]*
Broad mania	79.5 (7.5)	59.1 (7.2)	55.2 (11.9)	27.5 (3.7)	49.3 (12.0)	44.6 (15.4)	57.8 (7.4)	62.1 (13.8)	6.14(7,471872) [<0.01]*
Alcohol abuse or dependence	69.5 (8.1)	17.7 (3.0)	48.9 (10.6)	15.2 (2.3)	33.4 (6.0)	34.6 (7.9)	26.8 (7.0)	12.9 (6.8)	7.96(7,207551) [<0.01]*
Drug abuse or dependence	68.8 (9.6)	26.8 (9.6)	66.3 (10.5)	27.5 (5.7)	55.8 (15.2)	43.9 (10.1)	37.4 (10.1)	21.6 (8.7)	3.33(7,1465) [<0.01]*
Number of disorders									
Exactly one mental disorder	44.6 (5.6)	22.3 (1.9)	38.0 (4.3)	12.3 (1.5)	29.9 (4.3)	32.7 (5.1)	22.5 (3.2)	33.6 (4.5)	11.20(7,10032) [<0.01]*
Exactly two mental disorders	81.6 (4.7)	44.1 (3.5)	62.2 (6.2)	29.4 (3.9)	51.4 (6.1)	47.5 (7.1)	41.3 (4.6)	38.0 (6.5)	8.28(7,660) [<0.01]*
Three or more mental disorders	83.8 (5.1)	70.5 (6.1)	68.6 (8.4)	41.0 (6.0)	67.6 (8.0)	56.6 (10.4)	61.6 (9.9)	69.0 (10.9)	3.15(7,256) [<0.01]*

^aF-test to evaluate significant between-country difference in prevalence estimates. ndf, numerator degrees of freedom; ddf, denominator degrees of freedom.

3.3 | Mental disorders as predictors of severe role impairment

The results of the logistic regression models predicting any severe impairment adjusted by all our sociodemographic and college-related factors, as well as by country, are presented in Table 4. Model 1 shows that all disorders were significantly associated with the reporting of severe role impairment, major depression and GAD being those disorders with the highest odds of severe impairment (OR = 3.2 and OR = 3.0, respectively).

Models 2 and 3 include only the number of disorders, either considered as a continuous count (Model 2) or a 3-level categorical count (Model 3). Model 2 shows a significant OR = 2.5 of severe role impairment for each additional mental disorder, and Model 3 shows different effect for each of the categories considered (exactly one, exactly two, and three or more disorders), ORs ranging from 3.3 to 14.9. The joint effects corresponding to more than one disorder according to this model are subadditive.

Models 4 and 5 assess the association of 12-month mental disorders with any severe role impairment taking into account the number of disorders using a continuous (Model 4) and categorical (Model 5) approach. Both Models 4 and 5 show that estimates for individual disorders are even higher, when the disorder count is included, than they were in Model 1. In contrast with Models 2 and 3, once the presence a specific mental disorders is considered, additional disorders are associated with a subadditive effect (with OR significantly smaller than 1). Model AUCs were similar, ranging from 0.781 (Model 2) to 0.785 (Models 1 and 5) suggesting similar model performance. Model 5 was selected as the best fitting model.

The association of 12-month mental disorders with specific types of severe impairment when using a final logistic regression (Model 5 in Table 3) is presented in Table 5. All mental disorders are significantly associated with severe impairment in all four domains (with only two exceptions among 24 estimates). Major depressive episode is associated with the highest impairment (with ORs ranging from 4.4 for close relationships, to 2.2 for home management), closely followed by GAD (ORs ranging from 3.7 to 2.3). Social life and in close relationships were the domains consistently with the highest association with 12-month mental disorders, and home management, the domain with lowest but still high and significant ORs.

Table 5 also provides the population attributable risk proportion (PARP), estimated for severe impairment in each role domain. This can be interpreted as the theoretical proportion of impairment that could be avoided if there was no 12-month disorder. Overall, almost half (45.2%) of any severe role impairment (2nd column) could be avoided, with PARPs over 50% for social life, close relationship, and work impairment. AUCs indicated a similarly good performance for all models, with a range from 0.757 (work) to 0.803 (close relationships).

The association of 12-month mental disorders with any severe role impairment by country is presented in Table 6. For each row in the table, a separate model was built to determine the country-specific effects. All the models (and all the ORs shown) are also adjusted for sociodemographic and college-related variables, country membership, and variables-by-country interaction dummies. The second column

TABLE 4 Twelve-month mental disorders as predictors for any severe role impairment in the WMH-ICS surveys

	Model 1 aOR (95%CI)	Model 2 aOR (95%CI)	Model 3 aOR (95%CI)	Model 4 aOR (95%CI)	Model 5 aOR (95%CI)
Type of mental disorder					
Major depressive episode	3.2 (2.8–3.8)*			4.1 (3.4–4.9)*	4.0 (3.3–4.8)*
Generalized anxiety disorder	3.0 (2.6–3.6)*			4.1 (3.3–5.0)*	3.9 (3.1–4.8)*
Panic disorder	2.1 (1.5–2.9)*			2.8 (2.0–4.0)*	2.9 (2.1–4.2)*
Broad mania	1.9 (1.4–2.6)*			2.6 (1.9–3.7)*	2.7 (1.9–3.8)*
Alcohol abuse or dependence	1.4 (1.1–1.7)*			1.7 (1.3–2.2)*	1.7 (1.3–2.2)*
Drug abuse or dependence	1.6 (1.1–2.2)*			2.1 (1.5–3.1)*	2.1 (1.4–3.0)*
Number of disorders (continuous)		2.5 (2.4–2.7)*		0.8 (0.7–0.9)*	
Number of disorders (categorical)					
None			(ref)		(ref)
Exactly one mental disorder			3.3 (2.8–3.8)*		
Exactly two mental disorders			8.2 (6.9–9.8)*		0.7 (0.5–0.9)*
Three or more mental disorders			14.9 (11.4–19.5)*		0.4 (0.2–0.6)*
<i>F</i> (ndf,ddf) [<i>P</i> -value] ^a			272.76(3,2615) [<i><</i> 0.01]*		7.73(2,1992) [<i><</i> 0.01]*
Model fit					
Akaike information criterion	9416.8	9498.2	9474.4	9399.9	9398.2
Area under the curve	0.784	0.781	0.782	0.785	0.785

Note. Model 1 includes all the type of mental disorders; Model 2 includes the number of mental disorders (as a continuous variable); Model 3 Includes the number of mental disorders (as categorical variables); Model 4 includes all the types of mental disorders plus he number of disorders (continuous); and Model 5 includes all the types of mental disorders plus the number of disorders (categorical). All models are adjusted for sociodemographic (gender, age, parental educational level, parental marital status, place raised, religion, sexual orientation, and current living situation), college-related predictors (expected to work on a student job, academic performance in high school, most important reason to go to university), the predictors shown in the rows, and country membership. Significant findings are indicated in bold and marked with an asterisk *; α , 0.05.

aOR, adjusted odds ratio; CI, confidence interval.

^a*F*-test to evaluate joint significance of categorical predictor levels. *ndf*, numerator degrees of freedom; *ddf*, denominator degrees of freedom.

TABLE 5 Twelve-month mental disorders as predictors for types of severe role impairment in the WMH-ICS surveys, final model

	Any severe role impairment aOR (95%CI)	Home management aOR (95%CI)	College-related work aOR (95%CI)	Close relationships aOR (95%CI)	Social life aOR (95%CI)
Type of disorder					
Major depressive episode	4.0 (3.3–4.8)*	2.2 (1.6–3.0)*	3.6 (2.8–4.6)*	4.4 (3.4–5.5)*	4.3 (3.4–5.4)*
Generalized anxiety disorder	3.9 (3.1–4.8)*	2.3 (1.6–3.2)*	3.5 (2.7–4.7)*	3.1 (2.3–4.2)*	3.7 (2.9–4.8)*
Panic disorder	2.9 (2.1–4.2)*	1.9 (1.1–3.3)*	3.0 (2.0–4.4)*	2.4 (1.6–3.6)*	2.5 (1.7–3.7)*
Broad mania	2.7 (1.9–3.8)*	1.9 (1.1–3.2)*	2.3 (1.6–3.4)*	2.4 (1.6–3.6)*	2.4 (1.6–3.5)*
Alcohol abuse or dependence	1.7 (1.3–2.2)*	1.5 (1.0–2.3)*	1.4 (1.0–1.9)	1.6 (1.1–2.2)*	1.7 (1.3–2.4)*
Drug abuse or dependence	2.1 (1.4–3.0)*	1.7 (1.0–3.0)*	2.3 (1.5–3.5)*	2.4 (1.6–3.6)*	1.5 (1.0–2.3)
Number of disorders					
None or exactly one disorder					
Exactly two mental disorders	0.7 (0.5–0.9)*	1.0 (0.6–1.6)	0.7 (0.5–1.0)	0.7 (0.5–1.0)	0.6 (0.4–0.9)*
Three or more mental disorders	0.4 (0.2–0.6)*	0.8 (0.3–2.0)	0.4 (0.2–0.7)*	0.3 (0.2–0.6)*	0.3 (0.2–0.6)*
<i>F</i> (ndf,ddf) [<i>P</i> -value] ^a	7.73(2,1992) [<i><</i> 0.01]*	0.21(2,7336) [0.81]	5.31(2,946) [<i><</i> 0.01]*	6.14(2,690) [<i><</i> 0.01]*	6.95(2,751) [<i><</i> 0.01]*
PARP % (SE)	45.2 (1.8)	40.7 (3.2)	50.4 (2.2)	50.7 (1.9)	50.7 (1.6)
Area under the curve	0.785	0.757	0.775	0.803	0.798

Note. All models are adjusted for sociodemographic (gender, age, parental educational level, parental marital status, place raised, religion, sexual orientation, and current living situation), college-related predictors (expected to work on a student job, academic performance in high school, most important reason to go to university), the predictors shown in the rows, and country membership. Significant findings are indicated in bold and marked with an asterisk *; α , 0.05. aOR, adjusted odds ratio; CI, confidence interval.

^a*F*-test to evaluate joint significance of categorical predictor levels. *ndf*, numerator degrees of freedom; *ddf*, denominator degrees of freedom.

shows the overall effects for the pooled sample, while each country column represents deviation for this particular country sample, once adjusted by all sociodemographic and college-related factors. The vast majority of country-specific estimates were nonsignificantly different from one (i.e., no country effect); however, significant differences were found for Belgium (MDE, GAD, broad mania, and for three or more disorders), Australia (GAD and for exactly two disorders), and Spain (panic disorder), the odds ratios of country differences ranging from 1.4 to 2.4.

Analyses of the association of sociodemographic and college-related variables with impairment, adjusting for 12-month mental disorders are presented in Online Data Supplement Table 2. Results show that age, religion, sexual orientation, and country are associated with the likelihood of reporting any severe impairment adjusting by mental disorders. As shown in Online Data Supplement Table 2, adjusting by a large number of variables, including mental disorders, a significant association with the likelihood of reporting severe role impairment was found for country. Spain and Australia students consistently showed a lower likelihood while Northern Ireland in particular, but also South Africa and Germany showed a higher likelihood of severe role impairment, adjusting by mental disorders and other sociodemographic variables.

4 | DISCUSSION

To the best of our knowledge, this is the first study to report on the association between mental disorders and severe role impairment in first-year university students in a large, cross-national sample. Results indicate that severe role impairment is common among students with mental disorders and it encompasses broad domains of life, including social life, close personal relationships and work domains. Notably, PARP analyses indicated that if mental disorders were eliminated, nearly half of the reported severe role impairment in this population would be avoided, assuming that the observed association is causal (Krynska & Martin, 2009). When adjusting for sociodemographic and college-related factors as well as country membership, major depressive episode and GAD are most strongly associated with role impairment. Additionally, while comorbidity is linearly related to severe role impairment, after controlling for individual existing disorders, additional disorders show a subadditive association with impairment. Finally, we found that while there are cross-national differences in the frequency of severe role impairment, the association of mental disorders on severe role impairment is relatively similar across all the countries studied.

The finding that social life, close relationships, and college-related and other work domains are very frequently impaired among students with mental disorders is consistent with results of a previous study carried out in French universities ($n = 1,025$) (Verger et al., 2010). That study indicated comparable rates of impairment in the context of mental disorders. Our results are also in line with studies of adults in the general population, where individuals with mental disorders reported severe role impairment more frequently than those with physical conditions, due to a higher association with social and personal relationships (Ormel et al., 2008). The high frequency of impairment among

TABLE 6 Twelve-month mental disorders as predictors for any severe role impairment in the WMH-ICS surveys, country effect vs. overall effect

Type of disorder	Overall Effect aOR (95%CI)	Australia aOR (95%CI)	Belgium aOR (95%CI)	Germany aOR (95%CI)	Mexico aOR (95%CI)	Northern Ireland aOR (95%CI)	South Africa aOR (95%CI)	Spain aOR (95%CI)	United States aOR (95%CI)
Major depressive episode	3.8 (3.2-4.6)*	1.2 (0.8-1.9)	1.5 (1.2-2.0)*	0.8 (0.5-1.1)	0.8 (0.6-1.0)	1.0 (0.7-1.5)	0.8 (0.5-1.2)	0.8 (0.6-1.1)	1.4 (0.9-2.1)
Generalized anxiety disorder	3.9 (3.2-4.8)*	1.8 (1.1-2.8)*	1.4 (1.0-1.8)*	0.8 (0.5-1.3)	0.8 (0.6-1.1)	1.0 (0.7-1.4)	0.9 (0.5-1.4)	0.8 (0.6-1.1)	0.9 (0.6-1.4)
Panic disorder	3.1 (2.2-4.5)*	0.9 (0.4-1.7)	0.8 (0.5-1.4)	1.1 (0.5-2.6)	0.7 (0.4-1.3)	1.0 (0.4-2.6)	0.9 (0.3-3.0)	2.4 (1.2-4.8)*	0.8 (0.3-1.8)
Broad mania	2.6 (1.8-3.9)*	1.2 (0.5-2.8)	2.4 (1.2-4.6)*	0.8 (0.3-1.9)	0.7 (0.4-1.2)	0.5 (0.2-1.3)	0.4 (0.1-1.7)	1.4 (0.8-2.7)	2.3 (0.7-7.3)
Alcohol abuse or dependence	1.7 (1.3-2.3)*	1.5 (0.7-3.2)	1.0 (0.6-1.5)	1.7 (0.8-3.5)	0.9 (0.6-1.4)	1.0 (0.6-1.8)	1.1 (0.5-2.2)	0.9 (0.6-1.5)	0.4 (0.1-1.2)
Drug abuse or dependence	2.1 (1.4-3.0)*	1.1 (0.5-2.5)	1.1 (0.4-2.8)	1.7 (0.7-4.0)	1.1 (0.6-2.1)	0.9 (0.3-2.8)	1.0 (0.4-2.4)	0.8 (0.4-1.8)	0.6 (0.2-1.5)
Number of disorders									
None or exactly one disorder	(ref)	(ref)	(ref)	(ref)	(ref)	(ref)	(ref)	(ref)	(ref)
Exactly two mental disorders	0.7 (0.5-0.9)*	2.2 (1.2-3.9)*	1.2 (0.9-1.7)	0.9 (0.6-1.6)	1.0 (0.7-1.4)	0.9 (0.6-1.6)	0.8 (0.5-1.5)	0.8 (0.6-1.1)	0.6 (0.4-1.1)
Three or more mental disorders	0.3 (0.2-0.6)*	1.0 (0.5-2.0)	1.9 (1.0-3.4)*	0.9 (0.4-1.8)	0.7 (0.4-1.3)	1.0 (0.4-2.1)	0.7 (0.3-1.6)	1.2 (0.7-2.0)	1.1 (0.4-3.0)

Note: For each row in the table, a separate model was built to determine the country-specific effects. All the models (and all the ORs shown) are also adjusted for all other predictor variables, sociodemographic and college-related predictors, country membership, and predictor-by-country interaction dummies. The second column shows the overall adjusted predictor variable effect; the country columns show to what extent the country-specific adjusted predictor variable effect deviates from the overall adjusted predictor variable effect. For example, the country-specific effect for "Generalized anxiety disorder" in Australia can be obtained by multiplying aOR = 3.9 (the overall effect) by aOR = 1.8 (the country-specific deviation), i.e., aOR = 7.0. Significant findings are indicated in bold and marked with an asterisk*; α , 0.05. aOR, adjusted odds ratio; CI, confidence interval; SE, standard error.

college students with mental disorders suggests that screening for impairment may be critical to substantively improve health and associated academic functioning. Since role impairment mediates the impact of mental disorders on perceived health outcomes (Alonso et al., 2013), it is quite likely that diminishing or avoiding impairment on social relationships may improve students' health perceptions. It is nevertheless important to bear in mind that the data analyzed here are cross-sectional, precluding an interpretation of the directionality of the association between mental disorders and role impairment. Severe role impairment may have an effect on the incidence of new mental disorders, their impact and help-seeking behavior.

In our study, major depressive episode and GAD, the two most prevalent conditions, showed the strongest association with role impairment. Similar high prevalence of these two disorders have been previously reported (Auerbach et al., 2016; Farrer, Gulliver, Bennett, Fassnacht, & Griffiths, 2016; Ibrahim et al., 2013; Storrie, Ahern, & Tuckett, 2010; Vazquez et al., 2011). The combination of a high prevalence and high impact makes them a major source of role impairment among first-year university students (Ormel et al., 2008; Verger et al., 2010). Major depression has been associated with a higher degree of attrition of university failure (Auerbach et al., 2016). Screening for these two mental disorders among university students, in particular major depression, may diminish their burden. Online screening and intervention programs have been evaluated and shown to be feasible and effective for reducing symptoms (Andersson & Cuijpers, 2009; Spek et al., 2008) and prevent onset of full mental disorders (Buntrock et al., 2016). The impact of such strategies on academic outcomes should also be assessed.

We observed a monotonic relationship between number of mental disorders and frequency of severe role impairment in the bivariate analyses. However, a subadditive interaction between number of disorders and impairment was found in a more complex model that included terms for both the type and the number of comorbid conditions. That is, in a person with a given mental disorder, a comorbid disorder adds impairment but less so than it would be expected for the latter disorder if it happened alone. This general pattern has been replicated in other studies (Alonso et al., 2011; Bruffaerts et al., 2012). An important theoretical implication is that interventions that successfully treat only one disorder in a highly comorbidity set are unlikely to be successful in fully reducing the impairments associated with these disorders. The entire set of disorders needs to be targeted. The feasibility of the latter is increased by the fact that many comorbid mental disorders share similar symptoms and pathological pathways (34). Recognition of this fact and of the high prevalence of comorbidity among mental disorders has led to the development of transdiagnostic treatment approaches (Norton & Roberge, 2017). It has to be noted, that the subadditive interaction documented here is based on a logistic regression model. In such a model, a subadditive effect means that the joint effect of two conditions is lower than the product rather than the sum of their individual effects. Synergistic effects may still exist. And they might be important for prevention, since they might require innovative interventions beyond those directed to combat one particular disorder (Evans & Frank, 2004; Kessler et al., 2012; Lewinsohn, Rohde, & Seeley, 1995; Rothman, 2002; Scott et al., 2009). A comparative evaluation of

interventions targeted to specific disorders versus those targeted to more general manifestations is therefore necessary.

Differences across international sites were found for both the prevalence of mental disorders (Auerbach et al., 2018) and for the frequency of severe role impairment. But the association of mental disorder with severe role impairment was remarkably similar across sites (as shown in Table 6). This is consistent with reports about role impairment in the adult general population (Alonso et al., 2011), and suggests the generalizability of the impact of mental disorders. Additional analyses adjusting by many variables (Online Supplement Table 2) do suggest that university students in Australia and Spain would show a somewhat lower risk of impairment than average, while Northern Ireland shows a higher risk. No clear hypotheses emerge from our data. In future longitudinal analyses, we should test whether these differences persist. We also will assess the extent to which mental disorders are associated with academic outcomes internationally.

The findings reported here should be interpreted under the consideration of several limitations of our study. First, the response rates were low in several sites. However, these response rates compare favorably to those achieved in other large-scale prospective college student surveys (39–44%) (Eisenberg, Hunt, & Speer, 2013; Paul, Tsypes, Eidlitz, Ernhout, & Whitlock, 2015). While it has been shown that the empirical relationship between response rate and nonresponse bias is weak (Groves, 2006), recent findings warn of potential overestimation of mental disorders when response rates are low (Mortier, Cuijpers, et al., 2018). In addition, convenience samples rather than nationally representative samples of incoming first-year students were included in our study, which may limit the representativeness of the data. Second, we used an adapted version of the SDS. The original scale was targeted for individuals with mental disorders and asked specifically about the impairment caused by such conditions (Klemenc-Ketis et al., 2011). The modified scale used in our study did not differentiate between physical and mental health. This may have led us to attribute impairment related to physical health to the presence of mental disorders. But bias should be small given that impairment of mental disorders as measured by the SDS tends to be higher than that of physical conditions (Ormel et al., 2008). Nonetheless, additional analyses taking into account the presence of physical disorders could help better estimate those effects. Third, the assessment of mental disorders is based on self-report surveys as opposed to in-depth clinical interviews. While the survey uses well-validated screening scales, with good concordance with blinded clinical diagnoses (in the range $AUC = 0.70$ – 0.78 for mental disorders (Kessler et al., 2013) and 0.78 – 0.91 for alcohol use disorder (Reinert & Allen, 2002), this is not a substitute for diagnostic interviews, and replication of results using full standard diagnostic measures is needed. Fourth, we were limited to the use of cross-sectional data, and analyses only adjusted for a limited range of variables. We plan to undergo longitudinal analyses when data are gathered in our study to assess the extent to which we replicate our findings. Finally, monetary incentives were offered in our study. This might have augmented participation and therefore the representativity of the study samples. But, because the research on the use of incentives in web surveys is limited, the possibility of an associated bias cannot be ruled out.

In summary, this large online survey of first-year university students in eight countries has revealed that role impairment associated with mental disorders is very high, affecting social life, close personal relationships, and work domains. Major depressive episode and GAD are the disorders with the highest impact on role impairment. The combination of their high prevalence and substantial functional impairment should call for interventions that diminish their potential impact on academic achievement and future mental health decrements. Implementation of interventions addressed to major depression and GAD, specifically, should be strongly considered. The combination of high prevalence and substantial functional impairment should call for careful consideration of the potential benefit of screening for mental disorders in the university student population. Also, the implementation of interventions addressed to major depression and GAD, specifically, should be evaluated. There is evidence that online interventions are effective for the prevention and treatment of depression and anxiety disorders (Ebert, Cuijpers, Munoz, & Baumeister, 2017; Sander, Rausch, & Baumeister, 2016), but more research is needed to determine the potential of such approaches among university students, for example with regard to the reduction of role impairment and academic functioning. Also interventions targeted to the most frequent impairments associated with these disorders (i.e., social life, close personal relationships, and work) seem necessary.

ACKNOWLEDGMENTS

Funding to support this project was received from the National Institute of Mental Health (NIMH) R56MH109566 (RPA), and the content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health or NIMH; the Belgian Fund for Scientific Research (11N0514N/11N0516N/1114717N) (PM), the King Baudouin Foundation (2014-J2140150-102905) (RB), and Eli Lilly (IIT-H6U-BX-I002) (RB, PM); BARMER, a health care insurance company, for project StudiCare (DDE); ZonMw (Netherlands Organisation for Health Research and Development; grant number 636110005) and the PFGV (PFGV; Protestants Fonds voor de Geestelijke Volksgezondheid) in support of the student survey project (PC); South African Medical Research Council (DJS); Fondo de Investigación Sanitaria, Instituto de Salud Carlos III - FEDER (PI13/00343), ISCIII (Río Hortega, CM14/00125), ISCIII (Sara Borrell, CD12/00440), Ministerio de Sanidad, Servicios Sociales e Igualdad, PNSD (Exp. 2015I015); DIUE Generalitat de Catalunya (2017 SGR 452), FPU (FPU15/05728) (JA); Fondo de Investigación Sanitaria, Instituto de Salud Carlos III - FEDER (PI13/00506) (GV); European Union Regional Development Fund (ERDF) EU Sustainable Competitiveness Programme for Northern Ireland, Northern Ireland Public Health Agency (HSC R&D), and Ulster University (TB); The World Mental Health International College Student project is carried out as part of the WHO World Mental Health (WMH) Survey Initiative. The WMH survey is supported by the National Institute of Mental Health NIMH R01MH070884, the John D. and Catherine T. MacArthur Foundation, the Pfizer Foundation, the US Public Health Service (R13-MH066849, R01-MH069864, and R01-DA016558), the Fogarty International Center (FIRCA R03-TW006481), the Pan American Health

Organization, Eli Lilly and Company, Ortho-McNeil Pharmaceutical, GlaxoSmithKline, and Bristol-Myers Squibb. None of the funders had any role in the design, analysis, interpretation of results, or preparation of this paper (RK).

We thank the staff of the WMH Data Collection and Data Analysis Coordination Centres for assistance with instrumentation, fieldwork, and consultation on data analysis. A complete list of all within-country and cross-national WMH publications can be found at <https://www.hcp.med.harvard.edu/wmh/>.

WHO WMH-ICS Collaborators. *Australia:* Mark Boyes, School of Psychology & Speech Pathology, Curtin University; Glenn Kiekens, School of Psychology & Speech Pathology, Curtin University and RG Adult Psychiatry KU Leuven, Belgium; *Germany:* Harald Baumeister, University of Ulm; Fanny Kaehlke, Matthias Berking, Friedrich-Alexander University Erlangen Nuremberg; *Mexico:* Adrián Abrego Ramírez, Universidad Politécnica de Aguascalientes; Corina Benjet, Guilherme Borges, Instituto Nacional de Psiquiatría Ramón de la Fuente; Anabell Covarrubias Díaz, Universidad La Salle Noroeste; Ma. Socorro Durán, Universidad De La Salle Bajío; Rogaciano González González, Universidad De La Salle Bajío, campus Salamanca; Raúl A. Gutiérrez-García, Universidad De La Salle Bajío, campus Salamanca & Universidad Politécnica de Aguascalientes; Alicia Edith Hermsillo de la Torre, Universidad Autónoma de Aguascalientes; Kalina Isela Martínez Martínez, Universidad Autónoma de Aguascalientes, Departamento de Psicología, Centro Ciencias Sociales y Humanidades; María Elena Medina-Mora, Instituto Nacional de Psiquiatría Ramón de la Fuente; Humberto Mejía Zarazúa, Universidad La Salle Pachuca; Gustavo Pérez Tarango, Universidad De La Salle Bajío; María Alicia Zavala Berbena, Universidad De La Salle Bajío; *Northern Ireland:* Siobhan O'Neill, Psychology Research Institute, Ulster University; Tony Bjournson, School of Biomedical Sciences, Ulster University; *South Africa:* Christine Lochner, Janine Roos and Lian Taljaard, MRC Unit on Risk & Resilience in Mental Disorders, Department of Psychiatry, Stellenbosch University; Wylene Saal, Department of Psychology, Stellenbosch University; Dan Stein, Department of Psychiatry and MRC Unit on Risk & Resilience in Mental Disorders, University of Cape Town; *Spain:* The UNIVERSAL study group also includes Itxaso Alayo, IMIM (Hospital del Mar Medical Research Institute), CIBER en Epidemiología y Salud Pública (CIBERESP), Spain; José Almenara, Cadiz University; Laura Ballester, IMIM (Hospital del Mar Medical Research Institute), CIBER en Epidemiología y Salud Pública (CIBERESP), Spain; Gabriela Barbaglia, Pompeu Fabra University; María Jesús Blasco, Pompeu Fabra University; Pere Castellví, IMIM (Hospital del Mar Medical Research Institute); Ana Isabel Cebrià, Parc Taulí Hospital Universitari; Enrique Echeburúa, Basque Country University; Andrea Gabilondo, Osakidetza-Basque Health Service; Carlos García-Forero, CIBER en Epidemiología y Salud Pública (CIBERESP), Spain, Pompeu Fabra University; Álvaro Iruin, Hospital Universitario Donostia-Osakidetza; Carolina Lagares, Cadiz University; Andrea Miranda-Mendizábal, Pompeu Fabra University; Oleguer Parès-Badell, Pompeu Fabra University; María Teresa Pérez-Vázquez, Miguel Hernández University; José Antonio Piqueras, Miguel Hernández University; Miquel Roca, Illes Balears University; Jesús Rodríguez-Marín, Miguel Hernández University; Margalida Gili, Illes Balears University; Victoria

Soto-Sanz, Miguel Hernández University and Margarida Vives, Illes Balears University.

FUNDING INFORMATION

The funding sources had no role in the design and conduct of the study; collection, management, analysis, interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

CONFLICT OF INTEREST

In the past 3 years, Dr. Kessler received support for his epidemiological studies from Sanofi Aventis, was a consultant for Johnson & Johnson Wellness and Prevention, Sage Pharmaceuticals, Shire, Takeda, and served on an advisory board for the Johnson & Johnson Services Inc. Lake Nona Life Project. Kessler is a co-owner of DataStat, Inc., a market research firm that carries out healthcare research.

Dr. Ebert reports to have received consultancy fees/served in the scientific advisory board from several companies such as Minddistrict, Lantern, Schoen Kliniken and German health insurance companies (BARMER, Techniker Krankenkasse). He is also stakeholder of the Institute for health training online (GET.ON), which aims to implement scientific findings related to digital health interventions into routine care.

ORCID

Jordi Alonso MD, PhD  <http://orcid.org/0000-0001-8627-9636>

Pim Cuijpers PhD  <http://orcid.org/0000-0001-5497-2743>

David D. Ebert PhD  <http://orcid.org/0000-0001-6820-0146>

Ronald C. Kessler PhD  <http://orcid.org/0000-0003-4831-2305>

REFERENCES

- Abel, J. R., & Deitz, R. (2012). Do colleges and universities increase their region's human capital? *Journal of Economic Geography*, 12(3), 667–691.
- Alonso, J., Petukhova, M., Vilagut, G., Chatterji, S., Heeringa, S., Ustun, T. B., ... Kessler, R. C. (2011). Days out of role due to common physical and mental conditions: Results from the WHO World Mental Health surveys. *Molecular Psychiatry*, 16(12), 1234–1246. <https://doi.org/10.1038/mp.2010.101>
- Alonso, J., Vilagut, G., Adroher, N. D., Chatterji, S., He, Y., Andrade, L. H., ... Kessler, R. C. (2013). Disability mediates the impact of common conditions on perceived health. *Plos One*, 8(6), e65858. <https://doi.org/10.1371/journal.pone.0065858>
- Andersson, G., & Cuijpers, P. (2009). Internet-based and other computerized psychological treatments for adult depression: A meta-analysis. *Cognitive Behavior Therapy*, 38(4), 196–205. <https://doi.org/10.1080/16506070903318960>
- Auerbach, R. P., Alonso, J., Axinn, W. G., Cuijpers, P., Ebert, D. D., Green, J. G., ... Bruffaerts, R. (2016). Mental disorders among college students in the World Health Organization World Mental Health Surveys. *Psychological Medicine*, 46(14), 2955–2970. <https://doi.org/10.1017/S0033291716001665>
- Auerbach, R. P., Mortier, P., Bruffaerts, R., Alonso, J., Benjet, C., & Cuijpers, P. (2018). WHO World Mental Health Surveys International College Student Project: Prevalence and distribution of mental disorders. *Journal of Abnormal Psychology*, In press.
- Babor, T., Higgins-Biddle, J., Saunders, J., & Monteiro, M. (2001). *The Alcohol Use Disorders Identification Test. Guidelines for use in primary care*. Available at <http://apps.who.int/iris/handle/10665/67205>
- Blanco, C., Okuda, M., Wright, C., Hasin, D. S., Grant, B. F., Liu, S. M., & Olfson, M. (2008). Mental health of college students and their non-college-attending peers: Results from the national epidemiologic study on alcohol and related conditions. *Archives of General Psychiatry*, 65(12), 1429–1437. <https://doi.org/10.1001/archpsyc.65.12.1429>
- Bruffaerts, R., Mortier, P., Kiekens, G., Auerbach, R. P., Cuijpers, P., Demyttenaere, K., ... Kessler, R. C. (2018). Mental health problems in college freshmen: Prevalence and academic functioning. *Journal of Affective Disorders*, 225, 97–103. <https://doi.org/10.1016/j.jad.2017.07.044>
- Bruffaerts, R., Vilagut, G., Demyttenaere, K., Alonso, J., Alhazawi, A., Andrade, L. H., ... Kessler, R. C. (2012). Role of common mental and physical disorders in partial disability around the world. *British Journal of Psychiatry*, 200(6), 454–461. <https://doi.org/10.1192/bjp.bp.111.097519>
- Buntrock, C., Ebert, D. D., Lehr, D., Smit, F., Riper, H., Berking, M., & Cuijpers, P. (2016). Effect of a web-based guided self-help intervention for prevention of major depression in adults with sub-threshold depression: A randomized clinical trial. *Journal of the American Medical Association*, 315(17), 1854–1863.
- Cho, S. B., Llana, D. C., Adkins, A. E., Cooke, M., Kendler, K. S., Clark, S. L., & Dick, D. M. (2015). Patterns of substance use across the first year of college and associated risk factors. *Front Psychiatry*, 6, 152. <https://doi.org/10.3389/fpsy.2015.00152>
- Ebert, D. D., Cuijpers, P., Munoz, R. F., & Baumeister, H. (2017). Prevention of mental health disorders using internet- and mobile-based interventions: A narrative review and recommendations for future research. *Front Psychiatry*, 8, 116–131. <https://doi.org/10.3389/fpsy.2017.00116>
- Eisenberg, D., Golberstein, E., & Gollust, S. E. (2007). Help-seeking and access to mental health care in a university student population. *Medical Care*, 45(7), 594–601. <https://doi.org/10.1097/MLR.0b013e31803bb4c1>
- Eisenberg, D., Hunt, J., & Speer, N. (2013). Mental health in American colleges and universities: Variation across student subgroups and across campuses. *Journal of Nervous and Mental Disease*, 201(1), 60–67. <https://doi.org/10.1097/NMD.0b013e31827ab077>
- Evans, A. S., & Frank, S. J. (2004). Adolescent depression and externalizing problems: Testing two models of comorbidity in an inpatient sample. *Adolescence*, 39(153), 1–18.
- Farrer, L. M., Gulliver, A., Bennett, K., Fassnacht, D. B., & Griffiths, K. M. (2016). Demographic and psychosocial predictors of major depression and generalised anxiety disorder in Australian university students. *Bmc Psychiatry [Electronic Resource]*, 16, 241–249. <https://doi.org/10.1186/s12888-016-0961-z>
- Groves, R. M. (2006). Nonresponse rates and nonresponse bias in household surveys. *Public Opinion Quarterly*, 70(5), 646–675.
- Groves, R. M., & Couper, M. P. (1998). *Nonresponse in household interview surveys*. New York: Wiley.
- Ibrahim, A. K., Kelly, S. J., Adams, C. E., & Glazebrook, C. (2013). A systematic review of studies of depression prevalence in university students. *Journal of Psychiatric Research*, 47(3), 391–400. <https://doi.org/10.1016/j.jpsychires.2012.11.015>
- Inc., S. I. (2010). *SAS/STATR software*. Cary, NC: SAS Institute Inc.
- Kendler, K. S., Myers, J., & Dick, D. (2015). The stability and predictors of peer group deviance in university students. *Social Psychiatry and Psychiatric Epidemiology*, 50(9), 1463–1470. <https://doi.org/10.1007/s00127-015-1031-4>

- Kessler, R. C., Avenevoli, S., Costello, J., Green, J. G., Gruber, M. J., McLaughlin, K. A., ... Merikangas, K. R. (2012). Severity of 12-month DSM-IV disorders in the national comorbidity survey replication adolescent supplement. *Archives of General Psychiatry*, 69(4), 381–389. <https://doi.org/10.1001/archgenpsychiatry.2011.1603>
- Kessler, R. C., Foster, C. L., Saunders, W. B., & Stang, P. E. (1995). Social consequences of psychiatric disorders, I: Educational attainment. *American Journal of Psychiatry*, 152(7), 1026–1032.
- Kessler, R. C., McLaughlin, K. A., Green, J. G., Gruber, M. J., Sampson, N. A., Zaslavsky, A. M., ... Williams, D. R. (2010). Childhood adversities and adult psychopathology in the WHO World Mental Health Surveys. *British Journal of Psychiatry*, 197, 378–385.
- Kessler, R. C., Santiago, P. N., Colpe, L. J., Dempsey, C. L., First, M. B., Heeringa, S. G., ... Ursano, R. J. (2013). Clinical reappraisal of the Composite International Diagnostic Interview Screening Scales (CIDI-SC) in the army study to assess risk and resilience in service members (army STARRS). *International Journal of Methods in Psychiatric Research*, 22(4), 303–321.
- Kessler, R. C., & Ustun, T. B. (2004). The World Mental Health (WMH) survey initiative version of the World Health Organization (WHO) Composite International Diagnostic Interview (CIDI). *International Journal of Methods in Psychiatric Research*, 13(2), 93–121.
- Klemenc-Ketis, Z., Kersnik, J., Eder, K., & Colaric, D. (2011). Factors associated with health-related quality of life among university students. *Srpski Arhiv Za Celokupno Lekarstvo*, 139(3–4), 197–202.
- Krysinska, K., & Martin, G. (2009). The struggle to prevent and evaluate: Application of population attributable risk and preventive fraction to suicide prevention research. *Suicide & Life-Threatening Behavior*, 39(5), 548–557. <https://doi.org/10.1521/suli.2009.39.5.548>
- Leon, A. C., Olfson, M., Portera, L., Farber, L., & Sheehan, D. V. (1997). Assessing psychiatric impairment in primary care with the Sheehan Disability Scale. *International Journal of Psychiatry in Medicine*, 27(2), 93–105.
- Lewinsohn, P. M., Rohde, P., & Seeley, J. R. (1995). Adolescent psychopathology: III. The clinical consequences of comorbidity. *Journal of the American Academy of Child and Adolescent Psychiatry*, 34(4), 510–519.
- Mojtabai, R., Stuart, E. A., Hwang, I., Eaton, W. W., Sampson, N., & Kessler, R. C. (2015). Long-term effects of mental disorders on educational attainment in the National Comorbidity Survey ten-year follow-up. *Social Psychiatry and Psychiatric Epidemiology*, 50(10), 1577–1591. <https://doi.org/10.1007/s00127-015-1083-5>
- Mortier, P., Auerbach, R. P., Alonso, J., Bantjes, J., Benjet, C., Cuijpers, P., ... Collaborators, W. W.-I. (2018). Suicidal thoughts and behaviors among first-year college students: Results from the WMH-ICS project. *Journal of the American Academy of Child and Adolescent Psychiatry*, 57(4), 263–273. <https://doi.org/10.1016/j.jaac.2018.01.018>
- Mortier, P., Cuijpers, P., Kiekens, G., Auerbach, R. P., Demyttenaere, K., Green, J. G., ... Bruffaerts, R. (2018). The prevalence of suicidal thoughts and behaviours among college students: A meta-analysis. *Psychological Medicine*, 48(4), 554–565. <https://doi.org/10.1017/S0033291717002215>
- Nock, M. K., Borges, G., & Ono, Y. (2012). *Suicide: Global perspective from the WHO World Mental Health surveys*. New York: Cambridge University Press.
- Norton, P. J., & Roberge, P. (2017). Transdiagnostic therapy. *The Psychiatric Clinics of North America*, 40(4), 675–687. <https://doi.org/10.1016/j.psc.2017.08.003>
- OECD. (2016). Stat education and training. Education at a glance. educational attainment and labor-force status. Educational attainment of 25–64 year-olds. Accessed February 23, 2018.
- Ormel, J., Petukhova, M., Chatterji, S., Aguilar-Gaxiola, S., Alonso, J., Angermeyer, M. C., ... Kessler, R. C. (2008). Disability and treatment of specific mental and physical disorders across the world. *British Journal of Psychiatry*, 192, 368–375.
- Paul, E., Tsypes, A., Eidlitz, L., Ernhout, C., & Whitlock, J. (2015). Frequency and functions of non-suicidal self-injury: Associations with suicidal thoughts and behaviors. *Psychiatry Research*, 225(3), 276–282. <https://doi.org/10.1016/j.psychres.2014.12.026>
- Reinert, D. F., & Allen, J. P. (2002). The Alcohol Use Disorders Identification Test (AUDIT): A review of recent research. *Alcoholism, Clinical and Experimental Research*, 26(2), 272–279.
- Rothman, K. J. (2002). *Epidemiology: An introduction*. New York: Oxford University Press.
- Sander, L., Rausch, L., & Baumeister, H. (2016). Effectiveness of internet-based interventions for the prevention of mental disorders: A systematic review and meta-analysis. *JMIR Mental Health*, 3(3), e38. <https://doi.org/10.2196/mental.6061>
- Saunders, J. B., Aasland, O. G., Babor, T. F., de la Fuente, J. R., & Grant, M. (1993). Development of the alcohol use disorders identification test (AUDIT): WHO collaborative project on early detection of persons with harmful alcohol consumption—II. *Addiction*, 88(6), 791–804.
- Scott, K. M., Von Korff, M., Alonso, J., Angermeyer, M. C., Bromet, E., Fayyad, J., ... Williams, D. (2009). Mental-physical co-morbidity and its relationship with disability: Results from the World Mental Health Surveys. *Psychological Medicine*, 39(1), 33–43. <https://doi.org/10.1017/S0033291708003188>
- Skidmore, C. R., Kaufman, E. A., & Crowell, S. E. (2016). Substance use among college students. *Child and Adolescent Psychiatric Clinics of North America*, 25(4), 735–753. <https://doi.org/10.1016/j.chc.2016.06.004>
- Spek, V., Cuijpers, P., Nyklicek, I., Smits, N., Riper, H., Keyzer, J., & Pop, V. (2008). One-year follow-up results of a randomized controlled clinical trial on internet-based cognitive behavioural therapy for subthreshold depression in people over 50 years. *Psychological Medicine*, 38(5), 635–639.
- Storrie, K., Ahern, K., & Tuckett, A. (2010). A systematic review: Students with mental health problems—a growing problem. *International Journal of Nursing Practice*, 16(1), 1–6. <https://doi.org/10.1111/j.1440-172X.2009.01813.x>
- Van Buuren, S. (2012). *Flexible imputation of missing data*. Boca Raton: CRC Press (Taylor & Francis Group).
- Vazquez, F. L., Torres, A., Otero, P., & Diaz, O. (2011). Prevalence, comorbidity, and correlates of DSM-IV axis I mental disorders among female university students. *Journal of Nervous and Mental Disease*, 199(6), 379–383. <https://doi.org/10.1097/NMD.0b013e31821cd29c>
- Verger, P., Guagliardo, V., Gilbert, F., Rouillon, F., & Kovess-Masfetty, V. (2010). Psychiatric disorders in students in six French universities: 12-month prevalence, comorbidity, impairment and help-seeking. *Social Psychiatry and Psychiatric Epidemiology*, 45(2), 189–199.
- Wittchen, H. U., Nelson, C. B., & Lachner, G. (1998). Prevalence of mental disorders and psychosocial impairments in adolescents and young adults. *Psychological Medicine*, 28(1), 109–126.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Alonso J, Mortier P, Auerbach RP, et al. Severe role impairment associated with mental disorders: Results of the WHO World Mental Health Surveys International College Student Project. *Depress Anxiety*. 2018;35:802–814. <https://doi.org/10.1002/da.22778>