

## RESEARCH ARTICLE

## Open Access



# Adenosine relaxation in isolated rat aortic rings and possible roles of smooth muscle $K_v$ channels, $K_{ATP}$ channels and $A_{2a}$ receptors

Aryadi Arsyad<sup>1</sup> and Geoffrey P. Dobson<sup>2\*</sup>

## Abstract

**Background:** An area of ongoing controversy is the role adenosine to regulate vascular tone in conduit vessels that regulate compliance, and the role of nitric oxide (NO), potassium channels and receptor subtypes involved. The aim of our study was to investigate adenosine relaxation in rat thoracic aortic rings, and the effect of inhibitors of NO, prostanoids,  $K_v$ ,  $K_{ATP}$  channels, and  $A_{2a}$  and  $A_{2b}$  receptors.

**Methods:** Aortic rings were freshly harvested from adult male Sprague Dawley rats and equilibrated in an organ bath containing oxygenated, modified Krebs-Henseleit solution, 11 mM glucose, pH 7.4, 37 °C. Isolated rings were pre-contracted sub-maximally with 0.3  $\mu$ M norepinephrine (NE), and the effect of increasing concentrations of adenosine (1 to 1000  $\mu$ M) were examined. The drugs L-NAME, indomethacin, 4-aminopyridine (4-AP), glibenclamide, 5-hydroxydecanoate, ouabain, 8-(3-chlorostyryl) caffeine and PSB-0788 were examined in intact and denuded rings. Rings were tested for viability after each experiment.

**Results:** Adenosine induced a dose-dependent, triphasic relaxation response, and the mechanical removal of the endothelium significantly decreased adenosine relaxation above 10  $\mu$ M. Interestingly, endothelial removal significantly decreased the responsiveness (defined as % relaxation per  $\mu$ M adenosine) by two-thirds between 10 and 100  $\mu$ M, but not in the lower (1–10  $\mu$ M) or higher (>100  $\mu$ M) ranges. In intact rings, L-NAME significantly reduced relaxation, but not indomethacin. Antagonists of voltage-dependent  $K_v$  (4-AP), sarcolemma  $K_{ATP}$  (glibenclamide) and mitochondrial  $K_{ATP}$  channels (5-HD) led to significant reductions in relaxation in both intact and denuded rings, with ouabain having little or no effect. Adenosine-induced relaxation appeared to involve the  $A_{2a}$  receptor, but not the  $A_{2b}$  subtype.

**Conclusions:** It was concluded that adenosine relaxation in NE-precontracted rat aortic rings was triphasic and endothelium-dependent above 10  $\mu$ M, and relaxation involved endothelial nitric oxide (not prostanoids) and a complex interplay between smooth muscle  $A_{2a}$  subtype and voltage-dependent  $K_v$ ,  $SarcK_{ATP}$  and  $MitoK_{ATP}$  channels. The possible in vivo significance of the regulation of arterial compliance to left ventricular function coupling is discussed.

**Keywords:** Rat aorta, Adenosine, Vasodilation, Endothelium, Nitric oxide, Vascular tone

\* Correspondence: geoffrey.dobson@jcu.edu.au

<sup>2</sup>Heart, Trauma and Sepsis Research Laboratory, Australian Institute of Tropical Health and Medicine, College of Medicine and Dentistry, James Cook University, 1 James Cook Drive, Queensland 4811, Australia  
Full list of author information is available at the end of the article

## Background

Adenosine is a ubiquitous endogenous mediator that is activated in response to cellular ischemic/hypoxic/shear stress [1–5]. Adenosine exerts its cellular effects by binding to four major subtypes of the G-protein-coupled receptors;  $A_1$ ,  $A_{2a}$ ,  $A_{2b}$ , and  $A_3$  which activate intracellular survival kinase pathways in a cell- and tissue-specific manner [2, 3, 5, 6]. Through receptor-modulation and downstream signaling pathways adenosine alters coronary and peripheral vascular tone, cardiac function, brain and central nervous system signalling, sleep, the state of natural hibernation, ischemic preconditioning, post-conditioning, inflammation, coagulation, angiogenesis and cell proliferation and remodelling [4–7].

An area of ongoing controversy is the role adenosine to regulate vascular tone in the arterial tree, and the receptor subtypes involved. The subtype  $A_{2a}$  appears to be the predominate receptor in arterial vasodilation in mouse, rat, guinea pig, pigs and humans, however, the  $A_{2b}$  receptor has also been reported to dilate human coronary arteries [8], and possibly rat coronary arteries [6]. In the guinea pig,  $A_{2b}$  appears to predominate in the thoracic aorta to induce relaxation [9] and both  $A_{2a}$  and  $A_{2b}$  in the rat [10–12]. In addition, there is ongoing debate on the relative importance of an intact endothelium to adenosine relaxation in these vessels, and the role of nitric oxide (NO) and interplay between voltage-dependent transmembrane  $Na^+$ ,  $K^+$  and  $Ca^{2+}$  fluxes and signalling pathways. In the thoracic aorta, adenosine relaxation has been reported to be fully dependent [10, 13], partially dependent [14–17] or not dependent on the presence of an intact endothelium [10, 18–20]. Adenosine vasodilation has also been linked to  $A_1$  and  $A_{2a}$  receptor activation of endothelial production of NO and prostanoids [21], hyperpolarising factors [4], and a complex interplay between endothelial and smooth muscle mitochondrial and sarcolemmal  $K_{ATP}$  channels [16, 22, 23], and  $Na^+/K^+$  ATPase activation [4, 24].

The aim of the present study was to investigate adenosine relaxation in intact versus denuded rat thoracic aortic rings, and examine the effect of inhibitors of nitric oxide (NO), prostanoids,  $K_v$  channels,  $K_{ATP}$  channels, and adenosine  $A_{2a}$  and  $A_{2b}$  receptors. The rat thoracic aorta was chosen because of the ongoing debates about the mechanisms of adenosine relaxation, and its *in vivo* significance.

## Methods

### Animals

Male Sprague Dawley rats (300–350 g,  $n = 47$ ) were fed *ad libitum* and housed in a 12-h light/dark cycle. On the day of the experiment rats were anaesthetised with Na-thiopentone (100 mg/kg). Animals were treated in accordance with the Guide for the Care and Use of Laboratory Animals published by the US National Institutes of Health

(NIH Publication No. 85–23, revised 1996). The James Cook University (JCU) Animal Ethics Committee approval number for the present study was A1535. All other chemicals, drugs and inhibitors including adenosine (A9251 > 99 % purity) were purchased from Sigma Aldrich (Castle Hill, NSW).

### Aortic ring preparation and organ bath tension measurements

The thoracic cavity of anesthetized rats was opened and the thoracic aorta was harvested and placed in a modified ice-cold solution of Krebs-Henseleit (118 mM NaCl, 4.7 mM KCl, 1.2 mM  $Na_2PO_4$ , 0.5 mM  $MgCl_2$ , 1.12 mM  $CaCl_2$ , 25 mM  $NaHCO_3$ , 0.03 mM EDTA) pH 7.4 with 11 mM glucose. The aorta was carefully dissected from surrounding fat and connective tissue and cut into short transverse segments. Intact aortic rings were isolated from each rat and used without further processing. In those studies that required removal of the endothelium, intact rings were denuded by gently rubbing the intimal surface of the vessel segment with a smooth metal probe. Successful removal of the endothelium was assessed by testing the aortic ring for a vasodilatory response to 10  $\mu$ M acetylcholine (final concentration).

After preparation, intact or denuded aortic rings (3 to 4 mm long) were equilibrated in a standard 10 ml volume organ bath (Radnoti Glass, ADInstruments, NSW, AUS) containing modified Krebs-Henseleit (see above) and continuously bubbled with 95 %  $O_2$  and 5 %  $CO_2$  at 37 °C for 15 min (zero tension). The rings were vertically mounted on small stainless steel triangles, stirrups and connected to an isometric force transducer (PANLAB, distributed by ADInstruments as MLT 0201/RAD, NSW, AUS) coupled to a computer based data acquisition system (PowerLab, ADInstruments) and data recording software LabChart 7 (ADInstruments Pty Ltd., Castle Hill, Australia).

The ring tension was manually adjusted to 1.5 g and equilibrated for 60 min. A tension of 1.5 g was chosen from the literature for thoracic aortic rings [25, 26] and preliminary studies verified this tension. During equilibration, the solution was changed in 15 min intervals. The aortic rings were then washed with freshly prepared Krebs Henseleit buffer pH 7.4 and the tension was readjusted to 1.5 g tension. Each preparation was sub-maximally contracted using 3  $\mu$ l of 0.1 mM NE (0.3  $\mu$ M final concentration) [27, 28]. Those aortic rings that failed to contract were discarded. Ten microliters of 10 mM acetylcholine (10  $\mu$ M final concentration) was applied to confirm the presence or absence of an intact endothelium in all preparations. Acetylcholine will induce rapid relaxation of precontracted rings if the endothelium is intact and if the endothelium is removed (or denuded) the rings will remain in contracted

state [19]. Aortic rings were considered intact if the relaxation induced by 10  $\mu\text{M}$  ACh was greater than 80 %, and the aortic ring was assumed denuded if relaxation was less than 10 %.

Rings were contracted at least two more times before each experiment until a reproducible contractile response was obtained. Ten to 15 min after this state was achieved the experiment was commenced because preliminary studies showed that the increase in tension and plateau from 0.3  $\mu\text{M}$  of NE was reached at 10 min and remained at this plateau level for over 60 min, the time course of each experiment.

### Adenosine relaxation in intact and denuded rat aortic rings

Adenosine was added into the oxygenated organ bath containing Krebs-Henseleit solution to obtain 1, 5, 10, 50, 100, 500 and 1000  $\mu\text{M}$  adenosine concentrations. The change in tension of pre-contracted intact or denuded rings was measured. The inhibitors used in this study were incubated in organ bath 20–30 min before NE was administered followed by adenosine incremental administration. These included 1) 100  $\mu\text{M}$  N<sup>G</sup>-nitro-L-arginine Methyl Ester (L-NAME) (nitric oxide synthetase inhibitor) and 10  $\mu\text{M}$  indomethacin (cyclooxygenase or prostaglandin inhibitor e.g. prostacyclin). NO and prostacyclin are two major endothelial derived relaxation factors (EDRF), and the inhibitors were only applied in endothelium intact aortic rings, and 2) 1 mM 4-aminopyridine (4-AP) (Non-selective voltage-dependent K<sup>+</sup>-channel blocker of the Kv1 to Kv4 families rather than Kv7 channels) [29–31], 10  $\mu\text{M}$  glibenclamide (Non-selective SarcK<sub>ATP</sub> channel blocker) [32, 33] and 1 mM 5-hydroxydecanoate (5-HD) (Mito-K<sub>ATP</sub> channel blocker) [34], and Na<sup>+</sup>/K<sup>+</sup>-ATPase inhibitor (100  $\mu\text{M}$  ouabain) [24]. These inhibitors were applied to intact endothelium rings in the presence of L-NAME and indomethacin, and without the presence of L-NAME and indomethacin in denuded aortic rings. The adenosine A<sub>2a</sub> receptor inhibitor was 100  $\mu\text{M}$  8-(3-chlorostyryl) caffeine (CSC) [35, 36], and the A<sub>2b</sub> receptor inhibitor was 10  $\mu\text{M}$  8-(4-(4-(4-chlorobenzyl)piperazine-1-sulfonyl)-phenyl)-1-propylxanthine (PSB-0788) [37]. In rat striatal membranes, these antagonists have reported K<sub>i</sub> values of 24 nM for CSC [38] and 0.393 nM for PSB-0788 [37], and the micromolar concentrations used in the present study were based on previous published studies [39–41]. The inhibitors were applied in endothelium intact and denuded aortic rings in an oxygenated medium. At the end of each experiment, the rings were tested for viability (or patency) by being maximally dilated with 100  $\mu\text{M}$  papaverine, and relaxation was expressed as percentage of maximal relaxation to papaverine [24, 42].


### Statistics

Values are expressed as mean  $\pm$  SEM. Eight animals ( $n = 8$ ) were used for each group for seven measurement points using ANOVA analysis, and the number of rats was selected from a priori G-power analysis to achieve a level of 1.0. All data was tested for normality using *Kolmogorov-Smirnov* test. Relaxation responses to adenosine were analysed for homogeneity of variances followed by two-way ANOVA coupled with the *Bonferroni* post-hoc test for individual data point comparisons. The alpha level of significance for all experiments was set at  $p < 0.05$ .

### Results

#### Intact versus denuded aortic rings

In endothelium-intact rat aortic rings, adenosine led to 10, 21, 29, 60 and 81 % relaxation at 10, 50, 100, 500, and 1000  $\mu\text{M}$  adenosine concentrations respectively (Fig. 1). Adenosine relaxation in intact rings occurred in three linear phases (log scale); 0.96 % per  $\mu\text{M}$  from 1 to 10  $\mu\text{M}$  adenosine (Phase 1), 0.2 % per  $\mu\text{M}$  from 10 to 100  $\mu\text{M}$  adenosine (Phase 2), and 0.06 % per  $\mu\text{M}$  from 100 to 1000  $\mu\text{M}$  (Phase 3). After removing the endothelium, relaxation was reduced to 8, 10, 14, 45 and 67 % respectively, and was significant from 100 to 1000  $\mu\text{M}$ . In denuded rings, adenosine relaxation was 0.72 % per  $\mu\text{M}$  from 1 to 10  $\mu\text{M}$  adenosine, 0.07 % per  $\mu\text{M}$  from 10 to 100  $\mu\text{M}$  adenosine (Phase 1), and 0.06 % per  $\mu\text{M}$  in Phase 2 from 100 to 1000  $\mu\text{M}$ . Thus endothelial removal of rat aortic rings decreased the responsiveness (defined as % relaxation per  $\mu\text{M}$ ) to around one-third between 10



**Fig. 1** Concentration response curves to adenosine in intact and denuded isolated rat aortic rings. Relaxation is expressed as percent of maximal relaxation to 100  $\mu\text{M}$  papaverine. Points represent mean  $\pm$  S.E.M of aortic rings from a total of seven animals. \* $P < 0.05$  statistical difference in responses between the intact and denuded rings. Symbols (●) Intact rings (■) Denuded rings

and 100  $\mu\text{M}$ , but not in the lower (Phase 1) or higher (Phase 3) ranges (Fig. 1).

**Intact aortic rings**

**Effect of L-NAME and indomethacin**

Figure 2 shows that L-NAME and indomethacin significantly reduced adenosine relaxation at 50 to 1000  $\mu\text{M}$  adenosine. At 50  $\mu\text{M}$ , relaxation decreased from 26 to 11 % or 42 % ( $11/26 \times 100$ ) of the relaxation of intact controls. Thus at 50  $\mu\text{M}$  adenosine 59 % of relaxation was linked to L-NAME and indomethacin inhibition. At 100, 500 and 1000  $\mu\text{M}$  adenosine concentrations, L-NAME and indomethacin contribution to inhibition were 53, 33 and 19 % (Fig. 2). In addition, experiments with L-NAME alone showed a similar inhibition, indicating that indomethacin had little or no significant inhibition (Fig. 2). However, at 500  $\mu\text{M}$  and 1000  $\mu\text{M}$  adenosine there was a small difference of indomethacin from L-NAME but not significant (Fig. 2).

**Effect of  $K_v$ ,  $\text{SarcK}_{\text{ATP}}$ ,  $\text{MitoK}_{\text{ATP}}$  blockers and ouabain on adenosine relaxation**

The effect of  $K_v$ ,  $\text{SarcK}_{\text{ATP}}$ ,  $\text{MitoK}_{\text{ATP}}$  channels and  $\text{Na}^+/\text{K}^+$ -ATPase on adenosine relaxation in intact aortic rings is shown in Fig. 3a–d. In order to eliminate the effect of NO- and prostacyclin-induced relaxation in intact rings, 100  $\mu\text{M}$  L-NAME and 10  $\mu\text{M}$  indomethacin were included in the controls.

**$K_v$  inhibition**

Pre-incubating intact rings with 1 mM 4-aminopyridine (4-AP) on adenosine relaxation is shown in Fig. 3a. Percentage relaxation was 2.4, 5.1, 8.0, 32.1 and 52.5 % for 10, 50, 100, 500 and 1000  $\mu\text{M}$  adenosine, respectively. Expressed as a percentage contribution of adenosine relaxation relative to control intact rings, the  $K_v$  channel

was responsible for 78, 73, 72, 58.2 and 28 % for 10, 50, 100, 500 and 1000  $\mu\text{M}$  adenosine respectively, with greater between 10 to 100  $\mu\text{M}$  (Fig. 3a).

**$\text{SarcK}_{\text{ATP}}$  and  $\text{MitoK}_{\text{ATP}}$  inhibition**

The effect of glibenclamide on adenosine relaxation is shown in Fig. 3b. Glibenclamide was not as striking as 4-AP but significantly decreased adenosine relaxation at 50 and 100  $\mu\text{M}$  adenosine. The contribution of  $\text{SarcK}_{\text{ATP}}$  channel to adenosine relaxation was 63, 53 and 38 % at 10, 50 and 100  $\mu\text{M}$  adenosine (Fig. 3b).  $\text{MitoK}_{\text{ATP}}$  inhibitor, 5-hydroxydecanoate (5-HD), significantly led to a wider range of inhibition of adenosine relaxation compared to glibenclamide from 10 to 1000  $\mu\text{M}$ , but the differences between the two blockers were not significant (Fig. 3c). The contribution of  $\text{MitoK}_{\text{ATP}}$  channel to adenosine relaxation was 70, 63, 65, 40 and 27 % at 10, 50, 100, 500 and 1000  $\mu\text{M}$  adenosine level (Fig. 3c).

**$\text{Na}^+/\text{K}^+$ -ATPase inhibition**

Figure 3d showed that ouabain did not significantly change the inhibition produced by L-NAME and indomethacin in adenosine-induced relaxation at any given concentration, indicating that  $\text{Na}^+/\text{K}^+$ -ATPase contributed little extra to adenosine relaxation in endothelium intact aortic rings.


**Effect of  $A_{2a}$  and  $A_{2b}$  blockers in intact and denuded aortic rings**

**Intact rings**


L-NAME and indomethacin were not included in this experiment because it has been reported that NO or prostacyclin release are linked to adenosine  $A_{2a,b}$  receptor activation [43]. In the absence of any inhibitors, adenosine induced a rate of relaxation of about 10 % for every 50  $\mu\text{M}$  adenosine up to 100  $\mu\text{M}$ , and ~25 % relaxation per 50  $\mu\text{M}$  from 100 to 1000  $\mu\text{M}$  until 90 % full relaxation (Fig. 4a). Pre-incubating intact rings with adenosine  $A_{2a}$  receptor inhibitor, CSC, significantly reduced adenosine relaxation between 50 to 100  $\mu\text{M}$  (Fig. 4a). Although greater percentage falls in relaxation occurred at lower adenosine levels (e.g. 5 to 10  $\mu\text{M}$ ) these were not significantly different from controls (Fig. 4a). The  $A_{2a}$  receptor was responsible for 71, 66, 59 and 47 % adenosine relaxation at 5, 10, 50, and 100  $\mu\text{M}$  adenosine, respectively. In direct contrast, adenosine  $A_{2b}$  receptor inhibitor, PSB 0788, did not change relaxation at any adenosine concentration studied (Fig. 4b).

**Denuded rings**

In denuded rat aortic rings, incubation with  $A_{2a}$  blocker, CSC, showed a significant reduction of adenosine relaxation from 5 to 100  $\mu\text{M}$  (Fig. 4b). At 5, 10, 50 and 100  $\mu\text{M}$  adenosine, the  $A_{2a}$  receptor was responsible for



**Fig. 2** Concentration-response curves to adenosine with and without the presence of L-NAME alone and L-NAME + indomethacin in intact isolated rat aortic rings. Relaxation is expressed as percent of maximal relaxation to 100  $\mu\text{M}$  papaverine. Points represent mean  $\pm$  S.E.M of aortic rings from a total of eight animals. \* $P < 0.05$  statistically different in the presence of L-NAME alone ( $\blacktriangle$ ) and L-NAME + indomethacin ( $\blacksquare$ ) compared to control on intact rings ( $\blacklozenge$ )



**Fig. 3** Concentration-response curves to adenosine with and without the presence of some specific ion channel blockers in intact isolated rat aortic rings. **a** In the presence of 1 mM 4-aminopyridine (■). **b** In the presence of 1 mM 5-Hydroxydecanoate (■). **c** In the presence of 10 μM glibenclamide (■). **d** In the presence of 100 μM ouabain (■) compared to controls intact rings (◆). Relaxation is expressed as percent of maximal relaxation to 100 μM papaverine. Points represent mean ± S.E.M of aortic rings from a total of eight animals. \**P* < 0.05 statistical difference in responses between the presence and the absence of inhibitors on intact rings

72, 79, 66 and 55 % reduction in relaxation. Similar to 4-AP and 5-HD, the  $A_{2a}$  receptor blocker did not inhibit adenosine relaxation at 500 μM and 1000 μM. In contrast, adenosine  $A_{2b}$  blocker, PSB 0788, had no effect to reduce adenosine-induced relaxation (Fig. 4b).


**Effect of  $K_v$ ,  $SarcK_{ATP}$ ,  $MitoK_{ATP}$  blockers and ouabain on adenosine relaxation in denuded rings**

In the absence of endothelium and blockers, adenosine relaxed rat aortic rings in a dose-dependent manner and reaching 78 % relaxation at the highest 1000 μM adenosine concentration (Fig. 5). Pre-treatment with 4-AP significantly reduced relaxation from 1 to 500 μM adenosine but not at 1000 μM (Fig. 5a). 4-AP nearly completely abolished adenosine-induced relaxation up to 10 μM adenosine with

over 95 % inhibition. At 50, 100 and 500 μM adenosine, the  $K_v$  channel was responsible for 74 %, 62 %, 21 % of adenosine relaxation (Fig. 5a).

The  $SarcK_{ATP}$  channel blocker, glibenclamide, also significantly reduced relaxation at 10, 50 and 100 μM adenosine levels (Fig. 5b) indicating that the  $SarcK_{ATP}$  channel was responsible for 41, 38 and 22 % of adenosine relaxation, respectively. Mitochondrial  $K_{ATP}$  blocker, 5-HD, significantly reduced relaxation over a wider range than glibenclamide similar to intact rings (Figs. 3b, c and 5b, c). The greatest effect of 5-HD was found at 10 to 100 μM. The contributions of the  $MitoK_{ATP}$  channel to adenosine relaxation were 51, 48, 44 and 14 % at 10, 50, 100 and 500 μM adenosine levels, respectively. The  $Na^+/K^+$ -ATPase channel blocker ouabain, as in intact aortic rings, showed





**Fig. 4** Concentration-response curves to adenosine with and without the presence of adenosine  $A_{2ab}$  receptor blockers in intact (a) and denuded (b) isolated rat aortic rings. In the presence of 100  $\mu\text{M}$  8-(3-Chlorostyryl) caffeine (■) or 10  $\mu\text{M}$  PSB 0788 (■). Controls (intact and denuded rings) (◆). Relaxation is expressed as percent of maximal relaxation to 100  $\mu\text{M}$  papaverine. Points represent mean  $\pm$  S.E.M of aortic rings from a total of eight animals. \* $P < 0.05$  statistical difference in responses between the presence and the absence of inhibitors on intact rings

no significant effects to reduce adenosine-induced vasodilation at any given adenosine level (Fig. 5d).


### Discussion

Despite decades of investigation, the mechanisms of adenosine relaxation in large elastic arteries such as the rat thoracic aorta, and smaller muscular resistance arterioles are not fully understood [3, 4, 6, 19, 44]. We report in isolated rat thoracic rings that adenosine vasodilation was: 1) triphasic and partially dependent on an intact endothelium, 2) regulated predominately by endothelial NO, not prostanoids, 3) dependent on opening smooth muscle  $K_v$ ,  $\text{SarcK}_{\text{ATP}}$  and  $\text{MitoK}_{\text{ATP}}$  channels, 4) ouabain-insensitive ( $\text{Na}^+/\text{K}^+$  ATPase), and 5) activated by the  $A_{2a}$  subtype, not  $A_{2b}$ . We discuss the possible interplay between these potassium channels and adenosine relaxation in denuded and intact aortic rings, and the in vivo significance.

### Adenosine relaxation involves an NO-dependent pathway

Our study showed that L-NAME significantly reduced relaxation in intact rings and contributed up to 59 % of adenosine relaxation with little or no effect of indomethacin (Fig. 2). In the rat aorta, endothelial NO is believed to induce vasodilation via cGMP- and cAMP-dependent protein kinase mechanisms, and the inhibition of Rho-kinase constrictor activity [45]. The lack of a prostanoid effect in our study was surprising. In 2002, Ray and colleagues showed in an elegant series of studies, using a NO-sensitive electrode, that adenosine relaxation in the rat aorta produced a dose-dependent NO release from the endothelium [46]. They further showed that  $A_1$ -receptor NO release was linked to endothelial prostacyclin release via a common cyclic AMP signalling pathway [21].

In contrast to our study, Ray and colleagues used halothane- $\text{O}_2$  anesthetized, hypoxic, male 200-250 g Wistar



rats, and aortic conduits of 10 mm in length which were longitudinally opened and the NO-sensitive electrode directly in contact with the endothelial surface [21, 46]. Systemic hypoxia in their study was induced using 8 %  $O_2$  in  $N_2$  for 5 min prior to aorta harvest, but the group did not specify the  $pO_2$ ,  $pCO_2$  or temperature of their bathing media. This is an interesting contrast, as we harvested the thoracic aorta from normoxic, male 300-350 g Sprague Dawley rats under thiopentone anesthesia, and our isolated intact rings were 3–4 mm in length and fully oxygenated at all

times. It is possible that prostanoid production in rat aortic rings is not activated during normoxia but during hypoxia. In 2001, Verma and colleagues also reported in healthy humans that COX-2–selective inhibition did not result in significant changes in endothelial vasodilator responses [47]. Further work is required to examine these differences in different models.

**Role of the endothelium to adenosine relaxation**

In the present study, adenosine vasodilation was partially endothelium-dependent (Fig. 1), which is consistent with

earlier work of Yen and colleagues [14], Moritoki et al., [15], Headrick and Berne [16] and Rose-Meyer and colleagues [17] in rat and guinea pig thoracic aorta. However, we showed that adenosine relaxation was triphasic (Fig. 1), and that endothelial removal reduced ring relaxation 'responsiveness' between 10 to 100  $\mu\text{M}$  adenosine (Phase 2) with little or no change to denuded ring sensitivity from 1 to 10  $\mu\text{M}$  (Phase 1) or from 100 to 1000  $\mu\text{M}$  (Phase 3) compared to intact rings (Fig. 1). To our knowledge, this triphasic nature of adenosine relaxation has not been reported before, and although the underlying mechanisms for the different sensitivities are not known, they appear to involve differential endothelial-smooth muscle sensitivities to endothelial NO production, and smooth muscle  $A_{2a}$  receptor and voltage-dependent  $K_v$  and  $K_{ATP}$  channels (see below).

#### **Role of voltage-dependent $K_v$ channels in adenosine relaxation**

The 4-AP experiments (~70-95 % inhibition at 5 to 100  $\mu\text{M}$  adenosine) demonstrated that the  $K_v$  channel has the potential to be a potent activator of adenosine relaxation in rat aortic rings. A similar change in intact and denuded rings (Figs. 3a and 5a) suggests that 4-AP effect was independent of endothelial NO production, and was preferentially activated on vascular smooth muscle (Fig. 3a). Our data support the study of Tammaro and colleagues who reported the presence of smooth muscle  $K_v$  channels in rat aorta [48], and that of Heapes and Bowles in swine coronary arteries who showed 4-AP-sensitive  $K^+$  channels in adenosine relaxation [49]. In addition,  $K_v$  channels have also been widely reported in regulating tone in smaller resistance vessels of cerebral and mesenteric vascular beds [31, 50–52], and in vascular smooth muscle from larger rat pulmonary arteries [53]. In conclusion, our data indicate that adenosine relaxation in isolated NE-precontracted rat aortic rings involved  $K_v$  channels with higher sensitivities found at lower adenosine levels. Further studies are required using more specific  $K_v$  channel isoform inhibitors (and agonists), and their membrane voltage dependence on relaxation [54] at low and high adenosine levels.

#### **Contributions of $\text{SarcK}_{ATP}$ and $\text{MitoK}_{ATP}$ channels to adenosine relaxation, and $A_{2a}$ receptor activation**

We further showed that the  $\text{SarcK}_{ATP}$  channel contributed to 14 to 63 % of adenosine relaxation up to 100  $\mu\text{M}$  adenosine (Figs. 3b and 5b), and  $\text{MitoK}_{ATP}$  channels contributed to 22 to 70 % relaxation up to 1000  $\mu\text{M}$  adenosine in intact and denuded aortic rings (Figs. 3c and 5c). The wider range of adenosine inhibition with  $\text{MitoK}_{ATP}$  channel blocker 5-HD indicates that it shifted the control relaxation curve more to the right than glibenclamide (Figs. 3c and 5c). For example, at 10 and 100  $\mu\text{M}$  adenosine, 5-HD led to 50 % more

inhibition than glibenclamide in intact rings (Fig. 5c, b), and 17 and 29 % more inhibition in denuded rings (Fig. 5c, b). This difference may indicate differential contributions of the  $\text{MitoK}_{ATP}$  and  $\text{SarcK}_{ATP}$  channel activation to adenosine relaxation, however, 5-HD has been shown to exert effects independent of  $\text{MitoK}_{ATP}$  channels [55] which may influence that interpretation.

Our glibenclamide data showing significant relaxation reduction (Figs. 3b and 5b), albeit less potent than 5-HD (Figs. 3c and 5c), is in contrast to the study of Husken and colleagues who reported no effect in rat aorta [56]. However, their rings were bathed in a hypoxic, low-glucose medium. Similarly Kemp and Cocks reported lack of a glibenclamide effect in coronary artery rings prepared from cardiac surgery patients [8]. It appears therefore that glibenclamide-sensitive  $K_{ATP}$  channel activation and adenosine relaxation is dependent on the state of tissue oxygenation, prior disease states and possibly ischemia.

Furthermore, Kemp and Cocks found that adenosine relaxation in their discarded human coronary artery rings was mediated largely by  $A_{2b}$  receptors [8], unlike  $A_{2a}$  receptors we found in isolated rat aortic rings (Fig. 4a, b). Adenosine  $A_{2a}$  receptor activation and relaxation in rat aortic rings is consistent with the majority of studies in rabbit aorta and mesenteric and coeliac arteries [57], mouse hearts [58], and guinea pig, porcine and bovine coronary arteries [10, 59, 60]. However, Lewis and colleagues reported in Wistar rat isolated aortic preparations that  $A_{2a}$  adenosine relaxation was entirely endothelium-dependent [10], not smooth muscle-dependent as we found in the present study (Fig. 4a, b). In rat renal artery, Grbović and colleagues also showed that removal of the endothelium abolished  $A_{2a}$  adenosine relaxation, implicating endothelial relaxation factors such as NO for relaxation [42]. These contrasting results may be due to differences in species, age, prior disease state, aortic ring preparation, presence of an endothelium and the bathing media. Another difference may be the type of artery; studying the larger arterial conduits versus smaller arteriolar resistance vessels which have very different functions (see below '*Limitations of the Present Study and Future Studies*'). It is noteworthy that Leal and colleagues found that  $A_{2a}$  and  $A_{2b}$  subtypes were abundant in all three layers of Wistar rat thoracic aorta wall (intima, media, and adventitia) [61], again illustrating the deep complexity of receptor and channel expression in the thoracic aorta.

#### **Adenosine regulation of relaxation in rat aortic rings: a working hypothesis**

Although we did not investigate adenosine relaxation at different oxygenation states and pH, or from hypoxic animals, we propose the following scheme. Adenosine-linked NO



production appeared to be a major endothelium-derived relaxing factor in intact rat aortic rings, not prostanoids, which sets the stage for endothelial-smooth muscle coupling. In denuded aortic rings, adenosine appears to activate  $A_{2a}$  receptors and trigger downstream opening of  $K_v$  and  $K_{ATP}$  channels located on smooth muscle resulting in membrane hyperpolarization, and relaxation, which may have involved common protein kinase signalling transduction pathways and crosstalk [50, 57, 62–66]. Membrane hyperpolarization of only a few millivolts can lower cytosolic  $Ca^{2+}$  via reduced activity of membrane voltage-operated  $Ca^{2+}$  channels and reduced myofilament  $Ca^{2+}$  sensitivity [67], resulting in smooth muscle relaxation. Partial support for this hypothesis in denuded rings comes from reports showing adenosine activation of  $K_v$  channels in pig coronary arterioles occurs via cAMP-dependent protein kinase (PKA) activation and vasodilatation [68, 69], and from Kleppisch and Nelson who showed that adenosine  $A_{2a}$  (not  $A_{2b}$ ) activation opens  $K_{ATP}$  channels via the cAMP/PKA pathway in isolated rabbit mesenteric vascular smooth muscle cells [57]. More recently, Maimon and colleagues showed in skeletal muscle arterioles that PKA signalling varies with pre-exposure to adenosine, and that PKA activation alone was not sufficient to dilate these arterioles, and required other  $Ca^{2+}$ -dependent mechanisms to facilitate vasodilation to adenosine [66].

Another possible mechanism coupling adenosine  $A_{2a}$  receptor to opening  $K_v$  and  $K_{ATP}$  channels in rat denuded aorta rings may be via mitochondrial production of  $H_2O_2$  [70, 71].  $H_2O_2$  is a highly diffusible and signalling redox intermediate produced during mitochondrial phosphorylation of ADP to ATP, and is believed to trigger  $Ca^{2+}$  sparks that activate protein kinase pathways and adenosine relaxation [72, 73]. Dick and colleagues reported that  $H_2O_2$  activated  $K_v$  channels and led to coronary vasodilation along with increases in myocardial metabolism [72], and Sharifi-Saniani and colleagues showed that adenosine  $A_{2a}$  receptor activation in mouse aorta during reactive hyperemia was coupled to smooth muscle  $K_{ATP}$  channels via the production of  $H_2O_2$  [73]. It is possible that mitochondrial  $H_2O_2$  bursts may also facilitate crosstalk between mito- and sarc- $K_{ATP}$  channels in our model.

Lastly, activation of  $A_{2a}$  in rat aortic rings may also have occurred from adenosine's breakdown metabolite, inosine (via adenosine deaminase), which has recently been shown to be a functional agonist of the  $A_{2a}$  receptor [74]. It is possible therefore that adenosine engages  $A_{2a}$  receptor to generate initial relaxation followed by a dual agonist-mediated response from inosine to amplify or prolong  $A_{2a}$  activation in vivo. While inosine is known to relax aortic rings [75], its dual action with adenosine has only been studied in inflammatory/immune cells [74].

### Limitations of the present study and future studies

While all four major types of  $K^+$  channels ( $K_v$ ,  $K_{ATP}$ ,  $K_{IR}$  and  $K_{Ca}$ ) appear to be present in vascular endothelial and smooth muscle cells [50, 52, 64, 71, 76, 77], we limited our study to  $K_v$  and  $K_{ATP}$  channels in intact and denuded aortic rings. We also investigated aortic ring relaxation in a high  $pO_2$  environment and it would be of interest to investigate the effect of lowering  $pO_2$  and changing pH. In addition, adenosine receptor characterization may have been more robust with the use of more than one  $A_{2a}$  and  $A_{2b}$  antagonist at appropriate concentrations. The isolated aortic ring preparation also lacks sympathetic neurohumoral innervations and the vasa vasorum, which makes translation to the intact vessel challenging. The in vivo significance of our results may relate to regulating compliance of the thoracic aorta as part of ventricular-arterial coupling [78–80]. The thoracic aorta and other large arteries are compliance vessels and are continually subjected to different hemodynamic forces such as mechanical stretch due to pulsatile blood flow, and may adjust vascular tone by changing the balance of vasodilating and vasoconstricting factors and neurohumoral mechanisms [78–80]. In contrast, smaller peripheral and coronary arterioles supply vascular beds and regulate flow by changing resistance to maintain adequate tissue oxygenation. Further studies are required to investigate the possible role of adenosine (and possibly inosine) and its various receptor subtypes to regulate compliance versus resistance vessels (including venous capacitance vessels) in different regions and vascular beds in the body.

### Conclusions

It was concluded that adenosine relaxation in NE-precontracted rat thoracic aortic rings was triphasic and partially endothelium-dependent, and involved endothelial NO production with a complex interplay between smooth muscle  $A_{2a}$  subtype and voltage-dependent  $K_v$ , Sarc $K_{ATP}$  and Mito $K_{ATP}$  channels, but not a prostanoid-dependent pathway.

### Ethics approval and consent to participate

Animals were treated in accordance with the Guide for the Care and Use of Laboratory Animals published by the US National Institutes of Health (NIH Publication No. 85–23, revised 1996). The James Cook University (JCU) Animal Ethics Committee approval number for the present study was A1535.

### Consent for publication

Not Applicable

### Availability of data and materials

The datasets supporting the conclusions of this article can be made available by emailing the authors.

## Abbreviations

NO: nitric oxide; NE: norepinephrine; L-NAME: L-N<sup>G</sup>-Nitroarginine methyl ester; 4-AP: 4-aminopyridine; CSC: 8-(3-chlorostyryl) caffeine; PSB-0788: 8-(4-(4-(4-chlorobenzyl)piperazine-1-sulfonyl)phenyl)-1-propylxanthine; SarK<sub>ATP</sub>: sarcolemma K<sub>ATP</sub>; MitoK<sub>ATP</sub>: mitochondrial K<sub>ATP</sub> channels; 5-HD: 5-hydroxydecanoate; ACh: acetylcholine.

## Competing interests

There no financial and non-financial competing interests. Aryadi Arsyad has no conflicts to declare. Geoffrey Dobson is the sole inventor of the adenosine, lidocaine and magnesium concept for cardioplegia, surgery, infection and trauma.

## Authors' contributions

Both authors contributed equally to the design, implementation, literature and data analysis and the writing of the manuscript. Both authors read and approved the final manuscript.

## Acknowledgements

The authors would like to College of Medicine and AITHM and James Cook University (JCU) for support of the project, and to the Australian Government Endeavour Scholarship to Aryadi Arsyad to support his stay at JCU. Thanks also go to Dr Yulia Djahir and Hayley Letson for editorial and advice on statistical analysis.

## Funding

Research support was from internal research funds to GPD from College of Medicine and AITHM, James Cook University.

## Author details

<sup>1</sup>Physiology Department, Medical Faculty, Hasanuddin University, Jl. Perintis Kemerdekaan, Km. 10, Tamalanrea, Makassar 90213, Indonesia. <sup>2</sup>Heart, Trauma and Sepsis Research Laboratory, Australian Institute of Tropical Health and Medicine, College of Medicine and Dentistry, James Cook University, 1 James Cook Drive, Queensland 4811, Australia.

Received: 25 February 2016 Accepted: 29 April 2016

Published online: 23 May 2016

## References

- Ely SW, Berne RM. Protective effects of adenosine in myocardial ischaemia. *Circulation*. 1992;85:893–904.
- Fredholm BB, IJzerman AP, Jacobson KA, Klotz KN, Linden J. International Union of Pharmacology. XXV. Nomenclature and classification of adenosine receptors. *Pharmacol Rev*. 2001;53:527–52.
- Jacobson I, Gao Z-G. Adenosine receptors as therapeutic targets. *Nat Rev*. 2006;5:247–64.
- Mustafa SJ, Morrison RR, Teng B, Pelleg A. Adenosine receptors and the heart: role in regulation of coronary blood flow and cardiac electrophysiology. In: Wilson CN, Mustafa SJ, editors. *Adenosine receptors in health and disease: handbook of experimental pharmacology*. Berlin Heidelberg: Springer; 2009. p. 160–88.
- Burnstock G, Ralevic V. Purinergic signaling and blood vessels in health and disease. *Pharmacol Rev*. 2013;66:102–92.
- Headrick JP, Ashton KJ, Rosemeyer RB, Peart JN. Cardiovascular adenosine receptors: expression, actions and interactions. *Pharmacol Ther*. 2013;140:92–111.
- Dobson GP, Faggian G, Onorati F, Vinten-Johansen J. Hyperkalemic cardioplegia in adult and pediatric cardiac surgery: end of an Era? *Front Clin Transl Physiol*. 2013;4:1–28.
- Kemp BK, Cocks TM. Adenosine mediates relaxation of human small resistance-like coronary arteries via A2B receptors. *Br J Pharmacol*. 1999;126:1796–800.
- Martin PL, Ueeda M, Olsson RA. 2-Phenylethoxy-9-methyladenine: an adenosine receptor antagonist that discriminates between A2 adenosine receptors in the aorta and the coronary vessels from the guinea pig. *J Pharmacol Exp Ther*. 1993;265:248–53.
- Lewis CD, Hourani SM, Long CJ, Collis MG. Characterization of adenosine receptors in the rat isolated aorta. *Gen Pharmacol*. 1994;25:1381–7.
- Lewis CD, Hourani SM. Involvement of functional antagonism in the effects of adenosine antagonists and L-NAME in the rat isolated heart. *Gen Pharmacol*. 1997;29:421–7.
- Ponnoth DS, Sanjani MS, Ledent C, Roush K, Krahn T, et al. Absence of adenosine-mediated aortic relaxation in A(2A) adenosine receptor knockout mice. *Am J Physiol Heart Circ Physiol*. 2009;297:H1655–60.
- Newman WH, Becker BF, Heier M, Nees S, Gerlach E. Endothelium-mediated coronary dilatation by adenosine does not depend on endothelial adenylate cyclase activation: studies in isolated guinea pig hearts. *Pflugers Arch*. 1988;413:1–7.
- Yen MH, Wu CC, Chiou WF. Partially endothelium-dependent vasodilator effect of adenosine in rat aorta. *Hypertension*. 1988;11:514–8.
- Moritoki H, Matsugi T, Takase H, Ueda H, Tanioka A. Evidence for the involvement of cyclic GMP in adenosine-induced, age-dependent vasodilatation. *Br J Pharmacol*. 1990;100:569–75.
- Headrick JP, Berne RM. Endothelium-dependent and -independent relaxations to adenosine in guinea pig aorta. *Am J Physiol*. 1990;259:H62–7.
- Rosemeyer RB, Hope W. Evidence that A2 purinoceptors are involved in endothelium-dependent relaxation of the rat thoracic aorta. *Br J Pharmacol*. 1990;100:576–80.
- De Mey JG, Vanhoutte PM. Role of the intima in cholinergic and purinergic relaxation of isolated canine femoral arteries. *J Physiol*. 1981;316:347–55.
- Furchgott RF. Role of endothelium in responses of vascular smooth muscle. *Circ Res*. 1983;53:557–73.
- Rubanyi G, Vanhoutte PM. Inhibitors of prostaglandin synthesis augment beta-adrenergic responsiveness in canine coronary arteries. *Circ Res*. 1985;56:117–25.
- Ray C, Marshall J. The cellular mechanism by which adenosine evokes release of nitric oxide from rat aortic endothelium. *J Physiol*. 2006;570:85–96.
- Sato T, Sasaki N, O'Rourke B, Marban E. Adenosine primes the opening of mitochondrial ATP-sensitive potassium channels: a key step in ischemic preconditioning? *Circulation*. 2000;102:800–5.
- Taylor SG, Southerton JS, Weston AH, Baker JR. Endothelium-dependent effects of acetylcholine in rat aorta: a comparison with sodium nitroprusside and cromakalim. *Br J Pharmacol*. 1988;94:853–63.
- Grbovic I, Radenkovic M. Analysis of adenosine vascular effect in isolated rat aorta: possible role of Na<sup>+</sup>/K<sup>+</sup> -ATPase. *Pharmacol Toxicol*. 2003;92:265–71.
- Kinoshita H, Iranami H, Kimoto Y, Dojo M, Hatano Y. Mild alkalization and acidification differentially modify the effects of lidocaine or mexiletine on vasorelaxation mediated by ATP-sensitive K<sup>+</sup> channels. *Anesthesiology*. 2001;95:200–6.
- Dogan M, Peker RO, Donmez S, Gokalp O. Magnesium and diltiazem relaxes phenylephrine-precontracted rat aortic rings. *Interact Cardiovasc Thorac Surg*. 2012;15:1–4.
- Zerkowski HR, Knocks M, Konerding MA, Doetsch N, Roth G, et al. Endothelial damage of the venous graft in CABG. Influence of solutions used for storage and rinsing on endothelial function. *Eur J Cardiothorac Surg*. 1993;7:376–82.
- Evans GR, Gherardini G, Gurlek A, Langstein H, Joly GA, et al. Drug-induced vasodilation in an in vitro and in vivo study: the effects of nicardipine, papaverine, and lidocaine on the rabbit carotid artery. *Plast Reconstr Surg*. 1997;100:1475–81.
- Rautureau Y, Toumaniantz G, Serpillon S, Jourdon P, Trochu J-T, et al. Beta 3-adrenoceptor in rat aorta: molecular and biochemical characterization and signalling pathway. *Br J Pharmacol*. 2002;137:153–61.
- Novakovic A, Bukarica LG, Kanjuh V, Heinle H. Potassium Channels-Mediated Vasorelaxation of Rat Aorta Induced by Resveratrol. *Basic Clin Pharmacol Toxicol* 2006;99:360-4.
- Mackie MR, Byron KL. Cardiovascular KCNQ (Kv7) potassium channels: physiological regulators and new targets for therapeutic intervention. *Mol Pharmacol*. 2008;74:1171–9.
- O'Rourke B. Mitochondrial K<sub>ATP</sub> channels in preconditioning. *Circ Res*. 2000;87:845–55.
- Toyoda Y, Friehs I, Parker RA, Levitsky S, McCully JD. Differential role of sarcolemmal and mitochondrial K(ATP) channels in adenosine-enhanced ischemic preconditioning. *Am J Physiol Heart Circ Physiol*. 2000;279:H2694–703.
- Gross ER, Gross GJ. Pharmacologic therapeutics for cardiac reperfusion injury. *Expert Opin Emerg Drugs*. 2007;12:367–88.
- Mathoôt RA, Soudijn W, Breimer DD, IJzerman AP, Danhof M. Pharmacokinetic-haemodynamic relationships of 2-chloroadenosine at adenosine A1 and A2a receptors in vivo. *Br J Pharmacol*. 1996;118:369–77.
- Allende G, Acevedo S. Evidence for a role of cyclic AMP and endothelium in rat aortic relaxation induced by R-PIA. *Open Circ Vasc J*. 2011;4:6–11.

37. Borrmann T, Hinz S, Bertarelli DCG, Li W, Florin NC, et al. 1-Alkyl-8-(piperazine-1-sulfonyl)phenylxanthines: development and characterization of adenosine A2B receptor antagonists and a New radioligand with subnanomolar affinity and subtype specificity. *J Med Chem*. 2009;52:3994–4006.
38. Van der Walt MM, TerreBlanche G, Petzer A, Lourens ACU, Petzer JP. The adenosine A2A antagonistic properties of selected C8-substituted xanthines. *Bioorganic Chem*. 2013;49:49–58.
39. Kalkan S, Hocaoglu N, Akgun A, Gidener S, Tuncok Y. Effects of adenosine receptor antagonists on amitriptyline-induced vasodilation in rat isolated aorta. *Clin Toxicol*. 2007;45:600–4.
40. Kataoka K, Furukawa K, Nagao K, Ishii N, Tsuru N. The participation of adenosine receptors in the adenosine 5-triphosphate-induced relaxation in the isolated rabbit corpus cavernosum penis. *Int J Urology*. 2007;14:764–8.
41. Schiedel AC, Lacher SK, Linnemann C, Knolle PA, Müller CE. Antiproliferative effects of selective adenosine receptor agonists and antagonists on human lymphocytes: evidence for receptor-independent mechanisms. *Purinergic Signal*. 2013;9:351–65.
42. Grbović L, Radenković M, Prostran M, Pesić S. Characterization of adenosine action in isolated rat renal artery. Possible role of adenosine A(2A) receptors. *Gen Pharmacol*. 2000;35:29–36.
43. Hein TW, Belardinelli L, Kuo L. Adenosine A2A receptors mediate coronary microvascular dilation to adenosine: role of nitric oxide and ATP-sensitive potassium channels. *J Pharmacol Exp Ther*. 1999;291:655–64.
44. Félétou M, Vanhoutte PM. Endothelium-dependent hyperpolarizations: past beliefs and present facts. *Ann Med*. 2007;39:495–516.
45. Chitaley K, Webb RC. Nitric oxide induces dilation of rat aorta via inhibition of rho-kinase signaling. *Hypertension*. 2002;39:438–42.
46. Ray CJ, Abbas MR, Coney AM, Marshall JM. Interactions of adenosine, prostaglandins and nitric oxide in hypoxia-induced vasodilatation: in vivo and in vitro studies. *J Physiol*. 2002;544:195–209.
47. Verma S, Raj SR, Shewchuk L, Mather KJ, Anderson TJ. Cyclooxygenase-2 blockade does not impair endothelial vasodilator function in healthy volunteers: randomized evaluation of rofecoxib versus naproxen on endothelium-dependent vasodilatation. *Circulation*. 2001;104:2879–82.
48. Tammaro P, Smith AL, Hutchings SR, Smirnov SV. Pharmacological evidence for a key role of voltage-gated K<sup>+</sup> channels in the function of rat aortic smooth muscle cells. *Brit J Pharmacol*. 2004;143:303–17.
49. Heaps CL, Bowles DK. Gender-specific K<sup>+</sup> channel contribution to adenosine-induced relaxation in coronary arterioles. *J Appl Physiol*. 2002;92:550–8.
50. Cole WC, Clément-Chomienne O, Aiello EA. Regulation of 4-aminopyridine-sensitive, delayed rectifier K<sup>+</sup> channels in vascular smooth muscle by phosphorylation. *Biochem Cell Biol*. 1996;74:439–47.
51. Albarwani S, Nemetz LT, Madden JA, Tobin AA, England SK, et al. Voltage-gated K<sup>+</sup> channels in rat small cerebral arteries: molecular identity of the functional channels. *J Physiol*. 2003;551:751–63.
52. Coleman HA, Tare M, Parkington HC. Endothelial potassium channels, endothelium-dependent hyperpolarization and the regulation of vascular tone in health and disease. *Clin Exp Pharmacol Physiol*. 2004;31:641–9.
53. Archer SL, Wu X-C, Thebaud B, Nsair A, Bonnet S, et al. Preferential expression and function of voltage-gated, O<sub>2</sub>-sensitive K channels in resistance pulmonary arteries explains regional heterogeneity in hypoxic pulmonary vasoconstriction. *Circ Res*. 2004;95:308–18.
54. Smirnov SV, Tammaro P, Hutchings SR, Smith AF. Role of voltage-gated K<sup>+</sup> (KV) channels in vascular function. *Neurophysiology*. 2003;35:234–47.
55. Li X, Rapedius M, Baukowitz T, Liu GX, Srivastava DK, et al. 5-Hydroxydecanoate and coenzyme A are inhibitors of native sarcolemmal KATP channels in inside-out patches. *Biochim Biophys Acta*. 2010;1800:385–91.
56. Hüsken BC, Pfaffendorf M, van Zwieten PA. ATP-sensitive potassium channels in isolated rat aorta during physiologic, hypoxic, and low-glucose conditions. *J Cardiovasc Pharmacol*. 1997;29:130–5.
57. Kleppisch T, Nelson MT. Adenosine activates ATP-sensitive potassium channels in arterial myocytes via A2 receptors and cAMP-dependent protein kinase. *Proc Natl Acad Sci U S A*. 1995;92:12441–5.
58. Zatta AJ, Headrick JP. Mediators of coronary reactive hyperaemia in isolated mouse heart. *Br J Pharmacol*. 2005;144:576–87.
59. Shryock JC, Belardinelli L. Adenosine and adenosine receptors in the cardiovascular system: biochemistry, physiology, and pharmacology. *Am J Cardiol*. 1997;79:2–10.
60. Conti A, Lozza G, Monopoli A. Prolonged exposure to 5'-N-ethylcarboxamidoadenosine (NECA) does not affect the adenosine A2A-mediated vasodilation in porcine coronary arteries. *Pharmacol Res*. 1997;35:123–8.
61. Leal S, Sá C, Gonçalves J, Fresco P, Diniz C. Immunohistochemical characterization of adenosine receptors in rat aorta and tail arteries. *Microsc Res Tech*. 2008;71:703–9.
62. Quayle JM, Nelson MT, Standen NB. ATP-sensitive and inwardly rectifying potassium channels in smooth muscle. *Physiol Rev*. 1997;77:1165–232.
63. Brayden JE. Functional roles of KATP channels in vascular smooth muscle. *Clin Exp Pharmacol Physiol*. 2002;29:312–6.
64. Ko EA, Han JYH, Jung ID, Park WS. Physiological roles of K<sup>+</sup> channels in vascular smooth muscle cells. *J Smooth Muscle Res*. 2008;44:65–81.
65. Berwick ZC, Payne GA, Lynch B, Dick GM, Sturek M, et al. Contribution of adenosine A(2A) and A(2B) receptors to ischemic coronary dilation: role of K(V) and K(ATP) channels. *Vascul Pharmacol*. 2010;17:600–7.
66. Maimon N, Titus PA, Sarelis IH. Pre-exposure to adenosine, acting via A(2A) receptors on endothelial cells, alters the protein kinase A dependence of adenosine-induced dilation in skeletal muscle resistance arterioles. *J Physiol*. 2014;592:2575–90.
67. Akata T. Cellular and molecular mechanisms regulating vascular tone. Part 2: regulatory mechanisms modulating Ca<sup>2+</sup> mobilization and/or myofilament Ca<sup>2+</sup> sensitivity in vascular smooth muscle cells. *J Anesth*. 2007;21:232–42.
68. Heaps CL, Jeffery EC, Laine GA, Price EM, Bowles DK. Effects of exercise training and hypercholesterolemia on adenosine activation of voltage-dependent K<sup>+</sup> channels in coronary arterioles. *J Appl Physiol*. 2008;105:1761–71.
69. Ko EA, Park WS, Firth AL, Kim N, Yuan JX, et al. Pathophysiology of voltage-gated K<sup>+</sup> channels in vascular smooth muscle cells: modulation by protein kinases. *Prog Biophys Mol Biol*. 2010;103:95–101.
70. Krenz M, Oldenburg O, Wimpee H, Cohen MV, Garlid KD, et al. Opening of ATP-sensitive potassium channels causes generation of free radicals in vascular smooth muscle cells. *Basic Res Cardiol*. 2002;97:365–73.
71. Bonnet S, Archer SL. Potassium channel diversity in the pulmonary arteries and pulmonary veins: implications for regulation of the pulmonary vasculature in health and during pulmonary hypertension. *Pharmacol Ther*. 2007;115:56–69.
72. Dick GM, Bratz IN, Borbouse L, Payne GA, Dincer UD, et al. Voltage-dependent K<sup>+</sup> channels regulate the duration of reactive hyperemia in the canine coronary circulation. *Am J Physiol Heart Circ Physiol*. 2008;294:H2371–81.
73. Sharif-Sanjani M, Zhou X, Asano S, Tilley S, Ledent C, et al. Interactions between A(2A) adenosine receptors, hydrogen peroxide, and KATP channels in coronary reactive hyperemia. *Am J Physiol Heart Circ Physiol*. 2013;304:H1294–301.
74. Welihinda AA, Kaur M, Greene K, Zhai Y, Amento EP. The adenosine metabolite inosine is a functional agonist of the adenosine A2A receptor with a unique signaling bias. *Cell Signal*. 2016;28:552–60.
75. Chinellato A, Ragazzi E, Pandolfo L, Frolidi G, Caparrotta L, et al. Purine- and nucleotide-mediated relaxation of rabbit thoracic aorta: common and different sites of action. *J Pharm Pharmacol*. 1994;46:337–41.
76. Chen TT, Luykenar KD, Walsh EJ, Walsh MP, Cole WC. Key role of Kv1 channels in vasoregulation. *Circ Res*. 2006;99:53–60.
77. Edwards G, Félétou M, Weston AH. Endothelium-derived hyperpolarising factors and associated pathways: a synopsis. *Eur J Physiol*. 2010;459:863–79.
78. Sandoo A, Veldhuijzen van Zanten JJCS, Metsios GS, Carroll D, Kitas GD. The endothelium and its role in regulating vascular tone. *Open Cardiovasc Med J*. 2010;4:302–12.
79. Triggle CR, Samuel SM, Ravishankar S, Marei I, Arunachalam G, et al. The endothelium: influencing vascular smooth muscle in many ways. *Can J Physiol Pharmacol*. 2012;90:713–38.
80. Jufri NF, Mohamedali A, Avolio A, Baker MS. Mechanical stretch: physiological and pathological implications for human vascular endothelial cells. *Vasc Cell*. 2015;7:8.