

ResearchOnline@JCU

This is the **Accepted Version** of a paper published in the
Australian Journal of Primary Health:

Callander, Emily J., Corscadden, Lisa, and Levesque, Jean-Frederic
(2017) *Out-of-pocket healthcare expenditure and chronic disease –
do Australians forgo care because of the cost?* Australian Journal of
Primary Health, 23 (1). pp. 15-22.

<http://doi.org/10.1071/PY16005>

1 **Out-of-pocket healthcare expenditure and chronic disease – do Australians forgo care**
2 **due to the cost**

3 **Abstract**

4 *Background*

5 While we do know that out-of-pocket healthcare expenditure is relatively high in Australia,
6 little is known about what health conditions are associated with the highest out-of-pocket
7 expenditure, and whether the cost of healthcare acts as a barrier to care for people with
8 different chronic conditions.

9 *Methods*

10 Cross sectional analysis using linear and logistic regression models applied to the
11 Commonwealth Fund International Health Policy survey of adults aged 18 years and over
12 conducted in 2013.

13 *Results*

14 Adults with asthma, emphysema and COPD had household out-of-pocket healthcare
15 expenditure 109% higher than those with no health conditions (95% CI: 50% – 193%); and
16 adults with depression, anxiety and other mental health conditions had household out-of-
17 pocket expenditure 95% higher (95% CI: 33% – 187%). People with chronic conditions were
18 also more likely to forego care due to cost. People with depression, anxiety and other mental
19 health conditions had odds 7.65 times as high of skipping healthcare (95% CI: 4.13 – 14.20),
20 and people with asthma, emphysema and COPD had odds 6.16 times as high of skipping
21 healthcare (95% CI: 3.30 – 11.50) compared to people with no health condition. People with
22 chronic health conditions in Canada, the United Kingdom, Germany, France, Norway,
23 Sweden and Switzerland were all significantly less likely to skip healthcare because of cost
24 than people with a condition in Australia.

1 *Conclusions*

2 The out-of-pocket cost of healthcare in Australia acts as a barrier to accessing treatment for
3 people with chronic health conditions, with people with mental health conditions being likely
4 to skip care. Attention should be given to the accessibility and affordability of mental health
5 services in Australia.

6 **KEYWORDS:** Patient contributions; chronic health conditions

7

1 **Summary statement**

2 *What is known about this topic?*

- 3 - Individuals contribute 17% of total healthcare spending in Australia.

4 *What does this paper add?*

- 5 - Individuals with chronic health conditions are more likely to forgo care due to the cost
6 in Australia than many other countries.

- 7 - Individuals with mental health conditions are particularly likely to skip care,
8 indicating a need to address potentially prohibitive costs in this area of care.

1 **Introduction**

2 Despite Australia having a universal healthcare system, individuals' out-of-pocket medical
3 expenditure is reported by the Australian Institute of Health and Welfare to account for 17%
4 of total health spending (Australian Institute of Health and Welfare 2014). This makes
5 individuals the third largest contributor to health spending - only behind the Australian
6 government, and the state and territory governments, and means they contribute more than
7 double that paid by private health insurance companies (Australian Institute of Health and
8 Welfare 2014). By way of comparison, individual out-of-pocket expenditure was reported to
9 account for 9% of total health expenditure from all sources in the United Kingdom, 8% in
10 France, and 13% in Germany but as high as 25% in Switzerland (Organisation for Economic
11 Co-operation and Development (OECD) 2014) - based upon reported aggregate figures.

12 When broken down to the household's perspective, Australian's out-of-pocket expenditure on
13 healthcare equated to an average of 9% of total household expenditure per year for an older
14 household and 5% for a younger household (Yusuf and Leeder 2013). It is also known that
15 out-of-pocket expenditure for those with certain chronic health conditions is particularly high.
16 For example, individual stroke survivors reported spending an average of \$1,110 per year on
17 healthcare, however out-of-pocket expenditure ranged from \$0 to \$32,411 in the twelve
18 months following their stroke (Dewey, Thrift et al. 2004); similarly, for an individual with
19 arthritis, the mean out-of-pocket expenditure directly attributable to arthritis was \$1,513 per
20 year, but with a range of \$49 to \$20,527 (Lapsley, March et al. 2002) – both exclude health
21 insurance premiums and only consist of expenses directly related to the specific condition.
22 The expenditure by individuals in rural areas is even higher, with cancer patients treated in
23 North Queensland spending an average \$4,311 since diagnosis (interquartile range \$563 –
24 6,231), while those who lived more than 100km from the hospital spent an average of \$7,752

1 (Gordon, Ferguson et al. 2009). This high and heterogeneous out-of-pocket expenditure is
2 mirrored in the international literature (Lehnert, Heider et al. 2011).

3 However, the literature regarding out-of-pocket expenditure is fragmented – with studies only
4 focusing upon one type of chronic health condition, one type of healthcare expenditure, or
5 one particular population (for example, older people) (Lapsley, March et al. 2002, Dewey,
6 Thrift et al. 2004, Gordon, Ferguson et al. 2009, Essue, Kelly et al. 2011, McRae, Yen et al.
7 2013). International comparisons have also been made for overall healthcare expenditure in
8 Australian and other countries (Squires 2012), and the recent Senate Inquiry into Out-of-
9 pocket Health Expenditure in Australia did report on patient contributions being a barrier to
10 accessing care (Senate Standing Committees on Community Affairs 2014), but to date no
11 studies have been able to investigate the reported out-of-pocket expenditure and the impact
12 this has on decisions to access care for patients with specific health conditions for the
13 Australian population.

14 Given the current heightened level of community discussion on increases to out-of-pocket
15 healthcare expenditure (Duckett, Boxall et al. 2014, Bourke 2015, Owler 2015) associated
16 with proposed General Practitioner co-payments (Bourke 2015), the Extended Medicare
17 Safety Net (Department of Human Services 2014), and changes to Pharmaceutical Benefits
18 Scheme co-payments (Department of Human Services 2014), it is imperative to understand
19 current levels of out-of-pocket healthcare expenditure and whether the cost of healthcare may
20 be discouraging people from accessing care. This paper aims to answer the following
21 research questions: what level of out-of-pocket expenditure is associated with different health

1 conditions in Australia¹? With which conditions are people more likely to forego healthcare
2 due to the out-of-pocket cost compared to people with no conditions? How does Australia
3 compare internationally, in terms of self-reported out-of-pocket expenditure, and having
4 people with chronic health conditions forego healthcare due to cost?

5 **Methods**

6 The Commonwealth Fund International Health survey was conducted in 2013 by Social
7 Science Research Solutions and country contractors, and consisted of computer-assisted
8 telephone interviews of random samples of adults aged eighteen and older using a common
9 questionnaire between February and June 2013 (Rapoport, Tipan et al. 2013). Samples were
10 collected in Australia and 10 other sponsor countries including; Canada, Germany, France,
11 Netherlands, New Zealand, Norway, Sweden, Switzerland, the United Kingdom and the
12 United States. In Australia, there were 2,200 respondents with assigned survey weights such
13 that they would be representative of the age, gender, education and regional distribution of
14 Australia. Of note, the state of New South Wales (NSW) being oversampled as the
15 Australian sponsor of the survey was the NSW Bureau of Health Information, which makes
16 the margin of sampling error for Australia +/- 3.4%, larger than other countries with similar
17 sample sizes.

18 The Commonwealth Fund International Health Policy Surveys (The Commonwealth Fund
19 2013) have been conducted since 1998, with questions that have been tested and revised in
20 many settings, over several cycles (Osborn, Moulds et al. 2014). The reported out-of-pocket

¹ This paper focuses upon household expenditure on the direct costs of care and does not cover any indirect costs of seeking health care, such as transport costs, carer costs and opportunity costs from spending time seeking care rather than engaging in other activities such as employment.

1 expenditure on the 2013 survey was validated by comparing mean values with administrative
2 data (Table 1). The levels and order of magnitude correspond well.

3 Missing data

4 The survey asked respondents about their household expenditure on medical services with the
5 following two questions:

- 6 1. In the past 12 months, about how much have you and your family spent out-of-pocket
7 for medical treatments or services that were not covered by insurance? This would
8 include what you paid for prescription medicines, medical and dental care.
- 9 2. If you can't recall exactly how much you and your family spent out-of-pocket for
10 medical treatments or services, please provide your best estimate.

11 The first question was a continuous variable, with respondents able to record the dollar value
12 of their expenditure. The second question was a categorical variable, with respondents able to
13 select the following ranges: less than \$100; \$100 to less than \$500; \$500 to less than \$1,000;
14 \$1,000 to less than \$2,000; \$2,000 or more.

15 Of the 2,200 respondents, there were 212 respondents who stated they were not sure how
16 much their expenditure was or declined to answer to both questions. These were excluded
17 from the analysis.

18 For the respondents who stated they were unsure of the exact amount of their expenses in the
19 first question, but were able to identify which category their expenses fell into in the second
20 question, we assigned a continuous dollar value of expenditure based upon imputation from
21 the balance of the sample. For people with valid responses to questions 1 and 2, the mean
22 value of continuous household expenditure for those who stated in question 2 that they spent
23 less than \$100; \$100 to less than \$500; \$500 to less than \$1,000; \$1,000 to less than \$2,000;

1 \$2,000 or more was identified, as shown in Table 2. Respondents who were unsure of the
2 continuous value of their expenditure but were able to identify the range of their expenditure,
3 were assigned the average value shown in Table 2.

4 Exchange rates and currency

5 For all analysis that focuses only on Australia, Australian dollars are used as the currency.
6 For all analysis involving international comparison, United States dollars are used, with the
7 exchange rate set at \$1.04 US dollars to 1 Australian dollar, which was the exchange rate at
8 the time of the survey.

9 Australian Healthcare System

10 Australia does have a universal health care system, Medicare, which provides the population
11 with access to essential healthcare services at no charge to the end user or for a subsidised fee
12 (Department of Human Services 2015). In some areas of Australia many services are ‘bulk
13 billed’, meaning that the full cost of treatment is billed directly to Medicare and the patients
14 do not face any up-front or out of pocket charges. However, many services do have an ‘out
15 of pocket’ charge that is paid for by the patient, this occurs when the fee charged by the
16 service provider is higher than the rebate paid by Medicare. Similarly, for services that are
17 not covered by Medicare, or services which have a capped number of uses in a given time
18 period, the patient may have to pay the full fee.

19 Statistical methods

20 Initial univariate analysis was undertaken to document the distribution of stated out-of-pocket
21 healthcare expenditure in Australia. A series of logistic regression models were then
22 constructed to estimate the adjusted odds ratios of having any out-of-pocket healthcare
23 expenditure for people with different reported health conditions, compared to those with no

1 health conditions and adjusted for age, sex and highest level of education attainment
2 (secondary education or less; greater than secondary). As it was possible for respondents to
3 select multiple health conditions, separate models were constructed for each health condition.

4 The analysis was then limited to those who stated they had more than \$0 of healthcare
5 expenditure, and a series of linear regression models were then constructed on log
6 transformed data to estimate the adjusted difference of stated out-of-pocket expenditure for
7 people who reported having different health conditions, individually and in combination,
8 compared to those with no health conditions. The model was adjusted for age, sex and
9 highest level of education attainment.

10 The proportion of people who stated they skipped healthcare due to healthcare cost was then
11 compared between adults who had long term health conditions and those with no health
12 conditions. A series of logistic regression models were then constructed to assess the odds of
13 skipping care due to cost for those with each health condition compared to those with no
14 health conditions also adjusting for age, sex and highest level of education attainment.

15 To provide international context, the variation in reported expenditure on healthcare for
16 Australia and Canada, France, Germany, the Netherlands, New Zealand, Norway, Sweden,
17 Switzerland, the United Kingdom, and the United States is also provided. A logistic
18 regression model was constructed to compare the odds of skipping care due to the costs of
19 healthcare for all countries compared to Australia.

20 **Results**

21 There were 1,988 adults aged 18 and over from Australia with valid responses regarding
22 household out-of-pocket healthcare expenditure. Of these, 52% were female, 45% were aged
23 under 45 and 53% had more than a secondary level of education attainment (Table 3). When

1 asked if they had ever been told by a doctor they had: hypertension or high blood pressure,
2 heart disease, including heart attack, diabetes, asthma or chronic lung disease such as chronic
3 bronchitis, emphysema or chronic obstructive pulmonary disease, depression, anxiety or
4 other mental health problems, cancer, arthritis, or high cholesterol nearly half (49%) of the
5 adult population had at least one of these long term health conditions

6 Respondents reported spending an average of \$986 per year on household out-of-pocket
7 healthcare expenditure (median = AU\$250, range = \$0 - \$40,000). Table 4 shows the range
8 of expenditure per year. Around one quarter of people reported no out-of-pocket healthcare
9 expenditure, and a similar proportion reported between AU\$100 and AU\$500. Fourteen
10 percent reported expenditure over AU\$2,000.

11 Table 5 shows the distribution of this expenditure by gender, age and number of chronic
12 health conditions. Adults with one or more health conditions reported average expenditure
13 around double that reported by those who had no long-term health condition. Those in the
14 older age groups reported lower amounts of expenditure, as did those with multiple co-
15 morbidities.

16 Adults with different chronic health conditions had consistently higher average yearly
17 household out-of-pocket healthcare expenditure than those with the no chronic conditions.
18 People with Asthma, emphysema and COPD reported an average of AU\$1,642 per year in
19 household out-of-pocket healthcare expenditure, and those with high cholesterol reported an
20 average of AU\$1,423 per year (Table 6).

21 Australian adults with asthma, emphysema and COPD, depression, anxiety and other mental
22 health conditions, heart disease, hypertension and high cholesterol were all significantly more
23 likely to report any out-of-pocket expenditure compared to people with no chronic health
24 conditions (Table 6). People with Asthma, emphysema and COPD had 3.97 times the odds of

1 reporting out-of-pocket healthcare expenditure (95% CI: 2.21 – 7.14) and people with heart
2 disease had 3.93 times the odds (95% CI: 1.73 – 8.93) of reporting out-of-pocket healthcare
3 expenditure than people with no health condition, after adjusting for age, sex and education
4 attainment.

5 Looking only at people who had out-of-pocket expenditure, people with arthritis, asthma,
6 emphysema and COPD, depression, anxiety and other mental health conditions, diabetes,
7 hypertension and high cholesterol all had significantly higher out-of-pocket healthcare
8 expenditure than those with no health conditions, after adjusting for age, sex and education
9 attainment. People with asthma, emphysema and COPD had out-of-pocket healthcare
10 expenditure 109% higher than those with no health conditions (95% CI: 50% – 193%); and
11 people with depression, anxiety and other mental health conditions had out-of-pocket
12 expenditure 95% higher (95% CI: 33% – 187%).

13 Table 7 shows the proportion of people who stated they skipped healthcare treatment because
14 of the cost. An estimated 9% of Australians without a health condition stated they skipped
15 healthcare treatment because of the cost. Amongst those with one or more health conditions,
16 over 40% of people with depression, anxiety and other mental health conditions, and over one
17 third of people with asthma, emphysema and COPD stated they skipped healthcare treatment
18 because of the cost. After adjusting for age, sex and education attainment, people with
19 depression, anxiety and other mental health conditions had 7.65 times the odds of skipping
20 treatment (95% CI: 4.13 – 14.20), and people with asthma, emphysema and COPD had odds
21 of skipping care 6.16 times as high (95% CI: 3.30 – 11.50) as people with no health
22 condition.

23 Table 8 shows the average yearly reported household healthcare expenditure, in United States
24 dollars, for different countries. The average estimated yearly household healthcare

1 expenditure in Australia was US\$1,026, in the United States average spending was
2 US\$1,844, and in the United Kingdom average spending was US\$216 per year. Table 7 also
3 shows the odds ratio of skipping care due to costs amongst people with a chronic health
4 condition in different countries. People with chronic health conditions in Canada, the United
5 Kingdom, Germany, France, Norway, Sweden and Switzerland were all significantly less
6 likely to skip healthcare because of costs than people with chronic conditions in Australia.
7 People in the United States with conditions had twice the odds of skipping healthcare due to
8 costs compared to those in Australia (95% CI: 1.51 – 2.57). Figure 1 also shows the reported
9 annual expenditure of people with different numbers of co-morbidities across the different
10 countries.

11 **Discussion**

12 The Commonwealth Fund International Health Policy survey data provides a unique
13 opportunity to gain insights on the out-of-pocket healthcare expenditure of Australians, and
14 also allows comparisons with other countries to be made. The results of this analysis have
15 shown that respiratory disease (asthma, emphysema and COPD), mental health conditions
16 (depression, anxiety and other mental health conditions) and diabetes are the conditions with
17 high out-of-pocket healthcare expenditure in Australia (relative to people with no conditions).
18 For example, people with asthma, emphysema or COPD reported household spending of
19 \$1,600 per year on healthcare on average. The distribution analysis that shows people with,
20 multiple chronic conditions have lower out-of-pocket expenditure may be explained by a
21 cohort effect, where people with more chronic conditions are both more likely to be older and
22 more likely to receive some of their care in places where care is provided without an out-of-
23 pocket fee (such as within public hospitals as an inpatient, outpatients clinics, and Emergency
24 Departments). It could be a patient's trajectory to face significant expenditure for a condition
25 when followed in primary healthcare settings only and then see a reduction of costs, even for

1 that first disease, as more intense care is provided in parts of the hospital system. Similar
2 finding have been reported in other countries where older patients, once they got much sicker
3 with multiple chronic diseases, were returning to the public sector, not able to afford private
4 care anymore (Levesque 2008).

5 A high percentage of people with chronic conditions in Australia choose to forgo care
6 because of the cost. Even amongst those *without* a chronic health condition, 9% chose to
7 forgo care. However, the percentage of people who have a chronic health condition that
8 stated they are skipping care due to the cost is much greater. Over 40% of people with
9 depression, anxiety and other mental health conditions stated they skip treatment due to the
10 cost. Arthritis and asthma, emphysema and COPD were the other conditions with a high
11 percentage of people forgoing care due to the cost.

12 Those with mental health conditions were shown to have particularly large out-of-pocket
13 expenditure and be restively more likely to forgo care, which indicates that the costs of
14 mental health services may be prohibitively high. Other studies have noted that the Medicare
15 rebates for mental health are particularly low relative to the fees charged by mental health
16 practitioners (Farag, Sherrington et al. 2013). Other studies have also found that patients with
17 mental health problems were more likely to access GP services (which have lower out of
18 pocket charges on average) than mental health services (Lucas, Bayer et al. 2013). However,
19 this study was not able to specifically identify the exact treatments or services that were
20 foregone.

21 This study only reported the actual amount people spent on healthcare, and did not capture
22 the amount they would have spent had they been able to afford it. Given that a large
23 proportion of people with chronic conditions do forgo treatment and care due to the cost, their

1 actual healthcare expenditure could have been much higher but their household incomes may
2 have curtailed this.

3 Other studies have demonstrated that accessing healthcare and good condition management
4 are effective means for maintaining labour force participation amongst people with chronic
5 health conditions – particularly people with arthritis, diabetes and depression (Claxton 1999,
6 Yelin, Katz et al. 2001, Passey, Shrestha et al. 2012). The findings of this study show that,
7 ironically a large proportion of people with some chronic health conditions – particularly
8 people with depression and arthritis – are likely to forgo care due to the cost of healthcare.
9 Cost of healthcare has been an overlooked barrier to accessing healthcare, with previous
10 studies within Australia showing that lack of available services are a common barrier to
11 accessing care (Humphreys 2009). As such, policies aiming to promote affordable healthcare
12 to at risk and vulnerable populations (NSW Health 2015) are of vital importance to ensure
13 that out-of-pocket cost is not a barrier to treatment and do not widen the gap in health status
14 between those of high and low socioeconomic status.

15 It is well documented that people with chronic health conditions within Australia, and
16 internationally, have lower incomes, less wealth and are more likely to be in income poverty,
17 likely due to the impact that chronic health conditions have on ability to participate in the
18 labour force (Schofield, Shrestha et al. 2008, Schofield, Kelly et al. 2010, Callander,
19 Schofield et al. 2011, Hunter, Schofield et al. 2014). While the lower income of individuals
20 with chronic health conditions may act as a barrier to care, the out-of-pocket expenditure on
21 healthcare that people with chronic health conditions are able to afford will also add to the
22 financial burden of chronic illness leaving less disposable income available for other
23 expenses. Past studies assessing the disposable income of households with members who
24 have a chronic health condition have not taken into account the additional out-of-pocket

1 healthcare expenditure demonstrated in this study, which further reduces the actual
2 disposable income available to individuals with chronic conditions.

3 By way of international comparison, estimated self-reported out-of-pocket healthcare
4 expenditure in Australia is higher than other countries included in this analysis, with the
5 exception of the United States. This example demonstrates the caution that is need for the
6 interpretation of these international comparisons in light of the differences in healthcare
7 systems. People with chronic health conditions within Australia were also significantly more
8 likely to forgo care compared to most other countries. These findings are consistent with
9 administrative data for Australia showing 17% of all healthcare expenditure was attributed to
10 individuals' out-of-pocket expenditure in 2012-13, which is second only to Switzerland.
11 While those in the United States do have high out-of-pocket healthcare costs, total spending
12 on healthcare in the United States from all sources is also very high, and as a proportion of all
13 expenditure on healthcare, individuals from Australia and Switzerland contribute the highest
14 proportion of total expenditure (Bureau of Health Information 2014).

15 This study has a number of limitations that need to be acknowledged. First the study
16 population is designed to be representative of age, sex, regional and education based
17 populations of Australia. It is not designed to represent people with chronic conditions.
18 Therefore, it may under or over cover specific disease or population groups. Compared to
19 prevalence estimates from a national survey for the population Arthritis (15%), Asthma
20 (10%) and Diabetes (4%) estimates for the population are comparable but slightly lower than
21 the estimates for adults aged 18 and over from this international survey (17%, 12% and 9%
22 respectively) (Australian Institute of Health and Welfare 2014). In addition, older adults
23 living in nursing homes and other facilities were not sampled. This would reduce the
24 representativeness of the population with chronic conditions. Secondly, the survey did not
25 specifically ask respondents about over the counter medication, or aids and equipment, travel

1 for treatment, or health insurance premiums or excess amounts as a part of the question
2 regarding healthcare expenditure. As such, the figures represented in this study may
3 underestimate of the total amount spent by households on healthcare. Furthermore, the large
4 number of people stating they were unsure about the exact amount they spent on healthcare
5 indicates the difficulty individuals have with recalling exact amounts of expenditure,
6 particularly when the estimates are for an annual time period. There is no way to know how
7 these limitations influence international comparisons.

8 Despite these limitations, this study has shown the impact out-of-pocket healthcare
9 expenditure has on individuals with chronic health conditions within Australia, and that the
10 cost of healthcare does act as a barrier to receiving treatment, particularly those with mental
11 health conditions. This is the first time comparisons have been made to estimate the relative
12 amount of out-of-pocket expenditure between people with certain health conditions and
13 people without conditions, and has demonstrated that people with depression, anxiety and
14 other mental health conditions face particularly large costs, and are particularly deterred from
15 accessing care due to the cost. These findings come at a vital time when there has been much
16 discussion about the possibility of raising the cost of healthcare to individuals.

17 Conflict of interest

18 None.

1 **References**

- 2 Australian Institute of Health and Welfare (2014). *Australias Health, 2014*. Canberra, Australian
3 Government.
- 4 Bourke, L. (2015). Health Minister Sussan Ley backs away from plan to cut Medicare rebate. The
5 Sydney Morning Herald. Sydney, Fairfax Media.
- 6 Bureau of Health Information (2014). *Healthcare in Focus 2013: How well does NSW measure up?*.
7 Sydney, BHI.
- 8 Callander, E., D. Schofield and R. Shrestha (2011). "Multi-dimensional poverty in Australia and the
9 barriers ill health imposes on the employment of the disadvantaged." Journal of Socio Economics **40**:
10 736-742.
- 11 Claxton, A. J., Chawla, A.J., & Kennedy, S. (1999). "Absenteeism among workers treated for
12 depression." Journal of Occupational and Environmental Medicine **41**(7): 605-611.
- 13 Department of Human Services. (2014). "Budget 2013-14: Extended Medicare Safety Net."
14 Retrieved 23 Jan, 2015, from [http://www.humanservices.gov.au/corporate/publications-and-](http://www.humanservices.gov.au/corporate/publications-and-resources/budget/1314/measures/health-matters/20-60516)
15 [resources/budget/1314/measures/health-matters/20-60516](http://www.humanservices.gov.au/corporate/publications-and-resources/budget/1314/measures/health-matters/20-60516).
- 16 Department of Human Services. (2014). "Budget 2014-15: Pharmaceutical Benefits Scheme."
17 Retrieved 23 Jan, 2015, from [http://www.humanservices.gov.au/corporate/publications-and-](http://www.humanservices.gov.au/corporate/publications-and-resources/budget/1415/measures/health-matters-and-health-professionals/35-90114)
18 [resources/budget/1415/measures/health-matters-and-health-professionals/35-90114](http://www.humanservices.gov.au/corporate/publications-and-resources/budget/1415/measures/health-matters-and-health-professionals/35-90114).
- 19 Department of Human Services. (2015). "Medicare services." Retrieved 6 January, 2016, from
20 <http://www.humanservices.gov.au/customer/subjects/medicare-services>.
- 21 Dewey, H. M., A. G. Thrift, C. Mihalopoulos, R. Carter, R. A. Macdonell, J. J. McNeil and G. A. Donnan
22 (2004). "'Out of pocket' costs to stroke patients during the first year after stroke—results from the
23 North East Melbourne Stroke Incidence Study." Journal of clinical neuroscience **11**(2): 134-137.
- 24 Duckett, S., A. Boxall, F. Baum, M. Daube, P. Clarke and S. Javanparast (2014). *Federal budget 2014:*
25 *health experts react.* The Conversation, The Conversation Media Group.
- 26 Essue, B., P. Kelly, M. Roberts, S. Leeder and S. Jan (2011). "We can't afford my chronic illness!The
27 out-of-pocket burden associated with managing chronic obstructive pulmonary disease in western
28 Sydney." Journal of health services research & policy **16**: 226-231.

- 1 Farag, I., C. Sherrington, M. Ferreira and K. Howard (2013). "A systematic review of the unit costs of
2 allied health and community services used by older people in Australia." BMC health services
3 research **13**(1): 69.
- 4 Gordon, L. G., M. Ferguson, S. K. Chambers and J. Dunn (2009). Fuel, beds, meals and meds: out-of-
5 pocket expenses for patients with cancer in rural Queensland. Cancer Forum, The Cancer Council
6 Australia.
- 7 Humphreys, J. S. (2009). "Key considerations in delivering appropriate and accessible health care for
8 rural and remote populations: discussant overview." Australian Journal of Rural Health **17**(1): 34-38.
- 9 Hunter, D. J., D. Schofield and E. Callander (2014). "The individual and socioeconomic impact of
10 osteoarthritis." Nature Reviews Rheumatology **10**(7): 437-441.
- 11 Lapsley, H., L. March, K. Tribe, M. Cross, B. Courtenay and P. Brooks (2002). "Living with rheumatoid
12 arthritis: expenditures, health status, and social impact on patients." Annals of the rheumatic
13 diseases **61**(9): 818-821.
- 14 Lehnert, T., D. Heider, H. Leicht, S. Heinrich, S. Corrieri, M. Lupp, S. Riedel-Heller and H.-H. König
15 (2011). "Review: health care utilization and costs of elderly persons with multiple chronic
16 conditions." Medical Care Research and Review **68**(4): 387-420.
- 17 Levesque, J.-F. (2008). Inequalities in access to health care in urban south India. PhD, University of
18 New South Wales.
- 19 Lucas, N., J. K. Bayer, L. Gold, F. K. Mensah, L. Canterford, M. Wake, E. M. Westrupp and J. M.
20 Nicholson (2013). "The cost of healthcare for children with mental health difficulties." Australian and
21 New Zealand Journal of Psychiatry **47**(9): 849-858.
- 22 McRae, I., L. Yen, Y.-H. Jeon, P. M. Herath and B. Essue (2013). "Multimorbidity is associated with
23 higher out-of-pocket spending: a study of older Australians with multiple chronic conditions."
24 Australian journal of primary health **19**(2): 144-149.
- 25 NSW Health. (2015). "Chronic Disease." Retrieved 25 August, 2015, from
26 <http://www.health.nsw.gov.au/aboriginal/Pages/current-chronic-disease.aspx>.
- 27 Organisation for Economic Co-operation and Development (OECD) (2014). Out-of-pocket
28 expenditure on health 2014/15. Health: Key Tables from OECD Paris, OECD.

1 Osborn, R., D. Moulds, D. Squires, M. M. Doty and C. Anderson (2014). "International Survey Of
2 Older Adults Finds Shortcomings In Access, Coordination, And Patient-Centered Care." Health Affairs
3 **33**(12): 2247-2255.

4 Owler, R. (2015). AMA letter to Tony Abbott. T. Abbott. Sydney, SCRIBD.

5 Passey, M., R. Shrestha, M. Bertram, D. Schofield, T. Vos, E. Callander, R. Percival and S. Kelly (2012).
6 "The impact of diabetes prevention on labour force participation and income of older Australians: an
7 economic study." BMC Public Health **12**(16).

8 Rapoport, R., L. E. Tipan and E. Czyzewicz (2013). International Health Policy Survey 2013
9 Methodology Report. Washington, DC, The Commonwealth Fund.

10 Schofield, D., S. Kelly, R. Shrestha, E. Callander, M. Passey and R. Percival (2010). "Long term financial
11 impacts of CVD: Living standards in retirement." International Journal of Cardiology **155**(3): 406-408.

12 Schofield, D., R. Shrestha, M. Passey, A. Earnest and S. Fletcher (2008). "Chronic disease and labour
13 force participation among older Australians." Medical Journal of Australia **189**: 447-450.

14 Senate Standing Committees on Community Affairs (2014). Out-of-pocket costs in Australian
15 healthcare. Canberra, Commonwealth of Australia.

16 Squires, D. (2012). Multinational Comparisons of Health Systems Data, 2012, The Commonwealth
17 Fund.

18 The Commonwealth Fund (2013). 2013 Commonwealth Fund International Health Policy Survey.
19 Washington D.C., The Commonwealth Fund.

20 Yelin, E., P. Katz, D. Lubeck, L. Wanke and M. Buatti (2001). "Impact of etanercept (Enbrel®) on
21 health care use and employment in early RA." Arthritis & Rheumatism **44**(Supplement 9): 152.

22 Yusuf, F. and S. Leeder (2013). "Can't escape it: the out-of-pocket cost of health care in Australia."
23 MJA **199**(7): 475-478.

24

25

1 Table 1: Comparison of reported out-of-pocket costs from the Commonwealth Fund survey with
 2 administrative data

	HEALTHCARE EXPENDITURE PAID BY INDIVIDUALS OUT-OF-POCKET (\$ AUD PPP) SOURCE: OECD HEALTH STATS, 2013, AIHW) TOTAL HEALTHCARE EXPENDITURES PAID BY INDIVIDUALS DIVIDED BY NUMBER OF PEOPLE ALL AGES.	MEAN HOUSEHOLD OUT-OF-POCKET HEALTHCARE EXPENDITURE (AU\$) PPP) SOURCE: CWF SURVEY 2013 - BASED ON ADULTS 18+ BUT RESPONSES FOR 'FAMILY' IN THE PAPER
AUSTRALIA	1,185	990
CANADA	947	1160
FRANCE	421	550
GERMANY	987	720
NETHERLANDS	410	510
NEW ZEALAND	639	750
UNITED KINGDOM	488	330

3

1 Table 2: Average value of annual household out-of-pocket healthcare expenditure
2 corresponding to each expenditure group, from Commonwealth Fund International Health
3 Policy Survey, Australia, 2013

Response category of annual household out-of-pocket healthcare expenditure	Average continuous value of annual household out-of-pocket healthcare expenditure (US\$)
less than \$100	\$46
\$100 to less than \$500	\$249
\$500 to less than \$1,000	\$587
\$1,000 to less than \$2,000	\$1,196
\$2,000 or more	\$4,969

4

5

1 Table 3: Demographic and health characteristics of the weighted respondent population,
 2 Australia, 2013

Characteristic		N	%
		1988	
Sex	Female	1057	52
	Male	931	48
Age	under 45	883	45
	46 to 64	724	37
	65 plus	381	18
Education	Secondary or less	854	47
	More than secondary	1107	53
	Missing	27	
Chronic health conditions ^a	Arthritis	316	17
	Asthma, COPD or chronic lung disease	237	12
	Cancer	85	5
	Mental health	226	12
	Diabetes	150	9
	Heart disease	129	6
	High blood pressure	409	19
	High cholesterol	384	18
Yes, has any chronic condition	981	49	

3 ^arespondents may be in multiple categories of chronic conditions.

4

1 Table 4: Annual household out-of-pocket expenditure on healthcare (AU\$), Australia, 2013

Annual household out-of-pocket healthcare expenditure ranges by year (AU\$)	N	Percent
None	490	26
\$1 to less than \$100	110	5
\$100 to less than \$500	531	27
\$500 to less than \$1,000	322	17
\$1,000 to less than \$2,000	238	11
\$2,000 or more	297	14

2

3

1 Table 5: Annual out-of-pocket healthcare expenditure (AU\$) for Australians by age, gender
 2 and long term health condition

Characteristic	N	Mean annual household out-of-pocket healthcare expenditure (AU\$)	Median annual household out-of-pocket healthcare expenditure (AU\$)
	1990	990	250
<i>Gender</i>			
Male	930	690	250
Female	1060	1260	400
<i>Age group</i>			
Under 45	880	840	250
45 to 64	720	1110	480
65 and over	380	1100	300
<i>Long term health condition</i>			
No condition	960	660	200
Yes, has a condition	980	1340	500
Not sure/decline	40	730	100
<i>Number of conditions</i>			
One	440	1520	520
Two	280	1410	520
Three or more	260	1150	420

3

4

5

1 Table 6: Percentage difference in annual household out-of-pocket healthcare expenditure
 2 (AU\$) for Australians by chronic health condition

Long term health condition (with or without another condition)	Any expenditure (of 1,988 respondents who responded to expenditure question)					Of the 1,498 respondents with expenditure greater than zero		
	N	Mean per year (\$)	Median per year (\$)	Adjusted odds ratio of having any out-of-pocket expenditure*		N	% Difference in healthcare expenditure compared to reference group ^	
				OR (95% CI)	p-value		% difference (95% CI)	p-value
No health condition	964	660	200	REFERENCE		614	REFERENCE	
Arthritis	316	1,220	500	1.92 (0.93 – 3.96)	0.0786	270	59% (15 – 120)	0.0051
Asthma, emphysema, COPD	237	1,640	590	3.97 (2.21 – 7.14)	<.0001	206	109% (50 – 193)	<.0001
Cancer	85	810	250	0.86 (0.26 – 2.82)	0.8046	76	30% (-16 – 100)	0.2424
Depression, anxiety and other mental health conditions	226	1,350	590	2.34 (1.14 – 4.82)	0.0207	200	95% (33– 187)	0.0007
Diabetes	150	1,220	300	1.14 (0.58 – 2.22)	0.1382	124	53% (7 – 120)	0.0197
Heart disease	129	890	500	3.93 (1.73 – 8.93)	0.0011	115	21% (-29 – 107)	0.4799
Hypertension	409	1,030	400	2.12 (1.21 – 3.74)	0.0091	347	42% (6 – 91)	0.0186
High cholesterol	384	1,420	500	2.57 (1.39 – 4.71)	0.0025	338	50% (7 – 111)	0.02

3 Note: Respondents can have multiple conditions, each condition row represents a separate model, comparing people with
 4 that condition, with or without other conditions, to those with no condition.* adjusted for age, education and condition
 5 ^ adjusted for age, education and condition. Limited to those who had more than \$0 in health expenditure

6

1 Table 7: Percentage of Australian adults skipping healthcare treatment due to cost, by type of
 2 chronic health condition

Condition	Percentage skipping care due to cost	Adjusted odds of having skipped care due to cost*	
		OR (95% CI)	p-value
No health condition	9%	REFERENCE	
Arthritis	25%	5.35 (2.73 – 10.46)	<.0001
Asthma, emphysema, COPD	32%	6.16 (3.30 – 11.50)	<.0001
Cancer	21%	1.84 (0.81 – 4.16)	0.1451
Depression, anxiety and other mental health conditions	44%	7.65 (4.13 – 14.20)	<.0001
Diabetes	27%	4.28 (2.05 – 8.92)	0.0001
Heart disease	21%	3.88 (1.82 – 8.28)	0.0004
Hypertension	22%	3.45 (2.01 – 5.91)	<.0001
High cholesterol	22%	3.51 (1.96 – 6.26)	<.0001

3 * Each condition is modelled separately compared to people with no health conditions, with odds ratios adjusted
 4 for age, education and presence of that condition, with or without other conditions.

5


1 Table 8: Estimated yearly household expenditure on healthcare, Australia and 10 countries,
 2 2013 (US\$)

3

Country	Mean household out-of-pocket healthcare expenditure (AU\$ ppp)	Mean household out-of-pocket healthcare expenditure (US\$)	Median household out-of-pocket healthcare expenditure (US\$)	Percentage with a chronic health condition skipping care due to costs	Odds Ratio (95% CI) of skipping care due to costs amongst those with a chronic health condition	p-value
Australia	990	1030	260	25%	REFERENCE	
Canada	1160	760	200	17%	0.54 (0.41 – 0.70)	<.0001
New Zealand	750	490	170	23%	0.81 (0.57 – 1.17)	0.2632
United Kingdom	330	220	0	5%	0.16 (0.09 – 0.27)	<.0001
United States	2800	1840	500	42%	1.97 (1.51 – 2.57)	<.0001
Germany	720	480	130	18%	0.69 (0.49 – 0.97)	0.0327
Netherlands	510	340	260	23%	0.89 (0.63 – 1.25)	0.5068
France	550	360	130	20%	0.71 (0.52 – 0.98)	0.0348
Norway	1260	830	340	11%	0.34 (0.22 – 0.53)	<.0001
Sweden	470	310	150	8%	0.27 (0.19 – 0.38)	<.0001
Switzerland	1310	860	270	16%	0.57 (0.40 – 0.82)	0.0021

4

5



1

2 *Figure 1: Out of pocket expenditure by number of chronic conditions*