

ResearchOnline@JCU

This is the **Accepted Version** of a paper published in the journal Proceedings of the Royal Society of London Series B, Biological Science:

Gangur, Alexander, Smout, Michael, Liddell, Michael, Seymour, Jamie, Wilson, David, and Northfield, Tobin (2017) *Changes in predator exposure, but not diet induce phenotypic plasticity in scorpion venom*. Proceedings of the Royal Society of London Series B, Biological Sciences, 284 (1863).

<http://dx.doi.org/10.1098/rspb.2017.1364>

**Changes in predator exposure, but not diet induce
phenotypic plasticity in scorpion venom**

Journal:	<i>Proceedings B</i>
Manuscript ID	RSPB-2017-1364.R1
Article Type:	Research
Date Submitted by the Author:	14-Aug-2017
Complete List of Authors:	Gangur, Alexander; James Cook University, Centre for Tropical Environmental and Sustainability Studies, College of Science and Engineering Smout, Michael; James Cook University, Centre for Biodiscovery and Molecular Development of Therapeutics, Australian Institute for Tropical Health and Medicine Liddell, Michael; James Cook University, Centre for Tropical Environmental and Sustainability Studies, College of Science and Engineering Seymour, Jamie; James Cook University, Centre for Biodiscovery and Molecular Development of Therapeutics, Australian Institute for Tropical Health and Medicine Wilson, David; James Cook University, Centre for Biodiscovery and Molecular Development of Therapeutics, Australian Institute for Tropical Health and Medicine Northfield, Tobin; James Cook University, Centre for Tropical Environmental and Sustainability Studies, College of Science and Engineering
Subject:	Ecology < BIOLOGY, Biochemistry < BIOLOGY
Keywords:	plasticity, arms race, species interactions, predation
Proceedings B category:	Ecology




Figure 1. Averaged relative venom profiles for each of the four treatments taken: prior to treatments (A, $t = 0$ days), after the treatments ended (B, $t = 49$ days), and again after a recovery period (C, $t = 70$ days) divided into 14 fractions. Relative absorbance measures the absorbance units at 280 nm of any point along the venom profile relative to the point of maximum absorbance in the profile. Venom profiles obtained from scorpions subjected to the pressure for defensive venom (mouse exposure) treatment are given in dark blue (+ pressure for offensive venom, live cricket prey) and light blue (- pressure for offensive venom, dead cricket prey); profiles obtained from scorpions not subjected to this treatment are given in red (+ pressure for offensive venom) and orange (- pressure for offensive venom).








Figure 2. Loadings for principal components 1 and 2, separated into loadings for each scorpion (A) and loadings for each venom fraction (B, 14 total peaks) with fraction number indicated at the tips of each arrow. For clarity and consistency, colours are as per the curves presented in Figure 1; pressure for defensive venom treatment is indicated in dark (+ pressure for offensive venom) and light (- offensive pressure for offensive venom) blue, while data from scorpions not subjected to the pressure for defensive venom treatment are in red (+ pressure for offensive venom) and orange (- pressure for offensive venom). PC1 and PC2 described 53.6% and 25.4% of the overall variation, respectively.

238x132mm (96 x 96 DPI)



1 Changes in predator exposure, but not diet induce phenotypic plasticity in scorpion venom
2 Alex N. Gangur¹, Michael Smout², Michael J. Liddell¹, Jamie E. Seymour², David Wilson²,
3 Tobin D. Northfield¹

4 ¹ *Centre for Tropical Environmental and Sustainability Studies, College of Science and*
5 *Engineering, James Cook University, Cairns, QLD 4878, Australia.*

6 ² *Centre for Biodiscovery and Molecular Development of Therapeutics, Australian Institute*
7 *for Tropical Health and Medicine, James Cook University, Cairns, QLD 4878, Australia*

8

9 Abstract

10

11 Animals embedded between trophic levels must simultaneously balance pressures to deter
12 predators and acquire resources. Venomous animals may use venom toxins to mediate both
13 pressures, and thus changes in this balance may alter the composition of venoms. Basic
14 theory suggests that greater exposure to a predator should induce a larger proportion of
15 defensive venom components relative to offensive venom components, while increases in
16 arms races with prey will elicit the reverse. Alternatively, reducing the need for venom
17 expenditure for food acquisition, for example due to an increase in scavenging, may reduce
18 the production of offensive venom components. Here, we investigated changes in scorpion
19 venom composition using a mesocosm experiment where we manipulated scorpions'
20 exposure to a surrogate vertebrate predator and live and dead prey. After six weeks, scorpions
21 exposed to surrogate predators exhibited significantly different venom chemistry compared to
22 naïve scorpions. This change included a relative increase in some compounds toxic to
23 vertebrate cells, and a relative decrease in some compounds effective against their
24 invertebrate prey. Our findings provide, to our knowledge, the first evidence for adaptive

25 plasticity in venom composition. These changes in venom composition may increase the
26 stability of food webs involving venomous animals.

27

28 *1 Introduction*

29 Interspecific arms races are ubiquitous in ecological communities, and generally involve
30 reciprocal selection pressures that drive the evolution of adaptations and responses between
31 interacting organisms. Organisms investing in traits mediating these arms races need to
32 balance the fitness benefits of winning the race against the cost of maintaining those traits [1].
33 Some organisms defend themselves in arms races using chemical toxins, and these toxins are
34 often produced in low quantities unless induced by exposure to natural enemies to minimize
35 the cost of maintaining unnecessary defences [2]. Similarly, when predators are exposed to
36 prey with varying defensive adaptations, they may develop inducible chemical weapons [3,
37 4]. In venomous animals, the same delivery apparatus evolved for prey capture – such as
38 fangs or a stinger – can also be used to inject chemicals to deter enemies [5], and this dual-
39 purpose nature of the delivery apparatus also extends to the chemistry of the venom itself.
40 Animals generally need to balance arms races involving both predators and prey, and these
41 arms races drive the evolution of venom chemistry in both offensive and defensive contexts
42 [5]. However, physiological differences between predators and prey may necessitate different
43 toxins, and specificity of venom toxins to particular groups of animals has been identified in
44 many venom-users [6-11]. For example, sodium channel blocking α -toxins in scorpions
45 contain three separate subtypes of toxins that are effective against the voltage-gated sodium
46 channels of mammals only, insects only, and both [6]. The whole venom mixture can be
47 thought of as a cocktail of these different toxins, but whether the ‘recipe’ for this cocktail is
48 fixed or can exhibit plasticity in response to different environments and predator/prey
49 interactions remains unclear [12].

50 Broadly speaking, plasticity will be favourable when it enables an organism to have
51 higher fitness across multiple environments, or within a variable environment [13].
52 Theoretical work has shown that plasticity can be selected for when: (a) populations are
53 exposed to multiple environments or variability within an environment, (b) environments
54 produce reliable cues, (c) different phenotypes are favoured in each environment, and (d) no
55 single phenotype exhibits superior fitness across all environments [14-16]. Both plants and
56 animals can in principle exhibit ‘induced’ plasticity to calibrate their defences in response to
57 species interactions [17]. In this context, plasticity may provide a way of saving costs
58 associated with defences, allowing resources to be allocated towards growth and reproduction
59 instead [18]. For example, theory suggests that costly plant defensive compounds should be
60 constitutive (i.e. permanent) where the probability of herbivory is high, while an induced
61 defence in response to attack is optimal when the probability of herbivory is low but the
62 threat of injury from an attack is high [19, 20] Though less well-understood, induced
63 offensive traits enable a predator to capture certain prey more efficiently via plastic change in
64 response to cues signalling the presence of that prey [21]. Induced offences are more
65 favourable when a consumer can benefit from adapting to multiple resource (prey) species
66 with a variety of defences, or resources that can exhibit variable levels of a defence [22, 23].
67 For example, plastic induced ‘offensive traits’ can be seen in feeding-morphologies, such as
68 in snails from the genus *Lacuna*, which change the shape of their teeth to suit their prey [24],
69 and *Nephila pilipes* spiders can plastically modify the composition of their silk chemistry in
70 order to vary the architecture and physical properties of their webs to catch different prey
71 [25].

72 In venom-users, the high cost of chemical warfare has selected for a range of
73 behavioural ‘venom-metering’ strategies, and these plastic behaviours are used to minimize
74 the quantity of venom delivered. Spiders, for instance, may evaluate venom resistance in prey

75 based on olfactory cues and use their venom accordingly [26]. A choice of whether or not to
76 envenomate at all has been shown, based on the relative size and threat posed by the target
77 [27, 28]. Once the decision to envenomate is made, a range of other cues can influence the
78 delivery and volume of venom to minimize venom-use across venomous taxa [12]. By
79 employing similar cues, a venom-user is able to modify the ‘recipe’ of its venom cocktail,
80 thereby optimizing the fitness benefits of its costly venom in different environments
81 exhibiting differences in densities and types of predators and prey [4]. Analogous to induced
82 non-injected defences, a fixed, constitutive venom ‘recipe’ may be more favourable in
83 environments with higher rates of predator attack and lower variability in predator type, while
84 a plastic ‘recipe’ may be more favourable where rates of predator or prey encounters, or
85 predator or prey types, are variable. However, to our knowledge neither induced defensive
86 toxin production nor induced offensive toxin production have been demonstrated in
87 venomous animals.

88 Here, we present an experimental exploration of induced plasticity in the composition
89 of venom produced by scorpions in response to perceived predation risk, and reduced need
90 for venom-use for prey capture. We test the hypotheses that, if induced plasticity of venom
91 composition is exhibited by a model venom-user, higher predation risk will lead to higher
92 relative production of predator-active toxins; and that relative production of prey-active
93 toxins will increase in response to a prey-type that requires greater venom-expenditure to
94 ensure a meal. We did this by manipulating exposure to a surrogate predator and access to
95 live or dead prey, and evaluating changes in the relative concentrations of prey-specific
96 toxins, predator-specific toxins, and general venom compounds. To investigate whether
97 manipulated rates of predator-prey interactions would elicit the plastic changes in venom
98 composition we used the Australian rainforest scorpion *Liocheles waigiensis* (Gervais)
99 (Scorpionoidea: Hemisorpiidae). Our next goal was to evaluate the effects of the predator-

100 specific toxins present in *L. waigiensis* venom on mammalian cells and the prey-specific
101 toxins on crickets that represent their invertebrate prey. Finally, we evaluated whether the
102 relative proportions of vertebrate-toxic venom components would increase in response to a
103 higher rate of vertebrate predator-interactions, and whether the relative proportions of
104 invertebrate-toxic components would decrease in response to a lower need for venom in food
105 consumption, through increased scavenging behaviour.

106

107 ***2 Methods***

108 ***(a) Model organism***

109 The Australian rainforest scorpion *Liocheles waigiensis* (Gervais) (Scorpionoidea:
110 Hemisoriidae) used in our experiments is a common species found in the wet tropics of Far
111 North Queensland. *L. waigiensis* is a generalist predator of invertebrates, including crickets,
112 and is in turn preyed upon by a range of invertebrate and vertebrate predators [29]. Scorpions
113 sourced from rainforest areas around Cairns were individually held in 170 × 110 × 50 mm
114 650 ml plastic containers with one stone and moist organic soil (300 ml Searles Premium
115 Potting Mix brand potting mix) to provide a suitable microclimate for the animal and to aid
116 with moulting. These containers were randomly sorted and stacked two high, in two
117 Wisecube WGC-450 temperature and humidity chambers at 28 °C on a 14/10 light/dark
118 cycle. Relative humidity was maintained at 70%, and after 3 weeks of treatments all
119 scorpions were moved to new containers containing freshly autoclaved soil to reduce fungal
120 growth. All scorpions were maintained in the controlled environment for no more than 5 days
121 prior to the first venom extraction.

122

123 ***(b) Experimental treatments***

124 Given the general predation of invertebrates by *L. waigiensis* in the wild, we used the
125 common house cricket *Acheta domesticus* L. (Insecta: Orthoptera) as a surrogate prey
126 species. To simulate a vertebrate predator sometimes encountered by *L. waigiensis* in the
127 wild, a frozen feeder mouse, *Mus musculus* L. (Rodentia: Muridae), was taxidermied by
128 skinning, stuffing with cotton wool, and articulation with wire, then used as a simulated
129 model vertebrate predator (as in Digweed and Rendall [30]). We used a 2 x 2 factorial design
130 in which without and with pressure for induced offensive venom production (presence of live
131 versus dead prey) was crossed with pressure for defensive venom production (simulated
132 predator exposure). The number of replicates, accounting for scorpions that died during the
133 treatments and were therefore excluded from the analysis, were as follows: 15 (pressure for
134 offensive + defensive venoms), 14 (pressure for offense venom), 14 (pressure for defensive
135 venom), 13 (control).

136 For the prey treatment, scorpions were each fed either a live (pressure for offensive
137 venom) or dead cricket once per week. Live crickets were purchased 1-2 days prior to each
138 feeding, and were killed by freezing for approximately 12 hours. In doing so, the quality of
139 the diet was identical for both groups, but the pressure to use venom to obtain a meal was not.
140 Our taxidermied mouse was used to provoke defensive stings from scorpions in the defensive
141 venom pressure treatment three times a week, except for the first week to allow for
142 acclimation. The mouse was used to continuously probe on the cephalothorax of defensive
143 pressure treatment scorpions for 30 seconds. This stimulus readily stimulated anti-predator
144 responses in the scorpions, including alert and threat postures (with chelae extended and
145 open, and metasoma erect), grappling, pinching, stinging, squirming, and retreat [31, 32]. To
146 ensure that scorpions excluded from the defensive pressure treatment were otherwise equally
147 handled and exposed to laboratory conditions, the containers of these scorpions were opened
148 and exposed to laboratory conditions for 30 seconds. Six weeks after commencement of the

149 experiment, scorpions were subjected to the control (no offensive, no defensive pressures)
150 treatment for one week before venom was extracted. Although this may have diminished the
151 measured effects of the with-pressure treatments due to relaxing of any induced response, a
152 brief recovery time was necessary to ensure sufficient volume of venom had recovered to
153 perform chemical analyses.

154

155 *(c) Venom extraction*

156 We ran the treatments for 42 days so that the experiments lasted twice as long as the venom
157 regeneration time of 21 days, according to previous analyses [29]. Venom was first extracted
158 within 5 days of collection, and then again after a week of rest, following the end of the
159 experiment, 49 days later. Venom was then extracted from all scorpions a third time, 21 days
160 after the experimental treatments ceased, to assess how it had changed in the absence of
161 offensive and defensive pressures. By providing the scorpions with the full length of time
162 necessary to regenerate their venom, we ensured that there was ample time for the treatments
163 to elicit a response in the chemistry of the regenerated venom.

164

165 *(d) Venom analysis*

166 Venom was collected using an Arthur H. Thomas Co. Z789 Square Wave Stimulator to
167 electrostimulate the telson at approximately 25 volts (5.5 pulses/sec, for 15 milliseconds per
168 pulse). Extracted venoms were diluted in 150 μ L of degassed phosphate buffered solution
169 (PBS- Life Technologies), centrifuged for 10 minutes total at 32,000 RPM, and filtered
170 through a syringe-driven 4mm 0.22 μ m filter (Millipore). Venom profiles were obtained
171 using size-exclusion fast protein liquid chromatography (FPLC) using a Superdex™ 75
172 10/300 (Tricorn) GL Column (13 μ m, 10mm \times 300mm – GE Healthcare) at 4 °C with 100%
173 PBS buffer at 0.50 ml/minute with 0.5 ml elutions for 45 mL on an ÄKTA™ FPLC (GE

174 Healthcare). Venom components were detected by absorbance measured at a wavelength of
175 280 nm. Finally, using venom collected from the same scorpions, but three weeks after the
176 cession of the experimental treatments, we evaluated the toxicity of each venom fraction by
177 performing toxicity assays on a human cardiac cell line to test for vertebrate toxicity (see
178 section f), and by performing behavioural assays on crickets to test for effects on temporary
179 or permanent invertebrate paralysis (see section g).

180

181 *(e) Statistical analysis for profile changes*

182 To compare between the venom profiles from each treatment, we split the FPLC venom
183 profile into 14 different ‘fractions’, and differences in the amounts of each relative to the
184 other treatments could then be evaluated statistically. To identify the different fractions, we
185 first standardised each chromatogram to an area under the curve of 1, obtained the mean
186 chromatogram for each of the four treatments by averaging all the curves within each
187 treatment. We next fitted a spline curve to each of these mean chromatograms using the
188 smooth.spline function in R [33], with the smoothing parameter, $\lambda = 0.5$ [33, 34]. The local
189 minima in these splines were then designated as boundaries between two fractions. Local
190 minima within 1 ml from each other were averaged to create a single break between fractions,
191 with one exception: the local minima values of 38.64, 38.99, 39.78, 40.51, and 40.57 ml were
192 divided into the two groups: 38.64, and 38.99; and 39.78, 40.51, and 40.57 ml, for which
193 each was averaged to describe the combined fraction separation point. Principal component
194 analysis (PCA) was then used to describe these 14 fractions across the data set [34].
195 MANOVA and two separate, follow-up ANOVA analyses were performed to evaluate
196 treatment effects on the first two principal components. These analyses were conducted on
197 venom samples collected at three time points: prior to the initiation of treatments ($t = 0$ days),

198 at the cessation of the treatments ($t = 49$ days), and 21 days after the cessation of treatments (t
199 $= 70$ days).

200 To evaluate the effects of predator and prey main effects on particular peaks, we
201 calculated the mean and 95% confidence intervals for each peak evaluating the difference
202 between the predator-no predator treatment means, or the live prey – dead prey treatment
203 means. The 95% confidence intervals were calculated using non-parametric bootstrapping
204 with 10,000 simulations. For each simulation, we resampled with replacement the
205 absorbances for a particular treatment (e.g., with or without simulated predator exposure) and
206 chemical fraction. Chemical fractions five through eight were not easily distinguishable and
207 likely represent a number of compounds, and therefore, we also calculated the mean
208 treatment effects and 95% confidence intervals for the sum of these fractions (summed
209 individually for each scorpion). Treatment effects were considered significant for $\alpha = 0.05$
210 when 95% confidence intervals did not overlap zero.

211

212

213 ***(f) Predator cell assays***

214 The biological consequences of observed changes to the venom profiles were evaluated using
215 toxicity assays. Fraction concentrations were determined using the A280 method [35]. A
216 human cardiac cell line (ScienCell) was used as a vertebrate assay, following Schneider [29].
217 Vertebrate cells were maintained and assays were performed as previously described by
218 Andreosso, Smout [36] and Chaousis, Smout [37]. An xCELLigence SP RTCA system
219 (ACEA Biosciences) with an E-plate seeded with 150 μL cardiac media (ScienCell) and 5000
220 human cardiac cells were incubated overnight at 37 $^{\circ}\text{C}$ and 5 % CO_2 .

221 The cell response to each fraction (20 μL) and 100% PBS solution (control) was
222 measured by the xCELLigence system as changes to cell index. Cell response is a

223 combination measure of changes in media conductivity or cell contact/toxicity, which varies
224 as the cells deform in response to exposure to a chemical sample. The cell index readouts
225 were blanked against the PBS control, and the maximum drop value in 2 hours after venom
226 addition was deemed the predator cell response. The relative response to whole venom as a
227 percentage was then used to graph the activity level of the venom peaks. We used two-tailed
228 t-tests to compare the response of each venom fraction to the PBS control to identify peaks
229 that significantly altered media conductivity or cell contact/toxicity.

230

231 ***(g) Prey toxicity assays***

232 *Acheta domesticus* cricket assays were performed by evaluating whether a given venom
233 fraction was active towards immature crickets. To evaluate the effects of each fraction, 3 μ L
234 of one of the 14 chemical fractions was injected ventrally into the pronotum of an immature
235 cricket varying in mass from 0.1 to 0.2 grams. Immediately after injection, the cricket was
236 inserted into a clean, 9-dram clear styrene tube with snap-on lid and rolled onto its back every
237 10 seconds 18 times for a total of 3 minutes. A compromised righting response was recorded
238 when a cricket was unable to right itself within 60 seconds of being rolled onto its dorsal side.
239 Each fraction was replicated with 10 crickets. We used a 2 by 2 Fisher's exact test to compare
240 the cricket response from each venom fraction to a PBS control.

241

242 ***3 Results***

243 ***(a) Effects of predator-prey interactions on venom composition***

244 The experimental results were used to evaluate our hypotheses on venom plasticity using a
245 model animal the rainforest scorpion *Liocheles waigiensis*. The venom profiles obtained from
246 venom extraction before the experimental treatments began were not significantly different
247 from each other (Figure 1A, see Electronic Supplementary Material 1 for statistical analysis).

248 After treatment there was a difference between the venom profiles of the predator-treated and
249 the predator-excluded scorpions. These profiles varied greatly in the relative concentration of
250 multiple chemical fractions, with the greatest difference in treatments occurring in peak
251 fraction 12 (Figure 1B). From the principal component analysis we obtained two major
252 principal components, PC1 and PC2, which explained 53.6% and 25.4% of the overall
253 variation, respectively. Venom profiles obtained from scorpions that were and were not
254 subjected to the defensive pressure treatment were found to be significantly different using a
255 MANOVA to evaluate the treatments on the principal component weightings (Table 1), with
256 increased predator exposure leading to lower and higher values of PC1 and 2, respectively
257 (Figure 2). This was most clearly associated with changes in fraction 12, which was reduced
258 in the scorpions exposed to simulated predators (Figure 1B, 2), which significantly decreased
259 with predator exposure (Figure 3). Fraction 2 also varied strongly in both principal
260 components, but not in a way that was interpretable with the experimental treatments (Figure
261 2). There were no interaction effects, nor any significant effects from prey manipulation
262 treatment (live versus dead prey) on the venom profile principal components 1 and 2 (Table
263 1). Profiles obtained after a 21-day recovery period following cessation of treatments exhibited
264 similar patterns of difference between treatments (Figure 1B, C, ESM 1), but the magnitude
265 difference was reduced.

266

267 ***(b) Toxicity assays***

268 Higher activity towards mammalian cells (>60%) was generally found in the toxin fractions
269 containing larger proteins/peptides (fractions 2,3,4,5,7, and 8, Figure 3, ESM 1), which were
270 likely 3-25 kDa due to the Superdex™ 75 resin that was used [37]. One section of the profile
271 contained many fractions (5-8) that were not easily distinguishable, significantly increased in
272 response to simulated predator exposure (95% bootstrap confidence limits: 0.004, 0.083), as

273 did fractions 7 and 8 when evaluated individually (Figure 3). In addition, fractions 10 and 11
274 had some activity against mammalian cells, although the magnitude of these effects were
275 much lower than for other fractions (Figure 3, ESM 1). Toxicity towards crickets was
276 generally found in a fraction containing larger proteins/peptides (fractions 3-4), and the
277 fractions containing smaller compounds (fractions 8-14) (Figure 3A, ESM 1). Components of
278 each of these sections were reduced in response to simulated predator exposure (Figure 3). It
279 should be noted that many small molecules are not detectable at 280 nm and other detection
280 methods may be required. Undetected compounds were the likely source of activity against
281 crickets (>90%) exhibited by fractions 13 and 14, as the absorbance trace showed very
282 minimal contents. Example cell responses are provided in ESM 2.

283

284 *(c) Comparing treatment effects with toxicity assays*

285 Simulated predator exposure had the strongest effect on reducing the relative
286 production of fraction 12 that demonstrated activity on crickets, and to a lesser extent,
287 reduced the relative production of a section (fractions 3-4) that exhibited effects on both
288 crickets and mammalian cells (Figure 3). Fractions 5-8, which were not easily distinguishable
289 (Figure 1), significantly increased in response to simulated predator exposure (predator
290 treatment effect 95% limits: 0.003670351, 0.082905484) and portions of this section of the
291 profile exhibited activity on mammalian cells (Figure 3B). Fractions 1 and 11 slightly
292 increased in the presence of predators (Figure 3C), but neither of these had strong effects on
293 invertebrates or mammals (Figure 3A,B). The presentation of live versus dead prey had little
294 effect on the relative production of each chemical fraction, although it did slightly increase
295 the production of fraction 3 (ESM 1) that effected both crickets and mammalian cells (Figure
296 3A,B).

297

298 **4 Discussion**

299 Given current theory relating selection pressures to plastic changes in defence and
300 reproductive investment (e.g., Peacor, Peckarsky [38]), venomous mesopredators should shift
301 the balance of venom composition towards the defensive components when predator
302 exposure increases. In line with these predictions, we found evidence for a plastic change in
303 venom composition in response to increased perceived predation risk, showing for the first
304 time to our knowledge that organismal venom chemistry can change in response to a threat.
305 These changes imply a rerouting of resource expenditure, which may be nutritional or
306 energetic [12], to increase relative production of other venom fractions which are responsible
307 for toxicity to vertebrates. Overall, simulated predator exposure appeared to decrease relative
308 production of strong invertebrate toxins, while generally increasing the production of a
309 section of the venom profile with activity towards mammalian cells. These results suggest for
310 the first time to our knowledge that venoms can serve as inducible defences used against
311 predators. Inducible defence theory suggests that plastic defences are more likely to evolve in
312 highly variable or cyclic environments, where the fitness benefits of flexibility outweigh the
313 costs of maintaining this capacity for variability [2, 13] Venomous animals evolve vast,
314 complex armouries of peptides and proteins in their venoms [12], and it would appear that *L.*
315 *waigiensis* is able to modify the production of a subset of their complex venom cocktail to
316 suit a changing environment. The magnitude of the pressure to minimize venom cost and the
317 predatory pressure may also relate to how closely venom production tracks the rate of
318 ecological dynamics [17].

319 Resource type did not elicit a response in venom chemistry. This lack of effect, which
320 was probed through removal of the need for venom expenditure, may have been due to a)
321 insufficient variation in resource type, b) a time-lag in the scorpions' response which as a
322 result was not detected, or c) an absence of inducible offence. It is well-documented that the

323 magnitude of environmental variability can influence both the magnitude and the speed of a
324 plastic response [39]. For example, moderate levels of herbivore damage may only induce a
325 ‘primed’ state in plants rather than the immediate chemical response to high damage, while
326 low levels of damage may fail to provoke a plastic response at all [2]. In our experiment, we
327 introduced two resource ‘types’ (live or dead) to represent variation in the need for venom
328 during prey-capture. However, if the live resource type was not sufficiently different to the
329 dead resource type (i.e. often not requiring venom-use to obtain a meal) than any inducible
330 offence in the venom profile may not have been provoked. We tried to account for this by
331 feeding larger prey (i.e. larger in size than a scorpion’s chelae) to encourage the need for
332 envenomation following van der Meijden, Coelho [32], but scorpions were still occasionally
333 observed to be killing their prey without stinging. Secondly, there may have been a time-lag
334 in any potential response to the treatment. In plants, induced chemical defences can be
335 mounted in response to attack, followed by a substantially longer ‘relaxation’ period before
336 returning to a ground state. For example, *Trifolium repens* mounts a systemic chemical
337 defence within 51 hours of herbivory, but requires at least 28 days to relax (Gomez et al.
338 2010). Similarly, an induced offense in response to prey-type may exhibit a relaxation period.
339 For example, in snails from the genus *Lacuna*, the longer an individual fed on previous diet,
340 the slower its induced morphological offense switched to a new food source [24]. Finally,
341 there may be no plastic response to variation in resource-type. This may be due to either
342 insufficient variation in prey-type in the wild to drive the evolution of a plastic response
343 capability, resembling the conditions under which constitutive defences are favoured by
344 plants, or due to sufficiently high fluctuations in prey-type to favour a bet-hedging strategy
345 rather than plasticity [40]. When traits respond to a selective pressure, evolution balances this
346 response between optimising the trait for the maximum fitness benefit and over-investing in
347 the trait to compensate for the effect of environmental stochasticity [15, 41]. Such bet

348 hedging strategies are ubiquitous in arms races [42], and may also be seen in venom-users
349 [43]. In the absence of an alternate prey which doesn't require stinging to be subdued, it may
350 be favourable to delay a plastic response (or exhibit none at all) and continue producing
351 costly venom even in the absence of live prey to ensure success in future opportunities to
352 catch a meal. Future work may be able to distinguish between these competing explanations
353 by investigating the variability of food resources in the natural habitat of *L. waigiensis*.

354 Induced plastic defences can stabilize populations against fluctuating predatory
355 pressures [17], and as such in ecological communities where venomous animals provide an
356 important food resource (e.g. [44]) induced defences could act as an important stabilizing
357 force for the community and diminish trophic cascades in food webs. Adaptive plasticity can
358 mitigate the effects of sudden disturbances by allowing populations to evolve sufficiently
359 quickly to survive abrupt change [45]. Phenotypic plasticity permits more time for
360 evolutionary adaptation to occur, and may reduce the degree of evolutionary change
361 necessary to track a moving optimal trade-off between the costs and benefits of venom
362 production [45]. Indeed, populations which exhibit greater phenotypic plasticity are generally
363 able to evolve more under global change and thereby adapt to changing environments [46].
364 However, plasticity may also slow the rate of evolutionary pressure by reducing the selection
365 pressure for genetic change [16]; whether or not venom plasticity should facilitate or inhibit
366 adaptation by venom-users to modified predator-prey interactions driven by environmental
367 change remains an open question.

368 In our bioassays, we found some venom fractions (e.g., fraction 3) have activity
369 against both the scorpion's cricket prey and mammalian heart cells, suggesting they may
370 serve to improve both prey capture and defence against predators. This may lead to
371 complicated tritrophic interactions where phenotypic changes in response to one arms race
372 (e.g., with predators) can alter the investment in another arms race (e.g., with prey) (Gangur

373 et al. in review). Furthermore, we identified fractions of the venom profile (e.g. fraction 1)
374 that increased in response to simulated predator exposure, but in isolation did not demonstrate
375 activity against mammalian heart cells. These fractions may have effectiveness against
376 another vertebrate biological pathway (e.g., pain activation), may interact with other fractions
377 to improve potency, or may be increased incidentally due to physiological constraints in
378 venom production. Thus, further research to clarify the role of these venom fractions in
379 predator defence may shed light on adaptive advantage of the observed phenotypic changes
380 in response to simulated predator exposure. Furthermore, we have only evaluated the effects
381 of each venom fraction on two distantly related taxa (mammalian cells and arthropods). In
382 some cases, organisms can target phenotypic changes in defence to the specific threats e.g.
383 [47, 48]. Further research exposing these scorpions to a range of predator species and
384 evaluating changes in venom composition may elucidate the specificity of this phenotypic
385 plasticity.

386

387 Venom research has historically been intently focused on human toxicity, due to obvious
388 reasons, the prevention of mortalities and a strong interest in medical advancements [49]. The
389 ecological and evolutionary perspectives that have been increasingly explored, particularly in
390 the last decade, offer critical insights into venomous animal ecology that has improved health
391 outcomes as well as enriched our understanding of venom-use and production. Indeed, if
392 plastic responses are widespread in venomous animals, antivenom production may be
393 improved by accounting for this potential source of variation by ensuring live prey or
394 simulated predation [50]. Furthermore, the potential role of venom in stabilising ecological
395 dynamics needs to be further explored as in some cases this may be a substantial factor
396 controlling community structure.

397

398 *Data Accessibility: Venom extraction and assay data: Dryad doi:10.5061/dryad.sq2g4 [51].*

399 *Additional statistical analyses supporting this article have been uploaded as part of the*

400 *supplementary material.*

401 *Ethics: There were no live vertebrate animals used as experimental subjects in this study.*

402 *Authors' Contributions: Experiments were designed by ANG, TDN, and MJL, JES, and conducted by*

403 *ANG. Chemical analyses were conducted by ANG, MS, and DW. Mammalian and invertebrate*

404 *bioassays were conducted by MS and TDN, respectively. ANG and TDN conducted statistical*

405 *analyses with input from MJL and JES. The first draft was prepared by ANG and TDN, with all others*

406 *contributing extensively to subsequent drafts.*

407 *Funding: This study was supported by JCU internal research allocations to TN, ML, and JS. Funding*

408 *support for MS and DW was provided by Australian Institute of Tropical Health and Medicine*

409 *research support.*

410 *Competing Interests: We declare we have no competing interests.*

411 *Acknowledgements: We'd like to specially think Samantha Barnett for helping out with laboratory*

412 *protocols; Tim Jenkins, Dani Brewster-O'Brien, and David Clarke for helping with scorpion*

413 *collection and venom extraction, and Nalisa Neuendorf and John Doolan for helping with venom*

414 *extraction; and two anonymous referees for their comments, which greatly improved the manuscript.*

415 REFERENCES

- 416 1. Wade M.J. 2007 The co-evolutionary genetics of ecological communities. *Nat Rev Genet* **8**,
417 185-195.
- 418 2. Karban R. 2011 The ecology and evolution of induced resistance against herbivores. *Funct*
419 *Ecol* **25**, 339-347.
- 420 3. Kopp M., Tollrian R. 2003 Reciprocal phenotypic plasticity in a predator-prey system:
421 inducible offences against inducible defences? *Ecol Lett* **6**, 742-748.
- 422 4. Mougi A., Kishida O. 2009 Reciprocal phenotypic plasticity can lead to stable predator-prey
423 interaction. *J Anim Ecol* **78**, 1172-1181.
- 424 5. Casewell N.R., Wüster W., Vonk F.J., Harrison R.A., Fry B.G. 2013 Complex cocktails: the
425 evolutionary novelty of venoms. *Trends Ecol Evol* **28**, 219-229.
- 426 6. Quintero-Hernández V., Jiménez-Vargas J., Gurrola G., Valdivia H., Possani L. 2013 Scorpion
427 venom components that affect ion-channels function. *Toxicon* **76**, 328-342.
- 428 7. McClounan S., Seymour J. 2012 Venom and cnidome ontogeny of the cubomedusae
429 *Chironex fleckeri*. *Toxicon* **60**, 1335-1341.
- 430 8. Yang S., Liu Z., Xiao Y., Li Y., Rong M., Liang S., Zhang Z., Yu H., King G.F., Lai R. 2012 Chemical
431 Punch Packed in Venoms Makes Centipedes Excellent Predators. *Mol Cell Proteomics* **11**, 640-650.
- 432 9. Dutertre S., Jin A.-H., Vetter I., Hamilton B., Sunagar K., Lavergne V., Dutertre V., Fry B.G.,
433 Antunes A., Venter D.J. 2014 Evolution of separate predation-and defence-evoked venoms in
434 carnivorous cone snails. *Nat Commun* **5**, 3521.
- 435 10. Zlotkin E., Menashé M., Rochat H., Miranda F., Lissitzky S. 1975 Proteins toxic to arthropods
436 in the venom of elapid snakes. *J Insect Physiol* **21**, 1605-1611.
- 437 11. Nicholson G.M. 2007 Insect-selective spider toxins targeting voltage-gated sodium channels.
438 *Toxicon* **49**, 490-512.
- 439 12. Morgenstern D., King G.F. 2013 The venom optimization hypothesis revisited. *Toxicon* **63**,
440 120-128.

- 441 13. Bradshaw A.D. 1965 Evolutionary significance of phenotypic plasticity in plants. *Adv Genet*
442 **13**, 115-155.
- 443 14. Via S., Lande R. 1985 Genotype-environment interaction and the evolution of phenotypic
444 plasticity. *Evolution* **39**, 505-522.
- 445 15. Scheiner S.M. 1993 Genetics and evolution of phenotypic plasticity. *Annu Rev Ecol Syst* **24**,
446 35-68.
- 447 16. Ghalambor C.K., McKay J.K., Carroll S.P., Reznick D.N. 2007 Adaptive versus non-adaptive
448 phenotypic plasticity and the potential for contemporary adaptation in new environments. *Funct*
449 *Ecol* **21**, 394-407.
- 450 17. Cortez M.H. 2011 Comparing the qualitatively different effects rapidly evolving and rapidly
451 induced defences have on predator-prey interactions. *Ecol Lett* **14**, 202-209.
- 452 18. Stamp N. 2003 Out of the quagmire of plant defense hypotheses. *Q Rev Biol* **78**, 23-55.
- 453 19. Karban R., Agrawal A.A., Thaler J.S., Adler L.S. 1999 Induced plant responses and information
454 content about risk of herbivory. *Trends Ecol Evol* **14**, 443-447.
- 455 20. Ito K., Sakai S. 2009 Optimal defense strategy against herbivory in plants: Conditions
456 selecting for induced defense, constitutive defense, and no-defense. *J Theor Biol* **260**, 453-459.
- 457 21. Takatsu K., Kishida O. 2013 An offensive predator phenotype selects for an amplified
458 defensive phenotype in its prey. *Evol Ecol* **27**, 1-11.
- 459 22. Banerji A., Morin P.J. 2009 Phenotypic plasticity, intraguild predation and anti-cannibal
460 defences in an enigmatic polymorphic ciliate. *Funct Ecol* **23**, 427-434.
- 461 23. Karban R., Agrawal A.A. 2002 Herbivore offense. *Annu Rev Ecol Syst* **33**, 641-664.
- 462 24. Padilla D.K. 2001 Food and environmental cues trigger an inducible offence. *Evol Ecol Res* **3**,
463 15-25.
- 464 25. Blamires S.J., Piorkowski D., Chuang A., Tseng Y.-H., Toft S., Tso I.-M. 2015 Can differential
465 nutrient extraction explain property variations in a predatory trap? *R Soc Open Sci* **2015**.

- 466 26. Hostettler S., Nentwig W. 2006 Olfactory Information Saves Venom during Prey-Capture of
467 the Hunting Spider *Cupiennius salei* (Araneae: Ctenidae). *Funct Ecol* **20**, 369-375.
- 468 27. Edmunds M.C., Sibly R.M. 2010 Optimal sting use in the feeding behavior of the scorpion
469 *Hadrurus spadix*. *J Arachnol* **38**, 123-125.
- 470 28. Rein J.O. 1993 Sting use in two species of Parabuthus scorpions (Buthidae). *J Arachnol* **4**, 60-
471 63.
- 472 29. Schneider I. 2011 Venom Ecology in the Australian rainforest scorpion *Liocheles waigiensis*.
473 [Honours Thesis]. Cairns, QLD, James Cook University.
- 474 30. Digweed S.M., Rendall D. 2009 Predator-associated vocalizations in North American red
475 squirrels, *Tamiasciurus hudsonicus*: are alarm calls predator specific? *Anim Behav* **78**, 1135-1144.
- 476 31. Warburg M.R. 1998 Qualitative and quantitative analysis of intra-and interspecific
477 behavioural patterns among scorpions. *J Ethol* **16**, 115-121.
- 478 32. van der Meijden A., Coelho P.L., Sousa P., Herrel A. 2013 Choose Your Weapon: Defensive
479 Behavior Is Associated with Morphology and Performance in Scorpions. *PLoS one* **8**, e78955.
- 480 33. R Development Core Team. 2014 R: A language and environment for statistical computing.
481 (Vienna, Austria, R Foundation for Statistical Computing.
- 482 34. Wehrens R. 2011 *Chemometrics with R: multivariate data analysis in the natural sciences*
483 *and life sciences*, Springer.
- 484 35. Olson B.J.S.C., Markwell J. 2007 Assays for Determination of Protein Concentration. In
485 *Current Protocols in Protein Science*, 3.4.1-3.4.29, John Wiley & Sons, Inc.
- 486 36. Andreosso A., Smout M.J., Seymour J.E. 2014 Dose and time dependence of box jellyfish
487 antivenom. *J Venom Anim Tox incl Trop Dis* **20**, 34.
- 488 37. Chaousis S., Smout M., Wilson D., Loukas A., Mulvenna J., Seymour J. 2014 Rapid short term
489 and gradual permanent cardiotoxic effects of vertebrate toxins from *Chironex fleckeri* (Australian
490 box jellyfish) venom. *Toxicon* **80**, 17-26.

- 491 38. Peacor S.D., Peckarsky B.L., Trussell G.C., Vonesh J.R. 2013 Costs of predator-induced
492 phenotypic plasticity: a graphical model for predicting the contribution of nonconsumptive and
493 consumptive effects of predators on prey. *Oecologia* **171**, 1-10.
- 494 39. DeWitt T.J., Sih A., Wilson D.S. 1998 Costs and limits of phenotypic plasticity. *Trends Ecol*
495 *Evol* **13**, 77-81.
- 496 40. Tufto J. 2015 Genetic evolution, plasticity, and bet-hedging as adaptive responses to
497 temporally autocorrelated fluctuating selection: A quantitative genetic model. *Evolution* **69**, 2034-
498 2049.
- 499 41. Rosenheim J.A. 2011 Stochasticity in reproductive opportunity and the evolution of egg
500 limitation in insects. *Evolution* **65**, 2300-2312.
- 501 42. Beaumont H.J.E., Gallie J., Kost C., Ferguson G.C., Rainey P.B. 2009 Experimental evolution of
502 bet hedging. *Nature* **462**, 90-93.
- 503 43. Hayes W.K. 1992 Prey-handling and envenomation strategies of prairie rattlesnakes
504 (*Crotalus v. viridis*) feeding on mice and sparrows. *J Herpetol* **26**, 496-499.
- 505 44. Polis G. 1990 Ecology. In *The Biology of Scorpions* (ed. Polis G.), pp. 247-293. Palo Alto,
506 Stanford University Press.
- 507 45. Reed T.E., Schindler D.E., Waples R.S. 2011 Interacting effects of phenotypic plasticity and
508 evolution on population persistence in a changing climate. *Conserv Biol* **25**, 56-63.
- 509 46. Schaum C.E., Collins S. 2014 Plasticity predicts evolution in a marine alga. *Proc R Soc B* **281**,
510 20141486.
- 511 47. Relyea R.A. 2001 Morphological and behavioural plasticity of larval Anurans in response to
512 different predators. *Ecology* **82**, 523-540.
- 513 48. Voswamatjam D.V., Narwani A.J.T., Thaler J.S. Specificity in Induced Plant Responses Shapes
514 Patterns of Herbivore Occurrence on *Solanum dulcamara*. *Ecology* **86**, 886-896.

- 515 49. Fry B.G., Roelants K., Champagne D.E., Scheib H., Tyndall J.D., King G.F., Nevalainen T.J.,
516 Norman J.A., Lewis R.J., Norton R.S. 2009 The toxicogenomic multiverse: convergent recruitment of
517 proteins into animal venoms. *Annu Rev Genom Hum Genet* **10**, 483-511.
- 518 50. Richards D.P., Barlow A., Wüster W. 2012 Venom lethality and diet: Differential responses of
519 natural prey and model organisms to the venom of the saw-scaled vipers (*Echis*). *Toxicon* **59**, 110-
520 116.
- 521 51 Gangur A., Smout M., Liddell M.J., Seymour J.E., Wilson D., Northfield, T.D. 2017 Data from:
522 Changes in predator exposure, but not diet induce phenotypic plasticity in scorpion venom. Dryad
523 Digital Repository. (doi:10.5061/dryad.sq2g4).

524 **TABLES**

525

526 **Table 1:** MANOVA results from venom collected one week after the cessation of the
 527 experimental treatments, which demonstrated a significant overall difference between the
 528 fraction loadings from scorpions that were and were not subjected to the defensive pressure
 529 treatment. There was no significant interaction effect.

Source	d.f.	Pillai	<i>F</i>	Df den	<i>P</i>
Offence	1	0.040	0.978	2, 47	0.383
Defence	1	0.242	7.517	2, 47	0.001
Defence × Offence	1	0.036	0.887	2, 47	0.419
Residuals	48				

530

531 **Table 2:** ANOVA results from venom collected one week after cessation of experiments,
 532 demonstrating significantly different fraction loadings between the scorpions that were and
 533 were not subjected to the defensive pressure treatment along both PC1 and PC2. There were
 534 no significant interaction effects.
 535

Source	d.f.	M Sq	F	P
PC 1				
<i>Prey</i>	1	0.012	1.933	0.171
<i>Predator</i>	1	0.048	7.643	0.008
<i>Predator</i> × <i>Prey</i>	1	0.001	0.170	0.682
<i>Residuals</i>	48	0.006		
PC 2				
<i>Prey</i>	1	0.001	0.173	0.679
<i>Predator</i>	1	0.018	5.928	0.019
<i>Predator</i> × <i>Prey</i>	1	0.005	1.744	0.193
<i>Residuals</i>	48	0.003		

536

537

538 **FIGURE LEGENDS**

539 Figure 1. Averaged relative venom profiles for each of the four treatments taken: prior to
540 treatments (A, t = 0 days), after the treatments ended (B, t = 49 days), and again after a
541 recovery period (C, t = 70 days) divided into 14 fractions. Relative absorbance measures the
542 absorbance units at 280 nm of any point along the venom profile relative to the point of
543 maximum absorbance in the profile. Venom profiles obtained from scorpions subjected to the
544 pressure for defensive venom (mouse exposure) treatment are given in dark blue (+ pressure
545 for offensive venom, live cricket prey) and light blue (- pressure for offensive venom, dead
546 cricket prey); profiles obtained from scorpions not subjected to this treatment are given in red
547 (+ pressure for offensive venom) and orange (- pressure for offensive venom).

548 Figure 2. Loadings for principal components 1 and 2, separated into loadings for each
549 scorpion (A) and loadings for each venom fraction (B, 14 total peaks) with fraction number
550 indicated at the tips of each arrow. For clarity and consistency, colours are as per the curves
551 presented in Figure 1; pressure for defensive venom treatment is indicated in dark (+ pressure
552 for offensive venom) and light (- offensive pressure for offensive venom) blue, while data
553 from scorpions not subjected to the pressure for defensive venom treatment are in red (+
554 pressure for offensive venom) and orange (- pressure for offensive venom). PC1 and PC2
555 described 53.6% and 25.4% of the overall variation, respectively.

556 Figure 3. Invertebrate (A) and vertebrate (B) toxicity assay results. Invertebrate toxicity was
557 measured by evaluating the proportion of crickets (10 crickets per treatment) that were
558 paralysed for longer than 60 seconds. Statistically significant difference from the control was
559 evaluated using a Fisher's Exact Test for each peak (ESM 1). Vertebrate toxicity was
560 evaluated by measuring vertebrate cell response to venom fractions relative to whole venom
561 response using the xCELLigence platform. Due to small sample volume, it was not possible

562 to completely separate fraction 5 from fraction 6. Statistically significant difference from the
563 PBS control (blanked at 0) was evaluated using a two-tailed t-test for each peak (ESM 1).
564 Panel C presents mean (and 95% non-parametric bootstrap confidence intervals) for the
565 difference between the treatments with and without simulated predator exposure. Confidence
566 intervals entirely above (or below) zero suggest significant effects of increased (or decreased)
567 production after simulated predator exposure. Differences were calculated after the
568 treatments ended (t = 49 days). Asterisks represent confidence intervals that do not overlap
569 zero. Chemical fractions five through eight were not easily distinguishable, and likely
570 represent multiple similarly sized compounds. Therefore, we have also calculated this
571 confidence interval separately (predator treatment effect 95% limits: 0.004, 0.083). Panel D
572 presents the relationship between invertebrate and vertebrate toxicity for each peak. The
573 feature scaling function $x' = \frac{x - \min(x)}{\max(x) - \min(x)}$ was used to convert max drop value into a
574 normalized vertebrate assay score in the range [0,1] for ease of comparison (invertebrate
575 assays were already scored in this range as a proportion of crickets out of 10 replicates
576 experiencing paralysis for >60 seconds after toxin injection). Error bars indicate standard
577 error. Due to low yield volume, the vertebrate assay score for fraction 5 includes both
578 fractions 5 and 6. Asterisks indicate significance for alpha = 0.05.