

The Handbook of Microbial Metabolism of Amino Acids

EDITED BY
J.P.F. D'MELLO



CABI

1 Structural and Functional Properties of Glutamate Dehydrogenases

S. Brown^{1*} and D.C. Simcock^{1,2,3}


¹The Deviot Institute, Deviot, Tasmania, Australia; ²Institute of Food, Nutrition and Human Health, Massey University, Palmerston North, New Zealand; ³Faculty of Medicine and Bioscience, James Cook University, Cairns, Australia

1.1 Abstract

Glutamate dehydrogenases are homooligomeric enzymes that catalyse the reversible oxidative deamination of glutamate to 2-oxoglutarate, thereby linking nitrogen metabolism and the tricarboxylic acid cycle. Four different classes of the enzyme are known in microbes that differ in size and cofactor specificity. Despite some common features, the kinetic properties of the enzymes from different species vary considerably. We consider the relationship between the sequences, those structures that are available and the enzyme kinetics in the context of the metabolic role of the enzyme.

1.2 Introduction

Glutamate dehydrogenase (GDH) catalyses the reversible reductive amination of 2-oxoglutarate to glutamate:



The enzyme may be NAD⁺ specific (EC 1.4.1.2), NADP⁺ specific (EC 1.4.1.4) or able to use either cofactor (EC 1.4.1.3). Irrespective of the cofactor, the standard transformed Gibbs

free energy of reaction, $\Delta_r G'^0 \approx -40 \text{ kJ mol}^{-1}$ (Olson and Anfinsen, 1953; Strecker, 1953; Engel and Dalziel, 1967; Subramanian, 1978; Cook *et al.*, 1980) and the reaction operates in either direction depending on the relative magnitudes of

$$Q = \frac{[\text{Glutamate}][\text{NAD(P)}^+]}{[\text{2-Oxoglutarate}][\text{NAD(P)H}][\text{NH}_4^+]}$$

where Q is the reaction quotient and

$$K' = \exp(-\Delta_r G'^0/RT) \approx 10^{-7}$$

where K' is the apparent equilibrium constant, R is the gas constant and T is the absolute temperature. If $Q < K'$, the reaction will tend to form glutamate.

In bacteria, GDH can be involved in the assimilation of NH_4^+ (Kanamori *et al.*, 1987; Bruggeman *et al.*, 2005), in glutamate synthesis (Helling, 1994) and in controlling the intracellular concentration of 2-oxoglutarate to ensure the operation of 2-oxoglutarate dehydrogenase (ODH) and the whole tricarboxylic acid (TCA) cycle (Struhl and Maganasik, 1976; Janes *et al.*, 2001; Doucette *et al.*, 2011). We consider the structural and kinetic properties of the different forms of GDH, their taxonomic distribution and the roles of the enzyme in the metabolism of bacteria.

*Corresponding author: E-mail address: Simon.Brown@deviotinstitute.org

1.3 Enzyme Structure

Four broad classes of GDH, designated GDH1–4 (Andersson and Roger, 2003), have been distinguished using structural and functional criteria. Both GDH1 and GDH2 have monomers of about 50 kDa and are distinguished by their cofactor specificity: GDH1 is NAD⁺ specific, whereas GDH2 is NADP⁺-specific. In contrast, the monomers of GDH3 and GDH4 are of about 115 and 180 kDa, respectively (Fig. 1.1), but are not as well characterized.

Both GDH1 and GDH2 are about 450 residues in length, are highly conserved and are much more similar to one another than they are to GDH3 and GDH4 (Fig. 1.2). In the alignment on which Fig. 1.2 is based, the identity among the 14 GDH1 and GDH2 sequences shown is 21–89% (40–96% similarity; average overall distance = 0.80 substitutions per site), whereas the seven GDH3 sequences have 17–65% identity (32–79% similarity; mean overall distance = 1.22 substitutions per site) and the seven GDH4 sequences are more highly conserved (31–65% identity; 47–77% similarity; overall mean distance = 0.88 substitutions per site).

1.3.1 Substrate and cofactor binding

The residues involved in binding the substrates or the cofactor are mostly completely conserved among GDH1 and GDH2 sequences (Fig. 1.3),

and some of these are also conserved in GDH3 and GDH4 sequences. Of particular note among the latter are the lysine residue K126 and the aspartate residue D166, which, in concert with the α -amino group of the substrate glutamate and a water oxygen, form a tetrad capable of a wide range of hydrogen bonding patterns that may be catalytically significant (Fisher and Maniscalco, 2002).

The determinants of the cofactor specificity of GDH have not yet been identified unambiguously. It has been proposed that the Gly-X-Gly-X-X-Gly/Ala motif in the P-loop of the Rossmann fold is involved in determining the cofactor specificity of dehydrogenases (Bellamacina, 1996). An alanine in the final position (P6 or the alanine A244 in *Escherichia coli*) is supposed to favour NADP⁺ binding, whereas a glycine favours NAD⁺ binding, but the *Clostridium symbiosum* GDH1 has an alanine and the *Psychrobacter* sp. TAD1 GDH2 has a serine, so neither of these conform to this model. No consensus about P6 is apparent from the GDH3 and GDH4 sequences. Similarly, the presence of a negatively charged group near the 2'-OH and 3'-OH of adenosine (P7 or D263 in *E. coli*) has been suggested to favour NAD⁺ rather than NADP⁺ binding. However, as Sharkey *et al.* (2013) observed, there is an aspartate (or a glutamate in *Peptoniphilus asaccharolyticus*) in this position in most GDH sequences, with the exceptions of the *C. symbiosum* GDH1 and *Pseudomonas aeruginosa* GDH4. Based on the modelling of NADP⁺ binding to GDH2, it has been suggested that NADP⁺




Fig. 1.1. The domains of representative bacterial glutamate dehydrogenases (GDHs). The *Peptoniphilus asaccharolyticus* (AAA25611.1), *Escherichia coli* (P00370.1), *Desulfovibrio vulgaris* (ABM29040.1) and *Pseudomonas aeruginosa* (WP_053816968.1) enzymes are examples of GDH1, GDH2, GDH3 and GDH4, respectively. The domains identified using Pfam (the protein families database; Finn *et al.*, 2014) and CATH (class, architecture, topology, homology – a hierarchical protein domain classification system; Cuff *et al.*, 2011) are an NAD-binding Rossmann-like domain (use of 'NAD' intends no implication on cofactor specificity), a bacterial GDH domain (BAC) and a domain common to glu/leu/phe/val dehydrogenases (the substrate-binding domain, ELFV). For the *P. aeruginosa* sequence, Pfam identified a more extensive BAC domain (the dashed rectangle) than did CATH (the black rectangle).




Fig. 1.2. The bootstrap consensus phylogram derived by maximum likelihood from an alignment of 27 glutamate dehydrogenase (GDH) amino acid sequences (375 gap-free positions) using molecular evolutionary genetics analysis (MEGA5; Tamura *et al.*, 2011). The maximum likelihood was based on the JTT (Jones, Taylor, Thornton) matrix-based model (Jones *et al.*, 1992) using 500 bootstrap replicates (Felsenstein, 1985). The branch lengths (number of substitutions per site) are indicated below the branches unless the value was less than 0.1. The shape parameter of the discrete gamma distribution was 1.637.

specificity may be conferred by: (i) three positively charged residues (lysine K286, and arginines R289, R292 in the *E. coli* enzyme); and (ii) a serine (S264 in *E. coli*). In this model, these four residues are adjacent to the 2'-phosphate of NADP⁺ and stabilize the negative charge (Sharkey *et al.*, 2013). While K286 is almost completely conserved among GDH1 and GDH2, and S264 is also highly conserved (although neither is present in the *Burkholderia thailandensis* sequence), R289 and R292 are both absent from the NADP⁺-specific *Psychrobacter* enzyme (Di Fraia *et al.*, 2000). Overall, as Sharkey *et al.* (2013) suggest, cofactor specificity cannot be reliably inferred from primary sequence.

1.3.2 Tertiary and quaternary structure

As with other dehydrogenases, GDH has a substrate-binding domain (ELFV), which is similar to those of other amino acid dehydrogenases, and an NAD(P)⁺-binding domain (NAD), both of which can be identified from sequence homology alone (Fig. 1.2). Crystal structures are available for only GDH1 and GDH2, but these are similar to those of the human and bovine enzymes, except for the absence of the antenna domain. In each case, the ELFV domain consists of a series of β -strands sandwiched between α -helices and the NAD domain is a modified Rossmann fold. The nucleotide binds




Fig. 1.3. Summary of the structural properties of the NAD⁺-specific glutamate dehydrogenase (GDH) from *Burkholderia thailandensis* and the mutability of GDH1 and GDH2 sequences. The triangles (▲, ▼, △) indicate residues that are hydrogen bonded to glutamate, 2-oxoglutarate or NAD⁺ based on the crystal structures of the *B. thailandensis* (4XGI) and *Clostridium symbiosum* (1BGV) enzymes. The circles (●) indicate the approximate position of the residues involved in determining the NADP⁺-specificity of the *Escherichia coli* enzyme (Sharkey *et al.*, 2013). The secondary structure (4XGI) indicates only the positions of α -helices and β -strands (black and white rectangles, respectively). The mutability was calculated as described previously (Brown *et al.*, 2014) from the sequence alignment of the GDH1 and GDH2 sequences identified in Fig. 1.2. Abbreviations: ELFV, a domain common to glu/leu/phe/val dehydrogenases (the substrate-binding domain); NAD, a NAD-binding Rossmann-like domain (use of 'NAD' intends no implication on cofactor specificity).

in a cleft between the domains and is ligated by residues from each (Fig. 1.3).

The substrate and cofactor are necessarily close together. In both the bovine enzyme and in *B. thailandensis* GDH1, the substrate binds towards the narrow end of the cleft and NAD⁺ binds further towards the wider end of the cleft with the nicotinamide moiety oriented towards the substrate. In each case, they are so close that the 3'-OH of NAD⁺ and the carboxyl group of the substrate are both H bonded to a single asparagine residue.

Most microbial GDHs are homohexamers, although the GDH2 of *Streptomyces fradiae* (Vančurová *et al.*, 1989), the GDH3s of *Agaricus bisporus* and *Neurospora crassa* (Smith *et al.*, 1975) and the GDH4 of *P. aeruginosa* (Lu and Abdelal, 2001) are reportedly tetramers. Unusually, the GDH4 of *Psychrobacter* is dimeric and while some kinetic data have been reported (Camardella *et al.*, 2002), the primary sequence is not yet available. The crystal structures that are available exhibit D₃ symmetry in which monomers are located at the vertices of triangles and these are stacked in pairs with the monomers

approximately aligned. In each monomer the cleft between the domains opens away from, and the nicotinamide moiety of NAD⁺ is oriented towards, the threefold axis. It is inevitable that the tetrameric GDHs will have very different quaternary structures.

1.4 Enzyme Mechanism and Kinetics

The kinetics of GDH are complex, both because of the number of reactants and products (Eqn 1.1) and because of the variety of behaviours observed (Barton and Fisher, 1971; Bailey *et al.*, 1982; Syed and Engel, 1987). For some forms of the enzyme the usual rectangular hyperbolic relationship between initial rate (v_0) and substrate concentration (s) is observed, but cooperative kinetics have also been reported (Table 1.1). Reports of non-linear double reciprocal, Eadie-Hofstee or other transformed plots are common for both the mammalian (Barton and Fisher, 1971; Bailey *et al.*, 1982) and microbial enzymes (Table 1.1).

Table 1.1. Kinetic characteristics of some microbial glutamate dehydrogenases (GDHs). Where cooperativity is reported, the Hill coefficient (n_H) is given below the K_m or $s_{0.5}$ in parentheses.

Species	Cofactor	K_m or $s_{0.5}$ (n_H)				
		2-OG ^a (mM)	Glu ^b (mM)	NH ₄ ⁺ (mM)	NAD(P) ⁺ (μ M)	NAD(P)H (μ M)
GDH1						
<i>Bacillus cereus</i> DSM 31 (Jahns and Kaltwasser, 1993)	NAD ⁺	^c	7.4	96	560	33
<i>Bacillus subtilis</i> ISW1214 (Khan <i>et al.</i> , 2005)	NAD ⁺	0.65	0.34	55.6	80	70
<i>Bacteroides thetaiotaomicron</i> (Glass and Hylemon, 1980)	NAD ⁺		12.3		610	
	NADP ⁺	0.24	3.7	5.0	40	17
<i>Peptostreptococcus asaccharolyticus</i> (Hornby and Engel, 1984)	NAD ⁺	0.82	6.6	18.4	31	66
<i>Psychrobacter</i> sp. TAD1 (Camardella <i>et al.</i> , 2002)	NAD ⁺	2.36	28.6	24.6	500	40 (1.8)
<i>Pyrobaculum islandicum</i> (Kujo and Ohshima, 1998)	NAD ⁺	0.066	0.17	9.7	25	5.0
	NADP ⁺				240	270
GDH2						
<i>Escherichia coli</i> B/r (Sakamoto <i>et al.</i> , 1975)	NADP ⁺	0.64	1.3	1.1	42	40
<i>Psychrobacter</i> sp. TAD1 (Di Fraia <i>et al.</i> , 2000)	NADP ⁺	6.8 (3.6)	67.9 (1.2)	4.5 (1.1)	52 (0.8)	23 (2.3)
<i>Ruminococcus flavefaciens</i> (Duncan <i>et al.</i> , 1992)	NADP ⁺	0.41	62.19	19.22	–	34
<i>Saccharomyces cerevisiae</i> (DeLuna <i>et al.</i> , 2001)	NADP ⁺	1.27 (1.5)	6.36	5.00	10.5	331
<i>Saccharomyces cerevisiae</i> (DeLuna <i>et al.</i> , 2001)	NADP ⁺	0.29 (1.3)	9.79	5.96	14.1	113
GDH3						
<i>Benjaminiella poitrassi</i> (Joshi <i>et al.</i> , 2013)	NAD ⁺	3.12	33.3	27.7	170	60
<i>Agaricus bisporus</i> (Kersten <i>et al.</i> , 1999)	NAD ⁺	3.5	37.1	6.5	46	60
<i>Neurospora crassa</i> (Smith <i>et al.</i> , 1975)	NAD ⁺	4.6	5.5	17	330	550
GDH4						
<i>Janthinobacterium lividum</i> (Kawakami <i>et al.</i> , 2007, 2010)	NAD ⁺	9.1 (1.7)	7.1	20.9	2100	68 (1.8)
<i>Pseudomonas aeruginosa</i> (Lu and Abdelal, 2001)	NAD ⁺	–	4.4 (1.7)	–	130	–
<i>Streptomyces clavuligerus</i> (Miñambres <i>et al.</i> , 2000)	NAD ⁺	1.33 (1.5)	2.27 (1.39)	33.9	51.16 (1.48)	19.87 (1.52)

^a2-OG, 2-oxoglutarate.^bGlu, glutamate.^cResponse is sigmoidal, but no data are reported.

Rife and Cleland (1980b) proposed a model of the catalytic mechanism of GDH which is largely consistent with the crystal structures that have been found (Stillman *et al.*, 1993) and subsequent analyses of the kinetics (Fisher, 2005). The model shown in Fig. 1.4 is essentially that of Rife and Cleland (1980b) for bovine liver

GDH. The enzyme-bound 2-oxoglutarate is aminated, transiently forming a quaternary complex (ERNK in Fig. 1.4), to a 2-carbinolamine (2-hydroxy-2-iminoglutarate). This is dehydrated to 2-iminoglutarate which is then reduced to glutamate. At this point, the complex is protonated and then water is released




Fig. 1.4. Catalytic mechanism of bovine glutamate dehydrogenase (GDH) adapted from Rife and Cleland (1980b), Maniscalco *et al.* (1996) and Fisher (2005). The sequence of binding of NADH and 2-oxoglutarate is not shown. The structural fragments above each intermediate represent C1 and C2 of the ligand starting with 2-oxoglutarate (left) and ending with glutamate (right). The standard abbreviations used below the fragments are: C, 2-carbinolamine; E, enzyme; G, glutamate; I, 2-iminoglutarate; K, 2-oxoglutarate; N, NH_4^+ ; O, NAD^+ ; R, NADH.

from bovine liver GDH, although these steps are reversed in the *C. symbiosum* enzyme (Maniscalco *et al.*, 1996), before glutamate is released (Fig. 1.4).

The most commonly used initial rate (v_0) equation for GDH is the empirical expression

$$\frac{e}{v_0} = \phi_0 + \frac{\phi_1}{s_1} + \frac{\phi_2}{s_2} + \frac{\phi_3}{s_3} + \frac{\phi_{12}}{s_1 s_2} + \frac{\phi_{13}}{s_1 s_3} + \frac{\phi_{23}}{s_2 s_3} + \frac{\phi_{123}}{s_1 s_2 s_3} \quad (1.2)$$

or (see Eqn 1.3 at bottom of the page):

where e is the concentration of GDH, the s_i (s_1, s_2, s_3) are the concentrations of NAD(P)H , NH_4^+ and 2-oxoglutarate, and different combinations of rate and dissociation constants are distinguished by particular subscripts to ϕ . In Eqns 1.2 and 1.3, e/ϕ_0 is the maximum velocity (V_{\max}), the K_m for s_i is ϕ_i/ϕ_0 and all substrates are independent and have rectangular hyperbolic kinetics. This expression is based on a general random binding order model involving the formation of a quaternary complex, but it includes, as special cases, other models, including those with specific binding sequences (Dalziel, 1969). For example, some authors

have concluded that there was, in at least one direction, an ordered sequence of binding to the bovine enzyme (Frieden, 1959; Rife and Cleland, 1980a), whereas others have argued that the data are consistent with the random binding order and have obtained estimates of all eight parameters (Engel and Dalziel, 1970). The implication of the mechanism shown in Fig. 1.4 is that binding is at least partially ordered: NH_4^+ binds after NAD(P)H and 2-oxoglutarate and, in the opposite direction, glutamate binds after NAD(P)^+ .

The parameters of Eqn 1.2 are often obtained sequentially from linearized plots (Dalziel, 1969) which necessarily introduces bias into the estimates and may explain, in part, why it is sometimes difficult to obtain values for some parameters (Engel and Dalziel, 1970). Unsurprisingly, it is also clear that Eqn 1.2 cannot account for the reported cooperative kinetics (Di Fraia *et al.*, 2000; Miñambres *et al.*, 2000) of GDH and does not account for substrate inhibition (Sharkey and Engel, 2008) or the activation by high NAD^+ concentrations (Olson and Anfinsen, 1953; Engel and Dalziel, 1970). Other empirical expressions have been devised to account for some of these phenomena, and of these three

$$v_0 = \frac{s_1 s_2 s_3 e}{\phi_0 s_1 s_2 s_3 + \phi_1 s_2 s_3 + \phi_2 s_1 s_3 + \phi_3 s_1 s_2 + \phi_{12} s_3 + \phi_{13} s_2 + \phi_{23} s_1 + \phi_{123}} \quad (1.3)$$

are of particular interest. First, Engel and Ferdinand (1973) proposed a rational rate equation:

$$v_0 = \frac{\sum_{j=1}^q D_j s^j e}{N_0 + \sum_{i=1}^q N_i s^i}, \quad (1.4)$$

which can describe an almost unlimited range of behaviour, as has been considered in some detail (Bardsley and Childs, 1975; Bardsley *et al.*, 1980; Brown and Simcock, 2013). Second, Kurganov (2000) proposed that the K_m varies continuously with v_0/V_{\max} and modified the standard Michaelis–Menten expression accordingly:

$$v_0 = \frac{V_{\max} s}{\left(K_0 + (K_{\text{lim}} - K_0) \frac{v_0}{V_{\max}} \right) + s}, \quad (1.5)$$

where K_0 and K_{lim} are the effective values of K_m as v_0/V_{\max} approaches 0 and 1, respectively. This can be written as

$$s = (K_0 - K_{\text{lim}}) \frac{v_0}{V_{\max}} + K_{\text{lim}} \frac{v_0}{V_{\max} - v_0}, \quad (1.6)$$

so if $K_{\text{lim}} < K_0$, the linear v_0 - s regime dominates and the transition to the saturated regime occurs over a small range of s . If $K_{\text{lim}} > K_0$, a biphasic double reciprocal plot is generated which is reminiscent of those often reported for GDH (Barton and Fisher, 1971; Bailey *et al.*, 1982; Syed and Engel, 1987). Both Eqns 1.4 and 1.5 account for only one of the substrates and, like Eqn 2, their main purpose is to provide a means of summarizing particular aspects of the kinetics of GDH. This weakness is overcome, to some extent, by the third example, Eqn 1.6 – the expression devised by Barton and Fisher (1971) to account for the kinetics of the oxidative deamination reaction. Then again, this involves 13 parameters, few of which have any obvious mechanistic significance. A complete kinetic model is yet to be devised for the enzyme.

Despite rate equations such as shown in Eqns 1.2–1.5, the usual practice has been to estimate the K_m (or $s_{0.5}$ where there is cooperativity) and V_{\max} by independently varying the concentration of each reactant. The small selection of data in Table 1.1 prompts hypothesis that there might be some differences between GDHs. For example, it appears that the K_m values for both NH_4^+ and NAD(P)^+ may tend to be larger in

GDH1 than in GDH2 (Table 1.1). While this is merely speculation, careful analysis of the available data is warranted.

1.5 Genomics

The taxonomic distribution of the forms of GDH is uneven. Both GDH1 and GDH2 are widespread among bacteria and fungi, but GDH3 and GDH4 have more restricted distributions (Andersson and Roger, 2003). Based on the data available in Pfam (the protein families database; Finn *et al.*, 2014), homologues of GDH4 (based on the presence of the BAC domain shown in Fig. 1.1, Pfam accession code PF05088) can be found among the α -, β - and γ -proteobacteria (149, 43 and 146 species, respectively), and in the actinobacteria (84 species), as well as in a small number of fungi (11 species) and metazoa (five species). Among the bacteria that have GDH4 homologues are a number of *Mesorhizobium* spp., *Sinorhizobium* spp. and *Bradyrhizobium* spp., and some, such as *Sinorhizobium meliloti* (Fig. 1.2), also have other forms of GDH. However, rhizobia generally lack significant GDH activity (Ali *et al.*, 1981; Bravo and Mora, 1988; Rossi *et al.*, 1989), and several species of bacteria lack any GDH sequence (Andersson and Roger, 2003).

One of the many challenges of genomic analysis is the determination of the presence or absence of a gene (Cordwell, 1999; Osterman and Overbeek, 2003), but it is just one of many. Even if it can be confirmed that a gene occurs in a genome, that need not mean that it is expressed or, if it is, that the product has only the anticipated consequence. The identification of any homologue of a gene can be an even greater challenge if only because of insufficient data. For example, GDH4 is quite different from GDH1 (see Fig. 1.1) and so a database search may not identify all of the four forms of the enzyme. Depending on the circumstances, even the detection of a particular enzyme activity need not be unambiguous. For example, ODH was reported to be present in *Mycobacterium tuberculosis* (Cordwell, 1999) based on the presence of the genes and the oxidation of the substrate, but even if the genes are expressed they need not be incorporated into ODH (Tian *et al.*, 2005b) and the apparent ODH activity can be due to other enzymes entirely

(Tian *et al.*, 2005a). Further, there is the growing realization that an enzyme might have more than one function. For example, mammalian GDH may link lysosomes to microtubules (Rajas *et al.*, 1996) and bind mRNA (Preiss *et al.*, 1995), perhaps regulating the translation of proteins targeted to the mitochondria, while the aconitase apoprotein is involved in the regulation of iron metabolism (Beinert *et al.*, 1997; Brzóska *et al.*, 2006).

Nevertheless, the presence or absence of an enzyme such as GDH is biochemically and physiologically significant because it links carbon and nitrogen metabolism (Fig. 1.5). Notwithstanding, the TCA cycle has many variants (Weitzman, 1981; Huynen *et al.*, 1999) and so this metabolic intersection can be especially complex. Bearing in mind the intrinsic difficulties already described, several species do not have genes encoding any of the GDHs (Andersson and Roger, 2003), including *Aquifex aeolicus*, *Archaeoglobus*

fulgidus, *Borrelia burgdorferi*, *Methanobacterium thermoautotrophicum* and *Mycoplasma pneumoniae* (Cordwell, 1999). All five of the named species just listed also lack ODH activity, and two of them, *B. burgdorferi* and *M. pneumoniae*, do not have isocitrate dehydrogenase (IDH) activity either (Cordwell, 1999). This type of listing can be extended indefinitely (see Table 1.2). While such a list does not show the extent to which alternative reactions can substitute for any of these enzymes, it does prompt consideration of the possible metabolic variants.

1.6 Metabolic Context

GDH is an important link between amino acid and carbon metabolism through the TCA cycle (Fig. 1.5). Specifically, the 2-oxoglutarate produced by IDH (EC 1.1.1.42), is a substrate for




Fig. 1.5. Glutamate dehydrogenase (GDH) is the link between the tricarboxylic acid (TCA) cycle and nitrogen metabolism. The oxidative branch of the TCA cycle and several reactions involving glutamate and 2-oxoglutarate are shown, but the reactions that actually operate vary with the conditions and between species, as discussed in the text. Abbreviations: IDH, isocitrate dehydrogenase; ODH, 2-oxoglutarate dehydrogenase; OGC, 2-oxoglutarate decarboxylase; SCoAS, succinyl-CoA synthetase; SSADH, succinate semialdehyde dehydrogenase.

Table 1.2. The presence (+) or absence (–) of glutamate dehydrogenase (GDH), isocitrate dehydrogenase (IDH), 2-oxoglutarate dehydrogenase (ODH) and/or succinyl-CoA synthetase (SCoAS) in several species for which a complete genome sequence is available. The data for IDH, OGDH and SCoAS were adapted from Cordwell (1999).

Species	GDH	IDH	ODH	SCoAS
<i>Aquifex aeolicus</i>	–	+	–	+
<i>Archaeoglobus fulgidus</i>	–	+	–	+
<i>Bacillus subtilis</i>	+	+	+	+
<i>Borrelia burgdorferi</i>	–	–	–	–
<i>Chlamydia trachomatis</i>	–	–	+	+
<i>Corynebacterium glutamicum</i>	+	+	–	+
<i>Escherichia coli</i>	+	+	+	+
<i>Haemophilus influenzae</i>	+	–	+	+
<i>Helicobacter pylori</i>	+	+	–	–
<i>Methanobacterium thermoautotrophicum</i>	–	+	–	+
<i>Methanococcus jannaschii</i>	?	+	–	+
<i>Mycobacterium tuberculosis</i>	+	+	–	?
<i>Mycoplasma genitalium</i>	?	–	–	–
<i>Mycoplasma pneumoniae</i>	–	–	–	–
<i>Pyrococcus horikoshii</i>	+	–	–	–
<i>Synechocystis</i> sp. PCC6803	+	+	–	+
<i>Treponema pallidum</i>	?	–	–	–

both GDH and the ODH complex, which consists of multiple copies of a thiamine pyrophosphate-dependent dehydrogenase (E1; EC 1.2.4.2), dihydrolipoamide succinyltransferase (E2; EC 2.3.1.61) and dihydrolipoamide dehydrogenase (E3; EC 1.8.1.4). While GDH catalyses a reversible reaction, the citrate synthase, IDH and ODH reactions are irreversible. In fact, 2-oxoglutarate is among the key metabolites in the metabolic network of *E. coli* and glutamate is the most highly connected metabolite in this network (Wagner and Fell, 2001).

The interaction of the great variety of enzymes and reactions of the TCA cycle observed among microbes (Weitzman, 1981; Huynen *et al.*, 1999) with GDH has profound metabolic implications. Considered simply, the three enzymes directly involved in 2-oxoglutarate metabolism are IDH, ODH and GDH. In *E. coli*, IDH and GDH2 are NADP⁺ dependent with $K_m = 17 \mu\text{M}$ NADP⁺ (Hurley *et al.*, 1996) and $42 \mu\text{M}$ NADPH (Table 1.1), respectively, while ODH is NAD⁺ dependent with $K_m = 400 \mu\text{M}$ NAD⁺, and is also subject to product inhibition ($K_i = 1 \text{ mM}$) (Schmincke-Ott and Bisswanger, 1981). However, this difference in cofactor specificity is less significant than it might seem because *E. coli* has two transhydrogenases, one soluble (Cao *et al.*,

2011), and the other in the plasma membrane (Bizouarn *et al.*, 2000), that keep $[\text{NAD}^+]/[\text{NADH}] \approx 19$ and $[\text{NADP}^+]/[\text{NADPH}] \approx 1.2$ (Andersen and von Meyenburg, 1977). The K_m for 2-oxoglutarate is similar for GDH2 (0.64 mM; Table 1.1) and ODH (0.1 mM; Gupta and Dekker, 1980), but GDH2 also requires NH_4^+ ($K_m = 1.1 \text{ mM}$; Table 1.1), so the flux through ODH would tend to be greater than that through GDH2.


Under aerobic conditions, *E. coli* operates a complete TCA cycle, but in anaerobic conditions it lacks ODH activity and the pathway operates as two branches: an oxidative branch leading to the production of 2-oxoglutarate and a reductive branch yielding succinate (Hager and Kornberg, 1961; Amarasingham and Davis, 1965). *E. coli* only assimilates NH_4^+ under aerobic conditions and GDH is involved in this process in energy- or glucose-limited conditions, when the concentration of NH_4^+ is sufficiently high, whereas glutamine synthetase (GS; EC 6.3.1.2) and glutamate synthase (GOGAT, E. C. 1.4.1.13) are responsible in energy-rich conditions (Helling, 1994, 1998). In glucose-limited conditions, GDH and ODH share the available 2-oxoglutarate.

However, when ODH activity is low and 2-oxoglutarate accumulates, GDH2 can utilize the available NH_4^+ and the NADPH generated by


IDH to produce glutamate (if $Q < K'$). For example, the *E. coli* mutant 309-1, which constitutively lacks the capacity to oxidize 2-oxoglutarate, excretes glutamate when grown aerobically on glucose in the presence of succinate (Davis *et al.*, 1959; Shiiro *et al.*, 1961). A similar *Enterobacter aerogenes* mutant produces 2-oxoglutarate from glucose in anaerobic conditions (Stouthamer, 1966). Conversely, if the 2-oxoglutarate supply is low, then GDH2 can oxidatively deaminate glutamate (using NADP⁺; $K_m = 40 \mu\text{M}$) to produce more, providing that $Q > K'$. It should be noted though that while glutamate is an effective inducer of ODH activity (Amarasingham and Davis, 1965) and *E. coli* can use glutamate as a sole nitrogen source, the maximum growth rate is low whether or not glucose is limiting (Halpern and Umbarger, 1961; Senior, 1975).

Corynebacterium glutamicum (formerly *Brevibacterium flavum* and *B. lactofermentum*) has significant NADP⁺-dependent GDH activity (Shiiro *et al.*, 1959), which is presumably GDH2 (Shiiro and Ozaki, 1970), along with appreciable IDH activity (Shiiro *et al.*, 1961), but its ODH activity declines during growth (Kawahara *et al.*, 1997) and, in at least some circumstances, it can be undetectable (Shiiro *et al.*, 1961). During aerobic growth on glucose, 2-oxoglutarate, or in the presence of NH₄⁺, glutamate accumulates (Shiiro *et al.*, 1959), and this can be substantially reduced by biotin supplementation (Shiiro *et al.*, 1962). Deleting the gene for the E1 enzyme of ODH completely inactivates ODH and has little effect on the accumulation of glutamate (Asakura *et al.*, 2007), whereas deletion of the GDH gene reduces the intracellular concentration of glutamate and increases those of NH₄⁺ and 2-oxoglutarate (Müller *et al.*, 2006).

The oxidation of citrate and isocitrate by *M. tuberculosis* can be estimated from the rate of O₂ uptake, but this requires NADP⁺ and is independent of CoA (Murthy *et al.*, 1962). This is inconsistent with the operation of ODH, which requires NAD⁺ and CoA, and arises from the replacement of ODH and succinyl CoA synthetase (SCoAS; EC 6.2.1.4) by a combination of reactions catalysed by 2-oxoglutarate decarboxylase (OGC; EC 4.1.1.71)



and succinate semialdehyde dehydrogenase (SSADH; EC 1.2.1.16)



(Tian *et al.*, 2005a). The K_m for 2-oxoglutarate of OGC is 0.48 mM (Tian *et al.*, 2005a) is much lower than that of *M. smegmatis* (5 mM; Sarada *et al.*, 1980), the only GDH2 that has been characterized from *Mycobacterium* spp., which, if a similar situation pertains in *M. tuberculosis*, would tend to favour the operation of OGC/SSADH rather than GDH. While these, and other (Piston *et al.*, 1999), alternative reactions are interesting, what is remarkable is that *M. tuberculosis* has genes encoding the enzymes of ODH, but that they are used for other purposes (Maksymiuk *et al.*, 2015). This reinforces a point made previously: it is unwise to infer the presence of a gene product or an expected biological function from the presence of a gene.

1.7 Conclusions

Of the four forms of GDH in microbes, GDH1 and GDH2 have been well characterized, but GDH3 and GDH4 are less well understood. Crystal structures are available for GDH1 and GDH2 and much work has been done to analyse the kinetics of these enzymes, but there is little information on GDH3 or GDH4, and much remains to be done to generate a coherent whole from the rapidly growing volume of data. Some of the difficulty arises from the extraordinary metabolic and physiological flexibility of bacteria and fungi, the consequence of which is that genera (and sometimes species) have less in common than might be anticipated. This leads to the conclusion that more data are only part of the answer, and that it is also necessary to develop the means to integrate the data in meaningful ways.

References

- Ali, H., Niel, C. and Guillaume, J.B. (1981) The pathways of ammonium assimilation in *Rhizobium meliloti*. *Archives of Microbiology* 129, 391–394.
- Amarasingham, C.R. and Davis, B.D. (1965) Regulation of α -ketoglutarate dehydrogenase formation in *Escherichia coli*. *The Journal of Biological Chemistry* 240, 3664–3668.
- Andersen, K.B. and von Meyenburg, K. (1977) Charges of nicotinamide adenine nucleotides and adenylate energy charge as regulatory parameters of the metabolism in *Escherichia coli*. *The Journal of Biological Chemistry* 252, 4151–4156.
- Andersson, J.O. and Roger, A.J. (2003) Evolution of glutamate dehydrogenase genes: evidence for lateral gene transfer within and between prokaryotes and eukaryotes. *BMC Evolutionary Biology* 3:14.
- Asakura, Y., Kimura, E., Usuda, Y., Kawahara, Y., Matsui, K., Osumi, T. and Nakamatsu, T. (2007) Altered metabolic flux due to deletion of *odhA* causes L-glutamate overproduction in *Corynebacterium glutamicum*. *Applied and Environmental Microbiology* 73, 1308–1319.
- Bailey, J., Bell, E.T. and Bell, J.E. (1982) Regulation of bovine glutamate dehydrogenase. The effects of pH and ADP. *The Journal of Biological Chemistry* 257, 5579–5583.
- Bardsley, W.G. and Childs, R.E. (1975) Sigmoid curves, non-linear double-reciprocal plots and allosterism. *Biochemical Journal* 149, 313–328.
- Bardsley, W.G., Leff, P., Kavanagh, J. and Waight, R.D. (1980) Deviations from Michaelis–Menten kinetics. The possibility of complicated curves for simple kinetic schemes and the computer fitting of experimental data for acetylcholinesterase, acid phosphatase, adenosine deaminase, arylsulphatase, benzylamine oxidase, chymotrypsin, fumarase, galactose dehydrogenase, β -galactosidase, lactate dehydrogenase, peroxidase and xanthine oxidase. *Biochemical Journal* 187, 739–765.
- Barton, J.S. and Fisher, J.R. (1971) Nonlinear kinetics of glutamate dehydrogenase. Studies with substrates – glutamate and nicotinamide-adenine dinucleotide. *Biochemistry* 10, 577–585.
- Beinert, H., Holm, R.H. and Münck, E. (1997) Iron–sulfur clusters: nature's modular, multipurpose structures. *Science* 277, 653–659.
- Bellamacina, C.R. (1996) The nicotinamide dinucleotide binding motif: a comparison of nucleotide binding proteins. *FASEB Journal* 10, 1257–1269.
- Bizouarn, T., Fjellström, O., Mueller, J., Axelsson, M., Bergkvist, A., Johansson, C., Karlsson, B.G. and Rydström, J. (2000) Proton translocating nicotinamide nucleotide transhydrogenase from *Escherichia coli*. Mechanism of action deduced from its structural and catalytic properties. *Biochimica et Biophysica Acta* 1457, 211–228.
- Bravo, A. and Mora, J. (1988) Ammonium assimilation in *Rhizobium phaseoli* by the glutamine synthetase–glutamate synthase pathway. *Journal of Bacteriology* 170, 980–984.
- Brown, S., Muhamad, N., Walker, L.R., Pedley, K.C. and Simcock, D.C. (2014) An *in silico* analysis of the glutamate dehydrogenases of *Teladorsagia circumcincta* and *Haemonchus contortus*. *Journal of Biotechnology and Biotechnology* 3, 49–60.
- Brown, S. and Simcock, D.C. (2013) Confidence bands for rational rate equations. *International Journal of Emerging Sciences* 3, 15–27.
- Bruggeman, F.J., Boogerd, F.C. and Westerhoff, H.V. (2005) The multifarious short-term regulation of ammonium assimilation of *Escherichia coli*: dissection using an *in silico* replica. *FEBS Journal* 272, 1965–1985.
- Brzóska, K., Meczynska, S. and Kruszewski, M. (2006) Iron–sulfur cluster proteins: electron transfer and beyond. *Acta Biochimica Polonica* 53, 685–691.
- Camardella, L., Di Fraia, R., Antignani, A., Ciardiello, M.A., di Prisco, G., Coleman, J.K., Buchon, L. and Russell, N.J. (2002) The Antarctic *Psychrobacter* sp. TAD1 has two cold-active glutamate dehydrogenases with different cofactor specificities. Characterisation of the NAD⁺-dependent enzyme. *Comparative Biochemistry and Physiology* 131A, 559–567.
- Cao, Z., Song, P., Xu, Q., Su, R. and Zhu, G. (2011) Overexpression and biochemical characterization of soluble pyridine nucleotide transhydrogenase from *Escherichia coli*. *FEMS Microbiology Letters* 320, 9–14.
- Cook, P.F., Blanchard, J.S. and Cleland, W.W. (1980) Primary and secondary deuterium isotope effects on equilibrium constants for enzyme-catalyzed reactions. *Biochemistry* 19, 4853–4858.
- Cordwell, S.J. (1999) Microbial genomes and “missing” enzymes: redefining biochemical pathways. *Archives of Microbiology* 172, 269–279.
- Cuff, A.L., Sillitoe, I., Lewis, T., Clegg, A.B., Rentzsch, R., Furnham, N., Pellegrini-Calance, M., Jones, D., Thornton, J. and Orengo, C.A. (2011) Extending CATH: increasing coverage of the protein structure universe and linking structure with function. *Nucleic Acids Research* 39(Suppl. 1), D420–D426.

- Dalziel, K. (1969) The interpretation of kinetic data for enzyme-catalysed reactions involving three substrates. *Biochemical Journal* 114, 547–556.
- Davis, B.D., Kornberg, H.L., Nagler, A., Miller, P. and Mingioli, E. (1959) Formation and functions of succinate in *Escherichia coli*. *Federation Proceedings* 18: 211.
- DeLuna, A., Avendaño, A., Riego, L. and González, A. (2001) NADP-glutamate dehydrogenase isoenzymes of *Saccharomyces cerevisiae*. Purification, kinetic properties, and physiological roles. *The Journal of Biological Chemistry* 276, 43775–43783.
- Di Fraia, R., Wilquet, V., Ciardiello, M.A., Carratorre, V., Antignane, A., Camardella, L., Glansdorff, N. and di Prisco, G. (2000) NADP⁺-dependent glutamate dehydrogenase in the Antarctic psychrotolerant bacterium *Psychrobacter* sp. TAD1. Characterization, protein and DNA sequence, and relationship to other glutamate dehydrogenases. *European Journal of Biochemistry* 267, 121–131.
- Doucette, C.D., Schawb, D.J., Wingreen, N.S. and Rabinowitz, J.D. (2011) α -Ketoglutarate coordinates carbon and nitrogen utilization via enzyme I inhibition. *Nature Chemical Biology* 7, 894–901.
- Duncan, P.A., White, B.A. and Mackie, R.I. (1992) Purification and properties of NADP-dependent glutamate dehydrogenase from *Ruminococcus flavefaciens* FD-1. *Applied and Environmental Microbiology* 58, 4032–4037.
- Engel, P.C. and Dalziel, K. (1967) The equilibrium constants of the glutamate dehydrogenase systems. *Biochemical Journal* 105, 691–695.
- Engel, P.C. and Dalziel, K. (1970) Kinetics and mechanism of glutamate dehydrogenase. In: Sund, H. (ed) *Pyridine Nucleotide-dependent Dehydrogenases*. Springer, Heidelberg, Germany, pp. 245–256.
- Engel, P.C. and Ferdinand, W. (1973) The significance of abrupt transitions in Lineweaver–Burk plots with particular reference to glutamate dehydrogenase. Negative and positive co-operativity in catalytic rate constants. *Biochemical Journal* 131, 97–105.
- Felsenstein, J. (1985) Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39, 783–793.
- Finn, R.D., Bateman, A., Clements, J., Coggill, P., Eberhardt, R.Y., Eddy, S.R., Heger, A., Hetherington, K., Holm, L., Mistry, J. *et al.* (2014) Pfam: the protein families database. *Nucleic Acids Research* 42, D222–D230.
- Fisher, H.F. (2005) Transient-state kinetic approach to mechanisms of enzymatic catalysis. *Accounts of Chemical Research* 38, 157–166.
- Fisher, H.F. and Maniscalco, S.J. (2002) A close-packed planar 4-atom motif serves as a variable pathway mechanistic switching device in enzymatic catalysis. *Bioorganic Chemistry* 30, 199–210.
- Frieden, C. (1959) Glutamic dehydrogenase. III. The order of substrate addition in the enzymatic reaction. *The Journal of Biological Chemistry* 234, 2891–2896.
- Glass, T.L. and Hylemon, P.B. (1980) Characterization of a pyridine nucleotide-nonspecific glutamate dehydrogenase from *Bacteroides thetaiotaomicron*. *Journal of Bacteriology* 141, 1320–1330.
- Gupta, S.C. and Dekker, E.E. (1980) Evidence for the identity and some comparative properties of α -ketoglutarate and 2-keto-4-hydroxyglutarate dehydrogenase activity. *The Journal of Biological Chemistry* 255, 1107–1112.
- Hager, L.P. and Kornberg, H.L. (1961) On the mechanism of α -oxoglutarate oxidation in *Escherichia coli*. *Biochemical Journal* 78, 194–198.
- Halpern, Y.S. and Umbarger, H.E. (1961) Utilization of L-glutamic and 2-oxoglutaric acid as sole sources of carbon by *Escherichia coli*. *Journal of General Microbiology* 26, 175–188.
- Helling, R.B. (1994) Why does *Escherichia coli* have two primary pathways for synthesis of glutamate? *Journal of Bacteriology* 176, 4664–4668.
- Helling, R.B. (1998) Pathway choice in glutamate synthesis in *Escherichia coli*. *Journal of Bacteriology* 180, 4571–4575.
- Hornby, D.P. and Engel, P.C. (1984) Characterization of *Peptostreptococcus asaccharolyticus* glutamate dehydrogenase purified by dye–ligand chromatography. *Journal of General Microbiology* 130, 2385–2394.
- Hurley, J.H., Chen, R. and Dean, A.M. (1996) Determinants of cofactor specificity in isocitrate dehydrogenase: structure of an engineered NADP⁺ \rightarrow NAD⁺ specificity-reversal mutant. *Biochemistry* 35, 5670–5678.
- Huynen, M.A., Dandekar, T. and Bork, P. (1999) Variation and evolution of the citric-acid cycle: a genomic perspective. *Trends in Microbiology* 7, 281–291.
- Jahns, T. and Kaltwasser, H. (1993) Properties of the cold-labile NAD⁺-specific glutamate dehydrogenase from *Bacillus cereus* DSM 31. *Journal of General Microbiology* 139, 775–780.
- Janes, B.K., Pomposiello, P.J., Perez-Matos, A., Najarian, D.J., Goss, T.J. and Bender, R.A. (2001) Growth inhibition caused by overexpression of the structural genes for glutamate dehydrogenase (*gdhA*) from *Klebsiella aerogenes*. *Journal of Bacteriology* 183, 2709–2714.

- Jones, D.T., Taylor, W.R. and Thornton, J.M. (1992) The rapid generation of mutation data matrices from protein sequences. *Computer Applications in the Biosciences* 8, 275–282.
- Joshi, C.V., Pathan, E.K., Puneekar, N.S., Tupe, S.G., Kapadnis, B.P. and Deshpande, M.V. (2013) A biochemical correlate of dimorphism in a zygomycete *Benjaminiella poitrasii*: characterization of purified NAD-dependent glutamate dehydrogenase, a target for antifungal agents. *Antonie van Leeuwenhoek* 104, 25–36.
- Kanamori, K., Weiss, R.L. and Roberts, J.D. (1987) Role of glutamate dehydrogenase in ammonia assimilation in nitrogen-fixing *Bacillus macerans*. *Journal of Bacteriology* 169, 4692–4695.
- Kawahara, Y., Takahashi-Fuke, K., Shimizu, E., Nakamatsu, T. and Nakamori, S. (1997) Relationship between the glutamate production and the activity of 2-oxoglutarate dehydrogenase in *Brevibacterium lactofermentum*. *Bioscience, Biotechnology and Biochemistry* 61, 1109–1112.
- Kawakami, R., Sakuraba, H. and Ohshima, T. (2007) Gene cloning and characterization of the very large NAD-dependent L-glutamate dehydrogenase from the psychrophile *Janthinobacterium lividum*, isolated from cold soil. *Journal of Bacteriology* 189, 5626–5633.
- Kawakami, R., Oyama, M., Sakuraba, H. and Ohshima, T. (2010) The unique kinetic behavior of the very large NAD-dependent L-glutamate dehydrogenase from *Janthinobacterium lividum*. *Bioscience, Biotechnology and Biochemistry* 74, 884–887.
- Kersten, M.A.S.H., Müller, Y., Baars, J.J.P., Op den Camp, H.J.M., van der Drift, C., Van Griensven, L.J.L.D., Visser, J. and Schaap, P.J. (1999) NAD⁺-dependent glutamate dehydrogenase of the edible mushroom *Agaricus bisporus*: biochemical and molecular characterization. *Molecular and General Genetics* 261, 452–462.
- Khan, M.I.H., Ito, K., Kim, H., Ashida, H., Ishikawa, T., Shibata, H. and Sawa, Y. (2005) Molecular properties and enhancement of thermostability by random mutagenesis of glutamate dehydrogenase from *Bacillus subtilis*. *Bioscience, Biotechnology and Biochemistry* 69, 1861–1870.
- Kujo, C. and Ohshima, T. (1998) Enzymological characteristics of the hyperthermostable NAD-dependent glutamate dehydrogenase from the archaeon *Pyrobaculum islandicum* and effects of denaturants and organic solvents. *Applied and Environmental Microbiology* 64, 2152–2157.
- Kurganov, B.I. (2000) New approach to analysis of deviations from hyperbolic law in enzyme kinetics. *Biokhimiya* 65, 898–909.
- Lu, C.-D. and Abdellal, A.T. (2001) The *gdhB* gene of *Pseudomonas aeruginosa* encodes an arginine-inducible NAD⁺-dependent glutamate dehydrogenase which is subject to allosteric regulation. *Journal of Bacteriology* 183, 490–499.
- Maksymiuk, C., Balakrishnan, A., Bryk, R., Rhee, K.Y. and Nathan, C.F. (2015) E1 of α -ketoglutarate dehydrogenase defends *Mycobacterium tuberculosis* against glutamate anaplerosis and nitroxidative stress. *Proceedings of the National Academy of Sciences of the United States of America* 112(43), E5834–E5843.
- Maniscalco, S.J., Saha, S.K., Vicedomine, P. and Fisher, H.F. (1996) A difference in the sequence of steps in the reaction catalyzed by two closely homologous forms of glutamate dehydrogenase. *Biochemistry* 35, 89–94.
- Miñambres, B., Olivera, E.R., Jensen, R.A. and Luengo, J.M. (2000) A new class of glutamate dehydrogenases (GDH). Biochemical and genetic characterization of the first member, the AMP-requiring NAD-specific GDH of *Streptomyces clavuligerus*. *The Journal of Biological Chemistry* 275, 39529–39542.
- Müller, T., Strösser, J., Buchinger, S., Nolden, L., Wirtz, A., Krämer, R. and Burkovski, A. (2006) Mutation-induced metabolite pool alterations in *Corynebacterium glutamicum*: towards the identification of nitrogen control signals. *Journal of Biotechnology* 126, 440–453.
- Murthy, P.S., Sirsi, M. and Ramakrishnan, T. (1962) Tricarboxylic acid cycle and related enzymes in cell-free extracts of *Mycobacterium tuberculosis* H₃₇R_v. *Biochemical Journal* 84, 263–269.
- Olson, J.A. and Anfinsen, C.B. (1953) Kinetic and equilibrium studies on crystalline L-glutamic acid dehydrogenase. *The Journal of Biological Chemistry* 202, 841–856.
- Osterman, A. and Overbeek, R. (2003) Missing genes in metabolic pathways: a comparative genomics approach. *Current Opinion in Chemical Biology* 7, 238–251.
- Piston, S.M., Mendz, G.L., Srinivasan, S. and Hazell, S.L. (1999) The tricarboxylic acid cycle of *Helicobacter pylori*. *European Journal of Biochemistry* 260, 258–267.
- Preiss, T., Sang, A.E., Chrzanowska-Lightowlers, Z.M.A. and Lightowlers, R.N. (1995) The mRNA-binding protein COLBP is glutamate dehydrogenase. *FEBS Letters* 367, 291–296.
- Rajas, F., Gire, V. and Rousse, B. (1996) Involvement of a membrane-bound form of glutamate dehydrogenase in the association of lysosomes to microtubules. *The Journal of Biological Chemistry* 271, 29882–29890.
- Rife, J.E. and Cleland, W.W. (1980a) Kinetic mechanism of glutamate dehydrogenase. *Biochemistry* 19, 2321–2328.

- Rife, J.E. and Cleland, W.W. (1980b) Determination of the chemical mechanism of glutamate dehydrogenase from pH studies. *Biochemistry* 19, 2328–2333.
- Rossi, M., Defez, R., Chiurazzi, M., Lamberti, A., Fuggi, A. and Iaccarino, M. (1989) Regulation of glutamine synthetase isoenzymes in *Rhizobium leguminosarum* biovar *viceae*. *Journal of General Microbiology* 135, 629–637.
- Sakamoto, N., Kotre, A.M. and Savageau, M.A. (1975) Glutamate dehydrogenase from *Escherichia coli*: purification and properties. *Journal of Bacteriology* 124, 775–783.
- Sarada, K.V., Rao, N.A. and Venkatasubramanian, T.A. (1980) Isolation and characterisation of glutamate dehydrogenase from *Mycobacterium smegmatis* CDC 46. *Biochimica et Biophysica Acta* 615, 299–308.
- Schmincke-Ott, E. and Bisswanger, H. (1981) Dihydropyridine dehydrogenase component of the pyruvate dehydrogenase complex from *Escherichia coli* K12. Comparative characterization of the free and the complex-bound component. *European Journal of Biochemistry* 114, 413–420.
- Senior, P.J. (1975) Regulation of nitrogen metabolism in *Escherichia coli* and *Klebsiella aerogenes*: studies with the continuous-culture technique. *Journal of Bacteriology* 123, 407–418.
- Sharkey, M.A. and Engel, P.C. (2008) Apparent negative co-operativity and substrate inhibition in overexpressed glutamate dehydrogenase from *Escherichia coli*. *FEMS Microbiology Letters* 281, 132–139.
- Sharkey, M.A., Oliveira, T.F., Engel, P.C. and Khan, A.R. (2013) Structure of NADP⁺-dependent glutamate dehydrogenase from *Escherichia coli* – reflections on the basis of coenzyme specificity in the family of glutamate dehydrogenases. *FEBS Journal* 280, 4681–4692.
- Shiio, I. and Ozaki, H. (1970) Regulation of nicotinamide adenine dinucleotide phosphate-specific glutamate dehydrogenase from *Brevibacterium flavum*, a glutamate-producing bacterium. *The Journal of Biochemistry* 68, 633–647.
- Shiio, I., Ôsuka, S.-I. and Tsunoda, T. (1959) Glutamic acid formation from glucose by bacteria. II. Glutamic acid and α -ketoglutaric acid formation by *Brevibacterium flavum*, No. 2247. *The Journal of Biochemistry* 46, 1597–1605.
- Shiio, I., Otsuka, S. and Takahashi, M. (1961) Significance of α -ketoglutarate dehydrogenase on the glutamic acid formation in *Brevibacterium flavum*. *The Journal of Biochemistry* 50, 164–165.
- Shiio, I., Nariui, K., Yahaba, N. and Takahashi, M. (1962) Free intracellular amino acid of a glutamate-forming bacterium, *Brevibacterium flavum* No. 2247. *The Journal of Biochemistry* 51, 109–111.
- Smith, E.L., Austen, B.M., Blumenthal, K.M. and Nyc, J.F. (1975) Glutamate dehydrogenases. In P. Boyer (ed) *The enzymes*, vol. XI. Academic Press, New York pp. 293–367.
- Stillman, T.J., Baker, P.J., Britton, K.L. and Rice, D.W. (1993) Conformational flexibility in glutamate dehydrogenase. Role of water in substrate recognition and catalysis. *Journal of Molecular Biology* 234, 1131–1139.
- Stouthamer, A.H. (1966) Some properties of mutants of *Aerobacter aerogenes* blocked in the formation of succinyl-Co A. *Antonie van Leeuwenhoek* 32, 452.
- Strecker, H.J. (1953) Glutamic dehydrogenase. *Archives of Biochemistry and Biophysics* 46, 128–140.
- Struhl, K. and Maganasik, B. (1976) Ammonia-sensitive mutant of *Klebsiella aerogenes*. *Journal of Bacteriology* 126, 739–742.
- Subramanian, S. (1978) Thermodynamics of the glutamate dehydrogenase catalytic reaction. *Biophysical Chemistry* 7, 375–378.
- Syed, S.-E.H. and Engel, P.C. (1987) Glutamate dehydrogenase of *Clostridium symbiosum*: another example of an abrupt transition in the kinetic plots. *Biochemical Society Transactions* 15, 237–238.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. and Kumar, S. (2011) MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution* 28, 2731–2739.
- Tian, J., Bryk, R., Itoh, M., Suematsu, M. and Nathan, C. (2005a) Variant tricarboxylic acid cycle in *Mycobacterium tuberculosis*: identification of α -ketoglutarate decarboxylase. *Proceedings of the National Academy of Sciences of the United States of America* 102, 10670–10675.
- Tian, J., Bryk, R., Shi, S., Erdjument-Bromage, H., Tempst, P. and Nathan, C. (2005b) *Mycobacterium tuberculosis* appears to lack α -ketoglutarate dehydrogenase and encodes pyruvate dehydrogenase in widely separated genes. *Molecular Microbiology* 57, 859–868.
- Vančurová, I., Vančura, A., Volc, J., Kopecý, J., Neužil, J., Basařová, G. and Běhal, V. (1989) Purification and properties of NADP-dependent glutamate dehydrogenase from *Streptomyces fradiae*. *Journal of General Microbiology* 135, 3311–3318.
- Wagner, A. and Fell, D.A. (2001) The small world inside large metabolic networks. *Proceedings of the Royal Society of London* 268B, 1803–1810.
- Weitzman, P.D.J. (1981) Unity and diversity in some bacterial citric acid-cycle enzymes. *Advances in Microbial Physiology* 22, 185–244.