

Boar sperm DNA damage induced by tropical heat stress can be alleviated using antioxidants.

Santiago T. Peña, Jr.^{1,2,3}, Bruce Gummow^{2,4}, Anthony J. Parker⁵ and Damien B.B.P. Paris¹

¹Discipline of Biomedical Science, College of Public Health, Medical & Veterinary Sciences, James Cook University, Townsville, Queensland, Australia

²Discipline of Veterinary Science, College of Public Health, Medical & Veterinary Sciences, James Cook University, Townsville, Queensland, Australia

³College of Veterinary Medicine, Visayas State University, Baybay City, Leyte, Philippines

⁴Faculty of Veterinary Science, University of Pretoria, Onderstepoort, Pretoria, South Africa

⁵College of Food, Agricultural & Environmental Sciences, Ohio State University, Wooster, Ohio, USA

Seasonal heat stress is known to significantly diminish reproductive performance in pigs, particularly in the tropics, costing the industry millions in annual losses. The boar's reduced capacity to sweat, non-pendulous scrotum, and widespread use of European breeds in the tropics, makes this species particularly vulnerable to heat stress. While traditionally considered a sow problem, recent mouse studies demonstrate that heat stress-induced sperm DNA damage can result in arrested development and loss of early embryos. Our study investigated the impact of tropical summer heat stress on the quality and DNA integrity of boar sperm, and trialled antioxidant supplementation to alleviate the problem.

Motility of sperm obtained from $n = 5$ Large White boars housed in the dry tropics of Townsville, North Queensland, Australia was characterized by Computer-Assisted Sperm Analysis but did not differ between summer, winter or spring (total motility: 71.3 ± 8.1 vs. 90.2 ± 4.2 vs. $70.8 \pm 5.5\%$ respectively, $P > 0.05$; progressive motility: 35.4 ± 7.0 vs. 46.6 ± 4.0 vs. $41.7 \pm 2.8\%$ respectively, $P > 0.05$). Sperm DNA integrity in 20,000 sperm/boar/season, evaluated using TUNEL and flow cytometry, revealed 16-fold more DNA damaged sperm in summer than winter, and nearly 9-fold more than spring (16.1 ± 4.8 vs. 1.0 ± 0.2 vs. $1.9 \pm 0.5\%$ respectively, $P \leq 0.05$). However, boar feed supplemented with 100g/boar/day of proprietary custom-made antioxidants during summer significantly reduced sperm DNA damage to $9.9 \pm 4.5\%$ and $7.2 \pm 1.6\%$ ($P \leq 0.05$) after 42 and 84 days treatment respectively. Total and progressive motility were not altered by the supplement.

In summary, sperm DNA integrity is compromised in boars during summer, suggesting boar factors may contribute to seasonal embryo loss in sows. Moreover, such damage appears undetectable using traditional measures of sperm motility. Antioxidant supplementation during summer appears to mitigate the negative impact of heat stress on sperm DNA integrity.