

# Morphological characterization of virus-like particles in coral reef sponges

Cecília Pascelli<sup>1,2,3</sup>, Patrick W. Laffy<sup>1</sup>, Marija Kupresanin<sup>4</sup>, Timothy Ravasi<sup>4</sup> and Nicole S. Webster<sup>1,3,5</sup>

<sup>1</sup> Australian Institute of Marine Science, Townsville, Queensland, Australia

<sup>2</sup> James Cook University, Townsville, Queensland, Australia

<sup>3</sup> AIMS@JCU, Australian Institute of Marine Science and James Cook University, Townsville, Queensland, Australia

<sup>4</sup> KAUST Environmental Epigenetic Program (KEEP), Division of Biological and Environmental Sciences & Engineering, King Abdullah University of Science and Technology, Thuwal, Kingdom of Saudi Arabia

<sup>5</sup> Australian Centre for Ecogenomics, University of Queensland, Brisbane, Queensland, Australia

## ABSTRACT

Marine sponges host complex microbial consortia that vary in their abundance, diversity and stability amongst host species. While our understanding of sponge-microbe interactions has dramatically increased over the past decade, little is known about how sponges and their microbial symbionts interact with viruses, the most abundant entities in the ocean. In this study, we employed three transmission electron microscopy (TEM) preparation methods to provide the first comprehensive morphological assessment of sponge-associated viruses. The combined approaches revealed 50 different morphologies of viral-like particles (VLPs) represented across the different sponge species. VLPs were visualized within sponge cells, within the sponge extracellular mesohyl matrix, on the sponge ectoderm and within sponge-associated microbes. Non-enveloped, non-tailed icosahedral VLPs were the most commonly observed morphotypes, although tailed bacteriophage, brick-shaped, geminate and filamentous VLPs were also detected. Visualization of sponge-associated viruses using TEM has confirmed that sponges harbor not only diverse communities of microorganisms but also diverse communities of viruses.

**Subjects** Biodiversity, Marine Biology, Microbiology, Virology, Biological Oceanography

**Keywords** Transmission Electron Microscopy, Virus, TEM, VLP, Marine sponges, Great Barrier Reef, GBR, Red Sea

## INTRODUCTION

Sponges are abundant and ecologically important members of marine benthic communities (*Van Soest et al., 2012*). Most sponges are suspension filter feeders (*Thomassen & Røisgård, 1995*), with complex aquiferous systems capable of manipulating the seawater composition at both macro and micro scales (*Vacelet & Boury-Esnault, 1995; Patterson et al., 1997; De Goeij et al., 2013*). A unidirectional (ostia-chamber-atrium-oscula) water flow driven by flagellated choanocyte cells is responsible for capturing and retaining small eukaryotes, prokaryotic cells and viral particles (*Hadas et al., 2006*). Sponge filtration of large quantities of seawater represents an important nutrient link between the pelagic and benthic

Submitted 28 June 2018

Accepted 22 August 2018

Published 17 October 2018

Corresponding author

Cecília Pascelli,

ceciliapascelli@gmail.com

Academic editor

Mya Breitbart

Additional Information and  
Declarations can be found on  
page 15

DOI 10.7717/peerj.5625

© Copyright

2018 Pascelli et al.

Distributed under

Creative Commons CC-BY 4.0

## OPEN ACCESS

environments (*Pile & Young, 2006*), especially in oligotrophic ecosystems such as coral reefs (*De Goeij et al., 2013*).

Sponges form intimate partnerships with diverse microbial consortia, and these relationships range from mutualism to commensalism to parasitism (*Webster & Taylor, 2012; Thomas et al., 2016*). The sponge microbiome is often highly conserved across individuals of the same sponge species but varies considerably across species (*Thomas et al., 2016*). It is because of these functionally important symbiotic partnerships that sponges are considered a typical example of a marine 'holobiont', an organism comprised of various 'bionts', living in symbiogenesis (*Margulis & Fester, 1991; Webster & Thomas, 2016*). However, while the symbiotic association between sponges and their bacterial/archaeal symbionts has been extensively studied (*Schmitt et al., 2012; Thomas et al., 2016; Pita et al., 2018*), the role of viruses in the sponge holobiont remain largely unknown, despite TEM images from the 1970s alluding to viral-infected sponge cells (*Vacelet & Gallissian, 1978*), a demonstration of phage infection in a sponge-associated bacterium (*Lohr, Chen & Hill, 2005*), and a few recent metagenomic studies providing insights into sponge virus diversity and function (*Butina et al., 2015; Laffy et al., 2016; Laffy et al., 2018*).

Viruses are the most abundant biological agents in marine ecosystems, with about  $10^{10}$  viruses per liter of surface seawater and  $10^{10}$  per gram dry weight of marine sediment (*Suttle, 2007; Danovaro et al., 2011*). Importantly, viruses have the ability to regulate the prokaryotic and eukaryotic populations responsible for maintaining metabolic cycling in complex ecosystems such as coral reefs (*Seymour et al., 2005; Thurber & Correa, 2011; Mojica & Brussaard, 2014*). Viruses modulate microbial-driven processes through mortality, horizontal gene transfer and metabolic reprogramming by viral-encoded auxiliary metabolic genes (AMGs) (*Bergh et al., 1989; Rohwer & Thurber, 2009; Danovaro et al., 2011; Hurwitz et al., 2014; Breitbart et al., 2018*). Recent years have seen an increased focus on the diversity and function of viruses associated with reef invertebrates including sea anemones (*Wilson & Chapman, 2001*); starfish (*Hewson et al., 2018*); scleractinian corals and their associated microbial communities (*Patten, Harrison & Mitchell, 2008; Weynberg et al., 2014; Weynberg et al., 2017a; Laffy et al., 2018*). However, while viruses have been described as essential components of coral reef ecosystems, capable of controlling microbial community dynamics, playing a role in coral bleaching/disease, and mediating reef biogeochemical cycling (*Thurber et al., 2017*), there is a paucity of research exploring viruses associated with ecologically important reef sponges.

Metagenomic analysis of purified viral fractions (metaviromics) recently provided the first insights into the composition and function of viruses inhabiting reef sponges (*Laffy et al., 2016; Laffy et al., 2018*). Consistent with the pattern reported for sponge-associated microbial communities, the viral communities were found to be highly conserved within each sponge species, and displayed functional repertoires clearly distinct from viruses inhabiting the surrounding seawater (*Laffy et al., 2018*). Sequence analysis revealed that the metavirome assignments were dominated by viromes from the order *Caudovirales* but also contained representatives of the *Mimiviridae*, *Phycodnaviridae*, *Circoviridae*, *Parvoviridae*, *Bidnaviridae* and *Microviridae*. Unique viral adaptations to specific host

microenvironments were also evident, with viral auxiliary genes being differentially represented across sponge species (Laffy *et al.*, 2018).

While molecular approaches have substantially improved our understanding of viral-host interactions (Breitbart *et al.*, 2002; Rosario & Breitbart, 2011; Laffy *et al.*, 2016), biases associated with DNA/RNA extraction methods (Wood-Charlson *et al.*, 2015) and the limited genomic resources available for most environmental viruses (Roux *et al.*, 2015) can still constrain our understanding of host-associated viral ecology. Transmission electron microscopy (TEM) is a powerful approach that has helped to reveal the morphology and distribution of virus-like particles (VLPs) in many marine hosts as well as deciphering patterns of host-viral interactions (Wilson & Chapman, 2001; Patten, Harrison & Mitchell, 2008; Brum, Schenck & Sullivan, 2013; Pollock *et al.*, 2014; Weynberg *et al.*, 2017b). Here we use TEM to provide the first morphological characterization of viruses associated with 15 different coral reef sponge species and confirm the spatial localization of these VLPs within the sponge holobiont.

## MATERIALS AND METHODS

### Sponge collection and identification

Sampling was conducted on coral reefs of Orpheus Island, Great Barrier Reef, Australia (18°35'34"S, 146°28'53"E) and Al Fahal, Red Sea, Saudi Arabia (22°13'95"N, 39°01'81"E), between December 2015 and February 2016. Sampling in Australia was conducted under the Great Barrier Reef Marine Park Authority permit G12/35236.1, and sampling in Saudi Arabia was authorized by the Saudi Arabian coastguard as the study did not involve endangered or protected species.

Triplicate specimens of 15 sponge species were collected by scuba diving between three and 15 m depth. Two sponge species, *Stylissa carteri* and *Carteriospongia foliascens* were found at both locations, and sampling was performed in triplicate at both sites. Sponge specimens were photographed *in situ* before being individually placed within sterile Falcon<sup>®</sup> tubes and kept on ice until processing. All sampling materials were sterilized prior to and between each sampling. Morphological characterization of sponge species was performed as described in (Hooper & Van Soest, 2002) and DNA barcoding was additionally performed using mitochondrial cytochrome oxidase I (COI) gene primers and internal transcriber spacer 2 (ITS2) region of nuclear ribosomal DNA as described in (Erwin & Thacker, 2007; Andreakis, Luter & Webster, 2012; Wörheide *et al.*, 2012). Sponge species are described in Table 1 and can be seen in Fig. S1.

Three different sample preparation methods for TEM imaging of sponge-associated viruses were trialed: (i) ultrathin sectioning of sponge tissue (Cheville & Stasko, 2014); (ii) purification of viral fractions via density gradient ultracentrifugation (Lawrence & Steward, 2010; Weynberg *et al.*, 2014) and (iii) filtration of sponge mucus. All samples were examined using a Titan Cubed TEM and images were analyzed on the Cs-corrected Titan<sup>™</sup> 80–300 platform at the Imaging Characterization Core Lab in KAUST. TEM search time was standardized to 1 hr/sample.

**Table 1** Collection details for all sponge species examined by TEM. GBR refers to the Great Barrier Reef collection site and RS refers to the Red Sea collection site.

Sponge species	Location	Depth (m)
<i>Carteriospongia foliascens</i> , P.S. Pallas (1766)	GBR, RS	3–10
<i>Stylissa carteri</i> , A. Dendi (1889)	GBR, RS	10–15
<i>Xestospongia</i> sp.	GBR	5–15
<i>Lamellodysidea herbacea</i> , C. Keller (1889)	GBR	5–10
<i>Cymbastela marshae</i> , J.N.A. Hooper & P.R. Bergquist (1992)	GBR	10–15
<i>Cinachyrella schulzei</i> , C. Keller (1891)	GBR	3–7
<i>Pipestela candelabra</i> , B. Alvarez et al. (2008)	GBR	7–15
<i>Echinochalina isaaci</i> , J.N.A. Hooper (1996)	GBR	7–15
<i>Xestospongia testudinaria</i> , J.B.P. Lamarck (1815)	RS	7–15
<i>Amphimedon ochracea</i> , C. Keller (1889)	RS	7–15
<i>Hyrtilos erectus</i> , C. Keller (1889)	RS	5–15
<i>Crella (Grayela) cyathophora</i> , H.J. Carter (1869)	RS	7–15
<i>Mycale</i> sp.	RS	5–15

### Preparation of ultrathin sections of sponge tissue

Histological sections were prepared from fresh sponge tissue based on standard procedures for TEM (Cheville & Stasko, 2014). Briefly, each fragment of approximately 1 mm<sup>3</sup> was fixed in 2.5% glutaraldehyde in 0.2M cacodylate buffer and kept at 4 °C for 2–24 h. After fixation, samples were immersed in 1% osmium tetroxide in 100 mM phosphate buffer for 1–2 h, washed in distilled water and stained in the dark with 2% aqueous uranyl acetate for 2 h at 4 °C. Stained tissue was dehydrated through a series of ethanol and propylene oxide then embedded in epoxy resin. Ectosome-choanosome oriented sections (about 65 nm thick) were prepared using a Leica EM UC7 ultramicrotome and placed on TEM copper grids.

### Viral purification via density gradient solution

Viral purification was performed according to the fraction separation method by sedimentation in density gradients (Meselson, Stahl & Vinograd, 1957) following the pre-processing approach established to isolate viruses from coral and sponge tissue (Weynberg et al., 2014; Laffy et al., 2018). In order to eliminate contaminants present in the aquiferous system, sponges were partially dried via repeated gentle squeezing alternated with rinses of filtered (0.02 μm) seawater. Sponge tissue was then dissected into small pieces (~5 mm<sup>3</sup>) and covered with 15 μL of 0.02 μm filter-sterilized (Anotop, Whatman) SM buffer (100 mM NaCl, 8 mM MgSO<sub>4</sub>, 50 mM Tris pH 7.5), then homogenized with a Craig's HS30E homogenizer (Witeg, Germany) for 5 to 10 min (min). Tissue homogenate was filtered through a Falcon<sup>®</sup> 100 μm Cell Strainer (Corning, USA), then centrifuged at 500 g for 15 min at 4 °C to pellet the majority of cell debris. The supernatant was used to purify the VLP via centrifugation in Cesium Chloride solution, with density varying from 1.2 g/mL to 1.6 g/mL (Weynberg et al., 2014). After ultracentrifugation, sponge VLPs were collected from the fractions with densities between 1.2 g/mL and 1.5 g/mL. In order to exchange the buffer and remove CsCl salts, samples were loaded onto 30 KDa Amicon

centrifugal spin columns (Millipore, EUA) and centrifuged at 4,000 g for 30 min at 4 °C. This process was repeated four–six times per sample. Filter-sterilized SM Buffer was added to the concentrate and all flow-through was discarded. The concentrate was fixed in 0.5% glutaraldehyde and kept at 4 °C until TEM analysis. TEM preparation involved applying a droplet of sample onto a TEM Copper grid, rinsing with sterile water, staining with 1% uranyl acetate for one min, washing with sterile water, followed by removal of excess liquid from the grid by touching filter paper to the edge.

### ***Viral purification via filtration of sponge mucus***


To describe the VLPs associated with sponge mucus and the external ectoderm, the sponge surface was carefully scraped with a sterile scalpel blade followed by rinsing three times with filtered (0.02 μm) seawater. This TEM preparation method was based on a viral purification method described for marine hydras ([Grasis et al., 2014](#)). Extracted mucus was added to filtered (0.02 μm) Milli-Q<sup>®</sup> water (1:4) and centrifuged at 4,000 g for 10 min. Mucus supernatant was filtered through 0.45 μm filters (EMD Millipore, Burlington, CA, USA) and fixed in 1.5% glutaraldehyde. TEM imaging of mucus preparations was performed as described above for CsCl purified samples.

## **RESULTS**

### **Sponge associated viruses**


TEM analysis revealed that viral particles are diverse constituents of the sponge holobiont. Fifty VLP morphotypes ([Figs. 1–5](#); [Table S1](#); Morphotypes: M-I–M-L) were found in association with eight coral reef sponge species from the Great Barrier Reef: *Carteriospongia foliascens*, *Stylissa carteri*, *Xestospongia* sp., *Pipestela candelabra*, *Lamellodysidea herbacea*, *Cymbastella marshae*, *Echinochalina isaaci* and *Cinachyrella schulzei*; and seven sponge species from the Red Sea: *Carteriospongia foliascens*, *Stylissa carteri*, *Xestospongia testudinaria*, *Hyrtios erectus*, *Mycale* sp., *Amphimedon ochracea* and *Crella cyathophora*. VLPs were observed within sponge cells, in the extracellular mesohyal matrix, in the mucus/surface biofilm and within sponge-associated microbes. A diverse range of viral morphologies were observed, including hexagonal (tailed and non-tailed), spherical, filamentous, brick-shaped, beaded and geminate VLPs. While we detected numerous viral morphotypes, most were rare and often obscured by vesicles, cell debris and particulate organic matter.

Most sponge-associated VLP morphotypes possessed an icosahedral/polyhedral symmetry (~75%), ranging from 60–205 nm in diameter ([Figs. 1–3, 4A](#)). Tails were evident on some VLPs, confirming the presence of viruses from the bacteriophage order *Caudovirales*. Tailed VLPs were tentatively assigned to the three *Caudovirales* families based on their capsid symmetry and tail size/shape. VLPs characteristic of the *Podoviridae* presented a short tail attached to a non-enveloped icosahedral capsid and these VLPs were observed in the sponges *C. foliascens* ([Fig. 1A](#)), *Xestospongia* sp. ([Fig. 1B](#)), *E. isaaci* ([Fig. 1C](#)) and *S. carteri* ([Fig. 1D](#)). VLPs characteristic of the *Siphoviridae* presented an icosahedral head with a long non-contractile tail and these VLPs were detected in the surface biofilm of *C. schulzei* ([Fig. 1E](#)). VLPs characteristic of the *Myoviridae* presented an icosahedral head




**Figure 1** Representative morphotypes of virus-like particles associated with GBR and Red Sea sponges. GBR sponge species: (A, J, K, L) *C. foliascens*, (B) *Xestospongia* sp., (C, F, G) *E. isaaci*, (E) *C. schulzei*. Red Sea sponge species: (D, H) *S. carteri*, (K) *Amphimedon ochracea*. TEM preparation method: (A, J, K) ultrathin sections of sponge tissue, (B–I, L) viral purification via filtration of sponge mucus. Scale bar: 200 nm. Black arrows indicate the viral tail and white arrows indicate the VLPs.

Full-size  DOI: [10.7717/peerj.5625/fig-1](https://doi.org/10.7717/peerj.5625/fig-1)



**Figure 2** Representative morphotypes of virus-like particles associated with GBR sponges. Sponge species: (A) *C. foliascens*, (B, C) *Styliassa carteri*, (D) *Xestospongia* sp., (E–H) *Pipestela candelabra*, (I–K) *Lamellodysidea herbacea*, (L) *C. schulzei*. TEM preparation method: (A, H–K) viral purification via filtration of sponge mucus, (B–F, L) viral purification via CsCl gradient centrifugation, (G) ultrathin sections of sponge tissue. Scale bar: 200 nm.


Full-size DOI: [10.7717/peerj.5625/fig-2](https://doi.org/10.7717/peerj.5625/fig-2)



**Figure 3** Representative morphotypes of virus-like particles associated with GBR and Red Sea sponges. GBR sponge species: (A, B) *C. schulzei*, (C) *Cymbastella marshae*. Red Sea sponge species: (D, E) *C. foliascens*, (F–H) *S. carteri*, (I) *Xestospongia testudinaria*, (J–L) *Hyrtios erectus*. TEM preparation method: (A, D–E, I–L) ultrathin sections of sponge tissue, (B, C, F–H) viral purification via filtration of sponge mucus. Scale bar: (A–C, E–L) 200 nm, (D) 500 nm. Black arrows indicate the VLPs.


Full-size  DOI: [10.7717/peerj.5625/fig-3](https://doi.org/10.7717/peerj.5625/fig-3)





**Figure 4** Representative morphotypes of virus-like particles associated with GBR and Red Sea sponges. GBR sponge species: (A) *Mycale* sp., (B) *C. foliascens*, (C, D) *Xestospongia* sp., (E) *C. schulzei*. Red Sea sponge species: (F, G) *C. foliascens*, (H, I) *S. carteri*, (J–L) *Xestospongia testudinaria*. TEM preparation method: (A, B, D, E, H, I) viral purification via filtration of sponge mucus, (C) viral purification via CsCl gradient centrifugation, (F, G, J–L) ultrathin sections of sponge tissue. Scale bar: (A–E, H, I) 200 nm, (F, G, J–L) 500 nm.

Full-size DOI: [10.7717/peerj.5625/fig-4](https://doi.org/10.7717/peerj.5625/fig-4)



**Figure 5** Representative morphotypes of virus-like particles associated with GBR and Red Sea sponges. GBR sponge species: (C) *Lamellodysidea herbacea*, (I) *C. foliascens*. Red Sea sponge species: (A, B, H) *Crella cyathophora*, (D–G) *Amphimedon ochracea*, (J–L) *Hyrtios erectus*. TEM preparation method: (A, C, D) viral purification via filtration of sponge mucus, (B, E–L) ultrathin sections of sponge tissue. Scale bar: (D) 100 nm, (A–C, E, H–L) 200 nm, (F, G) 5  $\mu$ m. ECM: External Cell Matrix, om: outer membrane, im: inner membrane, cm: core membrane, lb: lateral bodies; c: core, e: external membrane; b: bacterium. Black arrows indicate the VLPs.

Full-size DOI: [10.7717/peerj.5625/fig-5](https://doi.org/10.7717/peerj.5625/fig-5)

and a long *contractile tail* and these VLPs were observed in the sponges *E. isaaci* (Figs. 1F, 1G), *S. carteri* (Fig. 1H) and *A. ochracea* (Fig. 1I).

Non-tailed icosahedral/polyhedral VLPs were observed using all three TEM preparation methods. Particle sizes ranged from 60 to 205 nm in diameter and some presented an electron dense core inside the viral capsid (35–124 nm in diameter). The majority of VLPs did not show an envelope outside the capsid, however an envelope was observed in association with a small proportion of VLPs. (Figs. 2E, 3K–3L, 5H). A typical example of an enveloped VLP was observed in *Hyrtios erectus* where a group of four virions were observed within a vacuole in the mesohyl matrix (Fig. 3L) and another free virion was captured merging its envelope into the cell membrane of the host (Fig. 3K).

In addition to the polyhedral VLPs, eight morphotypes of filamentous virus-like particles (FVLPs) were observed in the sponge mucus, mesohyl matrix, within sponge cells and associated with sponge-associated microorganisms (Figs. 4C–4L; 5A). These morphotypes varied greatly in size (100–1300 nm length, 12–60 nm width) and shape. Rod-shaped FVLPs were detected in the CsCl purified viral fraction of *Xestospongia* sp. (Figs. 4C, 4D) and the mucus of *S. carteri* (Fig. 4I). Although similar, the *S. carteri* bacilliform VLPs were longer than those observed in *Xestospongia* sp. (230 nm long, 19 nm wide in *S. carteri*; 120–130 nm long, 18 nm wide in *Xestospongia* sp.). In *C. foliascens*, a FVLP was frequently observed attached to cyanobacteria and within the sponge mesohyl, (Figs. 4F–4G). This FVLP resembled viruses of the family *Inoviridae* due to their shortened body (100–130 nm length, 50–60 nm width) and electron-translucent core with outer membrane structures consistent with a glycoprotein coat surrounding the entire membrane (Ploss & Kuhn, 2010). In *X. testudinaria*, a FVLP morphotype was observed within cells and dispersed throughout the mesohyl (Figs. 4J–4L). This thin, elongated FVLP (340–1,300 nm long and 15–30 nm wide) was observed at high abundance inside some choanocyte cells and lysed cells releasing virions were also evident (Figs. 4J–4L). Another distinct FVLP morphotype was evident in the sponge mucus of *C. cyathophora* (Fig. 5A). It presented a tube-like shape indicating helical symmetry and size ranging from 150–154 nm in length and 22–25 nm in width.

Geminate VLPs were observed in *C. cyathophora* mesohyl matrix (Fig. 5B), in *L. herbacea* mucus (Fig. 5C), and found infecting filamentous cyanobacteria associated with the sponge *A. ochracea* (Figs. 5D–5G). The cyanobacteria associated VLPs shared morphological traits with viruses from the family *Geminiviridae* (Li, Ou & Zhang, 2013) and were typically twinned (81–95 nm long, 37–48 nm wide), comprising two quasi-isometric particles (34–45 nm length). The VLPs were spread across the cytoplasm, thylakoid lumen, and vacuoles of the cyanobacterial cells and were often at high abundance surrounding the stellar bodies (Fig. 5F).

A brick-shaped VLP morphotype, closely resembling viruses from the *Poxviridae*, was observed in sections of *Crella cyathophora* (Fig. 5H). This morphotype had a complex structure comprising a biconcave core encased within a double layer membrane with two lateral bodies surrounded by an ovoid envelope (Buller & Palumbo, 1991). Three representatives of this morphotype were observed within the sponge mesohyl matrix, and a single non-enveloped VLP was also observed in close proximity to a lysed sponge cell.

A beaded VLP was observed in sections of the sponges *C. foliascens* (Fig. 5I) and *H. erectus* (Figs. 5J–5L). In *C. foliascens*, the branched VLP was 340 nm long, and was comprised of six beads, each measuring 30–35 nm in diameter. In *H. erectus*, the VLPs varied from 80 to 350 nm in length and were composed of 2–8 aligned beads with diameters ranging from 36–42 nm. This morphotype was observed as isolated VLPs, attached to extracellular vacuole membranes in the sponge mesohyl, and within intracellular vacuoles of archaeocyte cells.

## DISCUSSION

Sponges are complex holobionts that host a diverse array of bacteria, archaea, and eukaryotic microorganisms (Fan *et al.*, 2012; Fan *et al.*, 2013; Webster & Thomas, 2016). Whilst previous publications have alluded to the potential importance of viruses in sponges (Claverie *et al.*, 2009; Webster & Taylor, 2012; Laffy *et al.*, 2016; Laffy *et al.*, 2018), including in sponge disease (Luter, Whalan & Webster, 2010), this study provides the first visual evidence that viruses are diverse components of the sponge holobiont. The broad range of VLP morphologies visualised across the 15 different sponge species is consistent with recent molecular data showing sponges harbour diverse communities of viruses (Laffy *et al.*, 2016; Laffy *et al.*, 2018).

The frequent detection of multiple viral morphotypes within a single sponge species most likely reflects the large number of potential hosts within the sponge holobiont (sponge cells, bacteria, archaea, microeukaryotes). However, it is also possible that multiple viruses infect the same host, as has been observed in some bacterioplankton (Holmfeldt *et al.*, 2007) and corals (Thurber & Correa, 2011). Similarly, the same viral morphotype may infect multiple hosts within the holobiont, as recently highlighted from phage-bacteria network analyses (Flores *et al.*, 2011; Flores, Valverde & Weitz, 2013). This is particularly relevant considering the role of viruses in lateral gene transfer between hosts and their subsequent effects on host metabolism (Breitbart *et al.*, 2018). Observed viral morphotypes may also not be native to the holobiont, as some may have been extracted from the virioplankton by the sponge's aquiferous system. Although the isolation methods employed in this study unveiled a wide range of VLP morphotypes, no quantitative assessments were undertaken. To further our understanding of viral dynamics within the sponge holobiont, quantitative studies that count the number of VLPs per known tissue area, perform quantitative transmission electron microscopy (qTEM) (Brum, Schenck & Sullivan, 2013), flow cytometry (Brussaard, 2004; Pollock *et al.*, 2014) or fluorescent staining (Leruste, Bouvier & Bettarel, 2012; Pollard, 2012) should also be performed.

Morphology is an important feature for viral classification according to the International Committee on Taxonomy of Viruses (ICTV). However, there are also some limitations associated with using TEM to identify viruses. For instance, many viral groups lack morphological structures that characterize them as typical viral particles by TEM. Also, as many viruses are small and simple they can be mistaken for non-viral particles such as cellular vesicles or organelles. Although the assignment of viral-like particles in this study was made by comparison to morphologically characterised viruses, the possibility remains that some VLPs may not represent true viruses.

In this study, TEM analysis revealed a prevalence of polyhedral VLPs with characteristic bacteriophage morphology, consistent with what has been described for other marine invertebrates ([Wilson et al., 2005](#); [Davy et al., 2006](#); [Davy & Patten, 2007](#); [Patten, Harrison & Mitchell, 2008](#)). The presence of *Caudovirales*-like morphotypes highlights the potential for these VLPs to target sponge symbionts and ultimately control microbial population dynamics within the sponge holobiont. Amongst them, a *Siphoviridae* VLP detected in the surface biofilm of *C. schulzei* presented similar morphology, although slightly smaller, to the previously described sponge-associated Phage  $\Phi$ JL001 ([Lohr, Chen & Hill, 2005](#)).

Surprisingly, relatively few tailed bacteriophage were detected within the reef sponges, despite the dominance of *Caudovirales* within the assigned sponge viromes ([Laffy et al., 2018](#)). Although the dominance of tailed viruses in aquatic ecosystems is well characterised ([Mizuno et al., 2013](#); [Weynberg et al., 2017a](#); [Weynberg et al., 2017b](#); [Thurber et al., 2017](#); [Laffy et al., 2018](#)), results from morphological analysis of uncultivated viruses vary with respect to the relative dominance of tailed ([Cochlan et al., 1993](#); [Colombet et al., 2006](#); [Dutova & Drucker, 2013](#)) versus non-tailed ([Bergh et al., 1989](#); [Wommack et al., 1992](#); [Auguet, Montanié & Lebaron, 2006](#); [Brum, Schenck & Sullivan, 2013](#)) VLPs. The reduced number of tailed VLPs in morphological descriptions has been attributed to the destruction of the delicate VLP structures during centrifugation and TEM sample preparation ([Cochlan et al., 1993](#); [Proctor, 1997](#)). However, [Brum, Schenck & Sullivan \(2013\)](#) have shown that sample preservation and preparation do not alter the morphological characteristics of seawater derived VLPs ([Brum, Schenck & Sullivan, 2013](#)) and non-tailed VLP have therefore been proposed as the dominant viral group in aquatic ecosystems ([Brum, Schenck & Sullivan, 2013](#); [Kauffman et al., 2018](#)). Nevertheless, in this study, tailed VLPs were almost exclusively detected in samples purified via filtration of mucus or scraping of the external biofilm, the least disruptive of the three TEM preparation methods. This suggests that tailed VLPs are either more abundant on the external surface of the sponge or that the TEM preparation method could bias the detection of tailed VLPs in sponges by mechanically damaging or distorting viral structures.

Filamentous viral-like particles (FVLP) were detected in both prokaryotic and eukaryotic cells within the sponge holobiont. In *C. foliascens*, multiple individual *Inoviridae*-like VLPs were observed attached to the surface of cyanobacteria, although no virions were observed inside the cells. The absence of intracellular FVLPs combined with the absence of a dense core in these morphotypes provides further support for their classification as putative *Inoviridae*, as the replication mechanism of this viral family often relies on the virus injecting its DNA into the host cell and getting extruded without inducing cell lysis ([Bayer & Bayer, 1986](#); [Russel, 1991](#); [Ploss & Kuhn, 2010](#)). A previous study demonstrated that temperate viruses are relatively less abundant within host cells at high density ([McDaniel et al., 2002](#)).

FVLPs with helicoidal symmetry resembling *Spiraviridae* were detected in the sponge *C. cyathophora*, with this viral family known to infect Archaea ([Mochizuki et al., 2012](#)). FVLPs were also observed infecting eukaryotic cells in *X. testudinaria*. Abundant elongated and flexible FVLPs were also detected in the archaeocytes and extracellular mesohyl matrix of *X. testudinaria* (Figs. 4J–4L). The point of host cell lysis was captured with a recently burst

cell releasing virions into the extracellular matrix (Figs. 4J–4K), characteristic of typical lytic viral infection (Dyson *et al.*, 2015). Morphologically similar filamentous VLPs have been detected in coral mucus and associated *Symbiodinium* and were characterised as a coral-infecting RNA virus (Davy *et al.*, 2006; Weynberg *et al.*, 2017b). There is a general lack of studies investigating filamentous viruses in marine invertebrates, although metaviromic sequencing recently detected sequences assigned as filamentous viruses of the family Inoviridae in Great Barrier Reef sponges (Laffy *et al.*, 2018).

VLPs morphologically consistent with viruses from the family *Geminiviridae* were observed in association with cyanobacteria in the sponge *A. ochracea*. *Geminiviridae*-like viruses have been isolated from infected freshwater cyanobacteria (Li, Ou & Zhang, 2013), and, with the exception of being slightly smaller ( $79 \pm 5$  nm in length,  $28 \pm 3$  nm in diameter), the geminate VLPs from *A. ochracea* were morphologically similar. Most infected cyanobacterial cells had dense populations of these VLPs (Figs. 5D–5G), although no lysed cells or free geminate VLPs were observed in the sponge mesohyl. However, several extracellular vesicles containing VLPs were observed, indicating that VLPs could use cell extrusion as part of their reproductive cycle. A geminate VLP has previously been isolated from mucus secreted by scleractinian corals (Davy & Patten, 2007), however the morphology differs from the *A. ochracea* VLP, since it is notably bigger (about 145 nm in length, 82 nm width), with each isomer being wider than they are long, contrasting with the isomer dimensions in the *A. ochracea* VLP. Beaded VLPs were also detected in sponges and their non-isomeric particles comprising a flexible filament strongly resembled the beaded VLPs previously reported from scleractinian corals (Davy & Patten, 2007; Lawrence *et al.*, 2015).

Brick-shaped VLPs closely resembling viral morphotypes from the family *Poxviridae* were observed within the mesohyl of *Crella cyathophora*, a (Fig. 5H). Typical of enveloped viruses, poxviruses use their envelopes to connect and fuse with their host membrane so that the viral capsid is injected directly into the host cell (Moss, 2012). Poxviruses are notable pathogens, infecting a wide host range among vertebrate and invertebrate taxa (Bracht *et al.*, 2006; Grasis *et al.*, 2014; Haller *et al.*, 2014). In the marine environment, they have been reported associated with cetaceans and pinnipeds (Bracht *et al.*, 2006) and more recently, analysis of sponge metaviromes detected sequences affiliated to *Poxviridae* in *Amphimedon queenslandica* and *Ianthella basta* (Laffy *et al.*, 2018).

## CONCLUSION

In this study we validated the efficacy of three different methods for TEM imaging of sponge-associated viruses: (i) ultrathin sections of sponge tissue, (ii) purification via density gradient ultracentrifugation and (iii) ectoderm scraping and filtration of sponge mucus. While density gradient purification facilitated concentration and recovery of VLPs from different areas of the sponge holobiont, it also co-concentrated cellular debris, potentially masking many VLPs. Tissue sectioning enabled direct visualisation of spatial localisation and host-viral interactions but was labour intensive and some VLP structures were distorted during sectioning. Ectoderm scraping and collection of sponge mucus was

most effective at preserving delicate viral structures and minimizing the amount of cellular debris, however, it was restricted to recovering VLPs associated with the sponge mucus or ectoderm.

This first morphological characterisation of sponge-associated viruses revealed a wide diversity of VLPs infecting both the sponge cells and symbiont compartments of the holobiont. By confirming that viruses are a significant component of the sponge holobiont, this work paves the way for future metaviromic and cell culturing analyses that can characterise the taxonomy and function of the sponge viral community.

## ACKNOWLEDGEMENTS

The authors wish to acknowledge Dr. Karen Weynberg for support with the microscopy sample preparation and Dr. Rachid Sougrat and Dr. Ptissam Bergam for support with the operation of the TEM.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

Nicole S. Webster was funded through an Australian Research Council Future Fellowship FT120100480 and Cecília Pascelli was funded through a PhD Scholarship from The CAPES Foundation and Science without Borders. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:  
Australian Research Council Future Fellowship: FT120100480.  
The CAPES Foundation and Science.

### Competing Interests

The authors declare there are no competing interests.

### Author Contributions

- Cecília Pascelli conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Patrick W. Laffy analyzed the data, authored or reviewed drafts of the paper, approved the final draft.
- Marija Kupresanin performed the experiments, authored or reviewed drafts of the paper, approved the final draft.
- Timothy Ravasi contributed reagents/materials/analysis tools, authored or reviewed drafts of the paper, approved the final draft.
- Nicole S. Webster conceived and designed the experiments, analyzed the data, contributed reagents/materials/analysis tools, authored or reviewed drafts of the paper, approved the final draft.

### Field Study Permissions

The following information was supplied relating to field study approvals (i.e., approving body and any reference numbers):

Sampling in Australia was conducted under the Great Barrier Reef Marine Park Authority permit G12/35236.1

### Data Availability

The following information was supplied regarding data availability:

The raw data are provided in [Table S1](#).

### Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.5625#supplemental-information>.

## REFERENCES

- Andreakis N, Luter HM, Webster NS. 2012.** Cryptic speciation and phylogeographic relationships in the elephant ear sponge *Ianthella basta* (Porifera, Ianthellidae) from northern Australia. *Zoological Journal of the Linnean Society* **166**:225–235 DOI [10.1111/j.1096-3642.2012.00848.x](https://doi.org/10.1111/j.1096-3642.2012.00848.x).
- Auguet JC, Montanié H, Lebaron P. 2006.** Structure of virioplankton in the charente estuary (France): transmission electron microscopy versus pulsed field gel electrophoresis. *Microbial Ecology* **51**:197–208 DOI [10.1007/s00248-005-0043-0](https://doi.org/10.1007/s00248-005-0043-0).
- Bayer ME, Bayer MH. 1986.** Effects of bacteriophage fd infection on *Escherichia coli* HB11 envelope: a morphological and biochemical study. *Journal of Virology* **57**:258–266.
- Bergh Ø, Børsheim KY, Bratbak G, Haldal M. 1989.** High abundance of viruses found in aquatic environments. *Nature* **340**:467–468 DOI [10.1038/340467a0](https://doi.org/10.1038/340467a0).
- Bracht AJ, Brudek RL, Ewing RY, Manire CA, Burek KA, Rosa C, Beckmen KB, Maruniak JE, Romero CH. 2006.** Genetic identification of novel poxviruses of cetaceans and pinnipeds. *Archives of Virology* **151**:423–438 DOI [10.1007/s00705-005-0679-6](https://doi.org/10.1007/s00705-005-0679-6).
- Breitbart M, Bonnain C, Malki K, Sawaya NA. 2018.** Phage puppet masters of the marine microbial realm. *Nature Microbiology* **3**:754–766 DOI [10.1038/s41564-018-0166-y](https://doi.org/10.1038/s41564-018-0166-y).
- Breitbart M, Salamon P, Andresen B, Mahaffy JM, Segall AM, Mead D, Azam F, Rohwer F. 2002.** Genomic analysis of uncultured marine viral communities. *Proceedings of the National Academy of Sciences of the United States of America* **99**:14250–14255 DOI [10.1073/pnas.202488399](https://doi.org/10.1073/pnas.202488399).
- Brum JR, Schenck RO, Sullivan MB. 2013.** Global morphological analysis of marine viruses shows minimal regional variation and dominance of non-tailed viruses. *The ISME Journal* **7**:1738–1751 DOI [10.1038/ismej.2013.67](https://doi.org/10.1038/ismej.2013.67).
- Buller RM, Palumbo GJ. 1991.** Poxvirus pathogenesis. *Microbiological Reviews* **55**:80–122.
- Butina TV, Potapov SA, Belykh OI, Belikov SI. 2015.** Genetic diversity of cyanophages of the myoviridae family as a constituent of the associated community of the



- Baikal sponge *Lubomirskia baicalensis*. *Russian Journal of Genetics* **51**:313–317 DOI [10.1134/S1022795415030011](https://doi.org/10.1134/S1022795415030011).
- Chevillie NF, Stasko J. 2014.** Techniques in electron microscopy of animal tissue. *Veterinary Pathology* **51**:28–41 DOI [10.1177/0300985813505114](https://doi.org/10.1177/0300985813505114).
- Claverie JM, Grzela R, Lartigue A, Bernadac A, Nitsche S, Vacelet J, Ogata H, Abergel C. 2009.** Mimivirus and Mimiviridae: giant viruses with an increasing number of potential hosts, including corals and sponges. *Journal of Invertebrate Pathology* **101**:172–180 DOI [10.1016/j.jip.2009.03.011](https://doi.org/10.1016/j.jip.2009.03.011).
- Cochlan WP, Wikner J, Steward GF, Smith DC. 1993.** Spatial distribution of viruses, bacteria and chlorophyll a in neritic, oceanic and estuarine environments. *Marine Ecology Progress Series* **92**:77–87.
- Colombet J, Sime-Ngando T, Cauchie HM, Fonty G, Hoffmann L, Demeure G. 2006.** Depth-related gradients of viral activity in Lake Pavin. *Applied and Environmental Microbiology* **72**:4440–4445 DOI [10.1128/AEM.00021-06](https://doi.org/10.1128/AEM.00021-06).
- Danovaro R, Corinaldesi C, Dell'Anno A, Fuhrman JA, Middelburg JJ, Noble RT, Suttle CA. 2011.** Marine viruses and global climate change. *FEMS Microbiology Reviews* **35**:993–1034 DOI [10.1111/j.1574-6976.2010.00258.x](https://doi.org/10.1111/j.1574-6976.2010.00258.x).
- Davy JE, Patten NL. 2007.** Morphological diversity of virus-like particles within the surface microlayer of scleractinian corals. *Aquatic Microbial Ecology* **47**:37–44 DOI [10.3354/ame047037](https://doi.org/10.3354/ame047037).
- Davy S, Burchett S, Dale A, Davies P, Davy J, Muncke C, Hoegh-Guldberg O, Wilson W. 2006.** Viruses: agents of coral disease? *Diseases of Aquatic Organisms* **69**:101–110 DOI [10.3354/dao069101](https://doi.org/10.3354/dao069101).
- De Goeij JM, Van Oevelen D, Vermeij MJA, Osinga R, Middelburg JJ, De Goeij AFPM, Admiraal W. 2013.** Surviving in a marine desert: the sponge loop retains resources within coral reefs. *Science* **342**:108–110 DOI [10.1126/science.1241981](https://doi.org/10.1126/science.1241981).
- Dutova NV, Drucker VV. 2013.** Viral community of biofilms forming on different substrates under natural conditions of Lake Baikal. *Doklady Biological Sciences* **451**:238–240 DOI [10.1134/S0012496613030113](https://doi.org/10.1134/S0012496613030113).
- Dyson ZA, Tucci J, Seviour RJ, Petrovski S. 2015.** Lysis to kill: evaluation of the lytic abilities, and genomics of nine bacteriophages infective for *gordonia* spp. and their potential use in activated sludge foam biocontrol. *PLOS ONE* **10**:e0134512 DOI [10.1371/journal.pone.0134512](https://doi.org/10.1371/journal.pone.0134512).
- Erwin PM, Thacker RW. 2007.** Phylogenetic analyses of marine sponges within the order Verongida: a comparison of morphological and molecular data. *Invertebrate Biology* **126**:220–234 DOI [10.1111/j.1744-7410.2007.00092.x](https://doi.org/10.1111/j.1744-7410.2007.00092.x).
- Fan L, Liu M, Simister R, Webster NS, Thomas T. 2013.** Marine microbial symbiosis heats up: the phylogenetic and functional response of a sponge holobiont to thermal stress. *The ISME Journal* **7**:991–1002 DOI [10.1038/ismej.2012.165](https://doi.org/10.1038/ismej.2012.165).
- Fan L, Reynolds D, Liu M, Stark M, Kjelleberg S, Webster NS, Thomas T. 2012.** PNAS Plus: functional equivalence and evolutionary convergence in complex communities of microbial sponge symbionts. *Proceedings of the National Academy of Sciences of the United States of America* **109**:E1878–E1887 DOI [10.1073/pnas.1203287109](https://doi.org/10.1073/pnas.1203287109).

- Flores CO, Meyer JR, Valverde S, Farr L, Weitz JS. 2011. Statistical structure of host-phage interactions. *Proceedings of the National Academy of Sciences of the United States of America* **108**:E288–E297 DOI [10.1073/pnas.1101595108](https://doi.org/10.1073/pnas.1101595108).
- Flores CO, Valverde S, Weitz JS. 2013. Multi-scale structure and geographic drivers of cross-infection within marine bacteria and phages. *The ISME Journal* **7**:520–532 DOI [10.1038/ismej.2012.135](https://doi.org/10.1038/ismej.2012.135).
- Grasis JA, Lachnit T, Anton-Erxleben F, Lim YW, Schmieder R, Fraune S, Franzenburg S, Insua S, Machado G, Haynes M, Little M, Kimble R, Rosenstiel P, Rohwer FL, Bosch TCG. 2014. Species-specific viromes in the ancestral holobiont hydra. *PLOS ONE* **9**(10):e109952 DOI [10.1371/journal.pone.0109952](https://doi.org/10.1371/journal.pone.0109952).
- Hadas E, Marie D, Shpigel M, Ilan M. 2006. Virus predation by sponges is a new nutrient-flow pathway in coral reef food webs. *Limnology and Oceanography* **51**:1548–1550 DOI [10.4319/lo.2006.51.3.1548](https://doi.org/10.4319/lo.2006.51.3.1548).
- Haller SL, Peng C, McFadden G, Rothenburg S. 2014. Poxviruses and the evolution of host range and virulence. *Infection, Genetics and Evolution* **21**:15–40 DOI [10.1016/j.meegid.2013.10.014](https://doi.org/10.1016/j.meegid.2013.10.014).
- Hewson I, Bistolas KSI, Quijano Cardé EM, Button JB, Foster PJ, Flanzenbaum JM, Kocian J, Lewis CK. 2018. Investigating the complex association between viral ecology, environment, and northeast Pacific sea star wasting. *Frontiers in Marine Science* **5** DOI [10.3389/fmars.2018.00077](https://doi.org/10.3389/fmars.2018.00077).
- Holmfeldt K, Middelboe M, Nybroe O, Riemann L. 2007. Large variabilities in host strain susceptibility and phage host range govern interactions between lytic marine phages and their Flavobacterium hosts. *Applied and Environmental Microbiology* **73**:6730–6739 DOI [10.1128/AEM.01399-07](https://doi.org/10.1128/AEM.01399-07).
- Hooper JNA, Van Soest RWM. 2002. Systema Porifera. A guide to the classification of sponges. *Invertebrate Systematics* **18**:233–234 DOI [10.1007/978-1-4615-0747-5\\_1](https://doi.org/10.1007/978-1-4615-0747-5_1).
- Hurwitz BL, Westveld AH, Brum JR, Sullivan MB. 2014. Modeling ecological drivers in marine viral communities using comparative metagenomics and network analyses. *Proceedings of the National Academy of Sciences of the United States of America* **111**:10714–10719 DOI [10.1073/pnas.1319778111](https://doi.org/10.1073/pnas.1319778111).
- Kauffman KM, Hussain FA, Yang J, Arevalo P, Brown JM, Chang WK, Van Insberghe D, Elsherbini J, Sharma RS, Cutler MB, Kelly L, Polz MF. 2018. A major lineage of non-tailed dsDNA viruses as unrecognized killers of marine bacteria. *Nature* **554**:118–122 DOI [10.1038/nature25474](https://doi.org/10.1038/nature25474).
- Laffy PW, Wood-Charlson EM, Turaev D, Jutz S, Pascelli C, Botté ES, Bell SC, Peirce TE, Weynberg KD, van Oppen MJH, Rattei T, Webster NS. 2018. Reef invertebrate viromics: diversity, host specificity and functional capacity. *Environmental Microbiology* **20**(6):2125–2141 DOI [10.1111/1462-2920.14110](https://doi.org/10.1111/1462-2920.14110).
- Laffy PW, Wood-charlson EM, Turaev D, Weynberg KD, Botté ES, van Oppen MJH, Webster NS, Rattei T. 2016. HoloVir: a workflow for investigating the diversity and function of viruses in invertebrate holobionts. *Frontiers in Microbiology* **7**:1–15 DOI [10.3389/fmicb.2016.00822](https://doi.org/10.3389/fmicb.2016.00822).

- Lawrence JE, Steward GF. 2010.** Purification of viruses by centrifugation. In: Wilhelm SW, Weinbauer MG, Suttle CA, eds. *Manual of aquatic viral ecology*. American Society of Limnology and Oceanography, 166–181  
[DOI 10.4319/mave.2010.978-0-9845591-0-7.166](https://doi.org/10.4319/mave.2010.978-0-9845591-0-7.166).
- Lawrence S, Wilkinson S, Davy J, Arlidge W, Williams G, Wilson W, Aeby G, Davy S. 2015.** Influence of local environmental variables on the viral consortia associated with the coral *Montipora capitata* from Kaneohe Bay, Hawaii, USA. *Aquatic Microbial Ecology* 74:251–262 [DOI 10.3354/ame01743](https://doi.org/10.3354/ame01743).
- Leruste A, Bouvier T, Bettarel Y. 2012.** Enumerating Viruses in *Coral Mucus*. *Applied and Environmental Microbiology* 78:6377–6379 [DOI 10.1128/AEM.01141-12](https://doi.org/10.1128/AEM.01141-12).
- Li S, Ou T, Zhang Q. 2013.** Two virus-like particles that cause lytic infections in freshwater cyanobacteria. *Virologica Sinica* 28:303–305 [DOI 10.1007/s12250-013-3339-0](https://doi.org/10.1007/s12250-013-3339-0).
- Lohr JE, Chen F, Hill RT. 2005.** Genomic analysis of bacteriophage JL001: insights into its interaction with a sponge-associated alpha-proteobacterium. *Applied and Environmental Microbiology* 71:1598–1609 [DOI 10.1128/AEM.71.3.1598-1609.2005](https://doi.org/10.1128/AEM.71.3.1598-1609.2005).
- Luter HM, Whalan S, Webster NS. 2010.** Exploring the role of microorganisms in the disease-like syndrome affecting the sponge *Ianthella basta*. *Applied and Environmental Microbiology* 76:5736–5744 [DOI 10.1128/AEM.00653-10](https://doi.org/10.1128/AEM.00653-10).
- Margulis L, Fester R. 1991.** *Symbiosis as a source of evolutionary innovation: speciation and morphogenesis*. Cambridge: MIT press.
- McDaniel L, Houchin LA, Williamson SJ, Paul JH. 2002.** Lysogeny in marine *Synechococcus*. *Nature* 415:496–496 [DOI 10.1038/415496a](https://doi.org/10.1038/415496a).
- Meselson M, Stahl FW, Vinograd J. 1957.** Equilibrium sedimentation of macromolecules in density gradients. *Proceedings of the National Academy of Sciences of the United States of America* 43:581–588.
- Mochizuki T, Krupovic M, Pehau-Arnaudet G, Sako Y, Forterre P, Prangishvili D. 2012.** Archaeal virus with exceptional virion architecture and the largest single-stranded DNA genome. *Proceedings of the National Academy of Sciences of the United States of America* 109:13386–13391 [DOI 10.1073/pnas.1203668109](https://doi.org/10.1073/pnas.1203668109).
- Mojica KDA, Brussaard CPD. 2014.** Factors affecting virus dynamics and microbial host-virus interactions in marine environments. *FEMS Microbiology Ecology* 89:495–515 [DOI 10.1111/1574-6941.12343](https://doi.org/10.1111/1574-6941.12343).
- Moss B. 2012.** Poxvirus cell entry: how many proteins does it take? *Viruses* 4:688–707 [DOI 10.3390/v4050688](https://doi.org/10.3390/v4050688).
- Patten NL, Harrison PL, Mitchell JG. 2008.** Prevalence of virus-like particles within a staghorn scleractinian coral (*Acropora muricata*) from the Great Barrier Reef. *Coral Reefs* 27:569–580 [DOI 10.1007/s00338-008-0356-9](https://doi.org/10.1007/s00338-008-0356-9).
- Patterson MR, Chernykh VI, Fialkov VA, Savarese M. 1997.** Trophic effects of sponge feeding within Lake Baikal's littoral zone. 1. In situ pumping rates. *Limnology and Oceanography* 42:171–178 [DOI 10.4319/lo.1997.42.1.0171](https://doi.org/10.4319/lo.1997.42.1.0171).
- Pile AJ, Young CM. 2006.** The natural diet of a hexactinellid sponge: benthic—pelagic coupling in a deep-sea microbial food web. *Deep Sea Research Part I: Oceanographic Research Papers* 53:1148–1156 [DOI 10.1016/j.dsr.2006.03.008](https://doi.org/10.1016/j.dsr.2006.03.008).

- Pita L, Rix L, Slaby BM, Franke A, Hentschel U. 2018.** The sponge holobiont in a changing ocean: from microbes to ecosystems. *Microbiome* **6**:46 DOI [10.1186/s40168-018-0428-1](https://doi.org/10.1186/s40168-018-0428-1).
- Ploss M, Kuhn A. 2010.** Kinetics of filamentous phage assembly. *Physical Biology* **7**:45002 DOI [10.1088/1478-3975/7/4/045002](https://doi.org/10.1088/1478-3975/7/4/045002).
- Pollard PC. 2012.** Enumerating viruses by using fluorescence and the nature of the non-viral background fraction. *Applied and Environmental Microbiology* **78**:6615–6618 DOI [10.1128/AEM.01268-12](https://doi.org/10.1128/AEM.01268-12).
- Pollock F, Wood-Charlson E, van Oppen M, Bourne D, Willis B, Weynberg K. 2014.** Abundance and morphology of virus-like particles associated with the coral *Acropora hyacinthus* differ between healthy and white syndrome-infected states. *Marine Ecology Progress Series* **510**:39–43 DOI [10.3354/meps10927](https://doi.org/10.3354/meps10927).
- Proctor LM. 1997.** Advances in the study of marine viruses. *Microscopy Research and Technique* **37**:136–161 DOI [10.1002/\(SICI\)1097-0029\(19970415\)37:2<136::AID-JEMT3>3.0.CO;2-M](https://doi.org/10.1002/(SICI)1097-0029(19970415)37:2<136::AID-JEMT3>3.0.CO;2-M).
- Rohwer F, Thurber RV. 2009.** Viruses manipulate the marine environment. *Nature* **459**:207–212 DOI [10.1038/nature08060](https://doi.org/10.1038/nature08060).
- Rosario K, Breitbart M. 2011.** Exploring the viral world through metagenomics. *Current Opinion in Virology* **1**:289–297 DOI [10.1016/j.coviro.2011.06.004](https://doi.org/10.1016/j.coviro.2011.06.004).
- Roux S, Enault F, Hurwitz BL, Sullivan MB. 2015.** VirSorter: mining viral signal from microbial genomic data. *PeerJ* **3**:e985 DOI [10.7717/peerj.985](https://doi.org/10.7717/peerj.985).
- Russel M. 1991.** Filamentous phage assembly. *Molecular Microbiology* **5**:1607–1613 DOI [10.1111/j.1365-2958.1991.tb01907.x](https://doi.org/10.1111/j.1365-2958.1991.tb01907.x).
- Schmitt S, Tsai P, Bell J, Fromont J, Ilan M, Lindquist N, Perez T, Rodrigo A, Schupp PJ, Vacelet J, Webster N, Hentschel U, Taylor MW. 2012.** Assessing the complex sponge microbiota: core, variable and species-specific bacterial communities in marine sponges. *The ISME Journal* **6**:564–576 DOI [10.1038/ismej.2011.116](https://doi.org/10.1038/ismej.2011.116).
- Seymour J, Patten N, Bourne D, Mitchell J. 2005.** Spatial dynamics of virus-like particles and heterotrophic bacteria within a shallow coral reef system. *Marine Ecology Progress Series* **288**:1–8 DOI [10.3354/meps288001](https://doi.org/10.3354/meps288001).
- Suttle CA. 2007.** Marine viruses—major players in the global ecosystem. *Nature Reviews Microbiology* **5**:801–812 DOI [10.1038/nrmicro1750](https://doi.org/10.1038/nrmicro1750).
- Thomas T, Moitinho-Silva L, Lurgi M, Björk JR, Easson C, Astudillo-García C, Olson JB, Erwin PM, López-Legentil S, Luter H, Chaves-Fonnegra A, Costa R, Schupp PJ, Steindler L, Erpenbeck D, Gilbert J, Knight R, Ackermann G, Victor Lopez J, Taylor MW, Thacker RW, Montoya JM, Hentschel U, Webster NS. 2016.** Diversity, structure and convergent evolution of the global sponge microbiome. *Nature Communications* **7**:11870 DOI [10.1038/ncomms11870](https://doi.org/10.1038/ncomms11870).
- Thomassen S, Riisgård H. 1995.** Growth and energetics of the sponge *Halichondria panicea*. *Marine Ecology Progress Series* **128**:239–246 DOI [10.3354/meps128239](https://doi.org/10.3354/meps128239).
- Thurber RLV, Correa AMS. 2011.** Viruses of reef-building scleractinian corals. *Journal of Experimental Marine Biology and Ecology* **408**:102–113 DOI [10.1016/j.jembe.2011.07.030](https://doi.org/10.1016/j.jembe.2011.07.030).

- Thurber RV, Payet JP, Thurber AR, Correa AMS. 2017.** Virus—host interactions and their roles in coral reef health and disease. *Nature Reviews Microbiology* 15:205–216 DOI [10.1038/nrmicro.2016.176](https://doi.org/10.1038/nrmicro.2016.176).
- Vacelet J, Boury-Esnault N. 1995.** Carnivorous sponges. *Nature* 373:333–335 DOI [10.1038/373333a0](https://doi.org/10.1038/373333a0).
- Vacelet J, Gallissian M-F. 1978.** Virus-like particles in cells of the sponge *Verongia cavernicola* (demospongiae, dictyoceratida) and accompanying tissues changes. *Journal of Invertebrate Pathology* 31:246–254 DOI [10.1016/0022-2011\(78\)90014-9](https://doi.org/10.1016/0022-2011(78)90014-9).
- Van Soest RWM, Boury-Esnault N, Vacelet J, Dohrmann M, Erpenbeck D, De Voogd NJ, Santodomingo N, Vanhoorne B, Kelly M, Hooper JNA. 2012.** Global diversity of sponges (Porifera). *PLOS ONE* 7:e35105 DOI [10.1371/journal.pone.0035105](https://doi.org/10.1371/journal.pone.0035105).
- Webster NS, Taylor MW. 2012.** Marine sponges and their microbial symbionts: love and other relationships. *Environmental Microbiology* 14:335–346 DOI [10.1111/j.1462-2920.2011.02460.x](https://doi.org/10.1111/j.1462-2920.2011.02460.x).
- Webster NS, Thomas T. 2016.** The sponge hologenome. *mBio* 7:e00135–16 DOI [10.1128/mBio.00135-16](https://doi.org/10.1128/mBio.00135-16).
- Weynberg KD, Laffy PW, Wood-Charlson EM, Turaev D, Rattei T, Webster NS, van Oppen MJH. 2017a.** Coral-associated viral communities show high levels of diversity and host auxiliary functions. *PeerJ* 5:e4054 DOI [10.7717/peerj.4054](https://doi.org/10.7717/peerj.4054).
- Weynberg KD, Neave M, Clode PL, Voolstra CR, Brownlee C, Laffy P, Webster NS, Levin RA, Wood-Charlson EM, van Oppen MJH. 2017b.** Prevalent and persistent viral infection in cultures of the coral algal endosymbiont *Symbiodinium*. *Coral Reefs* 36:773–784 DOI [10.1007/s00338-017-1568-7](https://doi.org/10.1007/s00338-017-1568-7).
- Weynberg KD, Wood-Charlson EM, Suttle CA, van Oppen MJH. 2014.** Generating viral metagenomes from the coral holobiont. *Frontiers in Microbiology* 5:1–11 DOI [10.3389/fmicb.2014.00206](https://doi.org/10.3389/fmicb.2014.00206).
- Wilson WH, Chapman DM. 2001.** Observation of virus-like particles in thin sections of the plumose anemone, *Metridium senile*. *Journal of the Marine Biological Association of the UK* 81:879–880 DOI [10.1017/S0025315401004726](https://doi.org/10.1017/S0025315401004726).
- Wilson WH, Dale AL, Davy JE, Davy SK. 2005.** An enemy within? Observations of virus-like particles in reef corals. *Coral Reefs* 24:145–148 DOI [10.1007/s00338-004-0448-0](https://doi.org/10.1007/s00338-004-0448-0).
- Wommack KE, Hill RT, Kessel M, Russek-Cohen E, Colwell RR. 1992.** Distribution of viruses in the Chesapeake Bay. *Applied and Environmental Microbiology* 58:2965–2970.
- Wood-Charlson EM, Weynberg KD, Suttle CA, Roux S, van Oppen MJH. 2015.** Metagenomic characterization of viral communities in corals: mining biological signal from methodological noise. *Environmental Microbiology* 17:3440–3449 DOI [10.1111/1462-2920.12803](https://doi.org/10.1111/1462-2920.12803).
- Wörheide G, Dohrmann M, Erpenbeck D, Larroux C, Maldonado M, Voigt O, Borchiellini C, Lavrov DV. 2012.** Deep phylogeny and evolution of sponges (Phylum Porifera). *Advances in Marine Biology* 61:1–78 DOI [10.1016/B978-0-12-387787-1.00007-6](https://doi.org/10.1016/B978-0-12-387787-1.00007-6).