

# The Great Lakes Entomologist

---

Volume 18  
Number 2 - Summer 1985 *Number 2 - Summer*  
1985

Article 6

---

June 1985

## *Archilestes Grandis* in Wisconsin (Odonata: Lestidae)

Susan Sullivan Borkin  
Milwaukee Public Museum

Follow this and additional works at: <https://scholar.valpo.edu/tgle>



Part of the [Entomology Commons](#)

---

### Recommended Citation

Borkin, Susan Sullivan 1985. "*Archilestes Grandis* in Wisconsin (Odonata: Lestidae)," *The Great Lakes Entomologist*, vol 18 (2)

Available at: <https://scholar.valpo.edu/tgle/vol18/iss2/6>

This Peer-Review Article is brought to you for free and open access by the Department of Biology at ValpoScholar. It has been accepted for inclusion in *The Great Lakes Entomologist* by an authorized administrator of ValpoScholar. For more information, please contact a ValpoScholar staff member at [scholar@valpo.edu](mailto:scholar@valpo.edu).

**ARCHILESTES GRANDIS IN WISCONSIN (ODONATA: LESTIDAE)***Susan Sullivan Borkin*<sup>1</sup>

*Archilestes grandis* (Rambur) is the largest damselfly that occurs in America north of Mexico. Its distribution in the United States was reported by Muttkowski (1910) as Lower Sonoran, Washington to Baja California, and Texas. It was collected east of the Mississippi River for the first time in 1927 by E. B. Williamson (1931) in Ohio, and since that time has extended its range along the southern and mid-Atlantic states (Harwood 1973, Kennedy 1977, Tennessen 1979), and north of Texas to South Dakota (Gloyd 1980). With this history of recent expansion, it comes as no surprise to find *A. grandis* in Wisconsin.

I collected one pair of *A. grandis* in copula on 8 September 1978 in Milwaukee County, from vegetation along Honey Creek near its junction with the Menomonee River. No individuals were discovered at the site on 5 August 1982, but on 20 September 1984 after a two-hour search, one male was collected from a perch in Honey Creek and one female was taken from a tree trunk nearby. No individuals were found along the Menomonee River. The water was slow-moving and shallow in both the river and Honey Creek. Willow branches overhanging the water were examined for oviposition punctures but none were discovered. If *A. grandis* is able to overwinter here, presumably it hibernates in the egg stage (Ingram 1976).

## LITERATURE CITED

- Gloyd, L. K. 1980. The taxonomic status of the genera *Superlestes* and *Cyptolestes* Williamson, 1921. (Odonata: Lestidae). Occ. Pap. Mus. Zool. Univ. Michigan 694:1-3.
- Harwood, P. D. 1973. Notes on damselflies from West Virginia. Proc. West Virginia Acad. Sci. 45:167-169.
- Ingram, B. R. 1976. Life histories of three species of Lestidae in North Carolina (Odonata: Zygoptera). Odonatologica 5:231-244.
- Kennedy, J. H. 1977. The occurrence of *Archilestes grandis* Rambur (Zygoptera: Lestidae) in Virginia. Entomol. News 88:215-216.
- Muttkowski, R. A. 1910. Catalogue of the Odonata of North America. Bull. Public Mus., Milwaukee. No. 1 207 p.
- Tennessen, K. J. 1979. New records of Odonata for Alabama and Tennessee, with significant range extensions for several species. Entomol. News 90:118-120.
- Williamson, E. B. 1931. *Archilestes grandis* (Ramb.) in Ohio (Odonata: Agrionidae). Entomol. News 42:63-64.

---

<sup>1</sup>Invertebrate Zoology Section, Milwaukee Public Museum, Milwaukee, WI 53233.