



University of the Pacific
Scholarly Commons

Dugoni School of Dentistry Faculty Articles

Arthur A. Dugoni School of Dentistry

Fall 9-1-2008

Expanding circle of inhibition: small-molecule inhibitors of Bcl-2 as anticancer cell and antiangiogenic agents

Benjamin David Zeitlin

University of the Pacific, bzeitlin@pacific.edu

Isaac J. Zeitlin

University of Strathclyde

Jacques E. Nör

University of Michigan School of Dentistry, jenor@umich.edu

Follow this and additional works at: <https://scholarlycommons.pacific.edu/dugoni-facarticles>

 Part of the [Dentistry Commons](#)

Recommended Citation

Zeitlin, B. D., Zeitlin, I. J., & Nör, J. E. (2008). Expanding circle of inhibition: small-molecule inhibitors of Bcl-2 as anticancer cell and antiangiogenic agents. *Journal of Clinical Oncology*, 26(25), 4180–4188. DOI: [10.1200/JCO.2007.15.7693](https://doi.org/10.1200/JCO.2007.15.7693)
<https://scholarlycommons.pacific.edu/dugoni-facarticles/390>

This Article is brought to you for free and open access by the Arthur A. Dugoni School of Dentistry at Scholarly Commons. It has been accepted for inclusion in Dugoni School of Dentistry Faculty Articles by an authorized administrator of Scholarly Commons. For more information, please contact mgibney@pacific.edu.

Expanding Circle of Inhibition: Small-Molecule Inhibitors of Bcl-2 as Anticancer Cell and Antiangiogenic Agents

Benjamin D. Zeitlin, Isaac J. Zeitlin, and Jacques E. Nör

ABSTRACT

The specific targeting of diseases, particularly cancer, is a primary aim in drug development, as specificity reduces unwelcome effects on healthy tissue and increases drug efficacy at the target site. Drug specificity can be increased by improving the delivery system or by selecting drugs with affinity for a molecular ligand specific to the disease state. The role of the prosurvival Bcl-2 protein in maintaining the normal balance between apoptosis and cellular survival has been recognized for more than a decade. Bcl-2 is vital during development, much less so in adults. It has also been noted that some cancers evade apoptosis and obtain a survival advantage through aberrant expression of Bcl-2. The new and remarkably diverse class of drugs, small-molecule inhibitors of Bcl-2 (molecular weight approximately 400 to 800 Daltons), is examined herein. We present the activities of these compounds along with clinical observations, where available. The effects of Bcl-2 inhibition on attenuation of tumor cell growth are discussed, as are studies revealing the potential for Bcl-2 inhibitors as antiangiogenic agents. Despite an enormous body of work published for the Bcl-2 family of proteins, we are still learning exactly how this group of molecules interacts and indeed what they do. The small-molecule inhibitors of Bcl-2, in addition to their therapeutic potential, are proving to be an important investigative tool for understanding the function of Bcl-2.

J Clin Oncol 26:4180-4188. © 2008 by American Society of Clinical Oncology

INTRODUCTION

Evolution has adapted us more poorly to fight cancer compared with almost any other disease, and until recent years, approaches to treatment of cancer had changed little. Physical removal of the tumor by surgery remains an important first-line treatment but lacks effectiveness in the face of highly aggressive or invasive cancers, cancers that are difficult to detect, or those that have reached a metastatic stage. Radiotherapy and chemotherapy are the conventional second-line treatments; however, both are nonspecific to the tumor tissue. They are generally toxic to healthy tissue and, notably for DNA-directed drugs such as cisplatin (*cis*-diamminedichloroplatinum), may ultimately be carcinogenic in their own right.^{1,2}

This review deals with a novel class of targeted drugs developed by several independent groups over the last few years: small-molecule inhibitors of the antiapoptotic protein Bcl-2 (B-cell lymphoma/leukemia-2). The antitumor and antiangiogenic potential of these compounds will be discussed. In the literature, the term Bcl-2 inhibitor encompasses various drugs that bind the antiapoptotic Bcl-2 family members with more or less effi-

cacy. None of the compounds here discussed are specific to only one member of the Bcl-2 family.

BCL-2-RELATED PROTEIN FAMILY

Bcl-2 is a 26-kDa protein localized to the mitochondria and endoplasmic reticulum that acts as a key inhibitor of apoptosis. Expression of Bcl-2 is essential for growth of certain tumor cell lines in vitro and has been found upregulated in a variety of tumor types in vivo.³⁻⁵ The interactions of other Bcl-2 family members and their role in modulating Bcl-2 functions have been reviewed,^{6,7} and a detailed discussion of such is beyond the scope of this review. A brief description here may aid the appreciation of drug actions. Bcl-2 and other members of the family are primary regulators of apoptosis (Fig 1). They maintain a balance within the cell that is biased toward survival. Bcl-2 structure consists of five domains, Bcl-2 homology (BH) domains 1 to 4 plus a transmembrane domain (Fig 2). The other Bcl-2 family members bear four or fewer of these domains and display varying degrees of homology (Fig 2). The BH1-3 domains can form a labile hydrophobic groove, on tertiary folding, that allows binding of other molecules via their own BH3 domain. This

From the Department of Restorative Sciences, University of Michigan School of Dentistry, and University of Michigan Comprehensive Cancer Center, Ann Arbor, MI; and University of Strathclyde, Glasgow, Scotland, United Kingdom.

Submitted December 12, 2007; accepted April 29, 2008.

Supported by Grants No. R01-DE14601, R01-DE15948, and R01-DE16586 from the National Institutes of Health/National Institute of Dental and Craniofacial Research (J.E.N.) and a developmental project from the University of Michigan Head and Neck Specialized Program of Research Excellence (J.E.N.).

Terms in blue are defined in the glossary, found at the end of this article and online at www.jco.org.

Authors' disclosures of potential conflicts of interest and author contributions are found at the end of this article.

Corresponding author: Jacques E. Nör, PhD, Angiogenesis Research Laboratory, University of Michigan, 1011 N University Rm 2309, Ann Arbor, MI 48109-1078; e-mail: jenor@umich.edu.

© 2008 by American Society of Clinical Oncology

0732-183X/08/2625-4180/\$20.00

DOI: 10.1200/JCO.2007.15.7693




Fig 1. Role of Bcl-2 in tumor cell and endothelial cell apoptosis and survival. Diagram depicts the involvement of Bcl-2 in tumor cell and endothelial cell function, as well as in the regulation of cross-talks between the two cell types. Bcl-2 is a pivotal checkpoint in the apoptotic pathway, inhibiting the proapoptotic actions of Bax and Bak. Both Bcl-2 expression and hypoxia may result in vascular endothelial growth factor (VEGF) production, which is secreted by the tumor cell. Endogenous VEGF can bind to VEGF receptor 1 (VEGFR1) on tumor cells sending a pro-survival signal, upregulating Bcl-2 production, or can also bind to VEGFR2 on endothelial cells, promoting cell survival and stimulating proliferation and angiogenesis via Bcl-2-dependent pathways.

hydrophobic groove is called the BH3-binding domain. In general terms, these domains define the functionality of three groups within the Bcl-2 family: two groups of either proapoptotic or antiapoptotic effector molecules bearing multiple BH domains and one group of facilitator or potentiator molecules bearing only the BH3 domain (BH3-only proteins; Fig 2). The proapoptotic effectors Bax and Bak form oligomeric pores in the mitochondrial membrane that allow egress of apoptotic signaling molecules, whereas the antiapoptotic effectors Bcl-2, Bcl-x_L, Mcl-1, and Bfl-1/A1 bind to Bax or Bak and inhibit functional oligomerization. The proapoptotic BH3-only molecules may interact with the effector molecules to promote pore formation either by competitively inhibiting Bcl-2/Bcl-x_L/Mcl-1, binding to Bax/Bak, or by directly promoting Bax/Bak oligomerization.^{6,8} There are a number of models, variations on the above theme, proposed to explain the interaction of these proteins and how, or even whether, they induce apoptosis signaling via the mitochondria. Ex-

actly how this happens is unclear in real cellular models at this time (see Skommer et al⁶ for further discussion).

BCL-2: ROLE IN ANGIOGENESIS AND TUMOR STATUS

Angiogenesis is the mechanism by which blood vessels form from existing vasculature.⁹ It is fundamental to normal growth and development and as part of the physiologic response to wounding. Hyperplastic cell growth both encourages and requires formation of new blood vessels, as evidenced by the generally high vascularity of tumors. Indeed, beyond 1- to 2-mm size, tumors cannot sustain expansion without induction of angiogenesis. Simple diffusion of nutrients is clearly insufficient to maintain tumor growth.¹⁰ Three of the primary activators of angiogenesis are vascular endothelial growth factor (VEGF), CXCL1 (growth-related oncogene, GRO-α), and CXCL8 (interleukin-8), promoting endothelial cell survival and proliferation.^{11,12}

The involvement of Bcl-2 as a proangiogenic signaling molecule is only now becoming clearer for both tumor cells and vascular endothelial cells (Fig 1). Indeed it has long been recognized that both Bcl-2 and Bfl-1/A1 are mediators of protection for endothelial cells,¹³ although Bfl-1/A1 seems to perform a rapid, and more transient, function than Bcl-2, potentially related to inflammatory response.^{13,14} Although modulation of Bcl-2 in endothelial cells has been shown to affect angiogenesis in vitro and in vivo (discussed further in this paragraph and in Bcl-2 Inhibitors as Antiangiogenic Agents), little is known about the molecular mechanisms involved. Stimulation of the VEGF pathway results in increased expression of Bcl-2 in both tumor cells¹⁵ and endothelial cells.^{13,16,17} Bcl-2 upregulation in turn increases VEGF expression in endothelial cells and in tumor cells of various lineages.¹⁸⁻²⁰ Notably, Bcl-2 upregulation induces expression of the proangiogenic chemokines CXCL1 and CXCL8 through activation of the NF-κB signaling pathway in endothelial cells.^{16,21} It has recently been shown in human tumor biopsies that Bcl-2 is massively upregulated in head and neck squamous cell carcinoma-associated endothelial cells compared with endothelial cells in normal oral mucosa.²⁰ In this study, it was also demonstrated in vivo that factors secreted by endothelial cells, in response to modulation of Bcl-2 expression levels in neovascular endothelial cells, have a direct effect on tumor cell




Fig 2. Bcl-2 family of proteins. This is a nonexhaustive list of the Bcl-2 family of proteins, showing the conserved Bcl-2 homology (BH) domains, the transmembrane domains (TM), and the three primary classes of molecular activity. Bcl-2, Bcl-x_L, Mcl-1, and Bcl-w represent the major molecules in the antiapoptotic class of effector molecules. Bax, Bak, and Bok represent the proapoptotic effector molecules, lacking BH4 domain. The proapoptotic facilitator molecules bear only the BH3 domain and may lack the transmembrane domain. Sizes are representative and drawn to approximate scale relative to molecular weight.

BCL-2 INHIBITORS AS ANTITUMOR AGENTS

growth.²⁰ Additionally, the authors showed expression of VEGF to be significantly attenuated *in vitro* by small inhibitory RNA directed against Bcl-2, in both Bcl-2 overexpressing and control endothelial cells.²⁰ It is clear that disruption of the Bcl-2 pathway may be expected to affect angiogenesis, both directly by inhibiting function of endothelial cells and, as many cancers display increased Bcl-2 expression, by concomitant reduction in levels of endothelial-stimulating factors, such as VEGF, *CXCL1*, and *CXCL8* (Fig 1). Recently, endothelium-targeted Bcl-2 overexpression was shown to induce disruption of the blood vessel architecture and induce embryonic lethality in transgenic mice.²² This effect was limited to the microvasculature and related to a reduction in endothelial apoptosis, thus directly linking Bcl-2 levels during development to angiogenic function.²²

It should be noted that the clinical picture of Bcl-2 involvement in cancer progression or patient survival is unclear, with different studies finding varying degrees of correlation between Bcl-2 expression and disease severity or prognosis, both within and between cancer types.^{23,24} Moreover, there is no clear reason why Bcl-2 expression, as a negative or positive prognostic factor, should bear any relation to whether Bcl-2 is considered as a beneficial therapeutic target. One example of this dichotomy is that of small-cell lung cancer (SCLC) or non-small-cell lung cancer (NSCLC). Meta-analysis of the data from 28 clinical studies indicated that Bcl-2 expression was associated with good prognosis in NSCLC but had no correlation with disease state in SCLC.²⁵ Furthermore, some individual studies have indeed shown correlation between Bcl-2 or Bcl-x_L expression and poor disease prognosis in NSCLC.^{26,27} However, *in vitro* and *in vivo* studies consistently correlate increased Bcl-2 or Bcl-x_L expression with tumor aggression and tumor cell line resistance to chemotherapeutic drugs.²⁸⁻³⁰ They have further shown regression and reversal of drug resistance, by treatment with Bcl-2 family inhibitors, in SCLC tumors and others.^{29,30}

The disparity between tumor expression and therapeutic potential may result from varying methods of analysis, but may also be explained at the cellular level. This is illustrated by two studies. In the first, an acute lymphoblastic leukemia cell line treated with VEGF expressed a phosphorylated form of Bcl-2, and apoptosis was reduced.^{31,32} The expression of nonphosphorylatable Bcl-2 in these cells prevented this antiapoptotic effect. In contrast, the same authors³² also showed that VEGF-induced phosphorylation of Bcl-2 did not protect chronic lymphoblastic leukemia cells under similar conditions. Expression levels and post-translational modification of other pro- or antiapoptotic proteins, in particular the BH3-only proteins, may also account for the apparent differences.³³⁻³⁵ Furthermore, inactivation of Bcl-x_L by de-amidation occurs in response to common chemotherapeutic drugs, such as cisplatin.³⁶ With this number of variables, it is clear that simply looking at the expression level of Bcl-2 alone in a tumor is insufficient to predict the benefits of anti-Bcl-2 therapy. Profiling the expression of many specific markers within a tumor may provide a better indication for the suitability of therapies targeting individual pathways, but this analysis is not routinely performed. Accordingly, quantifiable RNA analysis of gene expression and antibody arrays or Western blot analysis of proteins could usefully supplement immunohistochemical data from tumor biopsies in future studies. Thus, at this stage, it is difficult to predict which cancers would benefit most from treatment with Bcl-2 inhibitors and in combination with which conventional therapies.

Both internal signaling and external stimuli may alter the complex balance toward apoptosis by activation of the proapoptotic BH3-only molecules and subsequent inactivation of Bcl-2/Bcl-x_L, thus tipping the balance in favor of apoptosis. Much research has been carried out attempting to exploit the BH3-binding permissive network toward development of Bcl-2/Bcl-x_L inhibitors that result in activation of Bax and Bak. A number of genetic and pharmacologic approaches have been used to this end. Targeting Bcl-2 family RNA has shown positive results, and *oblimersen* (Genasense; G3139; Genta Inc, Berkeley Heights, NJ), an anti-Bcl-2 antisense oligonucleotide, has reached phase III clinical trials in combination therapy.³⁷ Peptide-based drugs have also been shown both to attenuate Bcl-2 activity³⁸ and to activate Bax.⁸ However, nonpeptidic small-molecule inhibitors of signaling pathways are likely to remain developmental drugs of choice for the short to medium term, particularly because of the inherent low antigenicity of nonprotein organic compounds. In addition, the ease and plasticity of molecular modification allows ready tailoring of putative drugs. This last allows manipulation of compounds to aid route-specific delivery, increase bioavailability, increase target affinity, and, hopefully, to reduce patient drug load. This is a relatively new area for Bcl-2 inhibitors, and currently few compounds have been under developmental and clinical study since the start of the decade. Only within the last 6 to 12 months have publications appeared on a newer, larger group of compounds. The established compounds with available experimental data are discussed next in the context of cancer and angiogenesis. In this review, these compounds have been categorized as derived from natural sources (see "From Nature") or compounds that were designed in laboratories (see "From Desktop to Bedside").

From Nature

The first drugs to be examined for therapeutic potential in many diseases are generally pre-existing compounds that may be in use for ostensibly unrelated purposes. Even where they have been examined for anticancer activity, their molecular targets are rarely known in advance. This has certainly been true for Bcl-2 inhibitors. The health benefits of green and black tea polyphenols, in particular catechins and theaflavins, have been touted as almost the nearest equivalent to natural cure-alls. Indeed they do seem to have remarkably broad therapeutic properties according to the literature³⁹⁻⁴² and are commonly listed as small-molecule Bcl-2 inhibitors in cancer-related publications. Work from the Burnham Institute for Medical Research, using nuclear magnetic resonance structural data and binding displacement studies, provided clear evidence of binding between green or black tea polyphenols and Bcl-2 and Bcl-x_L.^{43,44} They also readily induce apoptosis in tumor cells *in vitro* and *in vivo*.⁴⁵⁻⁴⁸ However, tea polyphenols, and in particular the major active component epigallocatechin-3-gallate (EGCG; Fig 3F, Table 1), have been shown to act as highly specific ligands for a variety of cellular pathways other than those of Bcl-2. These include inhibition of platelet-derived growth factor receptor ligand binding,⁷¹ tyrosine phosphorylation of the Her/Neu epidermal growth factor receptor,⁷² and signaling by the metastasis-associated 67-kDa laminin receptor.⁷³ Expression of this last receptor conferred specific responsiveness on cancer cells to the growth inhibitory effects of EGCG and bound EGCG with a K_d of 39.9 nmol/L.⁷³ In addition to these extracellular targets, tea polyphenols




Fig 3. Small-molecule inhibitors of Bcl-2 family proteins. Chemical structures of the various small-molecule inhibitors of Bcl-2 discussed in this review. EGCG, epigallocatechin-3-gallate.

(EGCG in particular) have been shown to bind to, or directly affect, the pathways of a number of signaling molecules, including cyclooxygenase and retinoblastoma protein.^{74,75} These all may have equal claim, alongside Bcl-2, to be the cancer cell inhibitory targets for this class of compounds. Therefore, although tea polyphenols remain an exciting area for investigation as active antitumor-cell drugs, it is difficult to accurately describe them as primarily acting via the Bcl-2 pathway.

The other major natural derivative in this class, gossypol, was isolated from the cotton plant and has been well studied as a contra-

ceptive for human males.⁷⁶ However, gossypol, and in particular the (–)-gossypol enantiomer (AT-101; Ascenta Therapeutics Inc, Malvern, PA; Fig 3C, Table 1), was found to bind to Bcl-2 and Bcl-x_L through fluorescence-based ligand competition assay and nuclear magnetic resonance binding analysis.⁴⁹ In contrast with tea polyphenols, there is a preponderance of evidence to suggest that the Bcl-2 family is a specific target of gossypol for induction of apoptosis. Although gossypol does bind to other molecules, calcineurin, glutathione-s-transferase, and lactate dehydrogenase with good affinity, targeting the Bcl-2 pathways seems the most likely mechanism for

Table 1. Efficacies of Small-Molecule Inhibitors of Bcl-2

Small-Molecule Inhibitor	IC ₅₀ Range (μM)*	Binding Coefficients†			Reported Range In Vivo		Studies (first author)
		Bcl-2	Bcl-x _L	Mcl-1	Therapeutic	MTD (mg/kg qd)	
Gossypol	2.2-13.2	0.28-50	0.4-3.03	1.75	5-30 mg/kg	3 × 40	Kitada, ⁴⁹ Zhang, ⁵⁰ Mohammad, ⁵¹ Wolter, ⁵² Bauer, ²⁹ Xu, ⁵³ Zeitlin, ⁵⁵ Van Poznak ⁵⁷
HA14-1	5-20	0.9-100	NA	NA	400 nmol (approximately 136 μg)/mouse	NA	van Delft, ⁷ Wang, ⁵⁴ Lickliter, ⁵⁶ Chen, ⁵⁸ Manero ⁵⁹
TW-37	0.2-0.29	0.12-0.7	1.11	0.26	3-30 mg/kg qd	3 × 40	Wang, ⁶⁰ Mohammad, ⁶¹ Zeitlin, ⁵⁵ Verhaegen ⁶²
GX15-070	1.7	1.11‡	4.69‡	2.9‡	0.25-3 mg/kg/d iv	NA	Zhai, ⁴³ Campàs, ⁶³ Nguyen ⁶⁴
ABT-737	0.13-5	0.12	0.064	> 20	20-100 mg/kg/ip qd	NA	van Delft, ⁷ Oltersdorf, ³⁰ Konopleva, ⁶⁵ Zhai ⁴³
EGCG	20-65	0.45	0.59	0.92	1.5 mg/d ip	NA	Yang, ⁴⁵ Pan, ⁴⁶ Navarro-Perán, ⁴⁷ Jung ⁴⁸
Antimycin A and derivatives	0.37-0.7	0.82-100	2.7	2.51	2 mg/kg ip	NA	van Delft, ⁷ Zhai, ⁴³ Tzung, ⁶⁶ Manion, ⁶⁷ Kim, ⁶⁸ Cao, ⁶⁹ Guichard ⁷⁰

NOTE. Values are representative ranges of activities for drugs in a variety of cells and conditions. Lack of consistent and comparable studies between various compounds makes more exact reporting of values difficult to interpret.

Abbreviations: IC₅₀, half maximal inhibitory concentration; MTD, maximum-tolerated dose; qd, each day; NA, not available at time of writing; iv, intravenously; ip, intraperitoneally.

*Refers to reported IC₅₀, half maximal effective concentration (EC₅₀), or median lethal dose (LD₅₀).

†Determined using varying competitive binding assays.

‡Data for GX15, derivative of clinically tested GX15-070.

induction of apoptosis.⁷⁷⁻⁷⁹ Higher Bcl-2 expression levels in tumor cell lines have been correlated with increased apoptotic efficacy of gossypol.^{80,81} Notably, in addition to a direct Bcl-2 inhibitory action, gossypol modulates Bcl-2 family balance by downregulating the antiapoptotic members of the Bcl-2 family and upregulating the proapoptotic members in a concentration and cell-line specific manner.^{50,82,83} As a single agent, the (–)-gossypol enantiomer has shown encouraging efficacy *in vivo* in a head and neck cancer model.⁵² (–)-Gossypol has also been shown to overcome cisplatin resistance *in vitro* in a head and neck cancer line.²⁹ Two recent *in vivo* studies have reported a cooperative effect for administration of (–)-gossypol in conjunction either with chemotherapy (cyclophosphamide, doxorubicin, vincristine, and prednisolone regimen) or radiotherapy.^{51,53} A significant decrease in xenograft tumor volume (cells from a patient with clinically resistant diffuse large-cell lymphoma) was observed when animals were treated with (–)-gossypol/ cyclophosphamide, doxorubicin, vincristine, and prednisolone chemotherapy as compared with either individual treatment alone.

Mixed results were seen for racemic gossypol in phase I/II trials, but no severe toxicity was observed, except during determination of maximum-tolerated doses, where severe emesis was the dose-limiting toxicity.^{57,84-86} Currently, phase I/II clinical trials are underway for (–)-gossypol (AT-101) in single-agent treatment of relapsed B-cell malignancy. The compound is also being assessed in combination therapy with docetaxel and prednisone for hormone-refractory prostate cancer and also with an anti-CD20 antibody, rituximab, for chronic lymphocytic leukemia (see clinical trial listing at www.cancer.gov). Two further clinical trials have been approved for relapsed/refractory SCLC and newly diagnosed glioblastoma multiforme (www.cancer.gov).

Also under development are derivatives of Antimycin A, isolated from *Streptomyces*. Antimycin A (Fig 3G, Table 1) is an antifungal compound and a mitochondrial poison that targets complex III of the respiratory chain and that was shown to bind to the BH3 binding domain of Bcl-2/Bcl-x_L.^{43,66,67} It is highly toxic in mouse models, with a single median lethal dose of 1 mg/kg.^{66,68,87} Although a large body of literature can be found for the parent molecule, considerably less work has been published about the newer, less toxic derivatives, in particular, 2-methoxyantimycin.^{66,69,87,88} 2-Methoxyantimycin has been shown to function cooperatively with paclitaxel and docetaxel to induce apoptosis, whereas the parental compound, Antimycin A, displayed synergy with the topoisomerase inhibitor SN-38 in inhibiting cell growth in a human colon cancer line.^{70,87} If the less toxic antimycin analogs can retain activity displayed in the few available studies, they may prove to be a particularly interesting drug series.

From Desktop and Benchtop to Bedside

Inspired by the potential of natural Bcl-2 inhibitors, several research groups have developed specific inhibitors of Bcl-2. Primarily, various forms of *in silico* and *in vitro* screening of combinatorial libraries, for best-fit compounds, have resulted in Bcl-2 inhibitors of varying capacity and high affinity for one or more of the Bcl-2 family of proteins. HA14-1 (Fig 3D, Table 1) was the first Bcl-2-binding ligand to be discovered using these computer-based screening strategies using the predicted structure of Bcl-2.⁵⁴

Both synergistic and additive effects of HA14-1, with other chemotherapeutic compounds, contributed to reversal of drug resistance *in vitro* in a broad variety of cell lines and treatment regimens.^{56,58,89-92}

Notably, the order of treatment in a combination regimen made a profound difference in the final effect. HA14-1 acts as a radiosensitizer on LNCaP and PC3 prostate cancer cell lines with or without overexpression of Bcl-2. Synergy was only seen in the PC3 cells when HA14-1 was administered after exposure to gamma radiation, and not in the reverse order.⁹¹ In glioblastoma cells, a 1-hour pretreatment with HA14-1 sensitized the cells to radiation in a Bcl-2, but not Bcl-x_L, dependent manner.⁵⁹ Skommer et al⁹⁰ found synergy between HA14-1 and dexamethasone or doxorubicin, but not vincristine, in inhibiting B-cell lymphoma tumor cell proliferation. They observed that for HA14-1 and doxorubicin, synergistic inhibition was seen only when the Bcl-2 inhibitor was administered 24 hours before the conventional chemotherapeutic drugs, and antagonism was seen if they were administered simultaneously. In combination with the cyclin-dependent kinase (CDK) inhibitor flavopiridol, HA14-1 displays a similar synergy, but in this instance, only when the Bcl-2 inhibitor is administered after the CDK inhibitor.⁹³ It is noteworthy that the precise order of administration of HA14-1 in radiation, doxorubicin, or flavopiridol studies was important for enhancement of the inhibitory effect of HA14-1. Although not reported for all the Bcl-2 small-molecule inhibitors, most classes of drugs display administration order preferences for optimal cooperation with other therapeutic agents. The varying results, described above, for HA14-1 in combination therapy with radiation or chemotherapeutics may be due to tumor-specific differences in target expression (Bcl-2/Bcl-x_L/Mcl-1) or modulation of those proteins by whichever therapy is administered first.

Other small-molecule inhibitors of Bcl-2 began to appear after HA14-1, in particular TW-37 (TW37) and ABT-737. Both TW-37 and ABT-737 improve on HA14-1 in efficacy, having nanomolar affinities for Bcl-2. However, only TW-37 has nanomolar to low micromolar affinities for all of the major antiapoptotic members of the Bcl-2 family of proteins. TW-37 (Fig 3B, Table 1) represents a new class of pan-Bcl-2 protein family inhibitor. It is a BH3 mimetic developed by rational structural design using a computer-based modeling strategy.^{60,61} Apoptosis was shown to be the primary mechanism for the growth-inhibitory properties of TW-37 in the prostate cancer cell line PC3.⁶⁰ TW-37 has also been shown to effectively inhibit growth of Kaposi's sarcoma (SLK), breast cancer (MCF-7), and prostate cancer (LnCap) cell lines without inhibitory effects on human dermal fibroblasts.⁵⁵

Melanoma is particularly resistant to conventional chemotherapies.⁹⁴ TW-37 was found to be effective against melanoma-derived tumors when jointly administered with the mitogen-activated protein kinase kinase (MEK) inhibitors U0126 or CL-1040 and, to a lesser extent, as single agent both *in vivo* and *in vitro*.⁶² The rationale for coadministration was that some melanomas respond poorly to mitogen-activated protein kinase inhibition, possibly through ERK-mediated upregulation of the antiapoptotic Bcl-2 family of proteins. Using lentiviral-driven inhibitory RNA, the Soengas group produced a series of isogenic melanoma cell lines deficient in specific molecules of the apoptosis pathway. They identified Bcl-2, Mcl-1, and Bcl-x_L as integral components of resistance to MEK inhibition.⁶² They also showed synergistic inhibition of melanoma cell growth by TW-37 and the MEK inhibitors *in vitro*. Notably, *in vivo* coadministration significantly inhibited xenografted melanoma tumor growth when compared with either agent alone.⁶²

In an initial study by Oltersdorf et al,³⁰ ABT-737 (Fig 3E, Table 1) was shown to bind Bcl-2, Bcl-x_L, and Bcl-w with high affinity (K_i ≤ 1 nmol/L), but to have extremely low affinity for Mcl-1, Bcl-b, and A1. Indeed, the low affinity for Mcl-1 has defined part of the research into this compound, as the drug efficacy is greatly attenuated in the presence of that protein. ABT-737 was effective in delaying growth of tumors overexpressing Bcl-2. In contrast, tumors transduced with vectors inducing Mcl-1 overexpression were highly refractory to the drug.⁹⁵ Additionally, ABT-737 induced thrombocytopenia in mice in a transient and reversible manner but was nontoxic to hematopoietic progenitor cells, including megakaryocytes.^{65,96,97} This was shown to occur via direct inhibition of platelet Bcl-x_L by ABT-737, preventing binding of Bcl-x_L to Bak and inducing Bak-mediated platelet apoptosis.⁹⁷ Recently, Chen et al⁹⁸ have shown that selective downregulation of Mcl-1, either by CDK inhibitors such as roscovitine or by gene silencing, synergistically increases the sensitivity of human leukemia cells to ABT-737–induced apoptosis.

Given these issues, it is notable that ABT-737 is a highly active drug in several models. It has shown antitumor efficacy, as a single agent administered intraperitoneally, in two models of lung cancer using cell lines with known sensitivity to the compound and also in models of leukemia.^{30,65} In chronic myelogenous leukemia, the Bcr/Abl pathway promotes leukemogenesis, but the Bcr/Abl inhibitor (imatinib; Gleevec, Novartis Pharmaceuticals Corp, East Hanover, NJ) induces apoptosis in affected cells via a Bcl-2/Bcl-x_L–mediated pathway.⁹⁹ Coadministration of ABT-737 and imatinib to Bim/Bad double knockout Bcr/Abl-positive cells enhanced apoptosis as compared with either drug alone.¹⁰⁰ Thus ABT-737 seems to compensate for the loss of the BH3-only proteins Bim or Bad. It remains to be seen whether ABT-737 moves on to clinical trial, although a structurally related, orally available analog, ABT-263, is indeed entering phase I/II clinical trials for lymphocytic leukemia and SCLC¹⁰¹ (www.cancer.gov).

Another small-molecule inhibitor of all the major Bcl-2 antiapoptotic family proteins, GX15-070 (Obatoclox; Fig 3A; Table 1), was developed by Gemin X Biotechnologies Inc. (Montreal, Quebec, Canada). Although it was the first of the specifically designed Bcl-2 small-molecule inhibitor drugs to be approved in the United States for four clinical trials (www.cancer.gov) and has been under patent since 2002, little has yet been published on this compound. In one study by Campas et al,⁶³ GX15-070 displayed a mean half maximal effective concentration of 1.7 ± 0.1 μmol/L in isolated cell lines, ranging over a 48-hour period. However, this calculation of activity excluded two lines of 11, which were considered resistant to GX15-070. Therefore, reliable determination of the activity of this drug within this class of tumors must await other confirmatory studies at this time. Additive effects were reported in combination with fludarabine and chlorambucil administered with 0.5 μmol/L of GX15-070. However, synergy was observed for GX15-070 administered with the proteasome inhibitor, bortezomib, and the order of administration was not reported to alter synergistic efficacy.^{64,102} In three mantle-cell lymphoma lines studied, apoptotic concentrations of GX15-070 were able to inhibit binding of immunoprecipitated Mcl-1 and Bcl-x_L to Bak, resulting in mitochondrial depolarization and caspase-3 activation. The synergistic effects of GX15-070 with bortezomib on cytotoxicity in tumor cells were attributed in large part to inhibition of the Mcl-1/Bak interaction. Interestingly, they were also attributed to the in-

hibitory effects of GX15-070 on Mcl-1 expression, which may have countered the strong upregulation of Mcl-1 expression induced by bortezomib treatment.^{64,102}

BCL-2 INHIBITORS AS ANTIANGIOGENIC AGENTS

As discussed previously, Bcl-2 is intimately involved with angiogenesis through modulation of both endothelial and tumor cell signaling. Additionally, Bcl-2 inhibitors may directly induce apoptosis in angiogenically active endothelial cells in vitro, in similar fashion to their effect on tumor cells, and endothelial cell-specific apoptosis can disrupt neovascularization in vivo.^{55,103} Therefore, Bcl-2 and its related signaling network becomes a logical target for antiangiogenic cancer therapy (Fig 1). There is an emerging body of literature on the in vivo antiangiogenic effect of inhibition of the Bcl-2 family of proteins. Notably, most of what is available has been presented as unquantified immunohistochemical data. However, direct effects of Bcl-2 inhibition have been quantified in at least one study, which reported a significant reduction in microvessel density after TW-37 treatment in a humanized in vivo model of angiogenesis.⁵⁵ Of other studies, the antiangiogenic effects of EGCG have been demonstrated, including a significant reduction in microvessel density within xenografted ovarian cancer cell tumors and colon cancer tumors.^{48,104} However, as discussed previously, it may be pre-emptive to use EGCG as an example of an inhibitor inducing apoptosis solely through effects on Bcl-2. (–)-Gossypol has been shown to have potential antiangiogenic efficacy in vivo when administered in conjunction with radiotherapy, although without quantification of the reported changes in vessel density.⁵³ In total, these reports suggest that Bcl-2–dependent vasculature may indeed be targeted, in vivo, by small-molecule inhibitors of Bcl-2. Broadening the scope of literature search, two groups have published studies using antisense oligonucleotides to inhibit Bcl-2 expression in melanoma or prostate cancer cells transfected with Bcl-2.^{105,106} Anai et al¹⁰⁶ observed a reduction in microvessel density when the same cells were implanted as xenografts. Together, the work published to date suggests that Bcl-2 inhibition, whether by pharmacologic or genetic means, may potentially inhibit angiogenesis toward a therapeutic end.

SUMMARY AND IMPLICATIONS

There is an expanding industry devoted toward inhibiting various components of the Bcl-2 pathway using designer small-molecule drugs (mw approximately 400 to 800). These newly developed or identified compounds not only provide us with novel therapeutic approaches to cancer treatment but also deliver new tools for the analysis of the many functions of the Bcl-2 family of proteins. With our knowledge of the Bcl-2 family expanding rapidly, this specific avenue of pharmaceutical development is unlikely to slow down in the near future. It is strengthened by the proven therapeutic efficacy of both older and more recent pharmacologic agents in vivo, particularly (–)-gossypol and ABT-737, and the utility of the same compounds for dissection of apoptotic and angiogenic pathways.

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

The author(s) indicated no potential conflicts of interest.

AUTHOR CONTRIBUTIONS

Conception and design: Benjamin D. Zeitlin, Jacques E. Nör

Manuscript writing: Benjamin D. Zeitlin, Isaac J. Zeitlin, Jacques E. Nör

Final approval of manuscript: Benjamin D. Zeitlin, Isaac J. Zeitlin, Jacques E. Nör

REFERENCES

- Wang D, Lippard SJ: Cellular processing of platinum anticancer drugs. *Nat Rev Drug Discov* 4:307-320, 2005
- Rabik CA, Dolan ME: Molecular mechanisms of resistance and toxicity associated with platinating agents. *Cancer Treat Rev* 33:9-23, 2007
- Lee KH, Im SA, Oh DY, et al: Prognostic significance of bcl-2 expression in stage III breast cancer patients who had received doxorubicin and cyclophosphamide followed by paclitaxel as adjuvant chemotherapy. *BMC Cancer* 7:63, 2007
- Khor LY, Moughan J, Al-Saleem T, et al: Bcl-2 and bax expression predict prostate cancer outcome in men treated with androgen deprivation and radiotherapy on radiation therapy oncology group protocol 92-02. *Clin Cancer Res* 13:3585-3590, 2007
- Contu PC, Contu SS, Moreira LF: Bcl-2 expression in rectal cancer. *Arq Gastroenterol* 43:284-287, 2006
- Skommer J, Wlodkowic D, Deptala A: Larger than life: Mitochondria and the bcl-2 family. *Leuk Res* 31:277-286, 2007
- van Delft MF, Huang DC: How the bcl-2 family of proteins interact to regulate apoptosis. *Cell Res* 16:203-213, 2006
- Oh KJ, Barbuto S, Pitter K, et al: A membrane-targeted BID BCL-2 homology 3 peptide is sufficient for high potency activation of BAX in vitro. *J Biol Chem* 281:36999-37008, 2006
- Bikfalvi A: Angiogenesis: Health and disease. *Ann Oncol* 17:x65-70, 2006 (suppl 10)
- Carmeliet P, Jain RK: Angiogenesis in cancer and other diseases. *Nature* 407:249-257, 2000
- Mehrad B, Keane MP, Strieter RM: Chemokines as mediators of angiogenesis. *Thromb Haemostasis* 97:755-762, 2007
- Singh S, Sadanandam A, Singh RK: Chemokines in tumor angiogenesis and metastasis. *Cancer Metastasis Rev* 26:453-467, 2007
- Gerber HP, Dixit V, Ferrara N: Vascular endothelial growth factor induces expression of the antiapoptotic proteins bcl-2 and A1 in vascular endothelial cells. *J Biol Chem* 273:13313-13316, 1998
- Karsan A, Yee E, Kaushansky K, et al: Cloning of human bcl-2 homologue: Inflammatory cytokines induce human A1 in cultured endothelial cells. *Blood* 87:3089-3096, 1996
- Pidgeon GP, Barr MP, Harmey JH, et al: Vascular endothelial growth factor (VEGF) upregulates BCL-2 and inhibits apoptosis in human and murine mammary adenocarcinoma cells. *Br J Cancer* 85:273-278, 2001
- Nör JE, Christensen J, Mooney DJ, et al: Vascular endothelial growth factor (VEGF)-mediated angiogenesis is associated with enhanced endothelial cell survival and induction of bcl-2 expression. *Am J Pathol* 154:375-384, 1999
- Cai J, Ahmad S, Jiang WG, et al: Activation of vascular endothelial growth factor receptor-1 sustains angiogenesis and bcl-2 expression via the phosphatidylinositol 3-kinase pathway in endothelial cells. *Diabetes* 52:2959-2968, 2003
- Biroccio A, Candiloro A, Mottolese M, et al: Bcl-2 overexpression and hypoxia synergistically act to modulate vascular endothelial growth factor expression and in vivo angiogenesis in a breast carcinoma line. *FASEB J* 14:652-660, 2000
- Iervolino A, Trisciuglio D, Ribatti D, et al: Bcl-2 overexpression in human melanoma cells increases angiogenesis through VEGF mRNA stabilization and HIF-1-mediated transcriptional activity. *FASEB J* 16:1453-1455, 2002
- Kaneko T, Zhang Z, Mantellini M, et al: Bcl-2 orchestrates a cross-talk between endothelial and tumor cells that promotes tumor growth. *Cancer Res* 67:9685-9693, 2007
- Karl E, Warner K, Zeitlin B, et al: Bcl-2 acts in a proangiogenic signaling pathway through nuclear factor-kappaB and CXCL chemokines. *Cancer Res* 65:5063-5069, 2005
- Duval H, Johnson N, Li J, et al: Vascular development is disrupted by endothelial cell-specific expression of the anti-apoptotic protein bcl-2. *Angiogenesis* 10:55-68, 2007
- Hadar T, Shvero J, Yaniv E, et al: Expression of p53, ki-67 and bcl-2 in parathyroid adenoma and residual normal tissue. *Pathol Oncol Res* 11:45-49, 2005
- Daidone MG, Veneroni S, Benini E, et al: Biological markers as indicators of response to primary and adjuvant chemotherapy in breast cancer. *Int J Cancer* 84:580-586, 1999
- Martin B, Paesmans M, Berghmans T, et al: Role of bcl-2 as a prognostic factor for survival in lung cancer: A systematic review of the literature with meta-analysis. *Br J Cancer* 89:55-64, 2003
- Karczmarek-Borowska B, Filip A, Wojciorowski J, et al: Estimation of prognostic value of bcl-xL gene expression in non-small cell lung cancer. *Lung Cancer* 51:61-69, 2006
- Brooks KR, To K, Joshi MB, et al: Measurement of chemoresistance markers in patients with stage III non-small cell lung cancer: A novel approach for patient selection. *Ann Thorac Surg* 76:187-193, 2003; discussion 193
- Choi J, Choi K, Benveniste EN, et al: Bcl-2 promotes invasion and lung metastasis by inducing matrix metalloproteinase-2. *Cancer Res* 65:5554-5560, 2005
- Bauer JA, Trask DK, Kumar B, et al: Reversal of cisplatin resistance with a BH3 mimetic, (-)-gossypol, in head and neck cancer cells: Role of wild-type p53 and bcl-xL. *Mol Cancer Ther* 4:1096-1104, 2005
- Oltersdorf T, Elmore SW, Shoemaker AR, et al: An inhibitor of bcl-2 family proteins induces regression of solid tumours. *Nature* 435:677-681, 2005
- Wang L, Chen L, Benincosa J, et al: VEGF-induced phosphorylation of bcl-2 influences B lineage leukemic cell response to apoptotic stimuli. *Leukemia* 19:344-353, 2005
- Wang L, Coad JE, Fortney JM, et al: VEGF-induced survival of chronic lymphocytic leukemia is independent of bcl-2 phosphorylation. *Leukemia* 19:1486-1487, 2005
- Michels J, Johnson PW, Packham G: Mcl-1. *Int J Biochem Cell Biol* 37:267-271, 2005
- Nehls O, Okech T, Hsieh CJ, et al: Studies on p53, BAX and bcl-2 protein expression and microsatellite instability in stage III (UICC) colon cancer treated by adjuvant chemotherapy: Major prognostic impact of proapoptotic BAX. *Br J Cancer* 96:1409-1418, 2007
- Puthalakath H, Strasser A: Keeping killers on a tight leash: Transcriptional and post-translational control of the pro-apoptotic activity of BH3-only proteins. *Cell Death Differ* 9:505-512, 2002
- Deverman BE, Cook BL, Manson SR, et al: Bcl-xL deamidation is a critical switch in the regulation of the response to DNA damage. *Cell* 111:51-62, 2002
- Stein CA, Benimetskaya L, Mani S: Antisense strategies for oncogene inactivation. *Semin Oncol* 32:563-572, 2005
- Walensky LD, Kung AL, Escher I, et al: Activation of apoptosis in vivo by a hydrocarbon-stapled BH3 helix. *Science* 305:1466-1470, 2004
- Sang S, Hou Z, Lambert JD, et al: Redox properties of tea polyphenols and related biological activities. *Antioxid Redox Signal* 7:1704-1714, 2005
- Hsu S: Green tea and the skin. *J Am Acad Dermatol* 52:1049-1059, 2005
- Vita JA: Polyphenols and cardiovascular disease: Effects on endothelial and platelet function. *Am J Clin Nutr* 81:292S-297S, 2005
- Weinreb O, Mandel S, Amit T, et al: Neurological mechanisms of green tea polyphenols in Alzheimer's and Parkinson's diseases. *J Nutr Biochem* 15:506-516, 2004
- Zhai D, Jin C, Satterthwait AC, et al: Comparison of chemical inhibitors of antiapoptotic bcl-2-family proteins. *Cell Death Differ* 13:1419-1421, 2006
- Leone M, Zhai D, Sareth S, et al: Cancer prevention by tea polyphenols is linked to their

- direct inhibition of antiapoptotic bcl-2-family proteins. *Cancer Res* 63:8118-8121, 2003
45. Yang GY, Liao J, Kim K, et al: Inhibition of growth and induction of apoptosis in human cancer cell lines by tea polyphenols. *Carcinogenesis* 19:611-616, 1998
 46. Pan MH, Liang YC, Lin-Shiau SY, et al: Induction of apoptosis by the oolong tea polyphenol theasinensin A through cytochrome c release and activation of caspase-9 and caspase-3 in human U937 cells. *J Agric Food Chem* 48:6337-6346, 2000
 47. Navarro-Perán E, Cabezas-Herrera J, Garcia-Canovas F, et al: The antifolate activity of tea catechins. *Cancer Res* 65:2059-2064, 2005
 48. Jung YD, Kim MS, Shin BA, et al: EGCG, a major component of green tea, inhibits tumour growth by inhibiting VEGF induction in human colon carcinoma cells. *Br J Cancer* 84:844-850, 2001
 49. Kitada S, Leone M, Sareth S, et al: Discovery, characterization, and structure-activity relationships studies of proapoptotic polyphenols targeting B-cell lymphocyte/leukemia-2 proteins. *J Med Chem* 46:4259-4264, 2003
 50. Zhang M, Liu H, Guo R, et al: Molecular mechanism of gossypol-induced cell growth inhibition and cell death of HT-29 human colon carcinoma cells. *Biochem Pharmacol* 66:93-103, 2003
 51. Mohammad RM, Wang S, Aboukameel A, et al: Preclinical studies of a nonpeptidic small-molecule inhibitor of bcl-2 and bcl-X(L) [(–)-gossypol] against diffuse large cell lymphoma. *Mol Cancer Ther* 4:13-21, 2005
 52. Wolter KG, Wang SJ, Henson BS, et al: (–)-Gossypol inhibits growth and promotes apoptosis of human head and neck squamous cell carcinoma in vivo. *Neoplasia* 8:163-172, 2006
 53. Xu L, Yang D, Wang S, et al: (–)-Gossypol enhances response to radiation therapy and results in tumor regression of human prostate cancer. *Mol Cancer Ther* 4:197-205, 2005
 54. Wang JL, Liu D, Zhang ZJ, et al: Structure-based discovery of an organic compound that binds bcl-2 protein and induces apoptosis of tumor cells. *Proc Natl Acad Sci U S A* 97:7124-7129, 2000
 55. Zeitlin BD, Joo E, Dong Z, et al: Antiangiogenic effect of TW37, a small-molecule inhibitor of bcl-2. *Cancer Res* 66:8698-8706, 2006
 56. Lickliter JD, Wood NJ, Johnson L, et al: HA14-1 selectively induces apoptosis in bcl-2-overexpressing leukemia/lymphoma cells, and enhances cytarabine-induced cell death. *Leukemia* 17:2074-2080, 2003
 57. Van Poznak C, Seidman AD, Reidenberg MM, et al: Oral gossypol in the treatment of patients with refractory metastatic breast cancer: A phase I/II clinical trial. *Breast Cancer Res Treat* 66:239-248, 2001
 58. Chen J, Freeman A, Liu J, et al: The apoptotic effect of HA14-1, a bcl-2-interacting small molecule compound, requires bax translocation and is enhanced by PK11195. *Mol Cancer Ther* 1:961-967, 2002
 59. Manero F, Gautier F, Gallenne T, et al: The small organic compound HA14-1 prevents bcl-2 interaction with bax to sensitize malignant glioma cells to induction of cell death. *Cancer Res* 66:2757-2764, 2006
 60. Wang G, Nikolovska-Coleska Z, Yang CY, et al: Structure-based design of potent small-molecule inhibitors of anti-apoptotic bcl-2 proteins. *J Med Chem* 49:6139-6142, 2006
 61. Mohammad RM, Goustin AS, Aboukameel A, et al: Preclinical studies of TW-37, a new nonpeptidic small-molecule inhibitor of bcl-2, in diffuse large cell lymphoma xenograft model reveal drug action on both bcl-2 and mcl-1. *Clin Cancer Res* 13:2226-2235, 2007
 62. Verhaegen M, Bauer JA, Martin de la Vega C, et al: A novel BH3 mimetic reveals a mitogen-activated protein kinase-dependent mechanism of melanoma cell death controlled by p53 and reactive oxygen species. *Cancer Res* 66:11348-11359, 2006
 63. Campàs C, Cosialls AM, Barragan M, et al: Bcl-2 inhibitors induce apoptosis in chronic lymphocytic leukemia cells. *Exp Hematol* 34:1663-1669, 2006
 64. Nguyen M, Marcellus RC, Roulston A, et al: Small molecule obatocax (GX15-070) antagonizes MCL-1 and overcomes MCL-1-mediated resistance to apoptosis. *Proc Natl Acad Sci U S A* 104:19512-19517, 2007
 65. Konopleva M, Contractor R, Tsao T, et al: Mechanisms of apoptosis sensitivity and resistance to the BH3 mimetic ABT-737 in acute myeloid leukemia. *Cancer Cell* 10:375-388, 2006
 66. Tzung SP, Kim KM, Basanez G, et al: Antimycin A mimics a cell-death-inducing bcl-2 homology domain 3. *Nat Cell Biol* 3:183-191, 2001
 67. Manion MK, O'Neill JW, Giedt CD, et al: Bcl-XL mutations suppress cellular sensitivity to antimycin A. *J Biol Chem* 279:2159-2165, 2004
 68. Kim KM, Giedt CD, Basanez G, et al: Biophysical characterization of recombinant human bcl-2 and its interactions with an inhibitory ligand, antimycin A. *Biochemistry* 40:4911-4922, 2001
 69. Cao X, Rodarte C, Zhang L, et al: Bcl2/bcl-xL inhibitor engenders apoptosis and increases chemosensitivity in mesothelioma. *Cancer Biol Ther* 6:246-252, 2007
 70. Guichard SM, Hua ML, Kang P, et al: Short hairpin RNAs targeting bcl-xL modulate senescence and apoptosis following SN-38 and irinotecan exposure in a colon cancer model. *Cancer Chemother Pharmacol* 60:651-660, 2007
 71. Sakata R, Ueno T, Nakamura T, et al: Green tea polyphenol epigallocatechin-3-gallate inhibits platelet-derived growth factor-induced proliferation of human hepatic stellate cell line LI90. *J Hepatol* 40:52-59, 2004
 72. Pianetti S, Guo S, Kavanagh KT, et al: Green tea polyphenol epigallocatechin-3 gallate inhibits her-2/neu signaling, proliferation, and transformed phenotype of breast cancer cells. *Cancer Res* 62:652-655, 2002
 73. Tachibana H, Koga K, Fujimura Y, et al: A receptor for green tea polyphenol EGCG. *Nat Struct Mol Biol* 11:380-381, 2004
 74. Ahmad N, Adhami VM, Gupta S, et al: Role of the retinoblastoma (pRb)-E2F/DP pathway in cancer chemopreventive effects of green tea polyphenol epigallocatechin-3-gallate. *Arch Biochem Biophys* 398:125-131, 2002
 75. Peng G, Dixon DA, Muga SJ, et al: Green tea polyphenol (–)-epigallocatechin-3-gallate inhibits cyclooxygenase-2 expression in colon carcinogenesis. *Mol Carcinog* 45:309-319, 2006
 76. Coutinho EM, Athayde C, Atta G, et al: Gossypol blood levels and inhibition of spermatogenesis in men taking gossypol as a contraceptive: A multicenter, international, dose-finding study. *Contraception* 61:61-67, 2000
 77. Ford JM, Hait WN, Matlin SA, et al: Modulation of resistance to alkylating agents in cancer cell by gossypol enantiomers. *Cancer Lett* 56:85-94, 1991
 78. Yu Y, Deck JA, Hunsaker LA, et al: Selective active site inhibitors of human lactate dehydrogenases A4, B4, and C4. *Biochem Pharmacol* 62:81-89, 2001
 79. Baumgrass R, Weiwad M, Erdmann F, et al: Reversible inhibition of calcineurin by the polyphenolic aldehyde gossypol. *J Biol Chem* 276:47914-47921, 2001
 80. Oliver CL, Bauer JA, Wolter KG, et al: In vitro effects of the BH3 mimetic, (–)-gossypol, on head and neck squamous cell carcinoma cells. *Clin Cancer Res* 10:7757-7763, 2004
 81. Oliver CL, Miranda MB, Shangary S, et al: (–)-Gossypol acts directly on the mitochondria to overcome bcl-2- and bcl-X(L)-mediated apoptosis resistance. *Mol Cancer Ther* 4:23-31, 2005
 82. Wang X, Wang J, Wong SC, et al: Cytotoxic effect of gossypol on colon carcinoma cells. *Life Sci* 67:2663-2671, 2000
 83. Huang YW, Wang LS, Chang HL, et al: Molecular mechanisms of (–)-gossypol-induced apoptosis in human prostate cancer cells. *Anticancer Res* 26:1925-1933, 2006
 84. Bushnow P, Reidenberg MM, Wasenko J, et al: Gossypol treatment of recurrent adult malignant gliomas. *J Neurooncol* 43:79-86, 1999
 85. Flack MR, Pyle RG, Mullen NM, et al: Oral gossypol in the treatment of metastatic adrenal cancer. *J Clin Endocrinol Metab* 76:1019-1024, 1993
 86. Stein RC, Joseph AE, Matlin SA, et al: A preliminary clinical study of gossypol in advanced human cancer. *Cancer Chemother Pharmacol* 30:480-482, 1992
 87. Wang H, Li M, Rhie JK, et al: Preclinical pharmacology of 2-methoxyantimycin A compounds as novel antitumor agents. *Cancer Chemother Pharmacol* 56:291-298, 2005
 88. Schwartz PS, Manion MK, Emerson CB, et al: 2-methoxy antimycin reveals a unique mechanism for bcl-x(L) inhibition. *Mol Cancer Ther* 6:2073-2080, 2007
 89. Sinicrope FA, Penington RC, Tang XM: Tumor necrosis factor-related apoptosis-inducing ligand-induced apoptosis is inhibited by bcl-2 but restored by the small molecule bcl-2 inhibitor, HA 14-1, in human colon cancer cells. *Clin Cancer Res* 10:8284-8292, 2004
 90. Skommer J, Wlodkovic D, Matto M, et al: HA14-1, a small molecule bcl-2 antagonist, induces apoptosis and modulates action of selected anticancer drugs in follicular lymphoma B cells. *Leuk Res* 30:322-331, 2006
 91. An J, Chervin AS, Nie A, et al: Overcoming the radioresistance of prostate cancer cells with a novel bcl-2 inhibitor. *Oncogene* 26:652-661, 2007
 92. Dai Y, Rahmani M, Corey SJ, et al: A Bcr/Abl-independent, lyn-dependent form of imatinib mesylate (STI-571) resistance is associated with altered expression of bcl-2. *J Biol Chem* 279:34227-34239, 2004
 93. Pei XY, Dai Y, Grant S: The small-molecule bcl-2 inhibitor HA14-1 interacts synergistically with flavopiridol to induce mitochondrial injury and apoptosis in human myeloma cells through a free radical-dependent and jun NH2-terminal kinase-dependent mechanism. *Mol Cancer Ther* 3:1513-1524, 2004
 94. Hersey P: Apoptosis and melanoma: How new insights are effecting the development of new therapies for melanoma. *Curr Opin Oncol* 18:189-196, 2006
 95. van Delft MF, Wei AH, Mason KD, et al: The BH3 mimetic ABT-737 targets selective bcl-2 proteins and efficiently induces apoptosis via Bak/Bax if mcl-1 is neutralized. *Cancer Cell* 10:389-399, 2006

96. Trudel S, Stewart AK, Li Z, et al: The bcl-2 family protein inhibitor, ABT-737, has substantial antimyeloma activity and shows synergistic effect with dexamethasone and melphalan. *Clin Cancer Res* 13:621-629, 2007
97. Mason KD, Carpinelli MR, Fletcher JI, et al: Programmed anuclear cell death delimits platelet life span. *Cell* 128:1173-1186, 2007
98. Chen S, Dai Y, Harada H, et al: Mcl-1 down-regulation potentiates ABT-737 lethality by cooperatively inducing bak activation and bax translocation. *Cancer Res* 67:782-791, 2007
99. Ferrao PT, Frost MJ, Siah SP, et al: Overexpression of P-glycoprotein in K562 cells does not confer resistance to the growth inhibitory effects of imatinib (STI571) in vitro. *Blood* 102:4499-4503, 2003
100. Kuroda J, Puthalakath H, Cragg MS, et al: Bim and bad mediate imatinib-induced killing of Bcr/Abl(+) leukemic cells, and resistance due to their loss is overcome by a BH3 mimetic. *Proc Natl Acad Sci U S A* 103:14907-14912, 2006
101. Lock R, Carol H, Houghton PJ, et al: Initial testing (stage 1) of the BH3 mimetic ABT-263 by the pediatric preclinical testing program. *Pediatr Blood Cancer* 50:1181-1189, 2008
102. Pérez-Galán P, Roué G, Villamor N, et al: The BH3-mimetic GX15-070 synergizes with bortezomib in mantle cell lymphoma by enhancing noxa-mediated activation of bak. *Blood* 109:4441-4449, 2007
103. Dong Z, Zeitlin BD, Song W, et al: Level of endothelial cell apoptosis required for a significant decrease in microvessel density. *Exp Cell Res* 313:3645-3657, 2007
104. Spinella F, Rosano L, Di Castro V, et al: Green tea polyphenol epigallocatechin-3-gallate inhibits the endothelin axis and downstream signaling pathways in ovarian carcinoma. *Mol Cancer Ther* 5:1483-1492, 2006
105. Del Bufalo D, Trisciuglio D, Scarsella M, et al: Treatment of melanoma cells with a bcl-2/bcl-xL antisense oligonucleotide induces antiangiogenic activity. *Oncogene* 22:8441-8447, 2003
106. Anai S, Goodison S, Shiverick K, et al: Knockdown of bcl-2 by antisense oligodeoxynucleotides induces radiosensitization and inhibition of angiogenesis in human PC-3 prostate tumor xenografts. *Mol Cancer Ther* 6:101-111, 2007

Acknowledgment

We thank Shaomeng Wang for his continuous support of our studies and Chris Jung for the illustration of the model.

Glossary Terms

Bcl-2: First discovered as a translocated locus [t(14;22)] in B-cell leukemias, bcl-2 is an antiapoptotic protein that protects cells from programmed cell death by preventing the activation of proapoptotic caspase proteins.

Bcl-x_L: An antiapoptotic member of the Bcl-2 family of proteins, Bcl-x is a homolog of Bcl-2. Depending on splicing, Bcl-x may be present either as the short (Bcl-x_S) or the long (Bcl-x_L) form. Of these, Bcl-x_L has antiapoptotic properties similar to Bcl-2.

VEGF (vascular endothelial growth factor): VEGF is a cytokine that mediates numerous functions of endothelial cells, including proliferation, migration, invasion, survival, and permeability. VEGF is also known as vascular permeability factor. VEGF naturally occurs as a glycoprotein and is critical for angiogenesis. Many tumors overexpress VEGF, which correlates to poor prognosis. VEGF-A, -B, -C, -D, and -E are members of the larger family of VEGF-related proteins.

Angiogenesis: The process involved in the generation of new blood vessels. While this is a normal process that naturally occurs and is controlled by “on” and “off” switches, blocking tumor angiogenesis (antiangiogenesis) disrupts the blood supply to tumors, thereby preventing tumor growth.

BH3-binding domain: Hydrophobic groove present in Bcl-2, and some related proteins, that acts as a binding target for the BH3 domain of other proteins. The BH3-binding domain is the primary target for certain small molecule inhibitor drugs that prevent BH3-domain binding and induce apoptosis.

TW-37 (TW37): Small molecule inhibitor of Bcl-2 and Bcl-2 family of anti-apoptotic proteins, induces apoptosis in sensitive cells. TW-37 is an example of a new group of molecules designed to inhibit Bcl-2 and related proteins by targeting their BH3-binding domain. TW-37 is currently in pre-clinical trials.

(–)-Gossypol: Negative enantiomer of a polyphenolic molecule derived from the cotton plant. Having male contraceptive properties, it is a natural compound known to inhibit Bcl-2 and Bcl-x_L function via targeting their BH3-binding domain. (–)-Gossypol is currently entering clinical trial for cancer treatment.

Oblimersen (Genasense): Antisense oligonucleotide inhibiting Bcl-2 mRNA translation and therefore protein production. This is an example of a genetic inhibitor of Bcl-2 currently in clinical trial.