



Genomic Cloning and Sequence Analysis of *Trypanosoma brucei rhodesiense* Gene Encoding Putative N-glycosylation Enzyme

Waren Navarra Baticados¹, Noboru Inoue², Chihiro Sugimoto³, Hideyuki Nagasawa² & Abigail Morales Baticados¹

ABSTRACT

Background: *Trypanosoma brucei rhodesiense* is a haemoflagellate parasite of zoonotic significance. Aside from its public health importance, this parasite subspecies gained notoriety because of their effective system to circumvent the immune response of vertebrate host. The parasite cell surface is covered with millions of VSG dimers, which serve as an almost infinite repertoire of biomolecules needed for evasion of host immune system. Around two decades ago, it was resolved that all trypanosome VSG is associated with one or more N-linked oligosaccharides, with a range of structures including high mannose and complex types. This complex process of protein modification known as N-linked glycosylation is catalyzed by oligosaccharyl transferase (OST). In general, the incorporation of glycan structures can alter protein's antigenic properties and recently it was established that glycan molecules associated with VSG were found to be important in several aspects of trypanosome-host interaction, especially during parasite evasion of the host defense mechanisms. Therefore, our major interest is to clone and characterize the trypanosome OST.

Material, Methods and Results: The template genomic DNA for PCR amplification was extracted as described previously. In an attempt to clone *Trypanosoma brucei rhodesiense* putative oligosaccharyl transferase, an amplicon of ~2000 bp was obtained having an open reading frame of 2057 bp and deduced primary structure composed of 685 amino acid residues (TbrOST II). Comparison of TbrOST II ORF with annotated putative oligosaccharyl transferase in the genome of other organisms revealed sequence identity to other kinetoplastid. TbrOST II had high nucleotide (Ns) and amino acid (As) sequence similarity with the genomes of *T. brucei gambiense* (Ns:99%; As:78%) and *T. brucei* (Ns:95-98%; As:77%-98%). There was also significant nucleotide and amino acid sequence identity in the genomes of *T. cruzi* (Ns:74%; As:63%), *Leishmania infantum* (Ns:70-83%; As:46-57%), *L. braziliensis* (Ns:69-81%; As:46-55%) and *L. major* (Ns:69-80%; As:46-57%). Sequence similarity (71-77%) from other origins was also exhibited. The nucleotide sequence alignments and analysis were performed using the Oxford University Mac Vector 6.5 sequence analysis software and CLC Workbench 5.6 software.

Discussion: The nucleotide BLAST results indicate that sequence identity is higher between species of the same genus rather than of the same family. It is known that *T. brucei*, *T. gambiense* and *T. rhodesiense* are members of the *Brucei*-complex or *Brucei* group. Although *T. brucei brucei* has more similarities with *T. brucei rhodesiense* than *T. brucei gambiense*, these parasites are morphologically indistinguishable. This is the probable reason why high sequence identity was displayed by other subspecies of the *Brucei* group. In addition, the high percent identity possessed by TbrOST II with other trypanosomatids agrees with the evolutionarily conserved characteristics of the established OST. The DNA sequence data of TbrOST II showing similar sequences in the genome of other organisms further corroborate the previous reports regarding the ubiquitous nature of OST in other life forms. Based on the size of the amplicon and significant percentage of nucleotide and amino acid sequence identity to homologues within the genome of related species and various organisms, the results strongly indicate that TbrOST II is a trypanosome oligosaccharyl transferase gene candidate that should be fully characterized and subjected to functional genomic studies. The study reports the molecular cloning and sequencing of a potential oligosaccharyl transferase gene in *T. brucei rhodesiense* (TbrOST II). The sequence data has been deposited in the GenBank with accession number GU475126.

Keywords: *Trypanosoma brucei rhodesiense*, putative oligosaccharyl transferase, N-glycosylation.

INTRODUCTION

Trypanosoma brucei rhodesiense is a haemoflagellate parasite of zoonotic significance. A wide range of mammalian fauna, especially domestic livestock and wild bovids, serves as reservoir host [23,26]. Aside from its public health importance, this parasite subspecies gained notoriety because of their effective system to circumvent the immune response of vertebrate host. The parasite cell surface is covered with millions of VSG dimers, which serve as an almost infinite repertoire of biomolecules needed for evasion of host immune system. Unique VSG are alternately produced by sequential expression of about a thousand trypanosome VSG gene reservoir per parasite; a phenomenon described as antigenic variation [5,6]. Around two decades ago, it was resolved that all trypanosome VSG is associated with one or more N-linked oligosaccharides, with a range of structures including high mannose and complex types [7,19,21, 24]. This complex process of protein modification is generally known as glycosylations, one of which is N-linked glycosylation.

The central event in the N-linked glycosylation process is catalyzed by oligosaccharyl transferase (OST) [11]. It catalyzes the co-translational addition of preassembled oligosaccharide complexes (Dol-PP-GlcNAc₂Man₉Glc₃) to an asparagine residue in an Asn-Xaa-Ser/Thr consensus sequon (Xaa can be any amino acid excluding proline) of the growing nascent polypeptide chain being translocated into the endoplasmic reticulum through a structure called translocon [4].

OST appears ubiquitous among eukaryotes and conserved throughout eukaryotic evolution. OST has been molecularly isolated and purified from mammalian sources, avian species and yeast *Saccharomyces cerevisiae* [3,11-14,22] as a multimeric enzyme.

In general, the incorporation of glycan structures to different protein moieties is precedent towards proper protein folding and stability, intracellular targeting, intercellular recognition, hormone synthesis, anti-apoptotic response, control of salt/osmotic stress, and cell surface expression of some glycoproteins [8,9,15,17]. Glycans can alter protein's antigenic properties and recently it was established that, glycan molecules associated with VSG were found to be important in several aspects of trypanosome-host interaction, especially during parasite evasion of the host defense mechanisms [20,21,16]. Therefore our major interest is to clone and characterize the trypanosome OST.

MATERIALS AND METHODS

Laboratory animals

Female 8-week-old BALB/c mice¹ were used in the study. The animal room was maintained at 22 ± 3°C with a 12:12 hours of light-dark cycle. All experiments were conducted according to the guidelines for the care and use of laboratory animals, Obihiro University of Agriculture and Veterinary Medicine, Japan.

Trypanosomes and cultivation

Trypanosoma brucei rhodesiense IL2343 strain (Ivory Coast) stabilates were maintained in BALB/c mice. After reaching high parasitemia (~100 BSFs/field at magnification of x400), infected blood was collected by intracardiac puncture, cultured and maintained *in vitro* in HMI-9 medium supplemented with 10% fetal bovine serum² as previously described [10]. The HMI-9 *in vitro* cultivation medium was prepared using the following composition: Iscove's modified dulbecco's medium³ (IMDM), 10 mM bathocuproine disulfonic acid³, 100 mM pyruvic acid sodium salt³, 16 mM thymidine³, 40 mg/mL bovine serum albumin³, 1 mg/mL bovine holotransferrin, 100 mM hypoxanthine⁴, 150 mM L-cysteine hydro-chloride³, 14 mM 2-mercaptoethanol⁵, 25 mM HE-PES³ pH 7.2, 200 mM L-glutamine⁶, 10%v/v heat inactivated fetal bovine serum², 100 U/mL penicillin⁷ and 100 µg/mL streptomycin⁷.

DNA extraction

The template genomic DNA for PCR amplification was extracted as described previously [2]. Briefly, *T. brucei rhodesiense* IL2343 genomic DNA was extracted by adding 9 volumes of extraction buffer (0.2 M NaCl⁸, 10 mM Tris-HCl^{3,8} pH 8.0, 10 mM EDTA³ pH 8.0 and 1% SDS), proteinase K⁶ to a final concentration of 100 µg/mL and followed by 6 hours incubation at 55°C with gentle agitation. Overnight incubation was performed after additional proteinase K was placed. Genomic DNAs were phenol-chloroform-isoamyl alcohol³ extracted, ethanol precipitated, and resuspended in Tris-EDTA buffer, pH 8.0 or deionized water. The concentration of the sample DNA was determined by spectrophotometry.

Polymerase chain reaction amplification of putative OST gene

The primers were designed from the nucleotide sequences of *T. brucei* genomic clones as guided by EMBL-EBI Parasite Genomes WU-Blast

2 database search (www.ebi.ac.uk/blast/parasites.html) for African trypanosomes with *L. major* putative OST STT3 subunit sequence as the query [EMBL Q9U5N8 (AJ251127.1)]. PCR amplification was performed using a forward primer⁹ (5'-TGG TAC GAC TAC ATG AGC TGG TAC CCG CT-3') and a reverse primer⁹ (5'-TGG ATC TCC TTC GCT GGC GGG TAC TG-3'). Distilled water was used as template for negative control reaction. The samples were programmed to a temperature-step cycle of 94°C at 10 min, 94°C at 30 sec, 60°C at 30 sec for a total of 30 cycles followed by 4 min extension at 72°C. The PCR products were analyzed by electrophoresis on 1% TAE (Tris-acetate-EDTA) agarose gel. The PCR product was then processed for cloning after agarose gel extraction using a commercial kit¹⁰ according to the manufacturer's instructions.

Cloning and sequencing of PCR products

The PCR product was ligated into *EcoR* V site of pT7 bluescript plasmid vector¹¹ using Takara solution I ligation kit¹². Ligation reaction was transformed into DH5 α competent *E.coli* cells and plated on Luria Britani's-ampicillin (LB-amp) agar dishes. The presence of insert was confirmed by restriction digest against *Hind* III & *Xba* I site from the cloning site of the plasmid vectors flanking the PCR product.

Prior to sequencing, twenty-five cycles of Bigdye PCR was carried out in a total volume of 5 μ l and were performed using the following standard condition: 96°C at 2 min, 96°C at 10 sec, 50°C at 5 sec and 60°C at 4 minutes. Sequencing was started by the single strand dideoxynucleotide-chain-termination method using a cycle sequencing kit¹³, DNA sequence analyzer¹³ and T7 promoter primer and pUC/M13 reverse primers⁹. The second set⁹ [forward (5'-GAC ATA CAG CGT CAG TTT GC-3'); reverse (5'-GAT GAA TGT GAG TGA AGA GAG C-3')] and the third set⁹ [forward (5'-CGT TCG GAT TCT TCA AAC CTA CAG-3') and reverse (5'-AAT ACG GGC ATC TTC AGG CG-3')] primers were used to obtain the partial nucleotide sequence. The nucleotide sequence alignments and analysis were performed using sequence analysis softwares^{14,15}.

RESULTS

A putative trypanosome oligosaccharyl transferase gene was successfully amplified using *T. brucei rhodesiense* crude DNA as the template.

Genomic clone of approximately 2000 bp was obtained after PCR amplification (Figure 1). Nucleotide sequencing revealed that the amplified band was composed of 2057 bp partial nucleotide sequence (Figure 2). The deduced partial primary structure of the *T. brucei rhodesiense* putative oligosaccharyl transferase clone II (TbrOST II) was composed of 685 amino acid sequence (Figure 2). Determination of the sequence homology in other organisms was carried out using NCBI Basic Local Alignment Search Tool (**BLAST**) (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) [1].

Sequence analysis showed that TbrOST II had significant nucleotide (Ns) and amino acid sequence (As) percent identity to putative oligosaccharyl transferase subunit coming from other kinetoplastid genome in the NCBI public database.

When compared with the related trypanosome species, the DNA size of *T. brucei rhodesiense* putative oligosaccharyl transferase was 200 bp than *T. brucei gambiense* (1842 bp, FN554968.1). In addition, the sequence homologues from *T. brucei* (2466 bp, XM_839672.1) (2466 bp, XM_839671.1) (2406 bp, AC159432.1)(2406 bp, XM_839670.1), exceeded TbrOST II clone by >400 bp. The Latin American trypanosome species, *T. cruzi*, with percent identity to TbrOST II on the other hand had a molecular weight of 2397 bp. (XM_803446.1). Furthermore, the data showed that the annotated DNA size of the homologues under genus *Leishmania* was greater than that of genus *Trypanosoma* with




Figure 1. Polymerase Chain Reaction amplification of *Trypanosoma brucei rhodesiense* Putative Oligosaccharyl transferase clone II (TbrOST II). Lane (A), (B) & (C) represents 1 kbp DNA ladder, negative control reaction and ~ 2,000 bp PCR amplicon respectively.

a disparity ranging from ~300-500 bp (*L. infantum*) and ~200- 500bp (*L. major* & *L. braziliensis*).

Leishmania infantum displayed the greatest difference in terms of DNA molecular weight to TbrOST II (2592 bp, XM_001468890.1) (2385 bp, XM_001468892.1) (2355 bp, XM_001468891.1) followed by *L. braziliensis* (2580 bp, XM_001565753.1) (2319 bp, XM_001568165.1) (2565 bp, XM_001568167.1) (2472 bp, XM_001568166.1) and the least was *L. major* (2574 bp, XM_838130.1) (2322 bp, XM_838128.1) (2559 bp, XM_838127.1) (2373 bp, XM_838129.1) (2502 bp, AJ251127.1).

The molecular size of the amplicon showed akin to several homologues under order kinetoplastida. Taking into consideration the reported DNA size of the homologues from closely related species, the data suggest that the full-length nucleotide sequence of *T. brucei rhodesiense* putative oligosaccharyl transferase is also not more than 3000 bp.

Furthermore, the results using BLAST nucleotide (Ns) and amino acid (As) homology search

showed that the *T. brucei rhodesiense* putative oligosaccharyl transferase had a very high percent sequence identity to the genome of *T. brucei* (Ns:98%, XM_839670.1; As:98%, XP_844763.1). Other homologues found in the genome of *T. brucei* (Ns:95%, XM_839672.1; As:78%, XP_844765.1) (Ns:96%, XM_839671.1; As:77%, XP_844764.1) and *T. brucei gambiense* (Ns:99%, FN554968.1; As:78%, CBH10989.1) also showed very high DNA sequence identity but with only fairly high amino acid sequence identity to TbrOST II. A moderate sequence similarity in the genome of *T. cruzi* (Ns:74%, XM_803446.1; As:63%, XP_808539.1) was also observed.

Subsequently, percent sequence identity hits in other members of family Trypanosomatidae, genus *Leishmania*, were also obtained (Figure 3). All the homologues from genus *Leishmania* displayed moderate and low nucleotide and amino acid sequence identity to TbrOST II respectively. The following are the different *Leishmania* clones from the GenBank with nucleotide and amino acid

A	<pre> 1 TGGTACGACT ACATGAGCTG GTACCCGCTT GGCCGACCGG TGGGCACAAC CATCTTCCCC 61 GGAATGCAGC TTACCGGTGT AGCCATTTCAT CGTGTGCTGG AAATGCTCGG GCGAGGTATG 121 TCCATCAACA ATATCTGTGT GTACATTCCT GCATGGTTCG GTAGTATTGC CACTGTGTTG 181 GCTGCTCTCA TTGCGTACGA ATCATCTAAT TCGCTCAGTG TCATGGCGTT TACTCGGTAC 241 TTTTTTCCA TCGTACCTGC ACACCTGATG CGATCAATGG CTGGTGAATT TGACAAATGAG 301 TGTGTTGCAA TGGCGGCGAT GCTTCTGACG TTCTACATGT GGGTACGATC GTTACGCGAG 361 TCAAGTTCGT GGCCCATTTG CGCTTTAGCT GGTGTGGCAT ACGGGTACAT GGTGTCCAGC 421 TGGGGTGGTT ATATTTTCGT GCTGAAACATG GTAGCCTTCC ACGCTTCTGT ATGTGTACTG 481 CTTGATTGGG CTCGTGGGAC ATACAGCGTC AGTTTGTCTGA GGGCGTATTC ACTGTTTTTT 541 GTCAATTGGCA CTGCCCTTGC GATTTGCGTA CCGCCAGTGG AGTGGACGCC CTTCCGGTCC 601 CTGGAGCAAC TGACAGCATT GTTTGTCTTC GTTTTCATGT GGGCACCTCA CTACTCGGAA 661 TACCTGCGTG AGCGTGCCCG AGCGCCCATT CACTCTTCTA AAGCACTTCA GATCCGTGCC 721 CGCATTTTCA TGGGCACACT CTCCTTGCTG TTGATTGTGG CAAGTTTGTCT TGCCCCGTTT 781 GGATTCCTCA AACCTACAGC GTACCGCGTC CGTGGCTTGT TCGTGAACA TACGGGTACC 841 GGAATCCCCC TCGTGGATTG TGTGGCCGAG CATCGGCCGA CGACTGCCGG GGCATATCTG 901 CGCTACTTTC ATGTTTTGTA CCCTCTTTGG GGGTGCGGTG GGCTCTCTAT GTTGGTATTC 961 ATGAAAAAGG ACCGCTGGCG CGCCATTGTT TTTCTTGCTT CACTTTCCAC TGTTACGATG 1021 TATTTAGCGG CCGTATGTCT GCGATTACTT CTGTAGCGGG GTCCCGCAGC AACCGCTTGC 1081 GCCGGCATGT TCATAGGGGG GCTTTTTGAT CTGGCGCTGT CACAGTTTGT TGTATTGCGT 1141 AGCCCAAAGG ATGCCTCCGG CGATTCCGAT CCCGCGGGAG GGTGGAAGCG GGCAAAAGGG 1201 AAGTTTGTTA ATGAGCCTCC CAAAAGAGCC ATCTTTAGTC ACCGCTGGTT TCAACGTTTA 1261 TGCAATCGT TGCCCGTCCC GCTACGACGT GGTATCGCGG TTGTGGTGTCT CGTATGTCTC 1321 TTCGCCAATC CCATGAGACA CTCATTGCAA AAGTCCCTGG AGAAAAATGGC ACATGCACTT 1381 TCATCTCCAA GGATCATTGC CGTGCCTGAT CTACCCAATG GAGAGAGAGT CCTCGCCGAT 1441 GATTACTACG TGTGCTACTT GTGGCTGCGA AACCAATACG CTGAAGATGC CCGTATTCTC 1501 TCATGGTGGG ACTACGGGTA TCAAACTACT GGAATTGGCA ATCGCAACAC CTTGCGGAT 1561 GGTAAACAT GAGTCAACA GCACATAGCA ACTATTGGAAG AGATGCTTAC ATCCCCTGTG 1621 AAGGAGTCA ATGCTCTTAT ACGCCATCTC GCTGATTATG TGCTGATATG GTCTGGTCAA 1681 GATGGCAGCG ATTTACTTAA ATCGCCACAC ATGGCTCGGA TAGGCCACAG TGTATATCGC 1741 GATATGTGTT CAGAAGACGA TCCGCTGTGT AGACAGTTTC GGTTTTATAG TGGTACCTC 1801 AATAAGCCTA CGCCTATGAT GCAGCGGTCC CTATTATACA ATCTGCACAG GTTTGGTACG 1861 GATGGCGGGA AGACACAAC GGATAAGAAC ATGTTTTCAG TCGCCTACGT GTCAAAGTAT 1921 GGTGTTGGTA AGATCTCAA GGTGATGAAT GTGAGTGAAG AGAGCAAGGC GTGGTTGCA 1981 GACCAAAGA ACCGCGTATG CGACCCGCC GATCTTGGAA TATGCGCCGG CCACTACCCG 2041 CCAGCGAAGG AGATCCA </pre>
B	<pre> 1 WYDYSWYPL GRPVGTTIFP GMQLTGVAIH RVLEMLGRGM SINNICVYIP AWFSGIATVL 61 AALLAYESSN SLVMAFTAY FF SIVPAHLM RSMAGEFDNE CVAMAAMLLT FYMVRSLRS 121 SSSWPIGALA GVAYGYMVST WGGYIEVLM VAFHASVCVL LDWARGTYSV SLLRAYSLRF 181 VIGTALAI CV PPVEWTFPRS LEQLTALFVF VFMWALHYSE YLRERARAPI HSSKALQIRA 241 RIEMGTL SLL LIVASLLAPF GFFKPTAYRV RALFVKHIRT GNPLVD SVAE HRPTTAGAYL 301 RYFHVCYPLW GCGGLSMLVF MKKDRWRAIV FLASLSTVIM YFSARMSRLL LLAGPAATAC 361 AGMF IGGLFD LALSQF GDLR SPKDA S GDS PAGGSKRAG KVVNEPPKRA IF SHRWVQRL 421 VQSLPVPLRR GLAVVVLVCL FANPMRHSFE KSCCKMAHAL SSPRIILAVTD LPNGERVLAD 481 DYVYSYLWLR NNTPEDARIL SWWDYGYQIT GIGNRTILAD GNTWSHKHIA TIGKMLTSPV 541 KESHALIRHL ADYVLIWSSG DGSDDLKSPH MARI GNSVYR DMCSEDDPLC RQFGFYSGDL 601 NKPTMMQRS LLYNLHREFT DGGKTLQDKN MFQLAYVSKY GLVKIYKVMN VSEESKAWVA 661 DPKNRVCDPP GSWICAGQYP PAKEI </pre>

Figure 2. Partial Nucleotide (A) and Deduced Amino Acid (A) Sequence of *Trypanosoma brucei rhodesiense* Putative Oligosaccharyl transferase (TbrOST II) (GenBank: GU475126).

sequenced regions having local similarity to TbrOST II: *L. infantum* (Ns:70%, XM_001468891.1; As:57%, XP_001468928.1) (Ns:70%, XM_001468890.1; As:56%, XP_001468927.1) (Ns:83%, XM_00146892.1; As:46%, XP_001468929.1); *L. braziliensis* (Ns:69%, XM_001565753.1; As:54%, XP_001565803.1) (Ns:69%, XM_001568165.1; 55%,As:XP_001568215.1) (Ns:70%, XM_001568167.1; As:55%, XP_001568217.1) (Ns:81%, XM_001568166.1; As:46%, XP_001568216.1); *L. major* (Ns:69%, XM_838130.1; As:55%, XP_843223.1) (Ns:70%, XM_838128.1;As:57%, XP_843221.1) (Ns:70%, XM_838127.1; As:55%, XP_843220.1) (Ns:80%, XM_838129.1; As:46%, XP_843222.1) (Ns:80%,

```

TbrOSTII -----
Tbrucei MTKGGKVAVTKGSAQSDGAGEGGMASKAKSSTTFVATGGGSLPAWALKAVSTIVSAVILIY
Tbgam MTKGGKVAVTKGSAQSDGAGEGGMASKAKSSTTFVATGGGSLPAWALKAVSTIVSAVILIY

TbrOSTII -----WYDYMSWYPLGRPVG
Tbrucei SVHRAYDIRLTSVRLYGELEIHEFDPWFNYRATQYLSDNGWRAFFQWYDYMSWYPLGRPVG
Tbgam SVHRAYDIRLTSVRLYGELEIHEFDPWFNYRATQYLSDNGWRAFFQWYDYMSWYPLGRPVG

TbrOSTII TTIFFPGMQLTGVAIHRVLEMLGRGMSINNIVYIPAWFGSIATVLAALIAYESSNSLSVM
Tbrucei TTIFFPGMQLTGVAIHRVLEMLGRGMSINNIVYIPAWFGSIATVLAALIAYESSNSLSVM
Tbgam TTIFFPGMQLTGVAIHRVLEMLGRGMSINNIVYIPAWFGSIATVLAALIAYESSNSLSVM

TbrOSTII AFTAYFFSIVPAHLMRSMAGEFDNECVAMAAMLLTFYMWVRSLSRSSSSWPICALAGVAYG
Tbrucei AFTAYFFSIVPAHLMRSMAGEFDNECVAMAAMLLTFYMWVRSLSRSSSSWPICALAGVAYG
Tbgam AFTAYFFSIVPAHLMRSMAGEFDNECVAMAAMLLTFYMWVRSLSRSSSSWPICALAGVAYG

TbrOSTII YMVSTWGGYIFVLNMVAFHASVCVLLDWARGTYSVSLLRAYSLEFFVIGTALAICVPPVEW
Tbrucei YMVSTWGGYIFVLNMVAFHASVCVLLDWARGIYSVSLLRAYSLEFFVIGTALAICVPPVEW
Tbgam YMVSTWGGYIFVLNMVAFHASVCVLLDWARGTYSVSLLRAYSLEFFVIGTALAICVPPVEW

TbrOSTII TPFRSLEQLTALFVVFVMWALHYSEYLRERARAPIHSSKALQIRARIFMGTLSELLIVAS
Tbrucei TPFRSLEQLTALFVVFVMWALHYSEYLRERARAPIHSSKALQIRARIFMGTLSELLIVAS
Tbgam TPFRSLEQLTALFVVFVMWALHYSEYLRERARAPIHSSKALQIRARIFMGTLSELLIVAI

TbrOSTII LLAPFGFFKPTAYRVRALFVKHTRTGNPLVDSVAEHRPTTAGAYLRYFHVCYPLWCGGGL
Tbrucei LLAPFGFFKPTAYRVRALFVKHTRTGNPLVDSVAEHRPTTAGAYLRYFHVCYPLWCGGGL
Tbgam YLFSTGYFRSFSRVRALFVKHTRTGNPLVDSVAEHRPTTAGAFLRHLHVCYNGWIIGFF

TbrOSTII SMLVFMKKDRWRAIVFLASLSTVTMYFSARMSRLLLL LAGPAATACAGMFIGGLFDLALSQ
Tbrucei SMLVFMKKDRWRAIVFLASLSTVTMYFSARMSRLLLL LAGPAATACAGMFIGGLFDLALSQ
Tbgam FMSVSCFFHCTPGMSFLLLYSILAYYFSLKMSRLLLL SAPPVASILTGYVVGSIVDLAADC

TbrOSTII FGDLRSPKASGSDSPAGGSKRAKGVVNEPPKRAIF----SHRWFQRL-VQSLPVPLRR
Tbrucei FGDLRSPKASGSDSPAGGSKRAKGVVNEPPKRAIF----SHRWFQRL-VQSLPVPLRR
Tbgam FA-----ASGTEH--ADSKHQGKARGKQKRQITVECGCHNPFYKLWCNSFSSRLVV

TbrOSTII GIAVVVLVCLFANP--MRHSFEKSCEKMAHALSSPRIIAVTDLPNGERVLADDDYVSYLW
Tbrucei GIAVVVLVCLFANP--MRHSFEKSCEKMAHALSSPRIIAVTDLPNGERVLADDDYVSYLW
Tbgam GKFFVVVLAICGPTFLGSEFRAHCERFSLSVANPRIIS-SIRHSGKLVLADDDYVSYLW

TbrOSTII LRNNTPEDARILSWWDYGYQITGIGNRTTLADGNTWSHKHIATIGKMLTSPVKESHALIR
Tbrucei LRNNTPEDARILSWWDYGYQITGIGNRTTLADGNTWSHKHIATIGKMLTSPVKESHALIR
Tbgam LRNNTPEDARILSWWDYGYQITGIGNRTTLADGNTWNHEHIATIGKMLTSPVKESHALIR

TbrOSTII HLADYVLIWSGQDGSDDLKSPHMARIGNSVYRDMCSEDDPLCRQFGFYSGDLNKPTPMMQ
Tbrucei HLADYVLIWAGEDRGDLLKSPHMARIGNSVYRDMCSEDDPRCRQFGFEGDLNKPTPMMQ
Tbgam HLADYVLIWSGQDRGDLRKSRRHMARIGNSVYRDMCSEDDPLCRQFGFYSGDLKSKPTPMMQ

TbrOSTII RSLLYNLHRFGTDGGKTQLDKNMFQLAYVSKYGLVKIYKVMNVSEESKAWVADPKNRVCD
Tbrucei RSLLYNLHRFGTDGGKTQLDKNMFQLAYVSKYGLVKIYKVMNVSEESKAWVADPKNRVCD
Tbgam RSLLYNLHRFGTDGGKTQLDKNMFQLAYVSKYGLVKIYKVMNVSEESKAWVADPKNRKCD

TbrOSTII PPGSWICAGQYPPAKEI-----
Tbrucei PPGSWICAGQYPPAKEIQDMLAKRFHYE-----
Tbgam APGSWICAGQYPPAKEIQDMLAKRIDYEQLEDFNRRNRSDAHYRAYMRQMG
    
```

Figure 3. Amino Acid sequence identity of *Trypanosoma brucei rhodesiense* Putative Oligosaccharyl transferase (TbrOST II) within genome of genus *Trypanosoma*, family Trypanosomatidae. Aligned sequences were from *T. brucei* -Tbrucei (98%, XP_844763.1) and *T. brucei gambiense* -Tbgam (78%, CBH10989.1) species.

AJ251127.1; As:46% , CAB61 569.1).

DISCUSSION

The surface coat of trypanosome species were previously reported to have marked diversities in N-glycosylation [20]. This prompted us to hypothesize that trypanosomes also possess N-glycosylation enzyme. During the attempt to clone *T. brucei rhodesiense* putative oligosaccharyl transferase, a genomic clone of ~2000 bp was acquired.

Notably, *L. major* putative OST STT3 subunit sequence was used as the query during the primer design for PCR amplification of TbrOST II. The efficient amplification of the gene in genus *T. brucei rhodesiense* using primers designed from *L. major* stt3 gene as the query strongly indicate that TbrOST II seems to be conserved within the family Trypanosomatidae. The nucleotide BLAST results indicates that sequence identity is higher between species of the same genus rather than of the same family. It is known that *T. brucei*, *T. gambiense* and *T. rhodesiense* are members of the *Brucei*-complex or *Brucei* group. Although *T. brucei* *brucei* has more similarities with *T. brucei rhodesiense* than *T. brucei gambiense*, these parasites are morphologically indistinguishable [18,23]. This is the probable reason why high sequence identity was displayed by other subspecies of the *Brucei* group. This also indicates that the functional unit of this putative oligosaccharyl transferase is conserved within the *Brucei* complex trypanosomes. In addition, the high percent identity possessed by TbrOST II with other trypanosomatids agrees with the evolutionarily conserved characteristics of the established OST [27, 25]. Consequently, DNA identity searches within the public databases obtained homologues in the genome of mammalian, nematode, arthropod and algae species. Selected organisms include *Mus musculus* (Ns:71%, NM_024222.2; As:29%, NP_077184.2), *Schistosoma mansoni* (Ns:73%, XM_002577919.1; As:29%, XP_002577965.1), *Drosophila pseudoobscura pseudoobscura* (Ns:77%, XM_002134519.1; As:31%, XP_002134555.1), *Thalassiosira pseudonana* (Ns:77%, XM_002288187.1; As:46%, XP_002288223.1) and *Phaeodactylum tricorutum* (Ns:73%, XM_002185331.1; As:46%, XP_002185367.1). Moreover, obtained amino acid sequence identity in the genome of fly, nematode, mouse and algae homologues also exhibited lower sequence similarity (29-46%) than the previously documented percent identity (~50%) of reported

oligosaccharyl transferase in other eukaryotic species [25].

CONCLUSIONS

Finally, even currently regarded as putative, DNA sequence data of TbrOST II showing similar sequences in the genome of other organisms further corroborate the previous reports regarding the ubiquitous nature of OST in other life forms. Based on the size of the amplicon and significant percentage of nucleotide and amino acid sequence identity to homologues within the genome of related species and various organisms, the results strongly indicate that TbrOST II is a trypanosome oligosaccharyl transferase gene candidate that should be fully characterized and subjected to functional genomic studies. The study reports the molecular cloning and sequencing of a potential oligosaccharyl transferase gene in *T. brucei rhodesiense* (TbrOST II). The sequence data has been deposited in the GenBank with accession number of GU475126.

SOURCES AND MANUFACTURERS

¹CLEA Japan, Inc., Tokyo, Japan.

²Biosource International Inc., Camarillo, CA, USA.

³Sigma, St. Louis, USA.

⁴Calbiochem, La Jolla, CA, USA.

⁵BDH Chemicals, Poole, England.

⁶Invitrogen Co., Carlsbad, CA, USA.

⁷Meiji Seika Kaisha, Ltd., Tokyo, Japan.

⁸Wako Pure Chemical Industries Ltd., Osaka, Japan.

⁹Sigma-Genosys, Hokkaido, Japan.

¹⁰ GeneClean kit II, Q Biogene, Vista, CA, USA.

¹¹Novagen, Madison, WI, USA.

¹²TAKARA Shuzo, Japan.

¹³Applied Biosystems, Foster City, CA, USA.

¹⁴Mac Vector 6.5 sequence analysis software, Oxford University, England.

¹⁵CLC Workbench 5.6 software. USA.

Acknowledgment. This work was supported by the Japanese Ministry of Education, Culture, Sports, Science and Technology (Monbukagakusho).

REFERENCES

- 1 Altschul S.F., Madden T.L., Schäffer A.A., Zhang J., Zhang Z., Miller W. & Lipman D.J. 1997.** "Gapped BLAST and PSI-BLAST: a new generation of protein database search programs". *Nucleic Acids Research*. 25(17): 3389-3402.
- 2 Baticados W.N., Witola W.H., Inoue N., Kim J., Kuboki N., Xuan X., Yokoyama N. & Sugimoto C. 2005.** Expression of a Gene Encoding *Trypanosoma congolense* Putative Abc1 Family Protein is Developmentally Regulated. *Journal of Veterinary Medical Science*. 67(2): 157-164.
- 3 Breuer W. & Bause E. 1995.** Oligosaccharyl transferase is a constitutive component of an oligomeric protein complex from pig liver endoplasmic reticulum. *European Journal of Biochemistry*. 228(3): 689-696.
- 4 Brodsky J.L. 1997.** Translocation of proteins across the endoplasmic reticulum membrane. *International Review of Cytology*. 178: 277-328.
- 5 Cross G.A.M. 2002.** Antigenic variation in African trypanosomes and malaria. In: *Molecular Medical Parasitology*. Marr J.J., Nilsen T.W. & Komuniecki R.W. (Eds). Academic Press: New York, pp.89-110.
- 6 Donelson J.E., Hill K.L. & El-Sayed N.M.A. 1998.** Multiple mechanism of immune evasion by African trypanosomes. *Molecular & Biochemical Parasitology*. 91(1): 51-66.
- 7 Ferguson M.A.J. 1997.** The structure and biosynthesis of trypanosomatid glycosylphosphatidylinositols. In: Hide G. & Mottram J.C. (Eds). *Trypanosomiasis and Leishmaniasis*. U.K.: **Commonwealth Agricultural Bureaux** International, pp.65-77.
- 8 Garcia R.C., Cundell D.R., Tuomanen E.I., Kolakowski Jr. L.F., Gerard C. & Gerard N.P. 1995.** The role of N-glycosylation for functional expression of the human platelet-activating factor receptor. Glycosylation is required for efficient membrane trafficking. *Journal of Biological Chemistry*. 270(11): 25178-25184.
- 9 Goder V., Bieri C. & Spiess M. 1999.** Glycosylation can influence topogenesis of membrane proteins and reveals dynamic reorientation of nascent polypeptides within the translocon. *Journal of Cell Biology*. 147(2): 257-265.
- 10 Hirumi H. & Hirumi K. 1991.** *In vitro* cultivation of *Trypanosoma congolense* bloodstream forms in the absence of feeder cell layers. *Parasitology*. 102(2): 225-236.
- 11 Kaplan H.A., Welply J.K. & Lennarz W.J. 1987.** Oligosaccharyl transferase: the central enzyme in the pathway of glycoprotein assembly. *Biochimica et Biophysica Acta*. 906(2): 161-173.
- 12 Kelleher D.J. & Gilmore R. 1997.** DAD1, the defender against apoptotic cell death, is a subunit of the mammalian oligosaccharyltransferase. *Proceedings of the National Academy of Sciences*. 94(10): 4994-4999.
- 13 Knauer R. & Lehle L. 1999.** The oligosaccharyltransferase complex from *Saccharomyces cerevisiae*. *Journal of Biological Chemistry*. 274 (24): 17249-17256.
- 14 Kumar V., Heinemann F.S. & Ozols J. 1994.** Purification and characterization of avian oligosaccharyltransferase. Complete amino acid sequence of the 50-kDa subunit. *Journal of Biological Chemistry*. 269(18): 13451-13457.
- 15 Levin D. & Bartlett-Heubusch E. 1992.** Mutants in the *S. cerevisiae* *PKC1* gene display a cell cycle-specific osmotic stability defect. *Journal of Cell Biology*. 116(5): 1221-1229.
- 16 Lisowska E. 2002.** The role of glycosylation in protein antigenic properties. *Cellular and Molecular Life Sciences*. 59(3): 445-455.
- 17 Live D.H., Kumar R.A., Beebe X. & Danishefsky S.J. 1996.** Conformational influences of glycosylation of a peptide: A possible model for the effect of glycosylation on the rate of protein folding. *Proceedings of the National Academy of Sciences*. 93(23): 12759-12761.
- 18 Matthews K.R. 2005.** The developmental cell biology of *Trypanosoma brucei*. *Journal of Cell Science*. 118(2): 283-290.
- 19 Mehlert A., Bond C.S. & Ferguson M.A.J. 2002.** The glycoforms of a *Trypanosoma brucei* variant surface glycoprotein and molecular modeling of glycosylated surface coat. *Glycobiology*. 12(10): 607-612.
- 20 Parodi A.J. 1993.** N-glycosylation in trypanosomatid protozoa. *Glycobiology*. 3(3): 193-199.
- 21 Pays E. & Nolan D.P. 1998.** Expression and function of surface proteins in *Trypanosoma brucei*. *Molecular & Biochemical Parasitology*. 91(1): 3-36.
- 22 Silberstein S., Kelleher D.J. & Gilmore R. 1992.** The 48-kDa subunit of the mammalian oligosaccharyltransferase complex is homologous to the essential yeast protein WP1. *Journal of Biological Chemistry*. 267(33): 23658-23663.
- 23 Soulsby E.J.L. 1982.** Phylum: Sarcostigophora Honigberg and Balamuth, 1963, Subphylum: Mastigophora Dising, 1866. In: Soulsby E.J.L. (Ed). *Helminths, Arthropods and Protozoa of Domesticated Animals*. 7th edn. London: Baillière Tindall, pp.514-582.
- 24 Strickler J.E. & Patton C.L. 1980.** *Trypanosoma brucei brucei*: Inhibition of glycosylation of the major variable surface coat glycoprotein by tunicamycin. *Proceedings of the National Academy of Sciences*. 77(3): 1529-1533.

25 Yan Q. & Lennarz W.J. 2002. Studies on the function of oligosaccharyl transferase subunits. *Journal of Biological Chemistry*. 277(49): 47692-47700.

26 Zoller T., Fèvre E.M., Welburn S.C., Odiit M. & Coleman P.G. 2008. Analysis of risk factors for *T. brucei rhodesiense* sleeping sickness within villages in south-east Uganda. *BMC Infectious Diseases*. 8(88): 1-9.

27 Zufferey R., Knauer R., Burda P., Stagljar I., Heesen S., Lehle L. & Aebi M. 1995. STT3, a highly conserved protein required for yeast oligosaccharyl transferase activity *in vivo*. *EMBO Journal*. 14(20): 4949-4960.