

# Paediatric laser dentistry.

## Part 3: Dental trauma

C. Caprioglio \*  
G. Olivi\*\*  
M.D. Genovese\*\*\*  
M.C. Vitale\*\*\*\*

\*Visiting professor University of Pisa, Pisa, Italy

Private practice, Pavia, Italy

\*\*Visiting professor, Università Cattolica del Sacro Cuore, Rome, Italy

Private practice, Rome, Italy

\*\*\* Private practice, Rome, Italy

\*\*\*\*Assistant professor, Department of Clinical, Surgical, Diagnostic and Paediatric Sciences, Section of Dentistry, Unit of Orthodontics and Paediatric Dentistry, University of Pavia, Italy

e-mail: claudiagiulia.caprioglio@gmail.com - olivilaser@gmail.com

DOI: 10.23804/ejpd.2017.18.03.14

### ABSTRACT

Dental traumas are frequent and sometimes complex events, and at times real emergencies. There are no well-coded guidelines for laser applications in these clinical events. Laser-assisted therapy can offer new treatment possibilities, simplify dental procedures, reduce post-operative sensitivity and the need for post-operative medications because of the laser-induced biostimulating and anti-inflammatory effects.

**Keywords** Dental trauma; Laser dentistry; Low level laser therapy; Paediatric dentistry.

## Introduction

Approximately 20% of children suffer a traumatic injury to their primary teeth and more than 15% injure their permanent ones [Andreasen et al., 2007; Glendor, 2008; Kramer et al., 2016; Bagattoni, 2017]. Maxillary central incisors (50%) and maxillary lateral incisors (30%) are the teeth most frequently affected,

both in permanent and in primary dentition [Flores, 2002]. The literature shows that boys sustain more traumatic dental injuries than girls [Andreasen, 2007; Bani, 2016].

The revised classification of traumatic dental injuries of the World Health Organization includes injuries to teeth, supporting structures, and gingival and oral mucosa, and it is based on anatomical, therapeutic and prognostic considerations (Table 1) [WHO, 1995; Flores, 2001; Andreasen, 2007].

Careful collection of dental history, a good clinical examination, diagnostic imaging, photographic documentation, pulp testing are required for a complete medicolegal report.

## Laser in dental traumatology: hard tissues and pulp

Laser application offers multiple advantages in dental traumatology involving pulp and hard tissues.

- Pulp temperature increases only minimally during erbium laser treatment [Keller and Hibst, 1991; Hibst and Keller, 1996; Dostalova et al., 1997; Ozturk et al., 2004, Khouja, 2017].
- Laser irradiation provides high decontamination of

### 1. Traumatic injuries to the hard dental tissues and pulp

Crown infraction
Uncomplicated crown fracture
Complicated crown fracture
Uncomplicated crown-root fracture
Complicated crown-root fracture
Root fracture in the apical third
Root fracture in the middle third
Root fracture in the coronal third

### 2. Traumatic injuries to the periodontal tissues

Concussion
Subluxation
Extrusive luxation
Lateral luxation
Intrusive luxation
Avulsion

### 3. Injuries to the supporting bone

Comminution of the maxillary alveolar socket
Comminution of the mandibular alveolar socket
Fracture of the maxillary alveolar socket wall
Fracture of the mandibular alveolar socket wall
Fracture of the maxillary alveolar process
Fracture of the mandibular alveolar process
Fracture of the maxilla
Fracture of the mandible

### 4. Injuries to gingiva or oral mucosa

Laceration of gingival or oral mucosa
Contusion of gingival or oral mucosa

TABLE 1 Classification of traumatic injuries.

the exposed site (bactericidal effect) [Turkun et al., 2006; Moritz, 2006].

- Laser ablation removes the smear layer and debris, leaving the dentinal tubules open and thus allowing the adhesion process [Delmè and De Moor, 2007; Olivi et al., 2009].

The entire treatment can be performed with erbium laser, including preparation and conditioning for the reattachment of the fragment.

Laser therapy improves the psychological approach and the compliance of the patient, reducing discomfort and inducing laser analgesia by raising the pain threshold [Boj, 2005; Genovese and Olivi, 2008].

Crown infractions are rare; non-ablative laser energy in a defocused mode reduces the enamel-dentin permeability, the bacterial load and thus sensitivity. The procedure can be repeated after 7 to 15 days.

Enamel-crown fracture without pulp exposure (uncomplicated fracture) can be treated by using Erbium lasers [Caprioglio, 2010-2011; Olivi, 2010].

Erbium laser is used for enamel and dentin conditioning: signs of submorphological damage are related to the thermal vaporisation of the organic dentinal fibres and of the peripheral interprismatic enamel structure; acid etching is needed.

A complicated crown fracture exposes the pulp: treatment depends on the size of the exposure and the timeframe between the traumatic injury and treatment [AAPD, 2005-2006; Caprioglio, 2014; Smail-Faugeron, 2017]. The options include: pulp capping (Fig. 1), partial pulpotomy, pulpectomy and root canal therapy.

If the exposed area is very small (< 1 mm<sup>2</sup>) and the treatment is rapidly provided, pulp capping is the first choice of treatment: different laser wavelengths can be used for decontamination and coagulation of the exposed pulp.



**FIG. 1** Complicated crown fracture of the maxillary lateral incisor, treated with defocused treatment for coagulation and dentin melting (Erbium laser at 100 mJ).

### Pulp capping

Among the clinical trials and histological investigations in animal models, Santucci [1997] was the first to report the high success rate (90% after 6 months) *in vivo* using Nd:YAG laser and a glass ionomer cement for pulp capping. Later studies [Moritz, 1998] reported different success rates (89-93%) after 1 and 2 years using a CO<sub>2</sub> laser versus calcium hydroxide. Olivi et al. [2006-2007] have investigated *in vivo* different laser systems and calcium hydroxide, reporting success rates of 80% for Er,Cr:YSGG, 75% for Er:YAG and 63% for calcium hydroxide alone. Recently, Cengiz and Yilmaz [2016] have investigated Er,Cr:YSGG laser irradiation at low energy, with no agents and reported success rates after 6 months of 100% the laser, 73,3% for calcium hydroxide alone group and 66,6% for a resin-based tricalcium silicate material alone. Yazdanfor et al. [2015] have investigated a diode-laser-assisted method compared to conventional procedure, reporting 60% survival rate for the conventional method and 100% for the diode 808 nm procedure after one year. Several studies on different animal models have been performed by Jawardena et al. [2001] using Er:YAG laser; Hasheminia et al. [2010] reported better healing using Er:YAG laser with MTA in comparison with MTA alone. Suzuki et al. [2011] have investigated super-pulsed CO<sub>2</sub> laser preirradiation on exposed rat pulp in three groups, one irradiated and directly capped with a self-etching adhesive system, one capped with the self-etching adhesive system and the third was capped with commercially available calcium hydroxide, and the self-etching adhesive system was applied to the cavity, reporting no significant differences between the groups. However, the CO<sub>2</sub> laser irradiation showed a tendency to delay the formation of reparative dentin.

### Laser pulpotomy and pulpectomy

Pulpotomy is a very common technique in primary teeth, however several investigations using different wavelengths reported controversial results [Mendoza 2017].

Pulpotomy can be performed by using CO<sub>2</sub>, Nd:YAG, Erbium and diode lasers.

Two systematic reviews presented different results. De Coster et al. [2013] reviewed 7 articles and concluded that lasers (632/980 nm diode lasers, Nd:YAG, Er, YAG, CO<sub>2</sub>) were less successful than conventional pulpotomy techniques. Another study by Lin et al. [2012] including 37 studies and 22 meta-analyses highlighted that formocresol ferric sulfate and MTA had significantly better results in primary molar pulpotomy than calcium hydroxide and laser therapy in a 18-24 month follow-up study, with MTA as the first choice. More recent studies reported more positive results for lasers. Yadav et al. [2014] reported a better outcome of diode pulpotomy and electrosurgical pulpotomy versus ferric sulfate pulpotomy after 9 months. A randomised clinical



**FIG. 2** Complicated complete crown fracture in a 4-year-old patient. Gingivoplasty and tooth margin preparation are performed by using an Erbium laser.



**FIG. 3** Avulsion of the maxillary left central and lateral incisors in a 3-year-old girl. KTP laser irradiation (LLLT).

trial [Gupta et al., 2015] showed good clinical and radiographical results for laser pulpotomy performed on 30 primary molars. The importance of the sealing material was reported by different authors [Niranjani et al., 2015; Uloopi et al., 2016].

There are few studies on laser use for pulpectomy (Indexed on PubMed) in primary teeth (see Part 2).

## Laser in traumatic injuries to soft tissues

Traumatic injuries to supporting structures (alveolar bone, gingiva, ligaments, periodontium, fraenum and lips), defined as indirect traumas, can be effectively treated using lasers, especially the visible and near infrared types. These lasers are used for their ability to decontaminate the periodontal defect following a dental luxation or subluxation, for decontamination of the socket after an avulsion, for the ability to perform gingivectomy and/or gingivoplasty or surgical incisions (e.g. to remove a tooth fragment embedded in the soft tissue) (Fig. 2).

In all types of luxation injuries, lasers provide not only a bactericidal and detoxification effect (Er:YAG, Nd:YAG, and diode), but also favourable conditions for the attachment of the periodontal tissue, especially in the permanent dentition [Simunovic, 2000; Schindl, 2003; Carvalho et al., 2012; Matos et al., 2016]. In dental avulsion, which is often associated with injuries to the lips, a diode or Nd:YAG laser can be used to decontaminate the socket, promote healing of the oral mucosa and reduce pain. Some studies underlined the importance of removing the infected granulation tissue from the soft tissue walls [Schwarz, 2003; Crespi, 2007]. The biostimulating and pain control effects obtained with Low Level Laser Therapy (LLLT) are an integral part of the post-operative treatment of these patients. If oral mucosa and gingiva injuries are present, the LLLT can be used to help the tissue repair process through cell proliferation activation and reduction of inflammation (Fig. 3).

There is a large body of literature on this particular topic, even though, methodologically and in terms of doses, there is still considerable difference of opinion.

Semi-conductor diode type lasers (803 to 980 nm) are near-infrared lasers that at low power can be used for effective treatment of pain and inflammation and to promote tissue repair with great tissue penetration [Boj, 2011; Caprioglio, 2011; Ghaderi, 2017]. They can influence a large number of cell systems (fibroblasts, macrophages, lymphocytes, epithelial cells, endothelium) and can also have a series of benefits on the inflammatory mechanism, reducing the exudative phase and stimulating the reparative process [De Souza, 2016].

LLLT modifies nerve impulse stimulation and transmission and increases the metabolism of endorphins, acetylcholine, serotonin, and cortisol, resulting in reduced stimulation and perception of pain [Schindl, 2000; Simunovic, 2000; Pinheiro et al., 2004-2006]. LLLT increases production of adenosine triphosphate (ATP) as well as overall cell activity. Laser light increases mitochondrial ATP and can also react with beta growth factors [Karu, 1987-1988; Pinheiro, 2006]. LLLT modifies blood flow and induces angiogenesis, and the modification of lymph drainage reduces inflammation.

The study of these new treatments might lead to the definition of guidelines and protocols with specific doses and application sites.

## References

- > American Academy of Pediatric Dentistry. Guideline on pulp therapy for primary and young permanent teeth. *Pediatr Dent* 2005-2006; 27 (7 Suppl):130-134.
- > Application of the International Classification of Diseases and Stomatology. ICD-DA, ed 3. Geneva:World Health Organization, 1995.
- > Andreasen JO, Andreasen FM. Textbook and colour Atlas of Traumatic Injuries to the Teeth, ed 4. Ames, IA; Wiley-Blackwell, 2007:542-575.
- > Bagattoni S, Sadotti A, D'Alessandro G, Piana G. Dental trauma in Italian children and adolescents with special health care needs. A cross-sectional retrospective study. *Eur J Paediatr Dent* 2017;18(1):23-26.
- > Bani M, Bodur H, Kapci EG. Are behaviour risk factors for traumatic dental injuries in childhood different between males and females? *Eur J Paediatr Dent* 2016;16(1):25-32.
- > Boj JR, Galofre N, Espana A et al. Pain perception in pediatric patients undergoing laser treatment. *J Oral Laser Appl* 2005;5:85-89.
- > Boj JR, Poirier C, Hernandez M et al. Review: laser soft tissue treatments for paediatric dental patients. *Europ Arch Paed Dent* 2011;12:100-105.
- > Caprioglio C, Olivi G, Genovese MD. *I Laser in Traumatologia dentale*. Bologna, Italy: Martina Ed. 2009.
- > Caprioglio C. Lasers in dental Traumatology. *Laser* 2010;(2):12-18.
- > Caprioglio C, Caprioglio A. Dental Trauma in children. In Vitale MC, Caprioglio C (editors). *Lasers in Dentistry*, Bologna, Italy, Martina Ed.

- 2010; 105-138.
- › Caprioglio C, Olivi G, Genovese MD. Lasers in dental Traumatology and Low Level Laser Therapy (LLLT). *Europ Arch Paed Dentistry* 2011;12(2):79-84.
  - › Caprioglio C. Dental Trauma. In: Olivi G, Margolis F, Genovese MD. *Pediatric laser Dentistry: a user's guide*. Quintessence Pub: Chicago; 2011. Chapter 13, p. 181-190 and Chapter 14; p. 191-195.
  - › Caprioglio A, Conti V, Caprioglio C et al. A long term retrospective clinical study on MTA pulp potomies in immature permanent incisors with complicated crown fractures. *Eur J of Ped Dent* 2014;15(1):1-7.
  - › Carvalho EDS, Costa FTS, Campos MS et al. Root surface treatment using diode laser in delayed tooth replantation: radiographic and histomorphometric analysis in rats. *Dent Traumatol* 2012;28:429-
  - › Cengiz E, Yilmaz HG. Efficacy of Erbium, Chromium-doped: Yttrium, Scandium, Gallium, and Garnet Laser Irradiation Combined with Resin-based Tricalcium Silicate and Calcium Hydroxide on Direct Pulp Capping: a Randomized Clinical Trial. *J Endod* 2016;42(3):351-5. Epub 2015 Dec 23.
  - › Crespi R, Cappare P, Toscanelli I, Gherlone E, Romanos GE. Effects of Er:YAG laser compared to ultrasonic scaler in periodontal treatment: A 2-year follow-up split-mouth clinical study. *Periodontol* 2007;78:1195-1200.
  - › De Coster P, Rajasekharan S, Martens L. Laser-assisted pulpotomy in primary teeth: a systematic review. *Int J Paediatr Dent* 2013 Nov;23(6):389-99. Epub 2012 Nov 22.
  - › Delmè K, Deman PJ, Mieke AA, De Bruyne MMA, De Moor RJG. Influence of different Er:YAG laser energies and frequencies on the surface morphology of dentin and enamel. *J Oral Laser Application* 2006;6:43-52.
  - › Delmè KI, De Moor RJ. Scanning electron microscopic evaluation of enamel and dentin surfaces after Er:YAG laser and laser conditioning. *Photomed Laser Surg* 2007 Oct;25(5):393-401.
  - › De Souza Matos F, De Jesus Godolphim F et al. Effect of laser photobiomodulation on periodontal repair process of replanted teeth. *Dental Traumatol* 2016;32:402-408.
  - › Dostalova T, Jelinkova H, Krejsa O, Hamal K, Kubelka J, Prochazka S, Himmlova L. Dentin and pulp response to Erbium:YAG laser ablation: a preliminary evaluation of human teeth. *J Clin Laser Med Surg* 1997;15(3):117-121.
  - › Flores MT, Andreasen JO, Bakland LK, Feiglin B et al. Guidelines for the evaluation and management of traumatic dental injuries. *Dent Traumatol* 2001;17:1-2.
  - › Flores MT. Traumatic injuries in the primary dentition. *Dental Traumatol* 2002;3:287-98.
  - › Genovese MD, Olivi G. Laser in paediatric dentistry: Patient acceptance on hard and soft tissue therapy. *Eur J Paediatr Dent* 2008;9:13-17.
  - › Ghaderi F, Ghaderi R, Davarmanesh M, Bayani M, Arabzade Moghadam S. Pain management during needle insertion with low level laser. *Eur J Paediatr Dent* 2017;17(2):151-154.
  - › Glendor U. Epidemiology of traumatic dental injuries- A 12 year review of the literature. *Dent Traumatol* 2008;25:19-31.
  - › Gupta G, Rana V, Srivastava N, Chandna P. Laser Pulpotomy-An Effective Alternative to Conventional Techniques: A 12 Months Clinicoradiographic Study. *Int J Clin Pediatr Dent* 2015 Jan-Apr; 8(1):18-21. Epub 2015 Apr 28.
  - › Hasheminia SM, Feizi G, Razavi SM, Feizianfard M, Gutknecht N, Mir M. A comparative study of three treatment methods of direct pulp capping in canine teeth of cats: a histologic evaluation. *Laser Med Sci* 2010 Jan;25(1):9-15.
  - › Hibst R, Keller U. Effects of water spray and repetition rate on the temperature elevation during Er:YAG laser ablation of dentin. *Medical Application of Laser III, Proc SPIE* 1996.
  - › Jawardena JA, Kato J, Moriya K et al. Pulpal response to exposure with Er:YAG laser. *Oral Surg Oral med Oral Pathol Oral Radiol Endod* 2001;91:222-229.
  - › Yadav P, Indushekar K, Saraf B, Sheoran N1, Sardana D. Comparative evaluation of Ferric Sulfate, Electrosurgical and Diode Laser on human primary molars pulpotomy: an "in-vivo" study. *Laser Ther* 2014 Mar 27;23(1):41-7.
  - › Yazdanfar I, Gutknecht N, Franzen R. Effects of a diode laser on direct pulp capping treatment: a pilot study. *Lasers Med Sci* 2015 May;30(4):1237-43. Epub 2014 Apr 23.
  - › Karu TI. Photobiological fundamentals of low-power laser therapy. *J Quant Electron* 1987;10:1703-1717.
  - › Karu TI. Molecular mechanism of therapeutic effect of low intensity laser irradiation. *Laser Life Sci* 1988;2:53-74.
  - › Keller U, Hibst R. Tooth pulp reaction following Er:YAG laser application. *Laser in Orthopedic, Dental and Veterinary medicine. Proc SPIE* 1991;1424:127-133.
  - › Khouja F, Abdelaziz M, Bortolotto T, Krejci I. Intra-pulpal and subsurface temperature rise during tooth irradiation with 808 nm diode laser: an in vitro study. *Eur J Paediatr Dent* 2017;18(1):56-60.
  - › Kramer PF, Onetto J, Flores MT, et al. Traumatic Dental Injuries in the primary dentition: a 15 year bibliometric analysis of Dental Traumatology. *Dental Traumat* 2016;32:341-346.
  - › Lin PY, Chen HS, Wang YH, Tu YK. Primary molars pulpotomy: a systematic review and network meta-analysis. *J Dent* 2014 Sep;42(9):1060-77. Epub 2014 Feb 7.
  - › Mendoza-Mendoza A, Caleza-Jiménez C, Solano-Mendoza B, Iglesias-Linares A. Are there any differences between first and second primary molar pulpectomy prognoses? A retrospective clinical study. *Eur J Paediatr Dent* 2017;18(1):41-44.
  - › Moritz A, Schoop U, Goharkhay K, Sperr W. Advantages of pulsed CO<sub>2</sub> laser in direct pulp capping: a long term in vivo study. *Laser Surg Med.* 1998;22(5):288-93.
  - › Moritz A, Schoop U, Goharkhay K, Sperr W. The CO<sub>2</sub> laser as an aid in direct pulp capping. *J Endod* 1998 Apr;24(4):248-51.
  - › Olivi G. *Oral Laser Application*. Berlin: Quintessence Verlags;2006. P. 258-277.
  - › Niranjani K, Prasad MG, Vasa AA, Divya G, Thakur MS, Saujanya K. Clinical Evaluation of Success of Primary Teeth Pulpotomy Using Mineral Trioxide Aggregate ® laser and Biodentine™-an in vivo study. *J Clin Res* 2015 Apr;9(4):ZC35-7. Epub 2015 Apr 1.
  - › Olivi G, Genovese MD. Erbium Chromium laser in pulp capping treatment. *J Oral Laser Appl* 2006;6:291-299.
  - › Olivi G, Genovese MD, Maturo P, Docimo R. Pulp capping Advantages of using laser technology. *Eur J Ped Dent* 2007;8:89-95.
  - › Olivi G, Genovese MD, Caprioglio C. Evidence-based dentistry on laser paediatric dentistry: Review and outlook. *Eur J Paediatr Dent* 2009;10:29-40.
  - › Olivi G, Caprioglio C, Genovese MD. Lasers in dental Traumatology. *Eur J Paediatr Dent* 2010;18(2):71-76.
  - › Ozturk B, Usumez A, Ozturk AN, Ozer F. In vitro assessment of temperature change in the pulp chamber during cavity preparation. *J Prosthet Dent* 2004;91(5):436-440.
  - › Pinheiro AL, Meireles GC, de Barros Viera AL et al. Phototherapy improves healing of cutaneous wounds in nonnourished and undernourished Wistar rats. *Braz Dent J* 2004;15:21-28.
  - › Pinheiro ALB, Gerbi ME. Photobiomodulation of bone repair process. *Photomed Laser Surg* 2006;24:169-178.
  - › Santucci PJ. Dycal versus Nd:YAG laser and Vitrebond for direct pulp capping in permanent teeth. *J Clin Laser Med Surg.* 1999 Apr;17(2):69-75.
  - › Schindl A, Schindl M, Pernerstofer-Schin H, Mossbacher U, Schindl L. Low-intensity laser therapy: A review. *J Invest Med* 2000;48:312-326.
  - › Schwarz F, Sculean A, Berakdar M, Georg T, Reich E, Beck. Periodontal treatment with an Er:YAG laser or scaling and planning. A 2-year follow up split-mouth study. *J Periodontol* 2003;74:590-596.
  - › Simunovic Z, Ivanovich AD, Depolo A. Wound healing of animal and human body sport and traffic accident injuries using low-level laser therapy treatment: A randomized clinical study of seventy-four patients with control group. *J Clin Laser Med Surg* 2000;18:67-71.
  - › Smail-Faugeron V, Porot A, Muller-Bolla M, Courson F. Indirect pulp capping versus pulpotomy for treating deep carious lesions approaching the pulp in primary teeth: a systematic review. *Eur J Paediatr Dent* 2017;17(2):107-112.
  - › Soares F, Varella CH, Pileggi R, Adewumi A, Guelmann M. Impact of Er,Cr:YSGG laser therapy on the cleanliness of the root canal walls of primary teeth. *J Endod* 2008 Apr;34(4):474-7.
  - › Suzuki M, Ogisu T, Kato C, Shinkai K, Katoh Y. Effect of CO<sub>2</sub> laser irradiation on wound healing of exposed rat pulp. *Odontology.* 2011 Jan;99(1):34-44.
  - › Turkun M, Turkun LFI, Celik EU, Atefi M. Bactericidal effect of Er,Cr:YSGG laser on *Streptococcus mutans*. *Dent Mater J* 2006;25(1):81-86.
  - › Uloopi KS, Vinay C, Ratnaditya A, Gopal AS, Mrudula KJ, Rao RC. Clinical Evaluation of Low Level Diode Laser Application For Primary Teeth Pulpotomy. *J Clin Diagn Res* 2016 Jan;10(1):ZC67-70.