

Research Article

Marine fouling communities from artificial and natural habitats: comparison of resistance to chemical and physical disturbances

Jasmine Ferrario^{1*}, Ignacio Gestoso^{2,7}, Patrício Ramalhosa^{2,3}, Eva Cacabelos^{2,4}, Bernardo Duarte⁵, Isabel Caçador⁵ and João Canning-Clode^{2,6,7}

¹Department of Earth and Environmental Sciences, University of Pavia, Via S. Epifanio 14, 27100, Pavia, Italy

²MARE – Marine and Environmental Sciences Centre, Quinta do Lorde Marina, Sítio da Piedade, 9200 - 044, Caniçal, Madeira, Portugal

³OOM – Oceanic Observatory of Madeira, Agência Regional para o Desenvolvimento da Investigação Tecnologia e Inovação, Edifício Madeira Tecnopolo, Piso 0, Caminho da Penteada, 9020-105, Funchal, Madeira, Portugal

⁴cE3c/ABG – Centre for Ecology, Evolution and Environmental Changes/Azorean Biodiversity Group, Ponta Delgada, Azores, Portugal

⁵MARE – Marine and Environmental Sciences Centre, Faculty of Sciences of the University of Lisbon, Campo Grande 1749-016, Lisbon, Portugal

⁶Centre of IMAR of the University of the Azores, Department of Oceanography and Fisheries. Rua Prof. Dr. Frederico Machado, 4, PT-9901-862, Horta, Azores, Portugal

⁷Smithsonian Environmental Research Center, 647 Contees Wharf Road, Edgewater, MD 21037, USA

Author e-mails: jasmine.ferrario@unipv.it (JF), igestoso@mare-centre.pt (IG), pramalhosa@mare-centre.pt (PR), ecacabelos@mare-centre.pt (EC), baduarte@fc.ul.pt (BD), micacador@fc.ul.pt (IC), jcanning-clode@mare-centre.pt (JCC)

*Corresponding author

Citation: Ferrario J, Gestoso I, Ramalhosa P, Cacabelos E, Duarte B, Caçador I, Canning-Clode J (2020) Marine fouling communities from artificial and natural habitats: comparison of resistance to chemical and physical disturbances. *Aquatic Invasions* 15(2): 196–216, <https://doi.org/10.3391/ai.2020.15.2.01>

Received: 12 July 2019

Accepted: 11 October 2019

Published: 11 December 2019

Thematic editor: Charles W. Martin

Copyright: © Ferrario et al.

This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International - CC BY 4.0).

OPEN ACCESS

Abstract

Assessing the resistance of fouling communities to anthropogenic disturbances is an important goal for the development of effective management and control strategies. In this context, we conducted a manipulative experiment on natural and artificial habitats to examine fouling communities that developed outside and inside a marina on Madeira Island (NE Atlantic Ocean) following the application of two types of stressors frequently observed in coastal habitats, namely chemical and physical disturbances. The tested fouling communities, dominated by native and non-indigenous species respectively, were in general strongly affected by the chemical but not by the physical disturbance applied, and a higher resistance to disturbance was observed in the communities outside the marina. This suggests higher capacities for communities richer in native species to tolerate anthropogenic disturbances, while non-indigenous species did not play a key role. Further research can assess the resilience of natural and artificial fouling communities when exposed to disturbances.

Key words: artificial habitats, mesocosms, antifouling paint, mechanical stress, Madeira, non-indigenous species

Introduction

The composition and structure of fouling communities are dependent on several factors, such as pollution, nutrient availability, sedimentation rate, water flow, turbulence, type of substrate and orientation (Glasby and Connell 1999; Bulleri and Chapman 2004; Dafforn et al. 2015; Simpson et al. 2017). Generally, artificial habitats (e.g., concrete docks and floating pontoons) provide novel hard substrates, a suitable condition for the proliferation and spread of non-indigenous species (NIS; Minchin et al.

2006; Mineur et al. 2012; Airoidi et al. 2015). In contrast, it is typical for fouling native species to dominate natural coastal areas (Glasby et al. 2007; Piola and Johnston 2008), and reflect communities that are mostly local in biodiversity (Lewis 1998; Chapman and Bulleri 2003; Karlson and Osman 2012; Gestoso et al. 2017; Simpson et al. 2017).

Globally, we are witnessing an impressive proliferation of artificial structures, the so-called “coastal urban sprawl”, that is considered one of the most important human threats to marine biodiversity and ecosystem services worldwide (Elmqvist et al. 2016; Firth et al. 2016). The loss of natural habitat in favor of coastal urbanization can change the structure and functioning of benthic communities, and studies generally conclude that the new artificial structures cannot be considered surrogates for the natural habitats they replace (e.g., Moschella et al. 2005; Firth et al. 2013). In addition, anthropogenic activities acting along artificial habitats, such as ports and marinas, can increase the connectivity among distant localities, breaking down geographical barriers and facilitating the “stepping-stone” spreading of species (Hewitt et al. 2009; Hulme 2009; Bishop et al. 2017). This in turn can enhance the biotic homogenization of marine fouling communities in artificial habitats (McKinney and Lockwood 1999; Glasby et al. 2007; Bulleri and Chapman 2010; Davidson et al. 2010; Seebens et al. 2013; Airoidi et al. 2015; Ferrario et al. 2017). Artificial fouling communities are usually dominated by highly competitive species (e.g., NIS and cosmopolitan species), which show greater tolerance to variable environmental conditions and are often thought to be more resilient to disturbance than native communities (e.g., Piola and Johnston 2008; Canning-Clode et al. 2011; Crooks et al. 2011).

Both artificial and natural habitats can be affected by anthropogenic disturbance events (either physical or chemical) that can influence resident fouling communities. A disturbance event can alter the structure of an ecosystem, community or population by changing the availability of resources and leading to opportunities for new settlers (Altman and Whitlatch 2007; Lockwood et al. 2007). Physical disturbances (e.g., vessel grounding procedures or maintenance interventions of port structures) can cause abrasions on the substrate itself (Hudson and Goodwin 2001). These types of incidents, caused by small vessels, are mostly unreported although are likely to occur more frequently than by large vessels [e.g., in coral reefs (Lutz 2006)]. Benthic organisms and seagrass beds can also be damaged by disturbances caused by recreational SCUBA diving or snorkeling activities. For example, organisms can be accidentally kicked or buried by fine particles suspended with the fins, or damaged by trampling, anchoring or propeller action in shallow waters (Garrabou et al. 1998; Milazzo et al. 2002; Davenport and Davenport 2006; Huff 2011; Mendez et al. 2017; Renfro and Chadwick 2017).

In addition, heavy metal pollution is considered one of the most significant chemical disturbances in ports and marinas, mostly derived from the use of antifouling paints (Hall et al. 1998; Piola and Johnston 2009; Canning-Clode et al. 2011). These paints have important chemical inhibitors such as metallic biocides (such as zinc oxide, ZnO) that are often used in combination with copper to increase their toxicity and facilitate the leaching process (Guardiola et al. 2012; Watermann et al. 2005). When applied to vessel hulls and to several other artificial fixed structures (e.g., pilings, pontoons, buoys), antifouling paints can inhibit the growth of fouling organisms, slowly release heavy metals and harmful biocides into the water [e.g., copper and zinc (Lewis 1998; Comber et al. 2002; Warnken et al. 2004; Chambers et al. 2006; Thomas and Brooks 2010; Neira et al. 2014), and consequently cause a certain degree of environmental contamination. This can impact local communities (Kinsella and Crowe 2016) leading to losses of sensitive species and reductions in native diversity (Piola and Johnston 2008). In contrast, some widespread fouling species are known to have a certain degree of resistance to antifouling paints, e.g., bryozoans in the genus *Watersipora* or *Bugula neritina* (Linnaeus, 1758) (Wisley 1962; Clark and Johnston 2005; Dafforn et al. 2008; Piola et al. 2009; Crooks et al. 2011; Ramalhosa et al. 2019); and the ascidian *Asciidiella aspersa* (Müller, 1776) (considered a NIS outside Europe; Kenworthy et al. 2018).

Moreover, establishment of NIS may be ameliorated by disturbance events because free niche areas become available for more tolerant colonizers (Occhipinti-Ambrogi and Savini 2003). An example is the successful and rapid colonization of the invasive macroalga *Caulerpa* spp. following the experimental removal of the native seagrass *Posidonia oceanica* (Linnaeus) Delile, 1813 in Mediterranean meadows (e.g., Ceccherelli et al. 2014; Uyà et al. 2018).

Fouling communities have been extensively used in ecological studies as a model system to investigate a variety of general questions of succession, invasion ecology and community resilience (e.g., Svensson et al. 2007; Canning-Clode et al. 2009; Marraffini et al. 2017). Previous research on fouling communities from port habitats have evaluated the responses to different anthropogenic disturbances, e.g., heavy metal pollution and mechanical creation of bare patches (Clark and Johnston 2005; Canning-Clode et al. 2011; Crooks et al. 2011; Ramalhosa et al. 2019). However, most of the previous research has focused on the effects of disturbances acting separately, whereas the interactive effects of disturbances on communities from different habitats (artificial vs. natural) remains poorly explored.

In this context, we conducted a manipulative experiment to test the pulse of two types of disturbances on fouling communities composed of native and non-native organisms, in order to assess and compare their responses and resistance to anthropogenic disturbances. In particular, we hypothesize that fouling communities from natural habitats, expected to be




Figure 1. Map of Madeira Archipelago with the location of the marina of Quinta do Lorde (Caniçal) where 24 PVC plates were deployed at 5 meters depth inside and outside the marina.

richer in native species, would be less adapted to anthropogenic disturbances than communities from artificial habitats, whose dominance of NIS should better facilitate resistance to disturbance events.

Materials and methods

Study sites and community recruitment

The Madeira Archipelago is a group of volcanic islands located in the NE Atlantic Ocean 700 km off the Moroccan coast (Figure 1). The recruitment of fouling organisms was conducted in two areas: inside (artificial habitat) and outside (natural rocky habitat) the marina of Quinta do Lorde (Caniçal, Madeira Island: 32.741667N; -16.713333W, Figure 1).

In June 2017, a total of 24 grey polyvinylchloride PVC plates ($14 \times 14 \times 0.3$ cm) were deployed for colonization at 5 m depth both inside (12 plates) and outside (12 plates) the marina, known to host NIS (Canning-Clode et al. 2013). We constructed an experimental structure that was composed of a base attached to the rocky bottom (outside the marina) or to a brick (inside the marina), and PVC plates were placed on top, adequately separated from the base (only the side facing down was used for the experiment). The bricks inside the marina were used as a weight and laid approximately 6 cm from the soft bottom. In order to simplify its deployment in the field, plates were grouped in three different units per area, i.e. inside and outside marina (Figure S1). After a 4 month-period, all plates were retrieved and brought back to the laboratory for sampling (mid-October 2017). A preliminary analysis was performed to discard a potential “unit” effect on

the plate recruitment before starting the mesocosm experiment (PERMANOVA, non-significant effect of unit: Pseudo- $F_{2,18} = 1.975$, $P(\text{perm}) = 0.084$).

Mesocosm system and experimental design

A manipulative experiment was designed to examine the resistance of the two fouling communities in response to two types of disturbance (i.e. chemical vs. physical). This experiment was conducted inside the mesocosm system and laboratory facilities at the Madeira research unit of MARE – Marine and Environmental Research Centre, located at Quinta do Lorde Marina. Two independent tanks (350 L) were filled with 10 μm filtered seawater in a continuous water flow system of about 30 ml/sec, generating a complete turnover of water every 4 hours. Additionally, constant aeration systems were used in each tank, and artificial light was provided by fluorescent lamps (OSRAM- FH 14W/840 HE Lumilux Cool White 55 cm) under an approximate photoperiod of 12/12 h (light/dark). The use of filtered seawater excluded any possible new recruits, so that we tested the disturbance effects of only the original communities.

In this study, two types of pulse disturbance, *sensu* Glasby and Underwood (1996), were applied: physical and chemical. The “physical disturbance” consisted of pressing the PVC plates with approximately 7 kg weight for 30 seconds, as a proxy for any kind of lighter mechanic disturbances that can affect fouling communities, e.g., vessel docking, diver tourism, anchors (Clark and Johnston 2005). The “chemical disturbance” consisted of exposing the PVC plates to an antifouling paint for 48 hours to simulate the arrival of a boat freshly painted with antifouling coating (Canning-Clode et al. 2011; Crooks et al. 2011; Ramalhosa et al. 2019), considering that paints are designed to release an initial higher amount of biocides, followed by a rapid decline with slow and constant leaching rates (Schiff et al. 2004; Piola and Johnston 2008). Specifically, bare PVC plates were double coated with an antifouling paint (International® YBA067-Trilux 33 Black, containing as main metallic biocide Zinc Oxide 25 < 50%) and placed at the bottom of individual 10 L containers, with the relative manipulated plates hanged from the top. Chemical analysis for heavy metal concentrations of the water inside the containers was carried out in order to verify the effective release of the inhibitors proper of the antifouling paint used. In total, four disturbance types (three replicates per disturbance per area) were applied in this mesocosm experiment: control (C; no disturbance), physical (Phy), chemical (Che) and the combination of both physical and chemical disturbances on the same plates (mixed; Mix). Within the 12 PVC plates from inside and 12 from outside the marina, three plates were randomly selected for each of the four conditions. The manipulative experiment lasted 4 weeks (from October 12th to November 7th, 2017) and the resistance of fouling communities to disturbances was assessed over four scheduled sampling times (Figure 2). On Time_0 the PVC




Figure 2. Scheme of the experiment phases (divided in sampling times: Time_0, D, 1, 2) and the type of disturbance used. Grey bars represent the PVC plates inside a single container during the Time_D phase, showing the different disturbances: Control (C: PVC plate not treated and simply moved into a container), Physical (Phy: PVC plate firstly pressed with a weight and then moved into a container), Chemical (Che: PVC plate moved into a container with another PVC plate on bottom previously coated with an antifouling paint, represented in red color) and Mixed (Mix: PVC plates with combined disturbances).

plates were collected in the field, first analyzed and arranged in two tanks of the mesocosm system (from October 12th to 14th, 2017). After an acclimation period of 3 days in the two tanks, a pulse disturbance phase 48h long was carried out (Time_D; “D” stands for disturbance), based on the selected disturbances (on October 20th, 2017). Finally, fouling communities were evaluated after the first recovery week (Time_1; on October 27th, 2017) and the second recovery week (Time_2; on November 7th, 2017). At each sampling event, abundance and structure of the fouling communities were examined. The fouling species were identified under a stereomicroscope (Leica S8APO) using different taxonomic keys, existing literature or taxonomic experts (e.g., Hayward and Ryland 1995; Souto et al. 2015, 2018). During the sampling times Time_D-2, a semi-quantitative index on the “health status” of each species was also considered (values from 1 to 4): value “1” was assigned for totally dead specimens/colonies; value “2” was when the percentage of dead specimens/colonies were higher than live ones; value “3” was when the percentage of live specimens/colonies were higher than dead ones; and value “4” was for totally live specimens/colonies.

All plates per sampling event were photographed using a Olympus Stylus TG-4 camera and images were subsequently analyzed with the software CPCe [Coral Point Count with excel extensions (Kohler and Gill 2006)] to assess the abundance (in percent cover; % cover) of fouling species by considering 50 random points in each plate, and this information was then integrated with the “health status” of each species. Finally, during Time_D, all the plates were moved into single containers for 48-hours and, after each sampling time, all the 24 plates were completely randomized in two experimental tanks (12 plates each) to reduce potential “tank effect”.

Chemical-physical analyses

Dissolved oxygen, pH, salinity and temperature of the water were assessed by using a multi-parameter probe (YSI Professional Plus) during the sampling period. Water parameters were assessed i) in the field, during the sampling of the experimental units (inside and outside marina); ii) twice in 4 random containers, during the disturbance period and; iii) 7 times in the two tanks, during the entire duration of the experiment.

Finally, a total of 12 water samples (3 per disturbance level) were collected and analyzed from each container (during Time_D), for heavy metal concentration (zinc and copper), in order to verify water quality differences among conditions and estimate the concentration of these elements. For the water analysis, a total of 9 ml of seawater per sample was collected from the selected containers, with the use of 10 ml sterile syringes (B/Braun Injekt™); then, 1 ml of ultrapure HNO₃ (Trace SELECT®, Sigma-Aldrich) was added to each sample. Samples were subsequently refrigerated until the chemical analysis. Spiking assays with artificial and natural estuarine water were performed in order to test the accuracy of this method. Heavy metal recovery was always between 90–110% (Duarte et al. 2014).

Heavy metal concentrations were determined by Inductively Coupled Plasma-Atomic Emission Spectrometer [ICP-AES, Horiba Jobin-Yvon, France, Ultima, model equipped with a 40.68 MHz RF generator, Czerny-Turner monochromator with 1.00 m (sequential), autosampler AS500 and CMA (Concomitant Metals Analyzer)]. Blanks were always below the detection limits of ICP-AES (0.33 ppm for Zn and 0.03 ppm for Cu). The accuracy and efficiency of the results were checked by processing reference material CRM 145 and CRM 146, referent to sewage sludge from domestic origin and sludge, respectively. Trace metal concentrations in the reference materials determined by ICP-AES were not statistically different from their certified ones (t-student tests $p = 0.05$), indicating an accurate and total extraction of the heavy metals present in the samples.

Data analyses

Data on fouling abundance (% cover) was integrated with the “health status” visual assessment by multiplying the % cover value of each species with their correspondent “health status” factor; namely 0.1, 0.33, 0.66 and 0.99 factors, related to values “1”, “2”, “3” and “4” respectively. This integration was applied to limit the observed high % cover of species which, despite dead, maintained their calcareous structure on the plate (e.g., barnacles, encrusting bryozoans and serpulids). In all data analyses, NIS status was attributed only to those species with a confirmed non-indigenous category and verified by specific scientific literature (e.g., Canning-Clode et al. 2013; Chainho et al. 2015; Gestoso et al. 2017). Following a more conservative approach (Marchini et al. 2015) cryptogenic

species (i.e. unspecified origin, that is, unknown whether native or introduced; Carlton 1996) were included within the native species category.

Multivariate analyses were performed on the total fouling community structure, whereas univariate analyses were carried out considering the total % cover and species richness on i) the totality of species (total % cover and richness), ii) native/cryptogenic (native % cover and richness) and iii) NIS components (NIS % cover and richness).

First, different one-way permutational analyses of variance (PERMANOVAs) on Time_0 data were performed on the selected variables to compare the initial fouling communities from inside versus outside the marina, with “Area” as a fixed factor (2 levels: inside and outside; fixed, $n = 12$). A non-metric multidimensional scaling (nMDS) analysis was also used to graphically visualize differences detected between the selected communities. When significant, the structure of the two fouling communities was analyzed through a one-way SIMPER test (Similarity Percentages) in order to assess the main average dissimilarity between areas, and also which species mostly contribute to the similarity within the two areas (with a cut-off level of 90%; only species with a percentage contribution $\geq 2\%$ were reported; Clarke 1993).

As plates were repeatedly sampled, with the aim to assess the potential recovery of the fouling communities, uni- and multivariate PERMANOVA analyses were performed taking into account data from Time_D to Time_2 separately. The use of this approach was preferred to account for non-independence of samples. The design employed for these analyses included two orthogonal factors: “Disturbance” (4 levels: C, Phy, Che and Mix; fixed, orthogonal) and “Area” (2 levels: inside and outside; fixed, orthogonal). When significant differences among factors or their interactions were observed, post-hoc pairwise tests were performed. An additional nMDS was constructed to graphically visualize multivariate patterns of variation in two dimensions, showing the different responses of the communities to the disturbances.

All multivariate tests were run from Bray-Curtis similarity matrices, while for univariate analyses, the Euclidean distances matrices were preferred, using an approach similar to parametric ANOVA (Airoldi et al. 2015). P-values related to the pseudo-F ratios were calculated with 9999 random permutations (Underwood 1997). Whenever there were not enough possible permutations to get a reasonable test, the Monte Carlo p-values were used instead. Prior to analysis, PERMDISP was used to check data for heterogeneity of dispersions, and transformations were applied where necessary.

In order to identify those species with higher resistance capacity to the manipulative experiment, a SIMPER analysis was performed on live species from Time_2, i.e. considering only that ones with “health status” value from “2” to “4”. Statistical analyses were performed using the software

Table 1. Zinc (Zn) and copper (Cu) concentration ($\mu\text{g/l}$) in the water samples collected per condition (average values \pm SD, $n = 3$).

	C	Phy	Che	Mix
Zn	825.56 \pm 386.57	733.53 \pm 187.15	1322.90 \pm 6.37	1386.70 \pm 446.70
Cu	27.39 \pm 5.54	19.03 \pm 1.13	34.31 \pm 2.85	36.82 \pm 3.12

PRIMER version 6.1.13 (Clarke and Gorley 2006) with the add-on PERMANOVA + version 1.0.3 (Anderson 2005).

Results

Chemical-physical and heavy metal characterization

The chemical-physical parameters showed similar values between inside and outside the marina (in the field), and during all the experimental phases (both in the mesocosm tanks and in the containers used during Time_D; Table S1). The water samples collected in the containers for each condition, showed higher average concentrations (\pm SD, $\mu\text{g/l}$) of Cu (35.57 \pm 3.00) and Zn (1354.81 \pm 284.70) in samples treated with the antifouling paint (chemical and mixed disturbances), compared to the untreated treatments (control and physical disturbances): 23.16 \pm 5.77 (Cu) and 779.55 \pm 276.27 (Zn). Average values per condition (\pm SD) are presented in Table 1.

Initial structure of the fouling community inside and outside the marina (Time_0)

After 4 months of colonization, fouling communities inside and outside the marina were composed of a total of 38 and 43 species, respectively (average values \pm SD, 14.7 \pm 2.4 and 18.2 \pm 5.2 species), including 11 NIS inside and 9 NIS outside the marina (4.5 \pm 1.3 and 3.7 \pm 1.4 NIS) (Table 2; Figure S2). The main taxonomic groups observed on PVC plates inside the marina were ascidians (26%) and bryozoans (21%) while the outside community was dominated by bryozoans (41%).

At the local scale, total % cover and richness displayed significantly higher values outside the marina [PERMANOVAs: Pseudo- $F_{1,22} = 5.286$, $P(\text{perm}) = 0.031$, PERMDISP- $P(\text{perm}) = 0.365$; Pseudo- $F_{1,22} = 4.429$, $P(\text{perm}) = 0.049$, PERMDISP- $P(\text{perm}) = 0.057$ (Fourth root transformation), respectively], as well as for native species % cover and richness [PERMANOVAs: Pseudo- $F_{1,22} = 59.716$, $P(\text{perm}) = 0.0001$, PERMDISP- $P(\text{perm}) = 0.733$; Pseudo- $F_{1,22} = 7.283$, $P(\text{perm}) = 0.006$, PERMDISP- $P(\text{perm}) = 0.059$ (Square root transformation), respectively]. In contrast, the NIS % cover was significantly higher inside the marina [Pseudo- $F_{1,22} = 31.799$, $P(\text{perm}) = 0.0001$, PERMDISP- $P(\text{perm}) = 0.332$], while no difference was observed in NIS richness between the two areas [Pseudo- $F_{1,22} = 0.538$, $P(\text{MC}) = 0.4683$, PERMDISP- $P(\text{perm}) = 0.294$].

Table 2. Non-indigenous species recorded in the areas investigated (inside and outside the marina of Quinta do Lorde) at Time_0

Non-indigenous species	Potential native origin	References	Recorded inside the marina	Recorded outside the marina
Porifera				
<i>Paraleucilla magna</i> Klautau, Monteiro and Borojevic, 2004	Brazil	Klautau et al. 2004; Canning-Clode et al. 2013	•	•
Cnidaria				
<i>Ectopleura crocea</i> (Agassiz, 1862)	North-West Atlantic Ocean	Mills et al. 2007; Wirtz 2007		•
<i>Exaiptasia diaphana</i> (Rapp, 1829)	Mediterranean Sea	Riedl 1991; Canning-Clode et al. 2013	•	•
Polychaeta				
<i>Branchiommata bairdi</i> (McIntosh, 1885)	Gulf of Mexico	McIntosh 1885; Ramalhosa et al. 2014	•	
Crustacea: Cirripeda				
<i>Balanus trigonus</i> Darwin, 1854	Pacific Ocean	Carlton et al. 2011; Pilsbry 1916		•
Bryozoa				
<i>Beania maxilladentata</i> Ramalho, Muricy and Taylor, 2010	Brazil	Ramalho et al. 2008; Souto et al. 2015		•
<i>Celleporaria inaudita</i> Tilbrook, Hayward and Gordon, 2001	Red Sea – Indo-Pacific Ocean	Canning-Clode et al. 2013; Souto et al. 2018	•	
<i>Cradoscrupocellaria bertholletii</i> (Audouin, 1826)	Unknown	Canning-Clode et al. 2013; Vieira et al. 2013	•	
<i>Parasmittina alba</i> Ramalho, Muricy and Taylor, 2011	Brazil	Ramalho et al. 2011; Souto et al. 2018	•	•
<i>Schizoporella pungens</i> (Canu and Bassler, 1928)	Gulf of Mexico	Winston and Maturro 2009; Canning-Clode et al. 2013	•	•
<i>Tricellaria inopinata</i> d’Hondt and Occhipinti Ambrogi, 1985	North-East Pacific Ocean?	Marchini et al. 2007; Cook et al. 2013; Ramalhosa et al. 2019		•
Ascidiacea				
<i>Botrylloides niger</i> Herdman, 1886	Indian Ocean	Millar 1988; Gestoso et al. 2017	•	
<i>Botrylloides violaceus</i> Oka, 1927	North-West Pacific	Oka 1927; Canning-Clode et al. 2013	•	
<i>Botryllus schlosseri</i> (Pallas, 1766)	Unknown	Van Name 1945; Canning-Clode et al. 2008	•	
<i>Distaplia corolla</i> Monniot F., 1974	Caribbean	Monniot 1983; Canning-Clode et al. 2013	•	•

The two fouling communities at Time_0 revealed a significant difference in their structure (PERMANOVA: Pseudo- $F_{1,22} = 26.77$, $P(\text{perm}) = 0.0001$, PERMDISP- $P(\text{perm}) = 0.9833$; Figure 3A), also confirmed by an average dissimilarity of 82.87% (SIMPER analysis). The taxonomic groups that mostly contributed to the similarity of the community inside the marina were *Parasmittina alba* Ramalho, Muricy and Taylor, 2011 (44.36%) and spirorbids (20.59%); while outside the marina were *Spirobranchus triqueter* (Linnaeus, 1758) (37.07%), *Lithophyllum incrustans* Philippi, 1837 (11.46%) and *Puellina radiata* (Moll, 1803) (11.18%; Table S2).

Mesocosm experiment: effects of the disturbances on the fouling communities

Uni- and multivariate analyses of total richness in Time_1,2, of native richness in Time_1 and of fouling community structure along all times (Time_D-2) detected a significant interaction between the factors “Area” and “Disturbance”, revealing how the effect of the disturbance was dependent




Figure 3. Non-metric multidimensional scaling (nMDS) graphs on fouling communities inside and outside the marina at (A) Time_0 and (B) considering the sampling times from Time_D to Time_2 per each type of disturbance. Filled and empty symbols correspond to fouling community plates placed inside and outside the marina respectively.

on the area (Table S3). Overall, fouling communities from outside the marina maintained greater values of total and native richness than those communities from the inside (Table S4, Figure 4). This also reflects the different abundances of native and NIS in the two communities (Figure 4). The PERMANOVA performed with the other variables (i.e. total, native and NIS cover; and NIS richness) revealed significant effects of the main factor “Disturbance” and/or “Area” (Table S3). A conspicuous decline in total, native and NIS % cover was observed in both communities. Outside the marina, native species recorded higher values of % cover for all tested conditions in comparison with the communities inside the marina; while an opposite trend was observed for NIS % cover, which had higher values inside than outside the marina (Table S3; Figure 4).

Different responses to disturbances were observed between the two communities (Table S4 in Time_D-2) but, generally, while chemical or mixed




Figure 4. Average number of percent cover and richness, with corresponding SD, on total species (white bars), native/cryptogenic species (grey bars) and non-indigenous species (black bars) divided per experimental times, from inside and outside the marina.

disturbances significantly modified the structure of the communities, the exposure to physical disturbance did not differ from the control plates in both cases (Table S4). Some exceptions from this general trend were observed in the communities outside the marina, where no significant differences were recorded among the different disturbances on the structure of the fouling communities, except for mixed disturbance in Time_1 (Table S4); while cases of significant differences were observed between control and physical disturbances, when considering the total and native richness, due to a slight increase of these values in control from Time_D to Time_1 (Table S4; Figure 4). The nMDS clearly showed the lower effect of the physical disturbance on the fouling communities, while the chemical and mixed disturbances appeared separated from control and physical replicates (Figure 3B).

When considering the live biota from Time_2, algae, bryozoans, sponges and anthozoans were the dominant groups in both fouling communities. Inside the marina, the non-indigenous bryozoan *Celleporaria inaudita* Tilbrook, Hayward and Gordon, 2001 was the species most resistant to chemical disturbance, while algae (*L. incrustans*, *Nemoderma* sp. and Polysiphoniae) and sponges (*Mycale* sp. and *Sycon* sp.) mostly contributed to the average similarities of all the other conditions, followed by the NIS *Exaiptasia diaphana* (Rapp, 1829) and *P. alba* (the latter only in the control;




Figure 5. Average number of live/dead percent cover and richness, with corresponding SD, on total species (white bars), native/cryptogenic species (grey bars) and non-indigenous species (black bars) on Time_2, from inside and outside the marina.

Table S5). With regards to plates from outside the marina, the species that mostly contributed to the similarity within different conditions were *E. diaphana*, *Favosipora purpurea* Souto, Kaufmann and Canning-Clode, 2015, *L. incrustans* and *Nemoderma* sp., followed by sponges (e.g., *Mycale* sp., *Sycon* sp.; Table S5). Finally, a decrease of *P. alba* and other encrusting bryozoans (except for *F. purpurea*), spirorbids and serpulids was observed on fouling communities from both areas affected by physical disturbance, together with a slight increase of sponges and *E. diaphana* (Table S5). The species that were most resistant to the experimental conditions (“health status” from 2 to 4) were natives, with higher values observed outside the marina (Figure 5).

The complete species list per each PVC panel on Time_0, and from Time_D to Time_2 with the relative “health status” assessment for each species, are shown in Table S6 and S7 respectively.

Discussion and conclusions

The present study is a pioneering manipulative experiment to assess the resistance of fouling communities colonizing PVC plates from different areas (namely inside and outside a marina) to selected anthropogenic disturbances. Initially, a comparable number of NIS was recorded in both communities, but the PVC plates from inside the marina showed a dominant cover of NIS, in contrast with a higher native species cover and richness in the plates from the outside. Generally, both fouling communities were strongly affected by the chemical and mixed disturbances, with a small or null effect of the physical disturbance. Despite this general trend, fouling communities outside the marina displayed some exceptions, perhaps due to a slight increase of total/native species richness in the plates associated to control condition. In fact, during the experiment, the random placement of all plates in the tanks could have facilitated new recruitments. Furthermore, differences observed under disturbances between the two fouling communities may depend on their structures, confirming the potential of artificial habitats to modify ecological responses to environmental disturbances (e.g., Connell and Glasby 1999; Bulleri and Chapman 2004; Goodsell et al. 2007; Airoidi et al. 2015), as well as facilitate the settlement, establishment and further spread of NIS (Corriero et al. 2015).

The initial fouling communities experienced a sharp decrease of their most abundant species when treated with the antifouling paint; e.g., the NIS *Parasmittina alba* inside the marina, and the native species *Spirobranchus triqueter* and *Puellina radiata* from outside. Both natives and NIS were drastically reduced in presence and abundance, a trend observed in previous studies (Tait et al. 2018; Ramalhosa et al. 2019). Some species, however, demonstrated a certain degree of tolerance to the selected disturbances (e.g., the NIS *Celleporaria inaudita* and *Exaiptasia diaphana* inside marina; and the native *Favosipora purpurea* from outside). In contrast to previous studies (Clark and Johnston 2005; Piola and Johnston 2008; Piola et al. 2009; Crooks et al. 2011), a higher resistance of the non-indigenous community was not observed at the end of the experiment. As reported by Dafforn et al. (2009), native species also have the potential to survive and re-colonize the substrate after a disturbance event, demonstrating tolerance capabilities similar to NIS.

In this experiment, the impact of physical disturbance was not as strong as the chemical disturbance, but slight effects were recorded in terms of reduction of the hard and fragile encrusting bryozoans (apart to *F. purpurea*) and calcareous tubeworms, while soft and flexible taxa, like anthozoans and sponges, were more resistant (Clark and Johnston 2005). This trend was similar to results found by Clark and Johnston (2005) in a previous experiment using a similar approach, yet in our case, the pressure used was quite lower, i.e. 36 g/cm² vs. 140 g/cm². Other studies on fouling communities

manipulated with a physical disturbance were mostly focused on the removal of organisms from the substrata, with the aim to assess the colonization processes and not considering the resistance of fouling organisms to a simulated physical pressure (e.g., Bram et al. 2005; Valdivia et al. 2005; Cifuentes et al. 2007).

Contrary to our initial hypothesis for a higher expected tolerance and resistance to disturbances for NIS (i.e., inside the marina; Piola and Johnston 2008; Canning-Clode et al. 2011; Crooks et al. 2011), fouling communities from outside the marina showed a greater tolerance. Similar to Canning-Clode et al. (2011) and Ramalhosa et al. (2019), the native component exhibited slightly higher levels of resistance, with a higher amount of live native species in most conditions at the end of the experiment. In the same coastal region as the present study, Gestoso et al. (2017) demonstrated a certain degree of biotic resistance of fouling communities from marine protected areas to the NIS settlement rate in port habitats. Nevertheless, the resilient capacity of natural habitats as buffer areas surrounding marinas or ports, e.g., in reducing the stepping-stone of NIS, or the consequences of variation in recruitment across time, is still poorly studied (Glasby et al. 2007; Ruiz et al. 2009; Simkanin et al. 2017).

It is important to note that the experiment conducted here was limited to a single area, lasted 4 months and the manipulated communities were at an early stage of development. However, it represents a starting point in understanding the ecological consequences of anthropogenic disturbances on fouling communities. In summary, and in contrast with expectations, we found evidence that NIS did not play a key role in fouling communities against the selected disturbances. This highlights the need of future studies focused on the response of natural and artificial communities to combined anthropogenic disturbances, simulating long-term consequences and assessing possible solutions against the uninterrupted ocean sprawl phenomenon.

Acknowledgements

This work was based on the Short Term Scientific Mission (#38255) experience carried out by J. Ferrario and promoted by the COST Action CA-15121 “Advancing Marine Conservation in the European and Contiguous Seas - MarCons”. Authors are grateful to: Léa Riera, João Monteiro and Susanne Schäfer for their assistance during field work; Gustavo Martins for the statistical advice and April Monica Houghton Blakeslee for the English revision of the text. Finally, we thank two anonymous reviewers for their comments and suggestions that greatly improved the paper. This is contribution 45 from the Smithsonian’s MarineGEO Network.

Funding declaration

This work was partially funded by the MARE mini-grants 2017 project “Development of a new methodology for the study of bioinvasions on coastal communities: experimental test across insular and continental systems” and projects MIMAR (MAC/4.6.d/066), MIMAR+ (MAC2/4.6d/249) and PLASMAR (MAC/1.1a/030), in the framework of INTERREG MAC 2014-2020 Programme. I. Gestoso and E. Cacabelos were financially supported by postdoctoral grants from ARDITI (Regional Agency for Development of Research, Technology and Innovation of Madeira) (Projects 002458/2015/132 and M1420 -09 -5369 -FSE FSE-000001

respectively). P. Ramalhosa was partially funded by the Project Observatório Oceânico da Madeira-OOM (M1420-01-0145-FEDER-000001), co-financed by the Madeira Regional Operational Programme (Madeira 14-20), under the Portugal 2020 strategy, through the European Regional Development Fund (ERDF). B. Duarte investigation was supported by FCT throughout a Postdoctoral grant (SFRH/BPD/115162/2016). J. Canning-Clode was supported by a starting grant in the framework of the 2014 FCT Investigator Programme (IF/01606/2014/CP1230/CT0001). This study had the support of “Fundação para a Ciência e Tecnologia (FCT)”, through the strategic project UID/MAR/04292/2019 granted to MARE.

Authors' contribution

Conceptualization: JF, IG and JCC; Formal analyses: JF, IG, EC; Funding acquisition: JF, IG, JCC; Investigation: JF, IG, PR, EC; Methodology: JF, IG, PR, EC, BD, IC, JCC; Resources: JCC; Supervision: IG, JCC; Visualization: JF, IG; Writing: JF, IG, PR, EC, BD, IC, JCC.

Declaration of interest: none

Ethics and permits: none

References

- Airoldi L, Turon X, Perkol-Finkel S, Rius M (2015) Corridors for aliens but not for natives: effects of marine urban sprawl at a regional scale. *Diversity and Distributions* 21: 755–768, <https://doi.org/10.1111/ddi.12301>
- Altman S, Whitlatch RB (2007) Effects of small-scale disturbance on invasion success in marine communities. *Journal of Experimental Marine Biology and Ecology* 342: 15–29, <https://doi.org/10.1016/j.jembe.2006.10.011>
- Anderson MJ (2005) PERMANOVA: a FORTRAN computer program for permutational multivariate analysis of variance. University of Auckland, Auckland, New Zealand, 24 pp
- Bishop MJ, Mayer-Pinto M, Airoldi L, Firth LB, Morris RL, Loke LHL, Hawkins SJ, Naylor LA, Coleman RA, Chee SY, Dafforn KA (2017) Effects of ocean sprawl on ecological connectivity: impacts and solutions. *Journal of Experimental Marine Biology and Ecology* 492: 7–30, <https://doi.org/10.1016/j.jembe.2017.01.021>
- Bulleri F, Chapman MG (2004) Intertidal assemblages on artificial and natural habitats in marinas on the north-west coast of Italy. *Marine Biology* 145: 381–391, <https://doi.org/10.1007/s00227-004-1316-8>
- Bulleri F, Chapman MG (2010) The introduction of coastal infrastructure as a driver of change in marine environments. *Journal of Applied Ecology* 47: 26–35, <https://doi.org/10.1111/j.1365-2664.2009.01751.x>
- Bram JB, Page HM, Dugan JE (2005) Spatial and temporal variability in early successional patterns of an invertebrate assemblage at an offshore oil platform. *Journal of Experimental Marine Biology and Ecology* 317: 223–237, <https://doi.org/10.1016/j.jembe.2004.12.003>
- Canning-Clode J, Kaufmann M, Wahl M, Molis M, Lenz M (2008) Influence of disturbance and nutrient enrichment on early successional fouling communities in an oligotrophic marine system. *Marine Ecology-An Evolutionary Perspective* 29: 115–124, <https://doi.org/10.1111/j.1439-0485.2007.00210.x>
- Canning-Clode J, Bellou N, Kaufmann MJ, Wahl M (2009) Local-regional richness relationship in fouling assemblages-effects of succession. *Basic and Applied Ecology* 10: 745–753, <https://doi.org/10.1016/j.baae.2009.05.005>
- Canning-Clode J, Fofonoff P, Riedel GF, Torchin M, Ruiz GM (2011) The effects of copper pollution on fouling assemblage diversity: A tropical-temperate comparison. *PLoS ONE* 6: e18026, <https://doi.org/10.1371/journal.pone.0018026>
- Canning-Clode J, Fofonoff P, McCann L, Carlton JT, Ruiz G (2013) Marine invasions on a subtropical island: Fouling studies and new records in a recent marina on Madeira island (Eastern Atlantic Ocean). *Aquatic Invasions* 8: 261–270, <https://doi.org/10.3391/ai.2013.8.3.02>
- Carlton JT (1996) Biological invasions and cryptogenic species. *Ecology* 77: 1653–1655, <https://doi.org/10.2307/2265767>
- Carlton JT, Newman WA, Pitombo FP (2011) Barnacle invasions: introduced, cryptogenic, and range expanding Cirripedia of North and South America. In: Galil BS, Clark PF, Carlton JT (eds), In the wrong place-Alien marine crustaceans: distribution, biology and impacts. Springer-Verlag, Berlin, Germany, pp 159–213, https://doi.org/10.1007/978-94-007-0591-3_5
- Ceccherelli G, Pinna S, Cusseddu V, Bulleri F (2014) The role of disturbance in promoting the spread of the invasive seaweed *Caulerpa racemosa* in seagrass meadows. *Biological Invasions* 16: 2737–2745, <https://doi.org/10.1007/s10530-014-0700-7>
- Chainho P, Fernandes A, Amorim A, Ávila SP, Canning-Clode J, Castro JJ, Costa AC, Costa JL, Cruz T, Gollasch S, Grazziotin-Soares C, Melo R, Micael J, Parente MI, Semedo J, Silva T, Sobral D, Sousa M, Torres P, Veloso V, Costa MJ (2015) Non-indigenous species

- in Portuguese coastal areas, coastal lagoons, estuaries and islands. *Estuarine Coastal and Shelf Science* 167: 199–211, <https://doi.org/10.1016/j.ecss.2015.06.019>
- Chambers LD, Stokes KR, Walsh FC, Wood RJK (2006) Modern approaches to marine antifouling coatings. *Surface and Coatings Technology* 201: 3642–3652, <https://doi.org/10.1016/j.surfcoat.2006.08.129>
- Chapman MG, Bulleri F (2003) Intertidal seawalls - new features of landscape in intertidal environments. *Landscape and Urban Planning* 62: 159–172, [https://doi.org/10.1016/S0169-2046\(02\)00148-2](https://doi.org/10.1016/S0169-2046(02)00148-2)
- Cifuentes M, Kamlah C, Thiel M, Lenz M, Wahl M (2007) Effects of temporal variability of disturbance on the succession in marine fouling communities in northern-central Chile. *Journal of Experimental Marine Biology and Ecology* 352: 280–294, <https://doi.org/10.1016/j.jembe.2007.08.004>
- Clark GF, Johnston EL (2005) Manipulating larval supply in the field: a controlled study of marine invasibility. *Marine Ecology Progress Series* 298: 9–19, <https://doi.org/10.3354/meps298009>
- Clarke KR (1993) Non-parametric multivariate analyses of changes in community structure. *Australian Journal of Ecology* 18: 117–143, <https://doi.org/10.1111/j.1442-9993.1993.tb00438.x>
- Clarke KR, Gorley RN (2006) PRIMER v6: user manual/ tutorial, Primer-E. ed. Plymouth, 190 pp
- Comber SDW, Gardner MJ, Boxall ABA (2002) Survey of four marine antifoulant constituents (copper, zinc, diuron and Irgarol 1051) in two UK estuaries. *Journal of Environmental Monitoring* 4: 417–425, <https://doi.org/10.1039/b202019j>
- Connell SD, Glasby TM (1999) Do urban structures influence local abundance and diversity of subtidal epibiota? A case study from Sydney Harbour, Australia. *Marine Environmental Research* 47: 373–387, [https://doi.org/10.1016/S0141-1136\(98\)00126-3](https://doi.org/10.1016/S0141-1136(98)00126-3)
- Cook EJ, Stehlíková J, Beveridge CM, Burrows MT, De Blauwe H, Faasse M (2013) Distribution of the invasive bryozoan *Tricellaria inopinata* in Scotland and a review of its European expansion. *Aquatic Invasions* 8: 281–288, <https://doi.org/10.3391/ai.2013.8.3.04>
- Corriero G, Pierri C, Accoroni S, Alabiso G, Bavestrello G, Barbone E, Bastianini M, Bazzoni AM, Bernardi Aubry F, Boero F, Buia MC, Cabrini M, Camatti E, Cardone F, Cataletto B, Cattaneo Vietti R, Cecere E, Cibic T, Colangelo P, De Olazabal A, D’Onghia G, Finotto S, Fiore N, Fornasaro D, Frascchetti S, Gambi MC, Giangrande A, Gravili C, Guglielmo R, Longo C, Lorenti M, Lugliè A, Maiorano P, Mazzocchi MG, Mercurio M, Mastrototaro F, Mistri M, Monti M, Munari C, Musco L, Nonnis-Marzano C, Padedda BM, Patti FP, Petrocelli A, Piraino S, Portacci G, Pugnetti A, Pulina S, Romagnoli T, Rosati I, Sarno D, Satta CT, Sechi N, Schiapparelli S, Scipione B, Sion L, Terlizzi A, Tirelli V, Totti C, Tursi A, Ungaro N, Zingone A, Zupo V, Basset A (2015) Ecosystem vulnerability to alien and invasive species: a case study on marine habitats along the Italian coast. *Aquatic Conservation: Marine and Freshwater Ecosystems* 26: 392–409, <https://doi.org/10.1002/aqc.2550>
- Crooks JA, Chang AL, Ruiz GM (2011) Aquatic pollution increases the relative success of invasive species. *Biological Invasions* 13: 165–176, <https://doi.org/10.1007/s10530-010-9799-3>
- Dafforn KA, Glasby TM, Johnston EL (2008) Differential effects of tributyltin and copper antifoulants on recruitment of non-indigenous species. *Biofouling* 24: 23–33, <https://doi.org/10.1080/08927010701730329>
- Dafforn KA, Glasby TM, Johnston EL (2009) Links between estuarine condition and spatial distributions of marine invaders. *Diversity and Distributions* 15: 807–821, <https://doi.org/10.1111/j.1472-4642.2009.00587.x>
- Dafforn KA, Glasby TM, Airoidi L, Rivero NK, Mayer-Pinto M, Johnston EL (2015) Marine urbanization: an ecological framework for designing multifunctional artificial structures. *Frontiers in Ecology and the Environment* 13: 82–90, <https://doi.org/10.1890/140050>
- Davenport J, Davenport JL (2006) The impact of tourism and personal leisure transport on coastal environments: A review. *Estuarine Coastal and Shelf Science* 67: 280–292, <https://doi.org/10.1016/j.ecss.2005.11.026>
- Davidson IC, Zabin CJ, Chang AL, Brown CW, Sytsma MD, Ruiz GM (2010) Recreational boats as potential vectors of marine organisms at an invasion hotspot. *Aquatic Biology* 11: 179–191, <https://doi.org/10.3354/ab00302>
- Duarte B, Silva G, Lino Costa J, Medeiros JP, Azeda C, Sá E, Metelo I, Costa MJ, Caçador I (2014) Heavy metal distribution and partitioning in the vicinity of the discharge areas of Lisbon drainage basins (Tagus Estuary, Portugal). *Journal of Sea Research* 93: 101–111, <https://doi.org/10.1016/j.seares.2014.01.003>
- Elmqvist T, Zipperer WC, Güneralp B (2016) Urbanization, habitat loss and biodiversity decline. Solution pathways to break the cycle. In: Seta K, Solecki WD, Griffith CA (eds), *Routledge Handbook of Urbanization and Global Environmental Change*. London, New York, pp 139–151
- Ferrario J, Caronni S, Occhipinti-Ambrogi A, Marchini A (2017) Role of commercial harbours and recreational marinas in the spread of non-indigenous fouling species. *Biofouling* 33: 651–660, <https://doi.org/10.1080/08927014.2017.1351958>
- Firth LB, Thompson RC, White FJ, Schofield M, Skov MW, Hoggart SP, Jackson J, Knights AM, Hawkins SJ (2013) The importance of water-retaining features for biodiversity on

- artificial intertidal coastal defence structures. *Diversity and Distributions* 19: 1275–1283, <https://doi.org/10.1111/ddi.12079>
- Firth LB, Knights AM, Bridger D, Evans AJ, Mieszkowska N, Moore PJ, O'Connor NE, Sheehan EV, Thompson RC, Hawkins SJ (2016) Ocean Sprawl: challenges and opportunities for biodiversity management in a changing world. *Oceanography and Marine Biology - An Annual Review* 54: 193–269, <https://doi.org/10.1201/9781315368597-5>
- Garrabou J, Sala E, Arcas A, Zabala M (1998) The impact of diving on rocky sublittoral communities: a case study of a bryozoan population. *Conservation Biology* 12: 302–312, <https://doi.org/10.1046/j.1523-1739.1998.96417.x>
- Gestoso I, Ramalhosa P, Oliveira P, Canning-Clode J (2017) Marine protected communities against biological invasions: A case study from an offshore island. *Marine Pollution Bulletin* 119: 72–80, <https://doi.org/10.1016/j.marpolbul.2017.03.017>
- Glasby TM, Connell SD (1999) Urban Structures as Marine Habitats. *Ambio* 28: 595–598
- Glasby TM, Underwood AJ (1996) Sampling to differentiate between pulse and press perturbations. *Environmental Monitoring and Assessment* 42: 241–252, <https://doi.org/10.1007/BF00414371>
- Glasby TM, Connell SD, Holloway MG, Hewitt CL (2007) Nonindigenous biota on artificial structures: Could habitat creation facilitate biological invasions? *Marine Biology* 151: 887–895, <https://doi.org/10.1007/s00227-006-0552-5>
- Goodsell PJ, Chapman MG, Underwood AJ (2007) Differences between biota in anthropogenically fragmented habitats and in naturally patchy habitats. *Marine Ecology Progress Series* 351: 15–23, <https://doi.org/10.3354/meps07144>
- Guardiola FA, Cuesta A, Meseguer J, Esteban MA (2012) Risks of using antifouling biocides in aquaculture. *International Journal of Molecular Sciences* 13: 1541–1560, <https://doi.org/10.3390/ijms13021541>
- Hall LW, Scott MC, Killen WD (1998) Ecological risk assessment of copper and cadmium in surface waters of Chesapeake Bay watershed. *Environmental Toxicology and Chemistry* 17: 1172–1189, <https://doi.org/10.1002/etc.5620170626>
- Hayward PJ, Ryland JS (1995) Handbook of the marine fauna of North-West Europe. Oxford University Press, UK, 800 pp
- Hewitt CL, Gollasch S, Minchin D (2009) The vessel as a vector - biofouling, ballast water and sediments. In: Rilov G, Crooks JA (eds), *Biological Invasions in Marine Ecosystems*. Berlin, pp 117–131, https://doi.org/10.1007/978-3-540-79236-9_6
- Hudson JH, Goodwin WB (2001) Assessment of vessel grounding injury to coral reef and seagrass habitats in the Florida Keys National Marine Sanctuary, Florida: protocol and methods. *Bulletin of Marine Science* 69: 509–516
- Huff TM (2011) Effects of human trampling on macro- and meiofauna communities associated with intertidal algal turfs and implications for management of protected areas on rocky shores (Southern California). *Marine Ecology* 32: 335–345, <https://doi.org/10.1111/j.1439-0485.2011.00467.x>
- Hulme PE (2009) Trade, transport and trouble: managing invasive species pathways in an era of globalization. *Journal of Applied Ecology* 46: 10–18, <https://doi.org/10.1111/j.1365-2664.2008.01600.x>
- Karlson RH, Osman RW (2012) Species composition and geographic distribution of invertebrates in fouling communities along the east coast of the USA: A regional perspective. *Marine Ecology Progress Series* 458: 255–268, <https://doi.org/10.3354/meps09767>
- Kenworthy JM, Rolland G, Samadi S, Lejeune C (2018) Local variation within marinas: Effects of pollutants and implications for invasive species. *Marine Pollution Bulletin* 133: 96–106, <https://doi.org/10.1016/j.marpolbul.2018.05.001>
- Kinsella CM, Crowe TP (2016) Separate and combined effects of copper and freshwater on the biodiversity and functioning of fouling assemblages. *Marine Pollution Bulletin* 107: 136–143, <https://doi.org/10.1016/j.marpolbul.2016.04.008>
- Klautau M, Monteiro L, Borojevic R (2004) First occurrence of the genus *Paraleucilla* (Calcarea, Porifera) in the Atlantic Ocean: *P. magna* sp. nov. *Zootaxa* 710: 1–8, <https://doi.org/10.11646/zootaxa.710.1.1>
- Kohler KE, Gill SM (2006) Coral Point Count with Excel extensions (CPCe): a Visual Basic program for the determination of coral and substrate coverage using random point count methodology. *Computers & Geosciences* 32: 1259–1269, <https://doi.org/10.1016/j.cageo.2005.11.009>
- Lewis JA (1998) Marine biofouling and its prevention on underwater surfaces. *Materials Forum* 22: 41–61
- Lockwood JL, Hoopes MF, Marchetti MP (2007) *Invasion ecology*. Wiley-Blackwell, Malden, USA, 304 pp
- Lutz SJ (2006) A thousand cuts? An assessment of small-boat grounding damage to shallow corals of the Florida Keys. In: Precht WF (ed), *Coral Reef Restoration Handbook*. CRC Press, Boca Raton, pp 25–38, <https://doi.org/10.1201/9781420003796.ch2>
- Marchini A, Cunha MR, Occhipinti-Ambrogi A (2007) First observations on bryozoans and entoprocts in the Ria de Aveiro (NW Portugal) including the first record of the Pacific

- invasive cheilostome *Tricellaria inopinata*. *Marine Ecology* 28: 154–160, <https://doi.org/10.1111/j.1439-0485.2007.00173.x>
- Marchini A, Galil BS, Occhipinti-Ambrogi A (2015) Recommendations on standardizing lists of marine alien species: lessons from the Mediterranean Sea. *Marine Pollution Bulletin* 101: 267–273, <https://doi.org/10.1016/j.marpolbul.2015.09.054>
- Marraffini ML, Ashton GV, Brown CW, Chang AL, Ruiz GM (2017) Settlement plates as monitoring devices for non-indigenous species in marine fouling communities. *Management of Biological Invasions* 8: 559–566, <https://doi.org/10.3391/mbi.2017.8.4.11>
- McIntosh WC (1885) Report on the Annelida Polychaeta collected by H. M. S. Challenger during the years 1873–76, *Report of the scientific results of the voyage of H. M. S. Challenger London, Zoology* 12: 1–554, <https://biodiversitylibrary.org/page/50688426>
- McKinney ML, Lockwood JL (1999) Biotic homogenization: a few winners replacing many losers in the next mass extinction. *Trends in Ecology & Evolution* 14: 450–453, [https://doi.org/10.1016/S0169-5347\(99\)01679-1](https://doi.org/10.1016/S0169-5347(99)01679-1)
- Mendez MM, Livore JP, Calcagno JA, Bigatti G (2017) Effects of recreational activities on Patagonian rocky shores. *Marine Environmental Research* 130: 213–220, <https://doi.org/10.1016/j.marenvres.2017.07.023>
- Milazzo M, Chemello R, Badalamenti F, Camarda R, Riggio S (2002) The impact of human recreational activities in marine protected areas: what lessons should be learnt in the Mediterranean Sea? *Marine Ecology* 23: 280–290, <https://doi.org/10.1111/j.1439-0485.2002.tb00026.x>
- Millar RH (1988) Ascidiens collected during the International Indian Ocean Expedition. *Journal of Natural History* 22: 823–848, <https://doi.org/10.1080/00222938800770541>
- Mills CE, Marques AC, Migotto AE, Calder DR, Hand C, Rees JTS, Haddock HDC, Dunn W, Pugh PR (2007) Hydrozoa: Polyps, Hydromedusae, and Siphonophora. In: Carlton JT (ed), *The Light and Smith manual: intertidal invertebrates from central California to Oregon*. University of California Press, Berkeley, pp 118–167
- Minchin D, Floerl O, Savini D, Occhipinti-Ambrogi A (2006) Small craft and the spread of exotic species. In: John D, Davenport JL (eds), *The Ecology of Transportation: Managing Mobility for the Environment*. Berlin, pp 99–118, https://doi.org/10.1007/1-4020-4504-2_6
- Mineur F, Cook EJ, Minchin D, Bohn K, Macleod A, Maggs CA (2012) Changing Coasts: Marine aliens and artificial structures. *Oceanography and Marine Biology - An Annual Review* 50: 189–234, <https://doi.org/10.1201/b12157-5>
- Monniot F (1983) Ascidiens littorales de Guadeloupe 5. Polycitoridae. *Bulletin du Muséum National D'histoire Naturelle, Paris, 4 sér., section A* 4: 999–1019
- Moschella PS, Abbiati M, Åberg P, Airoidi L, Anderson JM, Bacchiocchi F, Bulleri F, Dinesen GE, Frost M, Gacia E, Granhag L, Jansson PR, Satta MP, Sundelöf A, Thompson RC, Hawkins SJ (2005) Low-crested coastal defence structures as artificial habitats for marine life: using ecological criteria in design. *Coastal Engineering* 52: 1053–1071, <https://doi.org/10.1016/j.coastaleng.2005.09.014>
- Neira C, Levin LA, Mendoza G, Zirino A (2014) Alteration of benthic communities associated with copper contamination linked to boat moorings. *Marine Ecology* 35: 46–66, <https://doi.org/10.1111/maec.12054>
- Occhipinti-Ambrogi A, Savini D (2003) Biological invasions as a component of global change in stressed marine ecosystems. *Marine Pollution Bulletin* 46: 542–551, [https://doi.org/10.1016/S0025-326X\(02\)00363-6](https://doi.org/10.1016/S0025-326X(02)00363-6)
- Oka A (1927) Zur kenntnis der japanischen Botryllidae (vorläufige Mitteilung). *Proceedings of the Imperial Academy Japan* 3: 607–609, <https://doi.org/10.2183/pjab1912.3.607>
- Pilsbry HA (1916) The sessile barnacles contained in the collections of the U.S. National Museum: including a monograph of the American species. *Bulletin United States National Museum* 93: 1–366, <https://doi.org/10.5479/si.03629236.93.1>
- Piola RF, Johnston EL (2008) Pollution reduces native diversity and increases invader dominance in marine hard-substrate communities. *Diversity and Distributions* 14: 329–342, <https://doi.org/10.1111/j.1472-4642.2007.00430.x>
- Piola RF, Johnston EL (2009) Comparing differential tolerance of native and non-indigenous marine species to metal pollution using novel assay techniques. *Environmental Pollution* 157: 2853–2864, <https://doi.org/10.1016/j.envpol.2009.04.007>
- Piola RF, Dafforn KA, Johnson EL (2009) The influence of antifouling practices on marine invasions. *Biofouling* 25: 633–644, <https://doi.org/10.1080/08927010903063065>
- Ramalho LV, Muricy G, Taylor PD (2008) Taxonomy of *Beania* Johnston, 1840 (Bryozoa, Flustrina) from Arrial do Cabo, Rio de Janeiro State, Brazil. *Arquivos do Museu Nacional* 66: 499–508
- Ramalho LV, Muricy G, Taylor PD (2011) Taxonomic revision of some lepraliomorph cheilostome bryozoans (Bryozoa: Lepraliomorpha) from Rio De Janeiro State, Brazil. *Journal of Natural History* 45: 767–798, <https://doi.org/10.1080/00222933.2010.535917>
- Ramalhosa P, Camacho-Cruz K, Bastida-Zavala R, Canning-Clode J (2014) First record of *Branchiomma bairdi* McIntosh, 1885 (Annelida: Sabellidae) from Madeiran Island, Portugal

- (northeastern Atlantic Ocean). *BioInvasions Records* 3: 235–239, <https://doi.org/10.3391/bir.2014.3.4.04>
- Ramalhosa P, Gestoso I, Duarte B, Caçador I, Canning-Clode J (2019) Metal pollution affects native and non-indigenous biofouling recruitment in a subtropical island system. *Marine Pollution Bulletin* 141: 373–386, <https://doi.org/10.1016/j.marpolbul.2019.02.072>
- Renfro B, Chadwick NE (2017) Benthic community structure on coral reefs exposed to intensive recreational snorkeling. *PloS ONE* 12: e0184175, <https://doi.org/10.1371/journal.pone.0184175>
- Riedl R (1991) Flora e fauna del Mediterraneo. Franco Muzzio Editore, 777 pp
- Ruiz GM, Freestone AL, Fofonoff PW, Simkanin C (2009) Habitat distribution and heterogeneity in marine invasion dynamics: the importance of hard substrate and artificial structure. In: Wahl M (ed), *Marine hard bottom communities*. Springer, Berlin, Heidelberg, pp 321–332, https://doi.org/10.1007/b76710_23
- Seebens H, Gastner MT, Blasius B (2013) The risk of marine bioinvasion caused by global shipping. *Ecology Letters* 16: 782–790, <https://doi.org/10.1111/ele.12111>
- Schiff K, Diehl D, Valkirs A (2004) Copper emissions from antifouling paint on recreational vessels. *Marine Pollution Bulletin* 48: 371–377, <https://doi.org/10.1016/j.marpolbul.2003.08.016>
- Simkanin C, Davidson IC, Therriault TW, Jamieson G, Dower JF (2017) Manipulating propagule pressure to test the invasibility of subtidal marine habitats. *Biological Invasions* 19: 1565–1575, <https://doi.org/10.1007/s10530-017-1379-3>
- Simpson TJS, Smale DA, McDonald JI, Wernberg T (2017) Large scale variability in the structure of sessile invertebrate assemblages in artificial habitats reveals the importance of local-scale processes. *Journal of Experimental Marine Biology and Ecology* 494: 10–19, <https://doi.org/10.1016/j.jembe.2017.05.003>
- Souto J, Kaufmann MJ, Canning-Clode J (2015) New species and new records of bryozoans from shallow waters of Madeira Island. *Zootaxa* 3925: 581–593, <https://doi.org/10.11646/zootaxa.3925.4.7>
- Souto J, Ramalhosa P, Canning-Clode J (2018) Three non-indigenous species from Madeira harbors, including a new species of *Parasmittina* (Bryozoa). *Marine Biodiversity* 48: 977–986, <https://doi.org/10.1007/s12526-016-0592-0>
- Svensson JR, Lindegarth M, Siccha M, Lenz M, Molis M, Wahl M, Pavia H (2007) Maximum species richness at intermediate intensities of disturbance: consistency among levels of productivity. *Ecology* 88: 830–838, <https://doi.org/10.1890/06-0976>
- Tait L, Inglis G, Seaward K (2018) Enhancing passive sampling tools for detecting marine bioinvasions. *Marine Pollution Bulletin* 128: 41–50, <https://doi.org/10.1016/j.marpolbul.2018.01.015>
- Thomas KV, Brooks S (2010) The environmental fate and effects of antifouling paint biocides. *Biofouling* 26: 73–88, <https://doi.org/10.1080/08927010903216564>
- Underwood AJ (1997) *Experiments in ecology: their logical design and interpretation using analysis of variance*. Cambridge University Press, Cambridge, UK, 657 pp, <https://doi.org/10.1017/CBO9780511806407>
- Uyà M, Bulleri F, Gribben PE (2018) Propagules are not all equal: traits of vegetative fragments and disturbance regulate invasion success. *Ecology* 99: 957–965, <https://doi.org/10.1002/ecs.2168>
- Valdivia N, Heidemann A, Thiel M, Molis M, Wahl M (2005) Effects of disturbance on the diversity of hard-bottom macrobenthic communities on the coast of Chile. *Marine Ecology Progress Series* 299: 45–54, <https://doi.org/10.3354/meps299045>
- Van Name WG (1945) The North and South American Ascidiaceans. *Bulletin of the American Museum of Natural History* 84: 476 pp
- Vieira LM, Spencer Jones ME, Winston JE (2013) *Cradoscrupocellaria*, a new bryozoan genus for *Scrupocellaria bertholletii* (Audouin) and related species (Cheilostomata, Candidae): taxonomy, biodiversity and distribution. *Zootaxa* 3707: 1–63, <https://doi.org/10.11646/zootaxa.3707.1.1>
- Warnken J, Dunn RJK, Teasdale PR (2004) Investigation of recreational boats as a source of copper at anchorage sites using time-integrated diffusive gradients in thin film and sediment measurements. *Marine Pollution Bulletin* 49: 833–843, <https://doi.org/10.1016/j.marpolbul.2004.06.012>
- Watermann BT, Daehne B, Sievers S, Dannenberg R, Overbeke JC, Klijnstra JW, Heemken O (2005) Bioassays and selected chemical analysis of biocide-free antifouling coatings. *Chemosphere* 60: 1530–1541, <https://doi.org/10.1016/j.chemosphere.2005.02.066>
- Winston JE, Maturo Jr. FJ (2009) Bryozoans (Ectoprocta) of the Gulf of Mexico. In: Felder DL, Camp DK (eds), *Gulf of Mexico-Origins, Waters, and Biota*. Biodiversity. Texas A&M Press, College Station, Texas, pp 1147–1164
- Wirtz P (2007) On a collection of hydroids (Cnidaria, Hydrozoa) from the Madeira archipelago. *Arquipélago Ciências Biológicas e Marinhas* 24: 11–16
- Wisley B (1962) Effect of an antifouling paint on a bryozoan larva. *Nature, London* 193: 543–544, <https://doi.org/10.1038/193543a0>

Supplementary material

The following supplementary material is available for this article:

Figure S1. Pictures of the experimental units deployed inside (A) and outside (B) the marina.

Figure S2. Pictures of two PVC panels from inside (A) and outside (B) the marina at Time_0.

Table S1. Dissolved oxygen, pH, salinity and temperature values assessed during i) the sampling period, both inside and outside the marina; ii) the experiment period, in the two tanks and iii) the treatment period, in the buckets.

Table S2. Species or taxonomic group average abundances and percentage contributions to similarity within each area derived from SIMPER analysis at Time_0.

Table S3. Non-parametric multivariate and univariate PERMANOVA analyses examining variability in fouling community structure, total percent cover, total richness, native percent cover, native richness, NIS percent cover and NIS richness for the different sampling times.

Table S4. Pairwise tests on significant factors from Table S3 PERMANOVA analyses: A, B) on factor “Area x Disturbance” for pair of levels of “Treatment” and “Area”, C) on factor “Disturbance” and D) on “Area” for the different sampling times.

Table S5. Average abundances and percentage contribution to similarity resulting from SIMPER analyses of fouling communities within the four treatments over sampling Time_2.

Table S6. Species list on sampling Time_0 with relative percent cover.

Table S7. Species list from sampling Time_D-2 with relative percent cover and the “health status” category assessments.

This material is available as part of online article from:

http://www.reabic.net/aquaticinvasions/2020/Supplements/AI_2020_Ferrario_etal_SupplementaryTables.xlsx

http://www.reabic.net/aquaticinvasions/2020/Supplements/AI_2020_Ferrario_etal_SupplementaryFigures.pdf