Underwater Photogrammetry for Change Detection

Alessandro CAPRA, Cristina CASTAGNETTI, Francesco MANCINI, Paolo ROSSI, University of Modena and Reggio Emilia, Italy

Key words: Engineering survey, Photogrammetry, Change Detection

SUMMARY

Underwater surveying has been used for several applications in oil industry, archaeology and biology. The presentation will focus the attention on underwater photogrammetric surveying for change detection applications. Structure from Motion photogrammetry and underwater imagery allow the three-dimensional quantification of submersed structures characteristic at patch scale and structural complexity. High accuracy and resolution are required in order to guarantee the repeatability of surveys over time within the same reference system; a proper geodetic network and acquisition scheme are mandatory as well for 3D models generation.

The direct comparison of multi-temporal point clouds enables the evaluation of the main trends and modification with an accuracy of millimeters to centimeters accuracy.

Some examples of underwater photogrammetry for change detection will be shown and particularly with reference to a case study of multi-temporal underwater photogrammetric survey of a reef patch located in Moorea Island to detect a coral growth of about 10 mm/years.

Underwater Photogrammetry for Change Detection (10745) Alessandro Capra (Italy)

FIG Working Week 2020 Smart surveyors for land and water management Amsterdam, the Netherlands, 10–14 May 2020

Underwater Photogrammetry for Change Detection

Alessandro CAPRA, Cristina CASTAGNETTI, Francesco MANCINI, Paolo ROSSI, University of Modena and Reggio Emilia, Italy

Underwater photogrammetric surveying through images acquired by divers and unmanned underwater vehicle has been used for many applications in oil industry, archaeology and biology for change detection purposes at high accuracy level (less than 1 cm accuracy) depending on the camera settings used and the distance to the imaged object (Capra 1993, Capra et al. 2015 and 2017; Drap et al. 2013; Sarakinou 2016; Shortis 2015 and 2019).

Particularly useful for these activities has been the utilization of Structure-from-Motion (SfM) photogrammetry (Agüera-Vega et al. 2016; Eltner et al. 2016; Fonstad et al. 2013; Harwin et al. 2015; Mancini et al. 2013; Nex and Remondino 2014; Rupnik et al. 2014 among others). The absolute positioning of the 3D models is achievable at high accuracy level when a reliable reference frame is established using ground control points (GCPs) whose coordinates have been surveyed and determined with standard deviations ranging from several millimeters up to 1 cm. Information about the accuracy achieved by the underwater photogrammetric surveys is essential whenever these data are planned for deformation monitoring playing a crucial role to determine the rate of changes from multi-temporal underwater photogrammetric surveys.

The case study used for describing the strategy for a reliable underwater surveying for change detection purposes refers to a multi-temporal photogrammetric survey of a patch reef located in Moorea, French Polynesia. The main aim of the project is to provide a workflow for the 3D mapping and monitoring of coral reefs at the level of accuracy required to detect annual changes in the 3D structure of the reef.

The applied **methodology** lies in geodetic and photogrammetric measurements, data processing and multi-temporal analysis and is deeply described in Rossi et al., 2019; a short summary of the case study is here reported.

In the monitoring of growth or loss of coral 3D structure at millimeters to centimeters level of accuracy, an accurate geodetic network has been established to guarantee a common unambiguous reference frame for further comparison of the photogrammetric models reconstructed from underwater photographic surveying over time (Guo et al. 2016; Rossi et al. 2019).

In the mentioned case study, the **geodetic reference frame** was established by installing and surveying a high-quality underwater control network, which is composed of benchmarks in fixed positions; highly redundant collection of distance measurements and highly accurate leveling with laser points have been carried out. The local network adjustment produced GCP coordinates at sub-centimeter level of accuracy that were used as reference markers in the external orientation of photogrammetric models.

Several models of cameras either singly or in multiple camera combinations were used to acquire imagery in order to test camera performance in the underwater environment and identify optimized camera settings (Guo et al, 2015,Nocerino et al, 2019, Rossi et al., 2019). A Lumix DMC-GH4 and a Nauticam underwater housing equipped with a dome port to reduce the distortion effects generated by the aquatic medium (Menna et al. 2016) was used as the standard

Underwater Photogrammetry for Change Detection (10745) Alessandro Capra (Italy)

reference camera, with a nominal focal length as fixed in air of 20 mm in 2017 and 22 mm in 2018 and an exposure time of 1/90 s during 2017 survey and 1/125 s in 2018 survey in order to optimize image capture and reduce motion blur.

Rossi et al. 2019 provides the detailed description of the **photogrammetric processing for 3D models** generation (SfM algorithms by means of Agisoft Photoscan software suite) whose main steps are here listed: removal of all poor quality images (i.e., out of focus, motion blurred) from the processed dataset; image orientation with bundle adjustment; GCPs coded targets automatically identified in the images; refinement orientation through a non-linear, least squares minimization during the bundle adjustment; self-calibration procedure; depth map and creation of a high-density point cloud; validation of the generated point clouds and the estimation of their final accuracies (Toschi et al. 2013).

A **multi-temporal analysis** was performed by the comparison of 3D models obtained in two subsequent years to detect any significant changes in reef geometry over time by calculating the oriented distances between the two 3D models using the M3C2 algorithm (Lague et al. 2013; Rossi et al.,2019). The comparison procedure was accurate enough to precisely detect typical rates of changes occurring in the structural properties of corals on the order of 1 cm/year (Bessat and Buigues 2001; Burns et al., 2015; Skarlatos et al. 2017; Neyer et al. 2018); therefore the case study achieved the goal to detect a coral growth of less than 10 mm/years and it could be followed as general methodology for the high accuracy change detection purpose.

The use of the same GCP coordinates each year, the minimal constraint reference network, similar flight paths, together with the repetition of subsequent surveys under similar environmental conditions (summer months), are all strategies allowing managers to remove errors related to instrumentations, observers, sea currents, refraction indices, and seasonal variations.

The proposed, underwater photogrammetric methodology allows for a cost-effective approach to the quantitative assessment of a complex structure change with an estimated **accuracy of about 10 mm/year.**

REFERENCES

Agüera-Vega, F., Carvajal-Ramírez, F., Martínez-Carricondo, P., 2016. Accuracy of digital surface models and orthophotos derived from unmanned aerial vehicle photogrammetry. *J Surv Eng*, 04016025. https://doi.org/10.1061/(ASCE)SU.1943-5428.0000206

Bessat, F., Buigues, D., 2001. Two centuries of variation in coral growth in a massive Porites colony from Moorea (French Polynesia): a response of ocean-atmosphere variability from south central Pacific. *Palaeogeogr Palaeocl Palaeoeco*, 175(1): pp.381-392.

Burns, J. H. R., Delparte, D., Gates, R. D., Takabayashi, M., 2015. Integrating structure-frommotion photogrammetry with geospatial software as a novel technique for quantifying 3D ecological characteristics of coral reefs. *PeerJ*, 3: e1077.

Capra, A., 1993. Non-conventional system in underwater photogrammetry. Int Arch Photogramm Remote Sens, 29: 234-234.

Underwater Photogrammetry for Change Detection (10745) Alessandro Capra (Italy) Capra, A., Castagnetti, C., Dubbini, M., Gruen, A., Guo, T., Mancini, F., Neyer, F., Rossi, P., Troyer, M., 2017. High accuracy underwater photogrammetric surveying. In *3rd IMEKO international conference on metrology for archaelogy and cultural heritage-MetroArchaeo 2017*: 696-701.

Capra, A., Dubbini, M., Bertacchini, E., Castagnetti, C., Mancini, F., 2015. 3D reconstruction of an underwater archaelogical site: Comparison between low cost cameras. *ISPRS-International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences*, 40(5W5): 67-72.

Drap, P., Merad, D., Seinturier, J., Mahiddine, A., Peloso, D., Boï, J. M., Garrabou, J., 2013 (October). Underwater programmetry for archaeology and marine biology: 40 years of experience in Marseille, France. In *2013 Digital Heritage International Congress* (*DigitalHeritage*), Vol. 1: 97-104).

Eltner, A., Kaiser, A., Castillo, C., Rock, G., Neugirg, F., Abellán, A., 2016. Image-based surface reconstruction in geomorphometry – Merits, limits and developments. *Earth Surf Dynam*, 4(2): 359–389. https://doi.org/10.5194/ esurf-4-359-2016

Figueira, W., Ferrari, R., Weatherby, E., Porter, A., Hawes, S., Byrne, M., 2015. Accuracy and precision of habitat structural complexity metrics derived from underwater photogrammetry. *Remote Sens*, 7(12): 16883-16900.

Fonstad M A, Dietrich J T, Courville B C, Jensen J L, Carbonneau, P. E., 2013. Topographic structure from motion: A new development in photogrammetric measurement. *Earth Surf Proc Land*, 38(4): 421–430. https://doi.org/10.1002/esp.v38.4

Guo, T., Capra, A., Troyer, M., Gruen, A., Brooks, A. J., Hench, J. L., Schmitt, R. L., Holbrook, S. J., Dubbini, M., 2016. Accuracy assessment of underwater photogrammetric three dimensional modelling for coral reefs. *ISPRS-International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences*, 41

Harwin, S., Lucieer, A., Osborn, J., 2015. The impact of the calibration method on the accuracy of point clouds derived using unmanned aerial vehicle multi-view stereopsis. *Remote Sens*, 7(9): 11933–11953. https://doi.org/10.3390/rs70911933

Lague, D., Brodu, N., Leroux, J., 2013. Accurate 3D comparison of complex topography with terrestrial laser scanner: Application to the Rangitikei canyon (NZ). *ISPRS J Photogramm Remote Sens*, 82: 10-26.

Mancini, F., Dubbini, M., Gattelli, M., Stecchi, F., Fabbri, S., Gabbianelli, G., 2013. Using unmanned aerial vehicles (UAV) for high-resolution reconstruction of topography: The structure from motion approach on coastal environments. *Remote Sens*, 5(12): 6880–6898. https://doi.org/10.3390/rs5126880

Menna, F., Nocerino, E., Fassi, F., Remondino, F., 2016. Geometric and optic characterization of a hemispherical dome port for underwater photogrammetry. *Sensors*, 16(1): 48.

Nex, F., Remondino, F., 2014. UAV for 3D mapping applications: a review. *Appl Geomat*, 6(1): 1-15.

Neyer, F., Nocerino, E., Gruen, A., 2018. Monitoring coral growth- the dicotomy between underwater photogrammetry and geodetic control network. *ISPRS-International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences*, 42(2)

Nocerino, E.,Neyer ,F.,Gruen ,A., Troyer ,M., Menna , F.,Brooks , A.J.,Capra ,A., Castagnetti , C.,Rossi. P., (2019) Comparison of diver-operated underwater photogrammetry systems for coral reef monitoring DOI: <u>10.5194/isprs-archives-XLII-2-W10-143-2019</u>; 143-150.

Underwater Photogrammetry for Change Detection (10745) Alessandro Capra (Italy)

Rossi, P., Mancini, F., Dubbini, M., Mazzone, F., Capra, A., 2017. Combining nadir and oblique UAV imagery to reconstruct quarry topography: Methodology and feasibility analysis. *Eur J Remote Sens*, 50(1): 211-22

Rossi ,P, Castagnetti, C., Capra , A., Brooks, A.J., Mancini , F., 2019. Detecting changes coral reef 3D structure using underwater photogrammetry: critical issues and performance metrics. Applied Geomatics (2019) https://doi.org/10.1007/s12518-019-00263-w

Rupnik, E., Nex, F., Remondino, F., 2014. Oblique multi-camera systems-orientation and dense matching issues. *ISPRS-International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences*, 40(3): 107. https://doi.org/10.5194/isprsarchives-XL-3-W1-107-2014

Sarakinou, I., Papadimitriou, K., Georgoula, O., Patias, P., 2016. Underwater 3D modeling: image enhancement and point cloud filtering. *ISPRS-International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences*, 41.

Skarlatos, D., Agrafiotis, P., Menna, F., Nocerino, E., Remondino, F., 2017. Ground control networks for underwater photogrammetry in archaeological excavations. *Proceedings of the 3rd IMEKO International Conference on Metrology for Archaeology and Cultural Heritage*, MetroArcheo 2017 October 23-25, 2017, Lecce, Italy

Shortis, M. Calibration techniques for accurate measurements by underwater camera systems. Sensors 2015, 15, 30810–30826.

Shortis M. (2019) Camera Calibration Techniques for Accurate Measurement Underwater. In: McCarthy J., Benjamin J., Winton T., van Duivenvoorde W. (eds) 3D Recording and

Interpretation for Maritime Archaeology. Coastal Research Library, vol 31. Springer, Cham Toschi, I., Rivola, R., Bertacchini, E., Castagnetti, C., Dubbini, M., Capra, A., 2013. Validation tests of open-source procedures for digital camera calibration and 3D image-based modelling. ISPRS-International Archives of the Photogrammetry. *Remote Sensing and Spatial Information Sciences*, Volume XL-5/W2: 647-652. https://doi.org/10.5194/isprsarchives-XL-5-W2-647-2013

Trapon, M. L., Pratchett, M. S., Adjeroud, M., Hoey, A. S., Baird, A. H., 2013. Post-settlement growth and mortality rates of juvenile scleractinian corals in Moorea, French Polynesia versus Trunk Reef. Australia. *Mar Ecol Prog Ser*, 488: 157-170.

BIOGRAPHICAL NOTES

Full Professor of Geomatics at DIEF (Engineering Department "Enzo Ferrari") of University of Modena and Reggio Emilia (UNIMORE). Director of DIASS Dept. of Polytechnic of Bari 2003-2005. Director of DIMEC Dept. of UNIMORE from 2010 to 2012. Director of DIEF Dept. of UNIMORE from 2012 to 2018. Author and co-author of 187 scientific publications and he presented memories at national and international symposia. Responsible of research projects at national and international level. Responsible of applied research activities with public and private bodies. Coordinator of Italian geodetic observatory of PNRA (Italian Research Program on Antarctica) since 1998. Chief Officer of SCAR (Scientific Committee on Antarctic Research) Geosciences Standing Group from 2004 to 2012. Editor-in-chief of Applied geomatics(bSpringer Editor) since 2009. President of SIFET (Italian Society of Photogrammetry and Surveying) from 2010 to 2014. Member of AIOSS (Italian Association of Scientific Divers) Scientific Committee since 2019. President of China Project (UNIMORE) since November 2019. Delegate for Internationalization of UNIMORE since November 2019.

Underwater Photogrammetry for Change Detection (10745) Alessandro Capra (Italy)

CONTACTS

Prof. Alessandro Capra University of Modena and Reggio Emilia Via Vivarelli n.10, Modena, ITALY Ph. +39 0592056188 Email: alessandro.capra@unimore.it Web site: www.unimore.it

Underwater Photogrammetry for Change Detection (10745) Alessandro Capra (Italy)

FIG Working Week 2020 Smart surveyors for land and water management Amsterdam, the Netherlands, 10–14 May 2020