

## UNIQUENESS OF LIMIT CYCLES FOR QUADRATIC VECTOR FIELDS

J.L. BRAVO, M. FERNÁNDEZ, I. OJEDA, F. SÁNCHEZ

ABSTRACT. This article deals with the study of the number of limit cycles surrounding a critical point of a quadratic planar vector field, which, in normal form, can be written as  $x' = a_1x - y - a_3x^2 + (2a_2 + a_5)xy + a_6y^2$ ,  $y' = x + a_1y + a_2x^2 + (2a_3 + a_4)xy - a_2y^2$ . In particular, we study the semi-varieties defined in terms of the parameters  $a_1, a_2, \dots, a_6$  where some classical criteria for the associated Abel equation apply. The proofs will combine classical ideas with tools from computational algebraic geometry.

### 1. INTRODUCTION AND MAIN RESULTS

The number of periodic solutions of a quadratic polynomial planar system is an open problem and the first non-trivial case of the second part of Hilbert's *XVI*-th problem.

It is known that if a quadratic system has a limit cycle, i.e., a periodic solution that is isolated in the set of periodic solutions of the system, then it must surround a focus of the system. In particular, if one takes the focus to be at the origin, then the system can be written in the form (see [5])

$$(1.1) \quad \begin{aligned} x' &= a_1x - y - a_3x^2 + (2a_2 + a_5)xy + a_6y^2, \\ y' &= x + a_1y + a_2x^2 + (2a_3 + a_4)xy - a_2y^2. \end{aligned}$$

One way to study the periodic solutions of (1.1) is to analyse the  $2\pi$ -periodic positive solutions of the polar equation

$$(1.2) \quad \frac{dr}{d\theta} = \frac{a_1r + f(\theta)r^2}{1 + g(\theta)r},$$

---

2010 *Mathematics Subject Classification*. Primary 34C25. Secondary: 34A34, 37C27, 37G15.

*Key words and phrases*. Abel equation, closed solution, periodic solution, limit cycle.

where  $f$  and  $g$  are the cubic homogeneous trigonometric polynomials defined by

$$\begin{aligned} f(\theta) &= -a_3 \cos^3 \theta + (3a_2 + a_5) \cos^2 \theta \sin \theta \\ &\quad + (2a_3 + a_4 + a_6) \cos \theta \sin^2 \theta - a_2 \sin^3 \theta, \\ g(\theta) &= a_2 \cos^3 \theta + (3a_3 + a_4) \cos^2 \theta \sin \theta \\ &\quad - (3a_2 + a_5) \cos \theta \sin^2 \theta - a_6 \sin^3 \theta, \end{aligned}$$

or of the Cherkas-equivalent Abel differential equation (see [8])

$$(1.3) \quad \rho' = A(\theta)\rho^3 + B(\theta)\rho^2 + a_1\rho,$$

where

$$A(\theta) = g(\theta)(a_1g(\theta) - f(\theta)), \quad B(\theta) = f(\theta) - 2a_1g(\theta) - g'(\theta).$$

There are several results that establish upper bounds for the number of limit cycles of (1.3). The best known ones impose the condition that one of the functions  $A$  or  $B$  has definite sign, see [14, 15, 19, 21, 23], where a  $2\pi$ -periodic function  $F(\theta)$  has definite sign if  $F(\theta) \geq 0$  for all  $\theta \in [0, 2\pi]$  or  $F(\theta) \leq 0$  for all  $\theta \in [0, 2\pi]$ .

In the particular case of Equation (1.3), the criteria in [15, 23] give the following result.

**Theorem 1.1** ([15, 23]). *If  $A$  or  $B$  has definite sign, then Abel equation (1.3) has at most one positive limit cycle.*

In [9] the quadratic systems for which the above criteria applies are described taking into account their number of critical points and the directions  $\theta$  in which  $g(\theta) = 0$ .

To establish our main results, which determine the semi-varieties in the space of parameters where the above criteria apply, and, as a consequence, to obtain that at most one limit cycle surrounds the origin of (1.1), we shall need the following notation.

The study of whether  $A$  has definite sign can, by the change of variable  $x = \tan(\theta)$  (see Section 4), be reduced to the study of the common roots of the polynomials  $p_1(x)$  and  $p_3(x) := a_1p_1(x) - p_2(x)$ , where

$$\begin{aligned} p_1(x) &= a_2 + (3a_3 + a_4)x - (3a_2 + a_5)x^2 - a_6x^3, \\ p_2(x) &= -a_3 + (3a_2 + a_5)x + (2a_3 + a_4 + a_6)x^2 - a_2x^3. \end{aligned}$$

Let us denote by  $D_1, D_3, D'_1, D'_3$  the discriminants of the polynomials  $p_1, p_3, p'_1, p'_3$ , respectively. If  $\text{res}(p_1, p_3)$  denotes the resultant of  $p_1$  and  $p_3$  with respect to  $x$ , then it factorizes as

$$\text{res}(p_1, p_3) = R_1R_2,$$

where

$$\begin{aligned} R_1 &= (4a_2 + a_5)^2 + (3a_3 + a_4 + a_6)^2, \\ R_2 &= a_3a_6l_0^2 + a_2l_0l_1l_2 + a_2^2(l_1 + l_2)(l_1 + l_3), \end{aligned}$$

with  $l_0 = 2a_3 + a_4$ ,  $l_1 = 2a_2 + a_5$ ,  $l_2 = a_3 + a_6$  and  $l_3 = a_3 - a_6$ .

Let us write

$$R_{113} = \text{res}(p'_1, p_3), \quad R_{133} = \text{res}(p_1, p'_3).$$

If  $r_1$  (resp.  $r_3$ ) denotes the remainder of the polynomial division of  $p_1$  by  $p'_1$  (resp.  $p_3$  by  $p'_3$ ), we shall write

$$\bar{R}_{113} = \text{res}(r_1, p_3), \quad \bar{R}_{133} = \text{res}(p_1, r_3).$$

Note that  $D_1, D_3, D'_1, \dots, \bar{R}_{113}, \bar{R}_{133}$ , are defined “for the generic case”, i.e., they are obtained as expressions on  $a_1, \dots, a_6$  without imposing any condition. Some of the expressions are not included in the paper as they are gruesome.

The first result determines the quadratic systems such that  $A(\theta)$  has definite sign.

**Theorem A.** *The coefficient  $A$  has definite sign (and, in consequence, (1.1) has at most one limit cycle surrounding the origin) if and only if one of the following conditions holds:*

- (1)  $p_1$  or  $p_3$  is identically null, or, equivalently, one of the following conditions holds:
  - (a)  $a_6 = a_5 = 3a_3 + a_4 = a_2 = 0$ ,
  - (b)  $a_1a_6 - a_2 = a_1a_5 - a_3 + a_4 + a_6 = a_1(3a_3 + a_4) - 3a_2 - a_5 = a_1a_2 + a_3 = 0$ .
- (2)  $p_1$  has degree one,  $p_3$  has degree three (i.e.,  $a_6 = 3a_2 + a_5 = 0$  and  $(3a_3 + a_4)a_2 \neq 0$ ),  $R_2 = 0$ , and  $a_2^2 \leq 4a_3^2 + 4a_1a_2a_3$ .
- (3)  $p_3$  has degree one,  $p_1$  has degree three (i.e.,  $a_2 - a_1a_6 = 2a_3 + a_4 + a_1(3a_2 + a_5) + a_6 = 0$  and  $a_6(3a_2 - a_1(3a_3 + a_4) + a_5) \neq 0$ ), and one of the following conditions holds:
  - (a)  $R_2 = 0$ ,  $D_1 \leq 0$ ,  $R_{113} \neq 0$ ,
  - (b)  $4a_4 - 9a_6 = 4a_3 + 5a_6 = 9a_2 + a_5 = 9a_1a_6 + a_5 = 8a_1^2 - 1 = 0$ .
- (4)  $p_1, p_3$  have degree two (i.e.,  $a_2 = a_6 = 0$  and  $a_5(a_3 - a_1a_5) \neq 0$ ),  $3a_3 + a_4 = 0$ , and  $4a_3^2 - 4a_1a_3a_5 \geq a_5^2$ .
- (5)  $p_1, p_3$  have degree three (i.e.,  $a_6(a_2 - a_1a_6) \neq 0$ ),  $R_2 = 0$ , and one of the following conditions holds:
  - (a)  $D_1 < 0$ ,  $D_3 < 0$ ,  $(a_3 - a_6)(a_2^2 + (a_4 + 2a_3)^2) \neq 0$ ,
  - (b)  $D_1 = 0$ ,  $D_3 < 0$ ,  $D'_1R_{113} \neq 0$ ,
  - (c)  $D_1 = D'_1 = 0$ ,  $D_3 < 0$ ,
  - (d)  $D_3 = 0$ ,  $D_1 < 0$ ,  $D'_3R_{133} \neq 0$ ,
  - (e)  $D_3 = D'_3 = 0$ ,  $D_1 < 0$ ,
  - (f)  $D_1 = D_3 = 0$ ,  $D'_1D'_3\bar{R}_{113}\bar{R}_{133} \neq 0$ ,
  - (g)  $D_1 = D'_1 = D_3 = 0$ ,  $\bar{R}_{133} \neq 0$ ,
  - (h)  $D_1 = D_3 = D'_3 = 0$ ,  $\bar{R}_{113} \neq 0$ .

*Remark 1.2.* The codimension of the semi-varieties defined by the conditions of Theorem A are the following (Proposition 4.10):

- 5a) has codimension one.
- 5b), 5d) have codimension two.
- 2), 3a), 4) have codimension three.
- 1a), 1b) have codimension four.
- 3b) has codimension five.
- 5f) has codimension two or three.
- 5c), 5e), 5g), 5h) have codimension of at least two.

Note that in case 3b) the equations already imply  $a_2 - a_1a_6 = 2a_3 + a_4 + a_1(3a_2a_5) + a_6 = 0$ .

Next, we determine quadratic systems such that  $B(t)$  has definite sign.

**Theorem B.** *The coefficient  $B$  has definite sign (indeed, it is identically null) if and only if the parameters  $a_1, \dots, a_6$  belong to any of the two codimension-four regular varieties defined by the equations*

$$(1.4) \quad a_4 + 4a_6 = 4a_3 + a_4 = 4a_2 + a_5 = a_1 = 0,$$

or

$$(1.5) \quad a_6 = 3a_3 + a_4 = 4a_2 + a_5 = 3a_1a_5 + 2a_4 = 0.$$

Moreover, (1.1) has at most one limit cycle surrounding the origin.

*Remark 1.3.* The conditions (1.4), (1.5) in Theorem B imply that  $B$  is identically null. Therefore, (1.3) reduces to a Bernoulli equation, and it is possible to obtain the exact number of limit cycles surrounding the origin (zero or one).

The rest of the paper is organized as follows. Section 2 contains some known results on the number of limit cycles of Abel equations. Section 3 describes the algebraic geometry tools that will be required for the proofs of the main results. Section 4 contains the proofs of Theorems A and B. Finally, in Appendix A we include the SINGULAR code for the proofs of Section 4.

## 2. ABEL EQUATIONS WITH AT MOST ONE NON-TRIVIAL LIMIT CYCLE

In this section we collect known results about the number of limit cycles of the Abel equation (1.3) that we will use subsequently.

**Proposition 2.1** ([23, 15]). *Assume  $A(\theta)$  has definite sign. Then Equation (1.3) has at most one positive limit cycle.*

*Proof.* From [23], we have that (1.3) has at most three limit cycles. Moreover, notice that  $\rho = 0$  is always a periodic solution of (1.3). Since  $A(\theta + \pi) = A(\theta)$  and  $B(\theta + \pi) = -B(\theta)$ , we have that  $\rho(\theta)$  is a solution of (1.3) if and only if  $-\rho(\theta + \pi)$  also is. Thus the number of limit cycles is the same in regions  $\rho > 0$  and  $\rho < 0$ , and consequently Equation (1.3) has at most one positive limit cycle.  $\square$

**Proposition 2.2.** *Assume  $A(\theta)$  to be identically null. Then Equation (1.3) has no limit cycle.*

*Proof.* When  $A(\theta) \equiv 0$ , Equation (1.3) is the Ricatti equation  $\rho' = B(\theta)\rho^2 + a_1\rho$ . Since  $\int_0^{2\pi} B(t) dt = 0$ , when  $a_1 = 0$  it is a centre and if  $a_1 \neq 0$  it has no limit cycle.  $\square$

**Proposition 2.3.** *If  $B(\theta)$  has definite sign, it is identically null. Moreover, equation(1.3) has at most one positive limit cycle.*

*Proof.* Since  $B(\theta + \pi) = -B(\theta)$ , if  $B(\theta)$  has definite sign, it is necessarily identically null. Then (1.3) is the Bernoulli equation  $\rho' = A(t)\rho^3 + a_1\rho$  which has at most one positive limit cycle.  $\square$

*Remark 2.4.* The criterion  $\alpha A + \beta B$  has definite sign for some  $\alpha, \beta \in \mathbb{R}$ ,  $\alpha^2 + \beta^2 \neq 0$ , used in [1, 16] to obtain upper bounds for the number of limit cycles in Abel equations is not relevant in this context since if  $\alpha A + \beta B$  has definite sign then, by the change of variables  $t \rightarrow \pi + t$ ,  $\alpha A - \beta B$  has the same definite sign. Therefore  $2\alpha A = (\alpha A + \beta B) + (\alpha A - \beta B)$  has definite sign, and consequently  $A$  has definite sign if  $\alpha \neq 0$  and  $B(t) \equiv 0$  otherwise.

### 3. ALGEBRAIC GEOMETRY TOOLS

In this section, we summarize the computational algebraic geometry results to be used subsequently. In all cases, we will include references to the SINGULAR ([11]) commands necessary to perform the corresponding computation. Those readers interested in considering computational algebraic geometry techniques in more depth are encouraged to consult [10] for an introduction, or [4] for a fuller development. Furthermore, readers familiar with differential equations will enjoy [24] which includes a comprehensive introduction to the basic generalities of computational algebraic geometry in its first chapter.

Let us consider a system of polynomial equations in  $n$  variables  $x_1, \dots, x_n$  with coefficients in a field  $\mathbb{k}$ ,

$$(3.6) \quad \begin{aligned} f_1(x_1, \dots, x_n) &= 0, \\ &\vdots \\ f_s(x_1, \dots, x_n) &= 0. \end{aligned}$$

Clearly,  $(a_1, \dots, a_n) \in \mathbb{k}^n$  is a solution of (3.6) if and only if

$$\sum_{i=1}^s g_i(a_1, \dots, a_n) f_i(a_1, \dots, a_n) = 0$$

for every  $g_i$  in the ring  $\mathbb{k}[x_1, \dots, x_n]$  of polynomials in  $n$  variables with coefficients in  $\mathbb{k}$ . Thus, the set of solutions of (3.6) in  $\mathbb{k}^n$  matches the set of zeros in  $\mathbb{k}^n$  of the ideal  $\langle f_1, \dots, f_s \rangle$  of  $\mathbb{k}[x_1, \dots, x_n]$  generated by  $f_1, \dots, f_s$ . The set of zeros of  $I = \langle f_1, \dots, f_s \rangle$  in  $\mathbb{k}^n$  is called the (affine) variety of  $I$  in  $\mathbb{k}^n$ . It is denoted  $\mathcal{V}_{\mathbb{k}}(I)$ , or simply  $\mathcal{V}(I)$  when no confusion is possible.

Here, it is convenient to recall that all the ideals of  $\mathbb{k}[x_1, \dots, x_n]$  are finitely generated by the Hilbert Basis Theorem (see [4, Theorem 1.3.5]). Therefore, to study a system of polynomial equations is the same as to study the ideal generated by the polynomials of the system, and vice versa.

Furthermore, since  $f(a_1, \dots, a_n) = 0$  if and only if  $f^r(a_1, \dots, a_n) = 0$  for every positive integer  $r$ , one has that  $\mathcal{V}(I) = \mathcal{V}(\sqrt{I})$ , where

$$\sqrt{I} = \{f \in \mathbb{k}[x_1, \dots, x_n] \mid f^r \in I, \text{ for some } r \in \mathbb{Z}_+\}$$

is the radical of  $I$ .

This ideal-variety approach has two immediate advantages. On the one hand, the varieties in  $\mathbb{k}^n$  of the ideals of  $\mathbb{k}[x_1, \dots, x_n]$  form the closed sets of a topology on  $\mathbb{k}^n$  called the Zariski topology of  $\mathbb{k}^n$  (see [4, Lemma A.2.4]). And on the other, there exists a kind of factorization theory for ideals of  $\mathbb{k}[x_1, \dots, x_n]$  in which the intersection of ideals plays the role of the product: the so-called primary decomposition theory that we shall outline in the following.

Observe that because of the well-known property

$$\mathcal{V}(J_1 \cap J_2) = \mathcal{V}(J_1) \cup \mathcal{V}(J_2),$$

for  $J_i$ ,  $i = 1, 2$ , ideals of  $\mathbb{k}[x_1, \dots, x_n]$  (see [4, Lemma A.2.3, part (2)]), a decomposition of the ideal defined by the polynomials in (3.6) as an intersection of “simpler ideals” will mean splitting the system 3.6 into several easier-to-solve systems, hopefully!

Depending on the purpose, some systems of generators of a polynomial ideal are better than others. For example, minimal systems of generators (i.e., systems of generators such that no generator is an algebraic combination of the others) are preferred for a concise description of the variety. But Gröbner bases, which are far from being minimal in the above sense, are special systems of generators with good computational properties. Given a system of generators of an ideal  $I$  of  $\mathbb{k}[x_1, \dots, x_n]$ , one can compute a minimal system of generators or a Gröbner basis of  $I$  by using the SINGULAR commands `mres(I, 1) [1]` or `std(I)`, respectively.

The original aim of the Gröbner bases methods was to compute the remainder of a polynomial under division by a polynomial ideal, something that can be done with the command `reduce` in SINGULAR. Nowadays, Gröbner bases are used for more sophisticated tasks. Computing the dimension of a variety or eliminating variables are just two classic examples.

Given a system of generators of an ideal  $I$  of  $\mathbb{k}[x_1, \dots, x_n]$ , the problem of the computation of the dimension of  $\mathcal{V}(I)$  (equivalently, the Krull dimension of  $\mathbb{k}[x_1, \dots, x_n]/I$ ) may be reduced to a pure combinatorial problem after the computation of one (any) Gröbner basis of  $I$  (see [10, Chapter 9]). The SINGULAR command `dim(std(I))` will compute the dimension of  $\mathcal{V}_{\mathbb{C}}(I)$  for us. The precise notion of dimension will be defined at the end of this section. On other hand, the problem of the elimination of a variable, say  $x_n$ , from the ideal  $I$ , consists of determining a system of generators of  $I \cap \mathbb{k}[x_1, \dots, x_{n-1}]$ .

This can be easily computed from a Gröbner basis of  $I$  with respect to a suitable well-ordering of the monomials in  $\mathbb{k}[x_1, \dots, x_n]$ . Geometrically, the elimination of variables has the following meaning:

**Proposition 3.1.** *Let  $\mathbb{k}$  be algebraically closed, and let  $I$  be an ideal of  $\mathbb{k}[x_1, \dots, x_n]$ . If  $\pi : \mathbb{k}^n \rightarrow \mathbb{k}^{n-1}$  is the projection map that sends  $(a_1, \dots, a_n)$  to  $(a_1, \dots, a_{n-1})$  then the Zariski closure of  $\pi(\mathcal{V}(I))$  in  $\mathbb{k}^{n-1}$  is equal to  $\mathcal{V}(I \cap \mathbb{k}[x_1, \dots, x_{n-1}])$ .*

*Proof.* See [10, Theorem 3, Section 3.2]. □

The elimination of variables is computed in SINGULAR with the command `eliminate`.

Let us now briefly summarize the primary decomposition process for ideals of  $\mathbb{k}[x_1, \dots, x_n]$ . To do so, we shall first introduce the quotient operation and its most elementary properties.

**Definition 3.2.** Let  $I$  and  $J$  be ideals of  $\mathbb{k}[x_1, \dots, x_n]$ . The quotient of  $I$  and  $J$  is the ideal  $(I : J)$  of  $\mathbb{k}[x_1, \dots, x_n]$  defined as follows:

$$(I : J) = \{g \in \mathbb{k}[x_1, \dots, x_n] \mid gf \subseteq I, \text{ for every } f \in J\}.$$

It is not difficult to see that  $I \subseteq (I : J)$  and  $((I : J) : J) = (I : J^2)$ . Then, we have a chain of ideals  $I \subseteq (I : J) \subseteq \dots \subseteq (I : J^r) \subseteq \dots$  that necessarily stabilizes by the Noetherian property of  $\mathbb{k}[x_1, \dots, x_n]$ . If  $N$  is the smallest integer for which the above chain stabilizes, then the ideal  $(I : J^N)$  is called the saturation of  $I$  by  $J$  and is usually denoted  $(I : J^\infty)$ .

Both quotient and saturation can be computed using the SINGULAR commands `quotient` and `sat`, respectively (the latter from the `elim` library).

*Remark 3.3.* Observe that an elementary necessary and sufficient condition for  $J \subseteq I$  is  $(I : J) = \langle 1 \rangle$ . Moreover, one has that  $f \in \sqrt{I}$  if and only if  $(I : \langle f \rangle^\infty) = \langle 1 \rangle$ . So, the radical membership problem can be computationally solved by computing the saturation of  $I$  by  $\langle f \rangle$ .

Geometrically, when  $\mathbb{k}$  is algebraically closed, the quotient and the saturation of  $I$  by  $J$  have the same behaviour which is nothing but the Zariski closure of the difference of varieties. In particular, the following holds:

$$\mathcal{V}(I : J) = \overline{\mathcal{V}(I) \setminus \mathcal{V}(J)} = \overline{\mathcal{V}(I) \setminus \mathcal{V}(J^r)} = \mathcal{V}(I : J^r),$$

for every positive integer  $r$  (see [10, Theorem 7, section 4.4]).

The next result represents a first step for the decomposition of an ideal of  $\mathbb{k}[x_1, \dots, x_n]$ :

**Lemma 3.4. (Splitting tool).** *Let  $I$  be an ideal of  $\mathbb{k}[x_1, \dots, x_n]$ , and let  $g \in \mathbb{k}[x_1, \dots, x_n]$ . If  $N$  is the smallest integer such that  $(I : \langle g \rangle^\infty) = (I : \langle g^N \rangle)$ , then*

$$I = (I : \langle g \rangle^\infty) \cap (I + \langle g^N \rangle).$$

*Proof.* See [4, Lemma 3.3.6]. □

When  $\mathbb{k}$  is algebraically closed, an immediate consequence of the splitting tool is the formula

$$\mathcal{V}(I) = \mathcal{V}(I : f^\infty) \cup \mathcal{V}(I + \langle f \rangle) = \overline{\mathcal{V}(I) \setminus \mathcal{V}(f)} \cup (\mathcal{V}(I) \cap \mathcal{V}(f))$$

where the varieties in the union on the right-hand side can be carefully interpreted as the solutions of the system associated with  $I$  by imposing the conditions  $f(x_1, \dots, x_n)$  different from or equal to zero, respectively.

At this point, we are in a position to clarify what “simpler ideals” means in the context of primary decomposition theory.

**Definition 3.5.** An ideal  $P$  of  $\mathbb{k}[x_1, \dots, x_n]$  is said to be prime if  $fg \in P$  and  $g \notin P$  implies  $f \in P$ . An ideal  $Q$  of  $\mathbb{k}[x_1, \dots, x_n]$  is said to be primary if  $fg \in Q$  and  $g \notin Q$  implies  $f \in \sqrt{Q}$ .

Notice that every prime ideal  $P$  is primary: indeed, if  $P$  is prime,  $\sqrt{P} = P$ . Moreover, one can easily check that the radical of a primary ideal is prime. Here, it is important to emphasize that, if  $\mathbb{k}$  is algebraically closed, then  $P$  is a prime ideal of  $\mathbb{k}[x_1, \dots, x_n]$  if and only if  $\mathcal{V}(P)$  is Zariski irreducible (see [10, Corollary 4, Section 4.5]). So, in this case, the variety of a primary ideal is a Zariski irreducible subset of  $\mathbb{k}^n$ .

**Theorem 3.6.** Let  $I$  be an ideal of  $\mathbb{k}[x_1, \dots, x_n]$ . If  $I \neq \langle 1 \rangle$ , there exists a decomposition of  $I$  as the intersection of finitely many primary ideals.

*Proof.* If  $I$  is primary, there is nothing to prove. Otherwise, there exists  $g \notin \sqrt{I}$  such that  $(I : g^\infty) \supsetneq I$ . Thus, by Lemma 3.4,  $I$  decomposes as  $(I : g^\infty) \cap (I + \langle g^N \rangle)$ . Both ideals strictly contain  $I$ . If they are primary, we are done. Otherwise, we can repeat the same argument with  $(I : g^\infty)$  and  $(I + \langle g^N \rangle)$ , and so on and so forth. In so far as this process cannot continue indefinitely because of the Noetherian property of  $\mathbb{k}[x_1, \dots, x_n]$ , our claim follows.  $\square$

A decomposition of  $I$  into primary ideals,  $I = Q_1 \cap \dots \cap Q_r$ , is called a primary decomposition of  $I$ . Since  $\sqrt{I} = \sqrt{Q_1} \cap \dots \cap \sqrt{Q_r}$ , by removing redundancies if necessary, we obtain finitely many prime ideals,  $P_1, \dots, P_t$ , not contained one in another, such that

$$\mathcal{V}(I) = \mathcal{V}(P_1) \cup \dots \cup \mathcal{V}(P_t).$$

Therefore, when  $\mathbb{k}$  is algebraically closed, a primary decomposition of an ideal  $I$  yields a decomposition of  $\mathcal{V}(I)$  into Zariski irreducible varieties. In general, the prime ideals defining these varieties do not depend on the decomposition, and are called minimal associated primes of  $I$  ([4, Theorem 4.1.5]).

*Remark 3.7.* Let  $\{P_1, \dots, P_t\}$  be the set of minimal associated primes of an ideal  $I$  of  $\mathbb{k}[x_1, \dots, x_n]$ . If  $P'$  is a prime ideal such that  $I \subseteq P' \subseteq P_j$  for some  $j$ , then  $P' = P_j$ . Indeed, it suffices to note that  $\sqrt{I} = P_1 \cap \dots \cap P_t \subseteq P' \subseteq P_j$  implies  $P_i \subseteq P' \subseteq P_j$  for some  $i$ , and that necessarily  $i = j$ . Therefore, the minimal associated primes of  $I$  are the “smallest” prime ideals containing  $I$ .



In conclusion, there exists a computational method to write the set of solutions of a system of polynomial equations in several variables as the union of the solution of finitely many systems. Moreover, if  $\mathbb{k}$  is algebraically closed, the varieties associated with those systems are Zariski irreducible.

The minimal associated primes of an ideal of  $\mathbb{k}[x_1, \dots, x_n]$  can be computed by using the SINGULAR command `minAssGTZ` (library `primary`).

We end this section by defining the notion of dimension of an algebraic variety.

**Definition 3.8.** Let  $I$  be an ideal of  $\mathbb{k}[x_1, \dots, x_n]$ . The dimension of  $I$ ,  $\dim(I)$ , is the supremum of the lengths of all chains of prime ideals in  $\mathbb{k}[x_1, \dots, x_n]/I$ .

Equivalently,  $\dim(I)$  is supremum of the lengths of all chains of prime ideals in  $\mathbb{k}[x_1, \dots, x_n]$  containing  $I$  (because of the well-known correspondence between ideals of the quotient  $A/I$  and ideals of  $A$  containing  $I$ ). Observe that

$$\dim(I) = \max \{ \dim(P) \mid P \text{ is a minimal associated prime of } I \}$$

by Remark 3.7.

This notion of dimension does not depend on the base field  $\mathbb{k}$  in the sense that if  $\mathbb{k} \hookrightarrow \mathbb{K}$  is an extension of  $\mathbb{k}$ , then the dimension of  $I$  is the same regardless of whether  $I$  is an ideal of  $\mathbb{k}[x_1, \dots, x_n]$  or an ideal of  $\mathbb{K}[x_1, \dots, x_n]$  (see [4, Theorem 3.5.1]).

Since the dimension of  $\mathcal{V}(I)$  is the supremum of the lengths of the chains of its closed irreducible sets, when  $\mathbb{k}$  is algebraically closed, the dimension of  $\mathcal{V}(I)$  is the maximum of  $\dim(P)$  where  $P$  is any minimal associated prime of  $I$ .

The next result is a particular version of the General Jacobian criterion (see [4, Theorem 5.7.1]).

**Theorem 3.9.** Let  $I = \langle f_1, \dots, f_m \rangle \subset \mathbb{k}[x_1, \dots, x_n]$  be an ideal and  $P$  a minimal associated prime of  $I$ . If  $\mathbf{a} = (a_1, \dots, a_n) \in \mathcal{V}(P) \subseteq \mathbb{k}^n$ , then

$$(3.7) \quad \text{rank} \left( \frac{\partial f_i}{\partial x_j}(\mathbf{a}) \right) \leq n - \dim(P),$$

and  $\mathbf{a} = (a_1, \dots, a_n)$  is a regular point of  $\mathcal{V}(I)$  if and only if the equality holds.

*Proof.* This theorem is nothing but [4, Theorem 5.7.1], from taking into account that  $n - \dim(P)$  is the height of  $P$ ,  $\text{ht}(P)$ , by [4, Theorem 3.5.1(4)] and the definition of regular point given in [4, Definition A.8.7].  $\square$

The left hand side in (3.7) can be computed in SINGULAR with the following command `rank(reduce(jacob(I), std(m_a)))`, where  $\mathbf{m}_a$  is the maximal ideal associated with  $\mathbf{a}$ , i.e.,  $\mathbf{m}_a = \langle x_1 - a_1, \dots, x_n - a_n \rangle$ .

## 4. PROOF OF THE MAIN RESULTS

In this section, we shall prove Theorem A and Theorem B.

A first consideration is that the functions  $A$  and  $B$  are homogeneous trigonometric polynomials of degrees 6 and 3, respectively. Since  $\sin(\theta) = -\sin(\theta + \pi)$  and  $\cos(\theta) = -\cos(\theta + \pi)$ , then for all  $\theta \in (-\pi/2, \pi/2]$

$$A(\theta) = A(\theta + \pi), \quad B(\theta) = -B(\theta + \pi).$$

In particular,  $B$  has definite sign if and only if  $B(\theta) \equiv 0$  for all  $\theta \in (-\pi/2, \pi/2]$ , and  $A$  has definite sign if and only if  $A(\theta) \geq 0$  for all  $\theta \in (-\pi/2, \pi/2]$ , or  $A(\theta) \leq 0$  for all  $\theta \in (-\pi/2, \pi/2]$ .

By the changes of variables  $x = \tan(\theta)$ , we obtain that  $A$  has definite sign if and only if the rational function

$$A(\operatorname{atan}(x)) = \frac{p_1(x)(a_1 p_1(x) - p_2(x))}{(1+x^2)^3}$$

has definite sign, where (we recall)

$$\begin{aligned} p_1(x) &= a_2 + (3a_3 + a_4)x - (3a_2 + a_5)x^2 - a_6x^3, \\ p_2(x) &= -a_3 + (3a_2 + a_5)x + (2a_3 + a_4 + a_6)x^2 - a_2x^3, \end{aligned}$$

or equivalently, that  $p_1(x)(a_1 p_1(x) - p_2(x))$  has definite sign.

Analogously, by the change of variable  $x = \tan(\theta)$ ,  $B$  is identically null if and only if

$$B(\operatorname{atan}(x)) = \frac{q(x)}{(1+x^2)^{3/2}} \equiv 0, \quad \text{for all } x \in \mathbb{R},$$

where

$$\begin{aligned} q(x) &= -\left(2a_1a_2 + 4a_3 + a_4\right) + \left(12a_2 + 3a_5 - a_1(6a_3 + 2a_4)\right)x \\ &\quad + \left(8a_3 + 3a_4 + 4a_6 + a_1(6a_2 + 2a_5)\right)x^2 - \left(4a_2 + a_5 - 2a_1a_6\right)x^3. \end{aligned}$$

Again, that is equivalent to  $q(x) \equiv 0$ .

**4.1. Proof of Theorem A.** We divide the proof of Theorem A into several propositions. A first comment is that if  $p_1(x) \equiv 0$  or  $p_3(x) := a_1 p_1(x) - p_2(x) \equiv 0$  then  $A(\theta) \equiv 0$ . In Proposition 4.1 we characterize when one of the polynomials  $p_1$ ,  $p_3$  is identically null. Next, we distinguish cases in terms of the minimum of the degrees of  $p_1$  and  $p_3$ . When this minimum is zero, Theorem A is proved in Proposition 4.2; when it is one, in Proposition 4.3; when it is two, in Proposition 4.4; and when it is three, in Proposition 4.9.

**Proposition 4.1.** *The polynomial  $p_1p_3$  is identically null if and only if*

$$(4.8) \quad a_6 = a_5 = 3a_3 + a_4 = a_2 = 0,$$

or

$$(4.9) \quad \begin{aligned} a_1a_6 - a_2 &= a_1a_5 - a_3 + a_4 + a_6 = 0, \\ a_1(3a_3 + a_4) - 3a_2 - a_5 &= a_1a_2 + a_3 = 0. \end{aligned}$$

*Proof.* It suffices to consider the ideals generated by the coefficients of the polynomials, and then, for each of these ideals, compute a minimal system of generators. (See Appendix A).  $\square$

In the following, we assume that neither of  $p_1, p_3$  is identically null. In consequence,  $p_1p_3$  has definite sign if and only if the odd-multiplicity real roots of  $p_1$  and  $p_3$  coincide. We shall distinguish several cases depending on the minimum degree of  $p_1$  and  $p_3$ .

If the minimum degree of  $p_1$  and  $p_3$  is zero (and neither of  $p_1, p_2, p_3$  is identically null), then  $A$  does not have definite sign.

**Proposition 4.2.** *If  $p_1$  and  $p_3$  are not identically null and  $p_1$  or  $p_3$  is constant, then the odd-multiplicity real roots of  $p_1$  and  $p_3$  do not coincide.*

*Proof.* Assume  $p_1$  is constant, i.e.,  $p_1(x) = a_2$ . If the odd-multiplicity roots of  $p_1, p_3$  coincide, then  $p_3$  has even degree. Hence  $a_2 = 0$ , in contradiction with  $p_1$  not being null.

Conversely, if  $p_3$  is constant and not null, then  $p_3(x) = a_1a_2 + a_3$ . Arguing as above,  $p_1$  has an even degree, so  $a_6 = 0$ . Moreover, since  $p_3(x)$  is constant,  $a_2 = 0$  in particular, and

$$p_1(x) = x(3a_3 + a_4 - a_5x).$$

I.e.,  $x = 0$  is a root of  $p_1$ . If it is a simple root, it should be a root of  $p_3$ , in contradiction with  $p_3$  being constant, so that  $3a_3 + a_4 = 0$ . But in this case,

$$p_3(x) = a_3 - a_5x + (-2a_3 - a_4 - a_1a_5)x^2.$$

In particular,  $a_5 = 0$ , so  $p_1(x) \equiv 0$ , and with this contradiction we conclude the proof.  $\square$

Next, we consider that one of  $p_1, p_3$  has degree one, and the other has an equal or greater degree.

**Proposition 4.3.** *Assume that the minimum of the degrees of  $p_1$  and  $p_3$  is one. Then the odd-multiplicity real-roots of  $p_1, p_3$  coincide if and only if*

$$(4.10) \quad a_6 = 3a_2 + a_5 = R_2 = 0, \quad a_2^2 \leq 4a_3^2 + 4a_1a_2a_3, \quad a_2(3a_3 + a_4) \neq 0,$$

or

$$(4.11) \quad \begin{aligned} a_2 - a_1a_6 &= 2a_3 + a_4 + a_1(3a_2 + a_5) + a_6 = R_2 = 0, \\ D_1 \leq 0, \quad a_6(3a_2 - a_1(3a_3 + a_4) + a_5) &R_{113} \neq 0. \end{aligned}$$

or

$$(4.12) \quad 4a_4 - 9a_6 = 4a_3 + 5a_6 = 9a_2 + a_5 = 9a_1a_6 + a_5 = 8a_1^2 - 1 = 0, \quad a_6 \neq 0.$$

*Proof.* Assume the odd-multiplicity real-roots of  $p_1, p_3$  coincide. Then the possible degrees of  $p_1, p_3$  are one or three. The polynomials  $p_1$  and  $p_3$  can not be simultaneously linear, since in this case  $p_1(x) = a_3x$  and  $p_3(x) = a_3 + a_1a_3x$ .

CASE 1. Assume that  $p_1$  has degree one and  $p_3$  has degree three. Then  $a_6 = 0$ ,  $3a_2 + a_5 = 0$ ,  $3a_3 + a_4 \neq 0$ , and  $a_2 \neq 0$ . Moreover,

$$\begin{aligned} p_1(x) &= a_2 + (3a_3 + a_4)x, \\ p_3(x) &= a_1a_2 + a_3 + a_1(3a_3 + a_4)x - (2a_3 + a_4)x^2 + a_2x^3. \end{aligned}$$

Assume that the odd-multiplicity roots of  $p_1$  and  $p_3$  coincide. The root of  $p_1$  is  $x_1 = -a_2/(3a_3 + a_4)$ . Then

$$p_3(x_1) = -\frac{(-a_2^2 + a_3(3a_3 + a_4))(a_2^2 + (3a_3 + a_4)^2)}{(3a_3 + a_4)^3} = 0.$$

Hence

$$0 = a_2^2 - a_3(3a_3 + a_4) = \frac{R_2}{a_2^2}.$$

In consequence  $a_3 \neq 0$ . Replacing  $a_4$  by  $\frac{a_2^2 - 3a_3^2}{a_3}$ , we obtain

$$p_1(x) = \frac{a_2(a_3 + a_2x)}{a_3}, \quad p_3(x) = \frac{(a_3 + a_2x)(a_1a_2 + a_3 - a_2x + a_3x^2)}{a_3}.$$

The odd-multiplicity roots of  $p_1$  and  $p_3$  coincide if and only if  $a_1a_2 + a_3 - a_2x + a_3x^2$  has no simple roots, i.e.,

$$a_2^2 - 4a_1a_2a_3 - 4a_3^2 \leq 0.$$

The converse is obvious.

CASE 2. Assume that  $p_3$  has degree one and  $p_1$  has degree three, or equivalently

$$\begin{aligned} a_2 &= a_1a_6, \quad 2a_3 + a_4 + a_1(3a_2 + a_5) + a_6 = 0, \\ a_6 &\neq 0, \quad 3a_2 - a_1(3a_3 + a_4) + a_5 \neq 0. \end{aligned}$$

Assume that the odd-multiplicity real roots of  $p_1$  and  $p_3$  coincide. From  $a_2 = a_1a_6$ ,  $a_4 = -2a_3 - a_1a_5 - a_6 - 3a_1^2a_6$ , we obtain

$$p_3(x) = a_3 + a_1^2a_6 - (a_5 + a_1(-a_3 + a_1a_5 + 4a_6 + 3a_1^2a_6))x.$$

where  $a_5 + a_1(-a_3 + a_1a_5 + 4a_6 + 3a_1^2a_6) \neq 0$ . Therefore,  $p_3$  has the unique root

$$x_0 = \frac{a_3 + a_1^2a_6}{a_5 + a_1(-a_3 + a_1a_5 + 4a_6 + 3a_1^2a_6)}.$$

As  $p_1$  and  $p_3$  have the same odd-multiplicity real roots,  $x_0$  must be a root of  $p_1$ . Substituting, one has

$$p_1(x_0) = \frac{-R_2((a_5 + 4a_1a_6)^2 + (a_3 - a_1a_5 - 3a_1^2a_6)^2)}{a_6^2(a_5 + a_1(-a_3 + a_1a_5 + 4a_6 + 3a_1^2a_6))^3}.$$

Since  $a_5 - a_1(a_3 - a_1a_5 - 4a_6 - 3a_1^2a_6) \neq 0$ , then  $(a_5 + 4a_1a_6)^2 + (a_3 - a_1a_5 - 3a_1^2a_6)^2 > 0$ . Therefore  $x_0$  is a root of  $p_1$  if and only if  $R_2 = 0$ .

If  $D_1 < 0$ , we shall prove that  $R_{113} \neq 0$ , so that (4.11) holds. Assume by contradiction that  $R_{113} = 0$ . Consider the ideal generated by  $D_1 + x^2$  (which implies  $D_1 < 0$  if  $x \neq 0$ ),  $a_2 - a_1a_6$ ,  $2a_3 + a_4 + a_1(3a_2 + a_5) + a_6$ , and  $R_{113}$ . This ideal has three associated primes (see Appendix A - the computations take some time in this case). The first one contains the polynomial  $x$ , so that it corresponds to  $D_1 = 0$ . The second contains the polynomial  $3a_2 - a_1(3a_3 + a_4) + a_5$ . The third contains  $1 + a_1^2$  so that it has no real points. Therefore, the variety of the ideal is contained in  $3a_2 - a_1(3a_3 + a_4) + a_5 = 0$ . But  $3a_2 - a_1(3a_3 + a_4) + a_5 \neq 0$  by hypothesis. This contradiction proves that  $R_{113} \neq 0$ .

If  $D_1 = 0$ , the multiplicity of  $x_0$  as a root of  $p_1$  must be one or three. The multiplicity is two or more if and only if  $p_1'(x_0) = 0$ , but

$$R_{113} = 9a_6^2 (a_5 + a_1(-a_3 + a_1a_5 + 4a_6 + 3a_1^2a_6))^2 p_1'(x_0).$$

I.e., the multiplicity is one if and only if  $R_{113} \neq 0$ . Finally, if the multiplicity is three, then  $p_1(x) = -a_6(x - a)^3$  for a certain  $a$ . We consider the ideal generated by  $R_2$ ,  $a_2 - a_1a_6$ ,  $2a_3 + a_4 + a_1(3a_2 + a_5) + a_6$ , and the coefficients of  $p_1(x) + a_6(x - a)^3$ . Eliminating  $a$ , and computing the minimal associated primes, we obtain three ideals. The first one contains  $1 + a_1^2$  so that it has no real points in its variety. The second contains the polynomial  $a_6$ , and, since by hypothesis  $a_6 \neq 0$ , it has no real points in its variety. The third is

$$\begin{aligned} 8a_5^2 - 81a_6^2 &= 4a_4 - 9a_6 = 9a_1a_6 + a_5 = 8a_1a_5 + 9a_6 = 8a_1^2 - 1 = 0, \\ 3a_1^2a_6 + a_1a_5 + 2a_3 + a_4 + a_6 &= -a_1a_6 + a_2 = 0. \end{aligned}$$

Computing a minimal system of generators, we obtain

$$(4.13) \quad 4a_4 - 9a_6 = 4a_3 + 5a_6 = 9a_2 + a_5 = 9a_1a_6 + a_5 = 8a_1^2 - 1 = 0.$$

To conclude, note that if (4.13) holds then

$$p_1(x) = -a_6 \left( x \pm \frac{1}{\sqrt{2}} \right)^3, \quad p_3(x) = \frac{9\sqrt{2}}{8} a_6 \left( x \pm \frac{1}{\sqrt{2}} \right).$$

□

Now, we consider that either  $p_1$  or  $p_3$  has degree two (and the other degree is two or more).

**Proposition 4.4.** *Assume that the minimum of the degrees of  $p_1$  and  $p_3$  is two. Then the odd-multiplicity real roots of  $p_1, p_3$  coincide if and only if*

$$(4.14) \quad a_2 = a_6 = 3a_3 + a_4 = 0, \quad 4a_3^2 - 4a_1a_3a_5 \geq a_5^2 > 0, \quad a_3 - a_1a_5 \neq 0.$$

*Proof.* Assume that the minimum of the degrees of  $p_1$  and  $p_3$  is two and the real odd-multiplicity roots of  $p_1, p_3$  coincide. Note that this implies that

they are both of degree two. I.e.,  $a_6 = a_2 = 0$ ,  $a_5 \neq 0$ ,  $2a_3 + a_4 + a_1a_5 \neq 0$ , and

$$\begin{aligned} p_1(x) &= (3a_3 + a_4)x - a_5x^2, \\ p_3(x) &= a_3 + (a_1(3a_3 + a_4) - a_5)x - (2a_3 + a_4 + a_1a_5)x^2. \end{aligned}$$

The roots of  $p_1$  are then  $x_1 = 0$  and  $x_2 = (3a_3 + a_4)/a_5$ .

Assume that  $3a_3 + a_4 \neq 0$ . As the simple real roots of  $p_1$  must be roots of  $p_3$ , we have that  $p_3(0) = 0$  which implies  $a_3 = 0$ . Moreover, evaluating  $p_3$  at  $x_2$ , we obtain

$$p_3\left(\frac{3a_3 + a_4}{a_5}\right) = -\frac{a_4(a_4^2 + a_5^2)}{a_5^2} = 0.$$

I.e.,  $a_4 = 0$ . But this is contradictory with  $3a_3 + a_4 \neq 0$ .

If  $3a_3 + a_4 = 0$  then  $p_1(x) = -a_5x^2$  has no odd-multiplicity real roots. The discriminant of  $p_3$ , replacing  $a_4$  by  $-3a_3$ , is

$$\text{disc}(p_3) = -4a_3^2 + 4a_1a_3a_5 + a_5^2,$$

so that  $p_3$  has no simple real roots if and only if  $4a_3^2 - 4a_1a_3a_5 - a_5^2 \geq 0$ . Finally, note that if  $3a_3 + a_4 = 0$  then the condition  $2a_3 + a_4 + a_1a_5 \neq 0$  is equivalent to  $a_3 - a_1a_5 \neq 0$ .

Conversely, assume that (4.14) holds. Then  $p_1(x) = -a_5x^2$  and  $p_3(x) = a_3 - a_5x + (a_3 - a_1a_5)x^2$ . Since  $\text{disc}(p_3) < 0$ , both  $p_1$  and  $p_3$  have no odd real roots.  $\square$

In the remainder of this subsection, we shall consider that both  $p_1$  and  $p_3$  have degree three. In this case, the number of real odd-multiplicity roots is given by the discriminant, being three if the discriminant is strictly positive and one if the discriminant is negative. Note that if  $D_1 \leq 0$  and  $D_3 > 0$ , or  $D_1 > 0$  and  $D_3 \leq 0$ , then the odd-multiplicity roots of  $p_1, p_3$  do not coincide since one has three simple roots and the other has one root with odd-multiplicity. Consequently, we only need to consider the cases  $D_1, D_3 > 0$  or  $D_1, D_3 \leq 0$ .

Firstly, we consider the case when  $p_1, p_3$  have three simple roots, for which we prove that the real roots can not coincide. The following result is a little more general since we do not impose the condition that the real roots be simple. It will be used in proving other cases.

**Proposition 4.5.** *If  $p_1, p_3$  have three real roots then the roots do not coincide (with multiplicity).*

*Proof.* The polynomials  $p_1, p_3$  have three real roots if and only if  $a_6, a_2 - a_1a_6 \neq 0$  and their discriminants are positive.

The three real roots of  $p_1, p_3$  coincide (with multiplicity) if and only if there exists  $\lambda \in \mathbb{R}$  such that  $p_1(x) = \lambda p_3(x)$ . Equating the coefficients of the leading term, one obtains

$$\lambda = \frac{a_6}{-a_2 + a_1a_6}.$$

Replacing  $\lambda$  in the rest of the equations yields the system (we have multiplied by  $a_2 - a_1a_6 \neq 0$ )

$$a_2^2 + a_3a_6 = a_2(3a_3 + a_4 - 3a_6) - a_5a_6 = a_2(3a_2 + a_5) + a_6(2a_3 + a_4 + a_6) = 0.$$

Solving this, one obtains (note that it is a staggered solution)

$$a_3 = \frac{-a_2^2}{a_6}, \quad a_5 = \frac{a_2(a_4a_6 + 3a_2 - 3a_2^2)}{a_6^2}, \quad a_4 = \frac{3a_2^2 - a_6^2}{a_6}.$$

Substituting in  $D_1$  gives  $D_1 = -4(a_2^2 + a_6^2)^2 < 0$ , in contradiction with  $p_1$  having three real roots.  $\square$

Recall that  $\text{res}(p_1, p_3)$  factorizes as the product of two polynomials,  $R_1, R_2$ . We shall prove that if  $p_1, p_3$  have a real root in common then  $R_2$  must vanish.

**Lemma 4.6.** *Assume  $a_2 - a_1a_6, a_6 \neq 0$ .  $p_1, p_3$  have a real root in common if and only if  $a = (a_1, \dots, a_6) \in \mathcal{V}(R_2)$ .*

*Proof.* If  $p_1, p_3$  have a real root in common, then  $\text{res}(p_1, p_3) = R_1R_2 = 0$ . Hence  $R_1 = 0$  or  $R_2 = 0$ . Assume that  $R_1 = (4a_2 + a_5)^2 + (3a_3 + a_4 + a_6)^2 = 0$ , i.e.,  $a_5 = -4a_2$  and  $a_6 = -3a_3 - a_4$ . Then

$$\begin{aligned} p_1(x) &= (a_2 + (3a_3 + a_4)x)(1 + x^2), \\ p_3(x) &= (a_1a_2 + a_3 + (a_2 + 3a_1a_3 + a_1a_4)x)(1 + x^2). \end{aligned}$$

Therefore,  $p_1, p_3$  have a real root in common if and only if  $a_2^2 - 3a_3^2 - a_3a_4 = 0$ . Since  $R_1 = (4a_2 + a_5)^2 + (3a_3 + a_4 + a_6)^2 = 0$  then

$$R_2 = (a_2^2 - 3a_3^2 - a_3a_4)(4a_2^2 + (2a_3 + a_4)^2).$$

Thus, the real root coincide if and only if  $R_2 = 0$ .  $\square$

Next, we study the singular points of the variety defined by  $R_2$ . We shall show that they are the intersection of the variety with the hyperplane  $a_3 = a_6$ . Moreover, in the intersection, the odd-multiplicity real roots of  $p_1$  and  $p_3$  do not coincide.

**Lemma 4.7.** *The point  $a = (a_1, a_2, \dots, a_6) \in \mathcal{V}(R_2)$  is singular if and only if  $a_3 = a_6$  or  $a_2 = 2a_3 + a_4 = 0$ .*

*Moreover, if  $a \in \mathcal{V}(R_2)$  is singular, then the real odd-multiplicity roots of  $p_1, p_3$  do not coincide.*

*Proof.* The variety of singular points of  $\mathcal{V}(R_2)$  is defined by  $\mathcal{V}(\langle R_2, \nabla R_2 \rangle)$ . It has two minimal associated prime ideals (see the SINGULAR code in Appendix A),

$$(4.15) \quad \langle 2a_2^2 + a_2a_5 + a_4a_6 + 2a_6^2, a_3 - a_6 \rangle \quad \text{and} \quad \langle 2a_3 + a_4, a_2 \rangle.$$

If  $a_3 = a_6$ , then  $R_2 = (2a_2^2 + a_2a_5 + a_4a_6 + 2a_6^2)^2$ . Hence,

$$R_2 = 0, a_3 = a_6 \quad \text{if and only if} \quad R_2 = 0, \nabla R_2 = 0.$$

Let  $a \in \mathcal{V}(\langle R_2, a_3 - a_6 \rangle)$ . Then, parametrizing the variety by  $a_1, a_2, a_3, a_4$ , we obtain

$$p_1(x) = \frac{(a_3x + a_2)(a_2 + (2a_3 + a_4)x - a_2x^2)}{a_2},$$

$$p_3(x) = \frac{(a_2 + (2a_3 + a_4)x - a_2x^2)(a_3 - a_2x + a_1(a_2 + a_3x))}{a_2}.$$

I.e.,  $p_1, p_3$  have three real roots (as the quadratic factor has positive discriminant), and by Proposition 4.5 they do not coincide.

Finally, let  $a \in \mathcal{V}(\langle a_2, 2a_3 + a_4 \rangle)$ . Then

$$p_1(x) = x(a_3 - a_5x - a_6x^2), \quad p_3(x) = (1 + a_1x)(a_3 - a_5x - a_6x^2).$$

Since  $x = 0$  has different parity as root of  $p_1$  than it does as root of  $p_3$ , they do not have the same odd-multiplicity real roots.  $\square$

The next proposition considers the case of  $p_1$  and  $p_3$  having a unique simple real solution.

**Proposition 4.8.** *Assume  $p_1$  and  $p_3$  have one simple root and two complex conjugate roots. Then  $p_1$  and  $p_3$  have the same odd-multiplicity real root if and only if*

$$(4.16) \quad \begin{aligned} R_2 = 0, \quad D_1 < 0, \quad D_3 < 0, \\ a_6 \neq 0, \quad a_2 - a_1a_6 \neq 0, \quad a_3 \neq a_6, \quad a_2^2 + (a_4 + 2a_3)^2 \neq 0. \end{aligned}$$

*Proof.* If  $p_1$  and  $p_3$  have the same real root then  $R = 0$  and, by Lemma 4.6,  $R_2 = 0$ . Moreover, applying Lemma 4.7,  $a_3 \neq a_6$ , and either  $a_2 \neq 0$  or  $a_4 + 2a_3 \neq 0$ .

Conversely, suppose  $R_2 = 0$ ,  $a_3 \neq a_6$ , and either  $a_2 \neq 0$  or  $a_4 + 2a_3 \neq 0$ . We have to prove that the real root of  $p_1$  coincides with that of  $p_3$ .

Assume on the contrary that these real roots do not coincide. In that case, the complex conjugate roots of  $p_1$  and  $p_3$  must coincide. Then there exist some  $a, a', b, d \in \mathbb{R}$  such that

$$(4.17) \quad \begin{aligned} p_1(x) &= -a_6(x - a)((x - b)^2 + d^2), \\ p_3(x) &= (-a_1a_6 + a_2)(x - a')((x - b)^2 + d^2). \end{aligned}$$

Equating the coefficients, eliminating the variables  $a, a', b, d$ , and computing the minimal associated prime ideals (see Appendix A), we obtain the ideals in (4.15) (which do not satisfy that  $a_3 \neq a_6$ , and either  $a_2 \neq 0$  or  $a_4 + 2a_3 \neq 0$ ), and an ideal such that one of its generators is  $R_1$ . By Lemma 4.6, we conclude.  $\square$

The last case is  $p_1, p_3$  of degree three with a unique odd-multiplicity real root, and possible double roots.

**Proposition 4.9.** *Assume  $p_1, p_3$  have degree three (i.e.,  $a_6(a_2 - a_1a_6) \neq 0$ ) and one of them has a root of multiplicity two or more. Then  $p_1$  and  $p_3$*



have the same odd-multiplicity real root if and only if  $R_2 = 0$  and one of the following statements holds:

$$(4.18) \quad D_1 = 0, D_3 < 0, D'_1 \neq 0, R_{113} \neq 0,$$

$$(4.19) \quad D_1 = D'_1 = 0, D_3 < 0,$$

$$(4.20) \quad D_3 = 0, D_1 < 0, D'_3 \neq 0, R_{133} \neq 0,$$

$$(4.21) \quad D_3 = D'_3 = 0, D_1 < 0,$$

$$(4.22) \quad D_1 = D'_1 = D_3 = 0, R_{133} \neq 0,$$

$$(4.23) \quad D_1 = D_3 = D'_3 = 0, R_{113} \neq 0,$$

$$(4.24) \quad D_1 = D_3 = 0, D'_1 \neq 0, D'_3 \neq 0, \bar{R}_{113} \neq 0, \bar{R}_{133} \neq 0,$$

*Proof.* Since  $p_1, p_3$  have degree three, then  $a_6 \neq 0, a_2 - a_1 a_6 \neq 0$ . By Lemma 4.6,  $p_1, p_3$  have a real root in common if and only if  $R_2 = 0$ . In the following, we shall assume this to be the case.

Assume that  $p_1$  has a root  $x_1$  of multiplicity two or more, and that  $p_3$  has a simple real root,  $x_3$ , and two complex conjugate roots, i.e.,  $D_1 = 0, D_3 < 0$ . The multiplicity of  $x_1$  is two if and only if  $D'_1 \neq 0$ , and is three if and only if  $D'_1 = 0$ . In the former case of  $D'_1 \neq 0$ ,  $p_1$  has a simple root  $\bar{x}_1 \neq x_1$ . Therefore  $p_1, p_3$  have the same odd-multiplicity real roots if and only if  $x_3 = \bar{x}_1$ . As  $R_2 = 0$ , then either  $x_3 = \bar{x}_1$  or  $x_1 = \bar{x}_1$ . Moreover,  $x_1$  is a root of  $p'_1$ , while  $\bar{x}_1$  is not, so that  $x_3 = \bar{x}_1$  if and only if  $R_{113} \neq 0$ . In the latter case of  $D'_1 = 0$ ,  $x_1$  is the unique real root of  $p_1$  with multiplicity three, and, as  $R_2 = 0$ ,  $x_1 = x_3$ , so that the odd-multiplicity real roots of  $p_1, p_3$  coincide.

Assume that  $p_3$  has a root of multiplicity two or more, and  $p_1$  has a simple real root and two complex conjugate roots. Arguing analogously, we obtain that the odd-multiplicity real roots of  $p_1, p_3$  coincide if and only if (4.20) or (4.21) hold.

Assume that  $p_1$  and  $p_3$  have a root of multiplicity two or more, i.e.,  $D_1 = D_3 = 0$ . Firstly, by Proposition 4.5, if both  $p_1$  and  $p_3$  have a root of multiplicity three, then it can not be common.

If  $D'_1 = 0$ , then  $p_1$  has a triple root. As  $R_2 = 0$ , this root coincides with one of the roots of  $p_3$ . If  $R_{133} \neq 0$ , then it coincides with a simple root of  $p_3$ , and in any other case ( $R_{133} = 0$  and  $D'_3 \neq 0$ ), it coincides with the double root of  $p_3$ .

Analogously, if  $D'_3 = 0$ , then the triple root of  $p_3$  coincides with the odd-multiplicity real root of  $p_1$  if and only if  $R_{113} \neq 0$ .

If  $D'_1, D'_3 \neq 0$ , then  $p_1$  and  $p_3$  have a root of multiplicity two and a simple root. In this case, the greatest common divisor of  $p_1$  and  $p'_1$  is  $r_1$ , a degree-one polynomial, so that  $\bar{R}_{113}$  is zero if and only if the double root of  $p_1$  is a root of  $p_3$ . Analogously,  $\bar{R}_{133}$  is zero if and only if the double root of  $p_3$  is a

root of  $p_1$ . By Proposition 4.5, if  $p_1, p_3$  have a double root in common, then their simple root is distinct. So  $p_1, p_3$  have the same simple root if and only if  $\bar{R}_{113} \neq 0$  and  $\bar{R}_{133} \neq 0$ .  $\square$

Finally, we compute examples of points for some of the semi-varieties and their dimensions.

**Proposition 4.10.** *The codimensions of the semi-varieties defined by conditions of Theorem A are the following:*

- 5a) has codimension one.
- 5b), 5d) have codimension two.
- 2), 3a), 4), 5f) have codimension three.
- 1a), 1b) have codimension four.
- 3b) has codimension five.
- 5f) has codimension two or three.
- 5c), 5e), 5g), 5h) have codimension of at least two.

*Proof.* In Table 1 we give one point in each of the semi-varieties, such that if the definition of the semi-variety contains inequalities then the inequalities hold strictly.

In the same table, we include the codimension of the tangent space of the semi-variety at that point,  $c_p$ . To obtain it, we compute the rank of the Jacobian matrix of the equations (equalities) defining the semi-variety at that point. If the rank is maximum (the point is regular), then it coincides with the codimension of the variety at that point. (We set it to \* if the point is not singular.)

Finally,  $c_I$  denotes the (Krull) codimension of the defining ideal  $I$  of the smallest variety containing the corresponding semi-variety (i.e., considering the ideal generated only by the polynomials of the equalities). In symbols,  $c_I = \text{codim}(\mathcal{V}_{\mathbb{C}}(I)) := n - \dim(I)$ , where  $n$  is the number of indeterminates in the base ring (see Appendix A). By Theorem 3.9  $c_p \leq c_P$ , where  $P$  is a minimal prime of  $I$  vanishing at  $p$  and the equality holds if the point is regular. Therefore, since the dimension of  $I$  is the maximum of the dimensions of its associated prime ideals, if  $c$  denotes the (real) codimension of the variety, then  $c_p \geq c \geq c_I$  at the regular points.  $\square$

**4.2. Proof of Theorem B.** The trigonometric polynomial  $B(\theta)$  has definite sign if and only if  $q(x) \equiv 0$ . I.e., the parameters belong to the variety defined by the ideal obtained by equating the coefficients of  $q(x)$  to zero:

$$\begin{aligned} 2a_1a_2 + 4a_3 + a_4 &= 0, \\ 12a_2 + 3a_5 - a_1(6a_3 + 2a_4) &= 0, \\ 8a_3 + 3a_4 + 4a_6 + a_1(6a_2 + 2a_5), \\ 4a_2 + a_5 - 2a_1a_6 &= 0. \end{aligned}$$

Case	Point	$c_p$	$c_I$
1a)	$a_1 = 1, a_2 = 0, a_3 = 1, a_4 = -3, a_5 = 0, a_6 = 0.$	4	4
1b)	$a_1 = 1, a_2 = 1, a_3 = -1, a_4 = 2, a_5 = -4, a_6 = 1.$	4	4
2)	$a_1 = -1, a_2 = \sqrt{14}, a_3 = -2,$ $a_4 = -1, a_5 = -3\sqrt{14}, a_6 = 0.$	3	3
3a)	$a_1 = -1, a_2 = (201 + 2\sqrt{1509})/58,$ $a_3 = (-33 + 4\sqrt{1509})/58, a_4 = -1,$ $a_5 = -16, a_6 = (-201 - 2\sqrt{1509})/58)$	3	3
3b)	$a_1 = 0, a_2 = 0, a_3 = 0, a_4 = 1, a_5 = -2, a_6 = -1.$	5	5
4)	$a_1 = 1, a_2 = 0, a_3 = 1/3, a_4 = -1, a_5 = -1, a_6 = 0.$	3	3
5a)	$a_1 = 0, a_2 = 1, a_3 = -15/16,$ $a_4 = -53/16, a_5 = (-941 - 31\sqrt{7913})/512, a_6 = 1.$	1	1
5b)	$a_1 = 0, a_2 = (4096 - 7\sqrt{1726})/16384, a_3 = 0,$ $a_4 = -(58339673 + 28672\sqrt{1726})/94666752,$ $a_5 = -1, a_6 = -2889/16384.$	2	2
5c)	$a_1 = 1, a_2 = 4, a_3 = -12, a_4 = 30, a_5 = -15, a_6 = 1/2$	*	2
5d)	$a_1 = 0, a_2 = \sqrt{185}/32, a_3 = 0,$ $a_4 = -1, a_5 = -3\sqrt{185}/32, a_6 = -5/32$	2	2
5e)	$a_1 = 0, a_2 = 2\sqrt{2}, a_3 = -1, a_4 = 0, a_5 = -9\sqrt{2}, a_6 = 8$	*	2
5f)	$a_1 = 0, a_2 = 2/3, a_3 = 0, a_4 = -1, a_5 = -2, a_6 = -1/3$	3	2
5g)	$a_1 = 0, a_2 = 1, a_3 = -9/2, a_4 = 15/2, a_5 = -15, a_6 = 8$	*	2
5h)	$a_1 = 0, a_2 = 1, a_3 = -8, a_4 = 35/2, a_5 = -15, a_6 = 9/2$	*	2

TABLE 1. Codimensions of the semi-varieties.

Computing the minimal associated prime ideals and a minimal set of generators (see Appendix A), we obtain three minimal ideals. But the first one contains the polynomial  $a_1^2 + 4$ , so that the associated variety is empty. The other two prime ideals obtained are

$$\langle a_1, 4a_2 + a_5, a_3 - a_6, a_4 + 4a_6 \rangle,$$

and

$$\langle a_6, 3a_3 + a_4, 4a_2 + a_5, 3a_1a_5 + 2a_4 \rangle.$$

## APPENDIX A. SINGULAR CODES

```
// Proposition 4.1;
LIB "primdec.lib";
ring r = 0, (a1,a2,a3,a4,a5,a6,x), dp;
poly p1 = -a6*x^3 - 3*a2*x^2 - a5*x^2 + 3*a3*x + a4*x + a2;
poly p2 = -a2*x^3 + 2*a3*x^2 + a4*x^2 + a6*x^2 + 3*a2*x + a5*x - a3;
poly p3 = a1*p1-p2;
ideal i1 = coeffs(p1,x);
ideal i3 = coeffs(p3,x);
mres(i1,1)[1];
```

```

mres(i3,1)[1];
// Proposition 4.3 Case 2;
// D1<0 implies R113!=0;
LIB "primdec.lib";
ring r = 0, (a1,a2,a3,a4,a5,a6,x), dp;
poly p1 = -a6*x^3-(3*a2+a5)*x^2+(3*a3+a4)*x+a2;
poly p2 = -a2*x^3+(2*a3+a4+a6)*x^2+(3*a2+a5)*x-a3;
poly p3 = a1*p1-p2;
ideal R = resultant(p1,p2,x);
poly R2 = minAssGTZ(R)[1][1];
poly dp1 = diff(p1,x);
ideal j1 = coeffs(p3,x)[4,1],coeffs(p3,x)[3,1], resultant(dp1,p3,x);
ideal j = R2, resultant(dp1,p1,x)+x^2, j1;
j = sat(j,a6)[1];
list l = minAssGTZ(j); // Takes some time
reduce(x,std(l[1]));
reduce(coeffs(p3,x)[2,1],std(l[2]));
reduce(1+a1^2,std(l[3]));
// Proposition 4.3 Case 2;
// p1 with a root of multiplicity three;
LIB "primdec.lib";
ring r = 0, (a1,a2,a3,a4,a5,a6,x,a), dp;
poly p1 = -a6*x^3 - 3*a2*x^2 - a5*x^2 + 3*a3*x + a4*x + a2;
poly p2 = -a2*x^3 + 2*a3*x^2 + a4*x^2 + a6*x^2 + 3*a2*x + a5*x - a3;
poly p3 = a1*p1-p2;
poly R2 = minAssGTZ(resultant(p1,p2,x))[1][1];
poly p1d = p1 + a6*(x-a)^3;
ideal i3 = coeffs(p3,x);
poly p3l = i3[3]*x+i3[4];
ideal i1d = coeffs(p1d,x);
ideal i3d = coeffs(p3l,x);
ideal i13 = i1d,i3d,R2;
ideal ie=eliminate(i13,a);
list J=minAssGTZ(ie);
mres(J[3],1)[1];
// Lemma 4.7;
LIB "primdec.lib";
ring r = 0, (a1,a2,a3,a4,a5,a6,x), dp;
poly p1 = -a6*x^3 - 3*a2*x^2 - a5*x^2 + 3*a3*x + a4*x + a2;
poly p2 = -a2*x^3 + 2*a3*x^2 + a4*x^2 + a6*x^2 + 3*a2*x + a5*x - a3;
poly R2 = minAssGTZ(resultant(p1,p2,x))[1][1];
ideal sR2 = R2,jacob(R2);
minAssGTZ(sR2);
// Proposition 4.8;

```

```

LIB "primdec.lib";
ring r = 0, (a1,a2,a3,a4,a5,a6,x,a,ap,b,d), dp;
poly p1 = -a6*x^3 - 3*a2*x^2 - a5*x^2 + 3*a3*x + a4*x + a2;
poly p2 = -a2*x^3 + 2*a3*x^2 + a4*x^2 + a6*x^2 + 3*a2*x + a5*x - a3;
poly p3 = a1*p1-p2;
poly R2 = minAssGTZ(resultant(p1,p2,x))[1][1];
poly p1d = p1 + a6*(x-a)*((x-b)^2-d^2);
poly p3d = p3 + (a1*a6-a2)*(x-ap)*((x-b)^2-d^2);
ideal i1d = coeffs(p1d,x);
ideal i3d = coeffs(p3d,x);
ideal i13 = i1d,i3d,R2;
ideal ie=eliminate(i13,a*ap*b*d);
list J=minAssGTZ(ie);

// Proposition 4.10;
LIB "primdec.lib";
ring r = 0, (a1,a2,a3,a4,a5,a6,x), dp;
poly p1 = -a6*x^3 - 3*a2*x^2 - a5*x^2 + 3*a3*x + a4*x + a2;
poly p2 = -a2*x^3 + 2*a3*x^2 + a4*x^2 + a6*x^2 + 3*a2*x + a5*x - a3;
poly p3 = a1*p1-p2;
poly dp1 = diff(p1,x);
poly dp3 = diff(p3,x);
poly ddp1 = diff(dp1,x);
poly ddp3 = diff(dp3,x);
poly D1 = resultant(p1,dp1,x);
poly D1p = resultant(p1,ddp1,x);
poly D3 = resultant(p3,dp3,x);
poly D3p = resultant(p3,ddp3,x);
poly R2 = minAssGTZ(resultant(p1,p2,x))[1][1];
ideal i1a = a6,a5,3*a3+a4,a2;
ideal i1b = a1*a6-a2,a1*a5-a3+a4+a6,a1*(3*a3+a4)-3*a2-a5,a1*a2+a3;
ideal i2 = a6,3*a2+a5,R2;
ideal i3a = a2-a1*a6,2*a3+a4+a1*(3*a2+a5)+a6,R2;
ideal i3b = 4*a4-9*a6,4*a3+5*a6,9*a2+a5,9*a1*a6+a5,8*a1^2-1;
ideal i4 = a2,a6,3*a3+a4;
ideal i5a = R2;
ideal i5b = R2,D1;
ideal i5c = R2,D1,D1p;
ideal i5d = R2,D3;
ideal i5e = R2,D3,D3p;
ideal i5f = R2,D1,D3;
ideal i5g = R2,D1,D1p,D3;
ideal i5h = R2,D1,D3p,D1;
nvars(basering) - dim(std(i1a));
nvars(basering) - dim(std(i1b));

```

```

nvars(basering) - dim(std(i2));
nvars(basering) - dim(std(i3a));
nvars(basering) - dim(std(i3b));
nvars(basering) - dim(std(i4));
nvars(basering) - dim(std(i5a));
nvars(basering) - dim(std(i5b));
nvars(basering) - dim(std(i5c));
nvars(basering) - dim(std(i5d));
nvars(basering) - dim(std(i5e)); // Takes some time;
nvars(basering) - dim(std(i5f));
nvars(basering) - dim(std(i5g));
nvars(basering) - dim(std(i5h)); // Takes some time;

// Theorem B
LIB "primdec.lib";
ring r = 0, (a1,a2,a3,a4,a5,a6), dp;
poly c0 = 2*a1*a2 + 4*a3 + a4;
poly c1 = 12*a2 + 3*a5 - a1*(6*a3 + 2*a4);
poly c2 = 8*a3 + 3*a4 + 4*a6 + a1*(6*a2+2*a5);
poly c3 = 4*a2 + a5 - 2*a1*a6;
ideal iB = c0,c1,c2,c3;
list LB = minAssGTZ(iB);
mres(LB[1],1)[1];
mres(LB[2],1)[1];
mres(LB[3],1)[1];

```

#### ACKNOWLEDGMENTS

The first two authors were partially supported by AEI/FEDER UE grant number MTM 2011-22751 and Junta de Extremadura grant GR15055 (Junta de Extremadura/FEDER funds). The third author was partially supported by the research group FQM-024 (Junta de Extremadura/FEDER funds) and by the project MTM2015-65764-C3-1-P (MINECO/FEDER, UE). The fourth author was partially supported by Junta de Extremadura grant GR15055 (Junta de Extremadura/FEDER funds).

#### REFERENCES

- [1] M.J. Álvarez, A. Gasull, H. Giacomini, *A new uniqueness criterion for the number of periodic orbits of Abel equations*, J. Differential Equations **234**, (2007), 161–176.
- [2] M.A.M. Alwash, N.G. Lloyd, *Nonautonomous equations related to polynomial two-dimensional systems*, Proc. Roy. Soc. Edinburgh Sect. A **105**, (1987), 129–152.
- [3] A.A. Andronov, E.A. Leontovich, I.I. Gordon, A.G. Maier, *Qualitative theory of second-order dynamic systems*. A Halsted Press Book. New York etc.: John Wiley & Sons; Jerusalem-London: Israel Program for Scientific Translations, 1973.
- [4] O. Bachmann, G.-M. Greuel, C. Lossen, G. Pfister, H. Schönemann. *A Singular introduction to commutative algebra*, Springer, Berlin, 2007.

- [5] N.N. Bautin, *On the number of limit cycles which appear with the variation of the coefficients from an equilibrium position of focus or centre type*, Math. USSR-Sb. **100** (1954), 397–413.
- [6] J.L. Bravo, M. Fernández, A. Gasull, *Limit cycles for some Abel equations having coefficients without fixed signs*, Internat. J. Bifur. Chaos Appl. Sci. Engrg. **19**, (2009), 3869–3876.
- [7] J.L. Bravo, J. Torregrosa, *Abel-like equations with no periodic solutions*, J. Math. Anal. Appl. **342**, (2008), 931–942.
- [8] L.A. Cherkas, *Number of limit cycles of an autonomous second-order system*, Diff. Eq. **5**, (1976) 666–668.
- [9] B. Coll, A. Gasull and J. Llibre, *Some theorems on the existence, uniqueness and non existence of limit cycles for quadratic systems*, J. Differential Equations **67** (1987), 372–399.
- [10] D. Cox, J. Little, D. O’Shea. *Ideals, varieties, and algorithms. An introduction to computational algebraic geometry and commutative algebra*. Second edition. Undergraduate Texts in Mathematics. Springer-Verlag, New York, 1997.
- [11] W. Decker, G.M. Greuel, G. Pfister, H. Schönemann. *Singular 4-1-0* — A computer algebra system for polynomial computations. <http://www.singular.uni-kl.de> (2016).
- [12] G.F.D. Duff, *Limit-cycles and rotated vector fields*, Ann. of Math. **57**, (1953), 15–31.
- [13] H. Dulac, *Détermination et intégration d’une certaine classe d’équations différentielles ayant pour point singulier un centre*, Bull. Soc. Math. France **32** no. 2 (1908), 230i;  $\frac{1}{2}$ -252.
- [14] A. Gasull, A. Guillamon, *Limit cycles for generalized Abel equations*, Internat. J. Bifur. Chaos Appl. Sci. Engrg. **16**, (2006), 3737–3745.
- [15] A. Gasull, J. Llibre, *Limit cycles for a class of Abel equations*, SIAM J. Math. Anal. **21-5**, (1990), 1235–1244.
- [16] Jianfeng Huang, Yulin Zhaob, *Periodic solutions for equation  $x' = A(t)x^m + B(t)x^n + C(t)x^l$  with  $A(t)$  and  $B(t)$  changing signs*, J. Differential Equations **253** (2012), 73–99.
- [17] A. Lins Neto, *On the number of solutions of the equation  $\frac{dx}{dt} = \sum_{j=0}^n b_j(t)x^j$ ,  $0 \leq t \leq 1$ , for which  $x(0) = x(1)$* , Inv. Math. **59**, (1980), 67–76.
- [18] J. Llibre, Xiang Zhang, *Non-existence and uniqueness of limit cycles for quadratic polynomial differential systems*, preprint.
- [19] N.G. Lloyd, *A note on the number of limit cycles in certain two-dimensional systems*, J. London Math. Soc. **20**, (1979), 277–286.
- [20] D. Mumford. *Algebraic Geometry I: Complex Projective Varieties*, Reprint of the 1976 edition, Classics in Mathematics, Springer-Verlag, Berlin, 1995
- [21] A.A. Panov. *The number of periodic solutions of polynomial differential equations*, Math. Notes **64**, (1998), 622–628.
- [22] L.M. Perko, *Differential equations and dynamical systems*, New York [etc.]: Springer-Verlag, 2001, 3rd ed.
- [23] V.A. Pliss, *Non-local problems of the theory of oscillations*, Academic Press, New York, 1966.
- [24] V. G. Romanovski, D. S. Shafer. *The centre and cyclicity problems. A computational algebra approach*. Birkh $\ddot{u}$ ser, 2009.
- [25] J. Sotomayor, *Curvas definidas por equações diferenciais no plano*, IMPA, Rio de Janeiro, 1981.

J.L. BRAVO, DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE EXTREMADURA,  
06006 BADAJOZ, SPAIN

*E-mail address:* trinidad@unex.es

M. FERNÁNDEZ, DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE EXTREMADURA,  
06006 BADAJOZ, SPAIN

*E-mail address:* `ghierro@unex.es`

I. OJEDA, DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE EXTREMADURA, 06006  
BADAJOZ, SPAIN

*E-mail address:* `ojedamc@unex.es`

F.SÁNCHEZ, DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE EXTREMADURA,  
06006 BADAJOZ, SPAIN

*E-mail address:* `fsanchez@unex.es`