

## P202. Biocompounds recovery from *Spirulina* by conventional and ohmic heating methodologies: chemical and biological properties

Pedro Ferreira-Santos<sup>1</sup>, Rafaela Nunes<sup>1</sup>, Filomena De Biasio<sup>2</sup>, Cristina M.R. Rocha<sup>1</sup>, Domenico Gorgoglione<sup>3</sup>, Giorgia Spigno<sup>4</sup>, José A. Teixeira<sup>1</sup>

<sup>1</sup> CEB - Centre of Biological Engineering, University of Minho, Campus de Gualtar, Braga, Portugal

<sup>2</sup> Osun Solutions s.r.l., Località Galdo, Lauria (PZ), Italy

<sup>3</sup> Evra s.r.l., Località Galdo Zona Industriale Lotto 20, Lauria (PZ), Italy

<sup>4</sup> Institute of Oenology and Agro-Food Engineering, Università Cattolica del Sacro Cuore, Piacenza, Italy

**E-mail:** [pedrosantos@ceb.uminho.pt](mailto:pedrosantos@ceb.uminho.pt)

Extracting the totality of bio-compounds with industrial interest from Cyanobacterium is often prevented by the intrinsic rigidity of its cell wall. In this sense, the present study focuses on evaluating the influence thermal batch extraction (conventional extraction technologies) and ohmic heating (OH) assisted extraction (considered a greener alternative technology) in blue green microalgae *Arthrospira platensis* (*Spirulina*) cell disruption for bioactive fractions recovery.

The proximal composition of *Spirulina* was initially determined. The maximum protein content (i.e., C-Phycocyanin), total carbohydrates (TC) and total phenolic compounds (TPC) extracted in water at different times (30-120 min) and temperatures (30-51 °C) was quantified after the conventional and OH- assisted extraction. The freeze-thawing process was used as control. The antioxidant activity (i.e., FRAP and DPPH assays) of the obtained extracts was assessed.

Results showed that with the freeze-thawing process, traditionally used for the recovery of bio- compounds from *Spirulina*, the concentration of C-phycoerythrin was approx. 42 mg/g of *Spirulina*, 26 mgGlcE/g *Spirulina* of TC and 9 mgGAE/g *Spirulina* of TPC. Using OH-assisted extraction, the maximum of C-Phycocyanin content obtained was 45 mg/g of *Spirulina* (obtained at 37 °C, 30 min), the maximum carbohydrates' content was 40 mgGlcE/g *Spirulina* and the maximum TPC was 10 mgGAE/g *Spirulina*. On the other hand, using conventional thermal treatment it can be observed that, under the same conditions, the bioactive compounds recovery decreased to 35 mg/g, 20 mgGlcE/g *Spirulina* for C-phycoerythrin concentration and TC ( $p < 0.05$ ), respectively. The concentration in phenolic compounds is not so affected, but even so the ohmic heating potentiates the extraction of these secondary metabolites.

The antioxidant activity of the extracts there was not different between conventional treatments and OH.

Thus, the results indicated that OH is a good alternative to conventional methods aiming at the extraction of intracellular components with a decrease in processing time and energy costs associated with the extraction process, which together with an easy upscale make OH an interesting methodology for use in the industrial production of microalgae colorants and bioactive supplements.