



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Prosopagnosia or Prosopdysgnosia

Facing up to a change of concepts

Sørensen, Thomas Alrik; Overgaard, Morten

Published in:

2018 10th International Conference on Knowledge and Smart Technology

DOI (link to publication from Publisher):

[10.1109/KST.2018.8426086](https://doi.org/10.1109/KST.2018.8426086)

Creative Commons License

Unspecified

Publication date:

2018

Document Version

Accepted author manuscript, peer reviewed version

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Sørensen, T. A., & Overgaard, M. (2018). Prosopagnosia or Prosopdysgnosia: Facing up to a change of concepts. In *2018 10th International Conference on Knowledge and Smart Technology: Cybernetics in the Next Decades, KST 2018* (pp. 260-263). [8426086] IEEE Press. International Conference on Knowledge and Smart Technology (KST) <https://doi.org/10.1109/KST.2018.8426086>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- ? Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- ? You may not further distribute the material or use it for any profit-making activity or commercial gain
- ? You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Prosopagnosia or Prosopdysgnosia

Facing up to a change of concepts

Thomas Alrik Sørensen

Department of Communication, Aalborg University
Centre for Cognitive Neuroscience
Aalborg, Denmark
thomasalrik@gmail.com

Morten Storm Overgaard

Department of Clinical Medicine, Aarhus University
Cognitive Neuroscience Research Unit, CFIN, MindLab
Aarhus, Denmark

Abstract—The term ‘Prosopagnosia’ refers to the inability to identify people using visual facial cues as a result of brain injury [1]. A further term, ‘Developmental Prosopagnosia’, has been proposed to refer to a condition with a similar dysfunction, but here there is an absence of any external brain injury. In recent years, these terms have been used increasingly, despite the lack of evidence about which specific functions are involved, and how they are affected during the development of human facial feature processing. Furthermore, most of these studies seem to address a dysfunction, rather than a function that never developed or has been lost altogether.

We propose clarifying the terminology by distinguishing between whether the function is absent (prosopagnosia) or functioning at a suboptimal level (prosopdysgnosia). This distinction is particularly important as an increasing number of studies indicate varying degrees of function, rather than a function that is either present or not. Focusing on the variations of dysfunctions may help to form a better understanding of how facial features are processed and used by different observers. Moreover, studies describing functions that are lost can help us to understand which parts of a network are critical for face perception. One model that may prove particularly useful, both in the general understanding of the dysfunction and in the specific possibilities regarding rehabilitation is the Reorganization of Elementary Functions model (REF) [2]. This model has recently been applied both within consciousness studies [3] and blindsight [4].

Keywords-component; Prosopagnosia; Individual differences; Prosopdysgnosia; Visual Perception; Agnosia

I. INTRODUCTION

Visual Agnosia is a rare deficit where the observer is no longer able to recognize objects based on visual information [5]. This condition has even been reported to affect very specific domains, such as facial identification, referred to as Prosopagnosia [1]. People with prosopagnosia can often use cues other than visual, in the task of identifying other people (e.g. the sound of a person’s voice). Nevertheless, they seem to have a profound “[...] inability to identify known people by the face” [6, p. 489]. The term ‘Prosopagnosia’ is often used not only to refer to a lost function, but also to dysfunctional facial recognition; here we argue that there is a need to distinguish between whether a function is missing (prosopagnosia), or merely operating at a suboptimal level (prosopdysgnosia). This

clarification may ultimately yield a better understanding of the substrate underlying these cognitive functions and variations between observers, whereby we may progress towards a gradual understanding, where particular functions at one level of description may be achieved in different ways at the neuronal level [2,7].

Originally Bodamer [1] described patients who were able to identify individual facial features, while unable to bind these features together into a coherent whole; effectively rendering the patients unable to recognize people from facial information alone. A variety of prosopagnosia, seemingly not caused by injury or trauma, was later coined ‘Developmental Prosopagnosia’ [8]. Despite little initial attention, this dysfunction has since been reported with increasing frequency [9]. McConachie [8] described a 12-year old girl (AB) who was suffering from a severe deficit in facial identification and recognition. This became apparent when AB changed to a different school where the use of school uniforms prevented AB from using some of the feature cues she would normally rely on to identify people (i.e. clothing). Compensatory strategies potentially disguise cognitive deficits, as in the case of AB. Contrary to prior cases of prosopagnosia, AB did not seem to have any history or indication of traumatic injury, suggesting to McConachie [8] a developmental disorder. Also, AB’s mother reported that she herself had difficulties identifying faces, however, this deficiency was not as severe as in the case of AB [8]. Such findings, naturally give rise to speculations as to whether there may be a hereditary component in the deficit [10,11]. For our present purposes, Developmental Prosopagnosia could be considered as one end of the spectrum of in a function that can have varying degrees [12]. This is further substantiated by reports of observers who populate the opposite end of the spectrum, the so-called ‘super recognizers’ [13, see also 34].

In a follow-up study of AB, De Haan and Campbell [6] reported that her deficit may not be limited to a severe dysfunction of facial recognition. She also has milder deficits, both in general object naming and in object recognition. It also seemed that AB had either retained or had developed some functions of face perception, since her performance was higher on a face-matching task [6]. Most studies of Developmental

Prosopagnosia rarely describe a pure inability to process face-specific information, as otherwise seemingly suggested by the label 'agnosia'. Studies that report on diagnostic tools for prosopagnosia, typically demonstrate variations in performance between individual observers [e.g. 14,15, see also 35]. Often, a performance criterion of two standard deviations from the mean is used as a cut-off, which is implicitly indicative of a dysfunction, rather than a lacking function [16].

Additionally, recent studies have begun to use the slightly different term of 'Congenital Prosopagnosia'. This is treated synonymously with Developmental Prosopagnosia, and is defined as an "impairment in face processing that is apparent from birth in the absence of any brain damage" [17, p. 180]. Some authors may have preferred the term 'Congenital Prosopagnosia' over 'Developmental Prosopagnosia', as the latter term has also been used to refer to acquired Developmental Prosopagnosia [18]. Generally, both Developmental and Congenital Prosopagnosia are used more broadly, referring to both deficits and dysfunctions in facial recognition, that are not caused by head injury or trauma [9,19]. We believe both terms to be problematic when addressing cognitive functions that are not developed from birth. In particular, the latter term is highly questionable, as one may argue that all newborns have Congenital Prosopagnosia. In fact, studies have suggested that it is not possible to see normal perceptual processing biases until a few months after birth [e.g. 20]. Only after the visual system matures, may one see whether a child has a developmental disorder affecting the ability to recognize faces.

As well as Developmental Prosopagnosia and Congenital Prosopagnosia, a third term has emerged, namely, 'Hereditary Prosopagnosia' [e.g. 10,11]. This also seem to be used analogous to the two other types of prosopagnosia, nevertheless, here the causal emphasis is on genetics (or upbringing). Studies of families, for example [e.g. 11], only give indirect evidence for hereditary components. One may indeed, question the contribution of nature and nurture in these types of studies. Both could be confounded by a number of other factors, such as how different people relate to their experiential qualia. As an example, it has long been argued that studies of synaesthesia suggest a hereditary element based on a strong female gender bias [e.g. 21]. Recent studies have established that this gender bias is, in fact, due to diverging cultural practices between how males and females discuss how they experience their surrounding environment [22]. To the knowledge of the author's, explicit genetic studies of face recognition have yet to be performed. There are a number of studies of twins [23,24], however, where it has been questioned whether these findings relate specifically to face perception [25].

The decreased ability to recognize faces may reflect a function that is much more prone to individual variation than we usually assume [10,12]. Already in her original paper, McConachie [8] briefly speculated, as to whether the

phenomenon may be more widespread in the general population. In related areas, where we see individual perceptual differences, such as synaesthesia mentioned above, observers discover the fact that they perceive the world differently from other people only rather late in their development [26]. Similarly, to the case of AB, where her difficulties went largely unnoticed, until an external event (the use of school uniforms) revealed that AB did not perceive the world similarly to her peers. If no such external pressure is experienced, then it is easy to imagine that many observers could be unaware of their synaesthesia, or in the present case, decreased performance in identification based on visual facial cues. Studies often suggest using a lifelong report of poor face recognition as an additional diagnostic characteristic in addition to the two standard deviation cut-off on more objective tasks; however, in the light of the example above, such subjective criterion does not seem valid.

Our aim is to shift the emphasis from cause to function. We propose the more accurate term 'Prosopdysgnosia' (or 'Prosopodysgnosia') in order to emphasize a decrease in ability compared to a neurotypical observer, rather than Developmental, Congenital, or Hereditary Prosopagnosia; as all of these latter seem to carry connotations of cause. Naturally, if the decrease in function can be linked to a specific cause, such as a trauma, we find it relevant to add 'acquired', and if other causal links can be established (e.g. developmental) for prosopdysgnosia (or prosopagnosia) then it would make sense to use developmental prosopdysgnosia. However, it still seems that more work is needed before one can ascribe the cases to developmental disorders with any certainty. Even in the absence of any evidence of external trauma, it is still difficult to fully rule out the possibility that a dysfunction is not actually due to an external cause. Hence one may argue that McConachie [8] was premature in coining a term for something in the absence of evidence. Imaging studies have sometimes had difficulties distinguishing observers with Developmental Prosopagnosia from neurotypical observers [see 12]. Moreover, if face perception is a varying spectrum in the normal population, then Developmental Prosopagnosia at times would also be much more similar to neurotypical observers than would the very severe cases originally reported by Bodamer [1].

Also, work linking face perception to general expertise seems to further support the proposed distinction [e.g. 25,27]. Investigations of face perception often reveal that observers have a processing advantage for upright compared to inverted faces [e.g. 27]. In a group of colour-grapheme synaesthetes we have previously shown a decreased upright advantage, but only a few of our observers had complaints related to their ability to recognize faces [28]. Moreover, Gauthier et al. [25] recently linked poor performance on face-recognition tasks to a general decrease in object recognition (as also later reported in AB [6]); this allows for some speculation into the degree of difference between face perception and object perception.

In addition, studies of the so-called ‘own-race bias’ in perception – a bias in the perceptual processing of faces – further demonstrate that this process is not present from birth and this bias is only established when the child is at least three months old [20]. Moreover, a study by Bar-Haim, Ziv, Lamy, and Hodes [29] showed that the own-race bias is established through an interaction with the surrounding environment, supporting the idea that face perception is acquired, and is fine-tuned through environmental exposure and expertise.

If, in fact, face and object perception are modulated by a general domain non-specific function, as suggested by Gauthier et al. [25], which develops through environmental exposure and interaction [20,29], then it is even more important to be able to locate the underlying domain-specific substrates in face perception. This may prove important in the development of intervention strategies for people with suboptimal facial recognition. These in turn, may not be an effective method in treating more severe cases of prosopagnosia.

With some caution, one might also consider the theoretical consequences of the argument above. The debate between strong localization and the equipotentiality of the brain has historically been discussed with varying emphasis, reaching a compromise between the two extreme positions today [30]. If we accept that cognitive functions such as perception are represented in varying degrees [12, see also 31], rather than as cognitive modules [32]. Cognitive functions, such as face perception, may be described as processes of interacting strategies rather than separated, individual modules, hereby relating to different neural networks. This would mean for instance that face perception may be achieved in a number of different ways encompassing varying strategies from observer to observer [e.g. 2,30]. The proposal of a distinction between prosopagnosia and prosopodysgnosia seem to support this view. Which is further supported by the number of varying sub-strategies that can be employed to identify other people, as an example, some people have been reported who have difficulties in identifying others from their voice alone (phonagnosia) similar to prosopagnosia in vision [36,37].

One model of functional reorganization following brain damage, does in fact predict that a neural injury may not result in the permanent abolishment of particular functions, but, rather, in the development of different strategies to replace them. In the recently proposed Reorganization of Elementary Functions (REF) model, the orchestration of ‘elementary functions’ (specific functions localized to a particular neural region) form ‘algorithmic strategies’ in larger brain networks [2]. According to the model, such algorithmic strategies subserve behavioural and cognitive functions, but not in a 1 to 1 fashion [7]. Rather, in the case of an injury to some region in the brain, following which some elementary functions are damaged, the associated strategies and behaviour or cognition are also (temporarily) lost. However, in many cases, new strategies consisting of different or partially different sets of elementary functions realize the relevant behaviour or

cognition. Nevertheless, at the level of the algorithmic strategies, the same function will be realized in a different way. In this view, it is highly plausible that we may find intervention strategies that can alleviate milder problems like prosopodysgnosia, whereas the damage to elementary functions in prosopagnosia may be too severe to be rehabilitated.

Models, such as the REF-model unfold a more complex picture of different possible strategies with different degrees of efficiency. As such, they fit nicely into other developments in visual perception research, where conscious experience of objects contains several levels of representation [see 31,33]. Moreover, this corresponds to the different strategies used in identifying people, where we rely on a number of different cues (e.g. clothes as in the case of AB).

II. CONCLUSION

We believe that the emphasis on function should be explicit, similarly to the distinctions between alexia and dyslexia. Moreover, distinguishing pure cases of prosopagnosia from different kinds of prosopodysgnosia may lead to a better understanding of the neural mechanisms constituting the algorithmic strategies behind face perception. Furthermore, the difficulty in accurately gaging the cause of the developmental disorder, seem to warrant the more accurate term ‘prosopodysgnosia’, and therefore the term ‘prosopagnosia’ should be reserved for cases where the facial identification through visual cues is abolished, or severely limited.

ACKNOWLEDGMENT

Thomas Alrik Sørensen was supported by the Danish Council for Independent Research and Morten Storm Overgaard was supported by a starting grant from The European Research Council.

The authors would also like to thank M. W. Lindegaard for fruitful discussions, and Dr. T. W. Teasdale for comments and proofreading of the manuscript.

Correspondence concerning this article should be addressed to Thomas Alrik Sørensen, Department of Communication and Psychology, Aalborg University, Kroghstræde 3, DK-9220 Aalborg, Denmark. Email: thomasalrik@gmail.com.

REFERENCES

- [1] J. Bodamer. Die Prosop-Agnosie. *European Archives of Psychiatry and Clinical Neuroscience*, vol. 179 nr. 1, pp. 6–53, 1947.
- [2] J. Mogensen, & H. Malá. "Post-traumatic functional recovery and reorganization in animal models: A theoretical and methodological challenge". *Scandinavian Journal of Psychology*, vol. 50 nr. 6, pp. 561–573, 2009. doi:10.1111/j.1467-9450.2009.00781.x
- [3] M. Overgaard & J. Mogensen. "Visual perception from the perspective of a representational, non-reductionistic, level-dependent account of perception and conscious awareness". *Philosophical Transactions of the Royal Society B: Biological Sciences*, vol. 369 nr 1641), 1-12, 2014.
- [4] M. Overgaard & J. Mogensen. "Reconciling current approaches to blindsight". *Consciousness and Cognition*, vol. 32, 33-40, 2015.

- [5] D. F. Benson, & J. P. Greenberg. "Visual form agnosia: A specific defect in visual discrimination". *Archives of Neurology*, vol. 20 nr. 1, pp. 82–89, 1969.
- [6] E. H. F. De Haan, & R. Campbell. "A fifteen year follow-up of a case of developmental prosopagnosia". *Cortex*, vol. 27 nr. 4, pp. 489–509, 1991.
- [7] M. Overgaard, & J. Mogensen. "A Framework for the Study of Multiple Realizations: The Importance of Levels of Analysis". *Frontiers in Psychology*, vol. 2, 2011. doi:10.3389/fpsyg.2011.00079
- [8] H. R. McConachie. "Developmental prosopagnosia: A single case report". *Cortex*, vol. 12, pp. 76–82, 1976.
- [9] T. Kress, & I. Daum. "Developmental prosopagnosia: A review". *Behavioural Neurology*, vol. 14 nr. 3/4, pp. 109–122, 2003.
- [10] I. Kennerknecht, T. Grueter, B. Welling, S. Wentzek, J. Horst, S. Edwards, & M. Grueter. "First report of prevalence of non-syndromic hereditary prosopagnosia (HPA)". *American Journal of Medical Genetics Part A*, vol. 140 nr. 15, pp. 1617-1622, 2006.
- [11] M. Grueter, T. Grueter, V. Bell, J. Horst, W. Laskowski, K. Sperling, P. W. Halligan, H. D. Ellis, & I. Kennerknecht. "Hereditary Prosopagnosia: The First Case Series". *Cortex*, vol. 43, pp. 734–749, 2007.
- [12] N. Furl, L. Garrido, R. J. Dolan, J. Driver, & B. Duchaine. "Fusiform gyrus face selectivity relates to individual differences in facial recognition ability". *Journal of Cognitive Neuroscience*, vol. 23 nr. 7, pp. 1723-1740, 2011.
- [13] R. Russell, B. Duchaine, & K. Nakayama. "Super-recognizers: People with extraordinary face recognition ability". *Psychonomic Bulletin & Review*, vol. 16 nr. 2, 252-257, 2009.
- [14] B. Duchaine, & K. Nakayama. "The Cambridge Face Memory Test: Results for neurologically intact individuals and an investigation of its validity using inverted face stimuli and prosopagnosic participants". *Neuropsychologia*, vol. 44, pp. 576–585, 2006.
- [15] B. Duchaine, L. Germine, & K. Nakayama. "Family resemblance: Ten family members with prosopagnosia and within-class object agnosia". *Cognitive Neuropsychology*, vol. 24 nr. 4, pp. 419-430, 2007. doi: 10.1080/02643290701380491
- [16] D. C. Bowles, E. McKone, A. Dawel, B. C. Duchaine, R. Palermo, L. Schmalzl, D. Rivolta, C. E. Wilson & G. Yovel. "Diagnosing prosopagnosia: Effects of ageing, sex, and participant–stimulus ethnic match on the Cambridge Face Memory Test and Cambridge Face Perception Test". *Cognitive Neuropsychology*, vol. 26 nr. 5, pp. 423–455, 2009. doi:10.1080/02643290903343149
- [17] M. Behrmann, & G. Avidan. "Congenital prosopagnosia: Face-blind from birth". *Trends in Cognitive Sciences*, vol. 9 nr. 4, pp. 180–187, 2005.
- [18] T. Grüter, M. Grüter, V. Bell, & C.-C. Carbon. "Visual mental imagery in congenital prosopagnosia". *Neuroscience Letters*, vol. 453 nr. 3, pp. 135–140, 2009. doi:10.1016/j.neulet.2009.02.021
- [19] R. D. Jones, & D. Tranel. "Severe Developmental Prosopagnosia in a Child With Superior Intellect". *Journal of Clinical and Experimental Neuropsychology (Neuropsychology, Development and Cognition: Section A)*, vol. 23 nr. 3, pp. 265–273, 2001. doi:10.1076/jcen.23.3.265.1183
- [20] D. J. Kelly, P. C. Quinn, A. M. Slater, K. Lee, A. Gibson, M. Smith, L. Ge, & O. Pascalis. "Three-month-olds, but not newborns, prefer own-race faces". *Developmental Science*, vol. 8 nr. 6, pp. F31-F36, 2005.
- [21] S. Baron-Cohen. "Is there a normal phase of synaesthesia in development?". *Psyche*, vol. 2 nr. 27, pp. 223-228, 1996.
- [22] J. Simner, C. Mulvenna, N. Sagiv, E. Tsakanikos, S. A. Witherby, C. Fraser, K. Scott, & J. Ward. "Synaesthesia: the prevalence of atypical cross-modal experiences". *Perception*, vol. 35 nr. 8, 1024-1033, 2006.
- [23] J. B. Wilmer, L. Germine, C. F. Chabris, G. Chatterjee, M. Williams, E. Loken, K. Nakayama, & B. Duchaine. "Human face recognition ability is specific and highly heritable". *Proceedings of the National Academy of Sciences*, vol. 107 nr. 11, 5238-5241, 2010.
- [24] Q. Zhu, Y. Song, S. Hu, X. Li, M. Tian, Z. Zhen, Q. Dong, N. Kanwisher, & J. Liu. "Heritability of the specific cognitive ability of face perception." *Current Biology*, vol. 20 nr. 2, pp. 137-142, 2010.
- [25] I. Gauthier, R. W. McGugin, J. J. Richler, G. Herzmann, M. Speegle, A. E. Van Gulick. "Experience moderates overlap between object and face recognition, suggesting a common ability". *Journal of Vision*, vol. 14 nr. 8, pp. 1-12, 2014.
- [26] R. E. Cytowic & D. M. Eagleman. *Wednesday Is Indigo Blue: Discovering the Brain of Synesthesia*. Cambridge, Massachusetts: MIT Press, 2009.
- [27] K. M. Curby, K. Glazek, & I. Gauthier. "A Visual Short-Term Memory Advantage for Objects of Expertise". *Journal of Experimental Psychology: Human, Perception and Performance*, vol. 35 nr. 1, pp. 94-107, 2009. DOI: 10.1037/0096-1523.35.1.94
- [28] T. A. Sorensen. "Facial Identification in Observers with Colour-Grapheme Synaesthesia" *Perception*, vol. 42, 167-167, 2013.
- [29] Y. Bar-Haim, T. Ziv, D. Lamy, & R. M. Hodes. "Nature and nurture in own-race face processing". *Psychological Science*, vol. 17 nr. 2, pp. 159-163, 2006.
- [30] M. Catani, F. Dell'Acqua, A. Bizzi, S. J. Forkel, S. C. Williams, A. Simmons, G. M. Declan, & M. Thiebaut de Schotten. "Beyond cortical localization in clinico-anatomical correlation". *Cortex*, vol. 48 nr. 10, pp. 1262-1287, 2012.
- [31] M. Overgaard & K. Sandberg. "Kinds of access: different methods for report reveal different kinds of metacognitive access". *Philosophical Transactions of the Royal Society B: Biological Sciences*, vol. 367 nr. 1594, pp. 1287–1296, 2012. doi:10.1098/rstb.2011.0425
- [32] J. Fodor. *The Mind Doesn't Work That Way: The Scope and Limits of Computational Psychology*. Cambridge, MA: MIT Press, 2000.
- [33] S. Kouider, V. de Gardelle, J. Sackur, & E. Dupoux. "How rich is consciousness? The partial awareness hypothesis". *Trends in Cognitive Sciences*, vol. 14 nr. 7, pp. 301–307, 2010. doi:10.1016/j.tics.2010.04.006
- [34] J. B. Wilmer, L. Germine, C. F. Chabris, G. Chatterjee, M. Williams, E. Loken, K. Nakayama, & B. Duchaine. Human face recognition ability is specific and highly heritable. *Proceedings of the National Academy of Sciences*, vol. 107 nr. 11, pp. 5238-5241, 2010.
- [35] R. Le Grand, P. A. Cooper, C. J. Mondloch, T. L. Lewis, N. Sagiv, B. de Gelder, & D. Maurer. "What aspects of face processing are impaired in developmental prosopagnosia?" *Brain and Cognition*, vol. 61 nr. 2, pp. 139-158, 2006.
- [36] B. E. Shilowich, & I. Biederman. "An estimate of the prevalence of developmental phonagnosia". *Brain and Language*, vol. 159, pp. 84-91, 2016.
- [37] S. B. Herald, X. Xu, I. Biederman, O. Amir, & B. E. Shilowich. "Phonagnosia: A voice homologue to prosopagnosia". *Visual Cognition*, vol. 22 nr. 8, pp. 1031-1033, 2014.