



Aalborg Universitet

AALBORG UNIVERSITY  
DENMARK

## Biostimulation strategies to enhance manganese removal in drinking water biofilters

Breda, Inês Lousinha Ribeiro; Ramsay, Loren; Søborg, Ditte; Roslev, Peter

*Publication date:*  
2018

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

*Citation for published version (APA):*

Breda, I. L. R., Ramsay, L., Søborg, D., & Roslev, P. (2018). *Biostimulation strategies to enhance manganese removal in drinking water biofilters*. Poster presented at International Symposium on Biological Treatment, Austin, United States.

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- ? Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- ? You may not further distribute the material or use it for any profit-making activity or commercial gain
- ? You may freely distribute the URL identifying the publication in the public portal ?

### Take down policy

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.



AALBORG UNIVERSITET

SKANDERBORG  
FORSYNING

# BIOSTIMULATION STRATEGIES TO ENHANCE MANGANESE REMOVAL IN DRINKING WATER BIOFILTERS

Inês L. Breda <sup>a,b</sup>, Loren Ramsay <sup>c</sup>, Ditte Søborg <sup>c</sup>, Peter Roslev <sup>b</sup>

<sup>a</sup> Department of Chemistry and Bioscience, Aalborg University, Denmark

<sup>b</sup> Group of Project and Development, Skanderborg Forsyningsvirksomhed A/S, Denmark

<sup>c</sup> Research Group for Energy and Environment, VIA University College, Denmark


## ABSTRACT

Maturation of drinking water biofilters for removal of manganese can vary considerably. The aim of this study was to investigate biostimulation strategies to enhance manganese removal during start-up of biofilters for treatment of groundwater. Two major biostimulation strategies were investigated: biostimulation using different filter media (e.g. quartz, calcium carbonate, polystyrene, manganese oxide), and biostimulation using inoculation of virgin quartz filters with matured quartz. The onset and extent of manganese oxidation was determined spectrophotometrically, and the bacterial communities were characterized using qPCR, 16S rRNA pyrosequencing, and enrichments of manganese oxidizing bacterial consortia. The investigation suggested that when inoculating different filter media with an identical water source, the bacterial community formed during the start-up period is strongly influenced by the filter media type. Biostimulation of virgin media to enhance initial manganese removal should take place in the early stages of filter development whereas autocatalytic processes appear to become dominant with time. The complex interactions between biological and chemical oxidation processes should be considered when optimizing biofilters for efficient removal of manganese from drinking water.

## FOCUS

EFFECT OF START-UP STRATEGIES ON MANGANESE REMOVAL, AND ON THE BACTERIAL COMMUNITIES FORMED IN THE EARLY STAGES OF THE BIOFILTER.

## DRINKING WATER TREATMENT IN DENMARK



### GROUNDWATER

– main water source for drinking water production

### NO ADDITION OF CHEMICALS

– strong legislation controlling the addition of chemicals

### NO DESINFECTION

– no chlorine added to the drinking water


## BIOFILTERS START-UP PERIOD

A major disadvantage of biofilters is the long start-up period required for virgin filter medium to become fully functional.



Start-up of drinking water biofilters hinges on a set of interconnected physical, chemical and biological processes. When manganese is present, the duration of a start-up varies from weeks to more than a year.



## MANGANESE REMOVAL PHYSICAL, CHEMICAL AND BIOLOGICAL




# RESULTS

## Alternative filter materials


### Column assay with 5 filter materials in triplicates



**Figure 1** | Manganese concentration in the column outlets over time as a percentage of manganese concentration in the source water. Each datum point represents the average of three filter columns.



**Figure 2** | Heatmap of the 25 most abundant genera present in filter media coatings at Day 75 (average of three filter columns). Shadings are based on read abundance (%) and dots mark indicate genera with known manganese oxidizing bacteria (MnOBs).




**Figure 3** | Principal component analysis of bacterial communities present in the source water and on the coating of each filter medium type. Numbers 1 to 3 represent medium replicates. Grey dots illustrate the taxa detected in all samples. Red labels identify selected key taxa.



**Manganese oxidation and bacterial diversity on different filter media coatings during the start-up of drinking water biofilters.** Breda I. L., Ramsay L., Roslev P. *Journal of Water Supply: Research and Technology –AQUA*, 2017. DOI:10.2166/aqua.2017.084

## Inoculation of new filters with matured quartz


### Pilot scale filters with virgin quartz (non-inoculated filter) and with virgin and matured quartz (inoculated filter)




**Figure 4** | Experimental setup pilot scale.




**Figure 5** | Manganese removal over time (top) and filters depth (bottom) by non-inoculated (green) and inoculated (red) pilot-scale biofilters.



**Figure 6** | ATP abundance in the top 10cm of initially virgin quartz from the inoculated and non-inoculated filters during the filters maturation.



**Figure 7** | Detection of MnOBs by lecouberlin blue in Leptotrix medium plate cultures using aliquots of fresh filter media samples from several depths of both inoculated and non-inoculated filter collected at the end of the start-up



**Figure 8** | Rate coefficient of manganese removal according to batch experiments using duplicates of fresh filter media with and without azide from depths 10 cm and 20 cm of both inoculated and non-inoculated filters.

## CONCLUSIONS

- The bacterial community formed during the start-up period is strongly influenced by the filter media type.
- Management of bacterial communities may be possible by selecting specific media to enhance growth and activity of specific bacteria.
- The interactions between biological and chemical oxidation processes should be considered when optimizing biofilters for efficient removal of manganese from drinking water.