



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Impact of baseline anemia in patients with acute coronary syndromes undergoing percutaneous coronary intervention

a prespecified analysis from the VALIDATE-SWEDEHEART trial

Wester, Axel; Attar, Rubina; Mohammad, Moman Aladdin; Andell, Pontus; Hofmann, Robin; Jensen, Jens; Szummer, Karolina; Erlinge, David; Koul, Sasha

Published in:
Journal of the American Heart Association

DOI (link to publication from Publisher):
[10.1161/JAHA.119.012741](https://doi.org/10.1161/JAHA.119.012741)

Creative Commons License
CC BY-NC 4.0

Publication date:
2019

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):
Wester, A., Attar, R., Mohammad, M. A., Andell, P., Hofmann, R., Jensen, J., Szummer, K., Erlinge, D., & Koul, S. (2019). Impact of baseline anemia in patients with acute coronary syndromes undergoing percutaneous coronary intervention: a prespecified analysis from the VALIDATE-SWEDEHEART trial. *Journal of the American Heart Association*, 8(16), [e012741]. <https://doi.org/10.1161/JAHA.119.012741>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- ? Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- ? You may not further distribute the material or use it for any profit-making activity or commercial gain
- ? You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Impact of Baseline Anemia in Patients With Acute Coronary Syndromes Undergoing Percutaneous Coronary Intervention: A Prespecified Analysis From the VALIDATE-SWEDEHEART Trial

Axel Wester, MD; Rubina Attar, BMSc; Moman Aladdin Mohammad, MD; Pontus Andell, MD, PhD; Robin Hofmann, MD, PhD; Jens Jensen, MD, PhD; Karolina Szummer, MD, PhD; David Erlinge, MD, PhD; Sasha Koul, MD, PhD

Background—The impact of baseline anemia in a contemporary acute coronary syndrome (ACS) population undergoing percutaneous coronary intervention in the era of predominant radial artery access, potent P2Y12 inhibition, and rare use of glycoprotein IIb/IIIa inhibitors has not been adequately studied.

Methods and Results—ACS patients who underwent percutaneous coronary intervention between 2014 and 2016 in the VALIDATE-SWEDEHEART (Bivalirudin Versus Heparin in ST-Segment and Non-ST-Segment Elevation Myocardial Infarction in Patients on Modern Antiplatelet Therapy in the Swedish Web System for Enhancement and Development of Evidence-Based Care in Heart Disease Evaluated According to Recommended Therapies Registry) trial without missing values for hemoglobin were included (n=5482). Mortality, myocardial reinfarction, and major bleeding at 180 days were assessed using Cox regression models and propensity score matching. All studied comorbidities were more common in ACS patients who had anemia (n=792). ACS patients with anemia had higher rates of 180-day mortality (6.9% versus 2.1%; hazard ratio, 1.9; 95% CI, 1.3–2.7; $P<0.001$), myocardial reinfarction (4.3% versus 1.9%; hazard ratio, 1.7; 95% CI, 1.1–2.7; $P=0.013$), and major bleeding (13.4% versus 8.2%; hazard ratio, 1.3; 95% CI, 1.0–1.6; $P=0.041$). The results were most evident in patients with a hemoglobin value <100 g/L, who had a nearly 10 times higher mortality rate.

Conclusions—Baseline anemia in ACS patients undergoing percutaneous coronary intervention, treated according to current practice including routine radial artery access, constitutes a high-risk feature for both ischemic events, bleeding events, and mortality. A multidisciplinary approach is warranted to maximize benefit and minimize patient risk.

Clinical Trial Registration—URL: <http://www.clinicaltrials.gov>. Unique identifier: NCT02311231. (*J Am Heart Assoc.* 2019;8:e012741. DOI: 10.1161/JAHA.119.012741.)

Key Words: acute coronary syndromes • anemia • mortality • VALIDATE-SWEDEHEART

Despite advances in the field of acute coronary care, major bleedings following percutaneous coronary intervention (PCI) still constitute a major clinical problem.¹ It has been suggested that a major bleeding carries the same prognostic impact as a new myocardial infarction (MI) during the first year after an acute coronary syndrome (ACS).² The HORIZONS-AMI (Harmonizing Outcomes With Revascularization and Stents in Acute Myocardial Infarction) trial

demonstrated more than a decade ago that bivalirudin reduced mortality in patients with ST-segment-elevation myocardial infarction (STEMI) undergoing primary percutaneous coronary intervention, compared with heparin plus glycoprotein IIb/IIIa inhibitors (GPIs).³ This was in part due to a reduction in major bleedings.^{3,4} Both non-access site and access site major bleedings increase mortality; however, radial artery access essentially removes the risk of excess

From the Department of Cardiology, Clinical Sciences, Lund University, Skåne University Hospital, Lund, Sweden (A.W., R.A., M.A.M., P.A., D.E., S.K.); Department of Cardiology, Clinical Medicine, Aalborg University, Aalborg, Denmark (R.A.); Unit of Cardiology, Department of Medicine, Karolinska Institute, Stockholm, Sweden (P.A., K.S.); Heart and Vascular Division, Karolinska University Hospital, Stockholm, Sweden (P.A., K.S.); Division of Cardiology, Department of Clinical Science and Education, Södersjukhuset, Karolinska Institute, Stockholm, Sweden (R.H.); Department of Clinical Science and Education, Södersjukhuset, Karolinska Institute, Stockholm, Sweden (J.J.); Unit of Cardiology, Capio S:t Göran Hospital, Stockholm, Sweden (J.J.).

Accompanying Tables S1 and S2 and Figure S1 are available at <https://www.ahajournals.org/doi/suppl/10.1161/JAHA.119.012741>

Correspondence to: Axel Wester, MD, Department of Cardiology, Skåne University Hospital, 221 85 Lund, Sweden. E-mail: axel.wester@med.lu.se

Received April 5, 2019; accepted July 9, 2019.

© 2019 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

Clinical Perspective

What Is New?

- Baseline anemia in acute coronary syndrome patients who undergo percutaneous coronary intervention and are treated according to current practice with >90% use of radial artery access, >97% potent P2Y12 inhibitors during percutaneous coronary intervention, and <3% glycoprotein IIb/IIIa inhibitors was associated with higher rates of 180-day mortality, myocardial reinfarction, and major bleeding.
- Patients with severe anemia (a hemoglobin value <100 g/L) had a nearly 10 times higher mortality rate, 6 times higher rate of myocardial reinfarction, and 3 times higher rate of major bleeding at 180 days.

What Are the Clinical Implications?

- Our results indicate that anemia constitutes a high-risk feature in patients with acute coronary syndromes, despite treatment with contemporary therapies, including high use of radial artery access.
- In complex patients with anemia and concomitant coronary disease, a multidisciplinary approach is warranted to maximize benefit and minimize patient risk.

mortality associated with major access site bleedings.^{5,6} In parallel with the increased use of radial artery access, acute medical care of ACS patients has also improved with the advent of potent P2Y12 inhibitors.^{7,8}

Baseline anemia, defined according to the World Health Organization,⁹ is present in nearly 20% of ACS patients.^{10,11} ACS patients with anemia are older and more often present with previous cardiovascular disease, diabetes mellitus, a history of major bleeding, and malignancy, compared with those without anemia.^{10–19} Having anemia at baseline therefore identifies both patients at high risk of bleeding and those at high risk for ischemic events. In fact, anemia in ACS patients is not only associated with overall increased mortality and bleeding but in some studies also associated with increased rates of MI.^{10,11}

The vast majority of studies investigating the impact of baseline anemia on hard clinical events have been performed in an era of either extensive GPI use or with limited radial artery use. The recent VALIDATE-SWEDEHEART (Bivalirudin Versus Heparin in ST-Segment and Non-ST-Segment Elevation Myocardial Infarction in Patients on Modern Antiplatelet Therapy in the Swedish Web System for Enhancement and Development of Evidence-Based Care in Heart Disease Evaluated According to Recommended Therapies Registry) trial compared bivalirudin to heparin monotherapy in ACS patients undergoing PCI.²⁰ The patients enrolled in the trial constituted a modern ACS population with high use of radial artery access

(90.3%) as well as high use of modern P2Y12 inhibitors (ticagrelor 94.9%, prasugrel 2.1%, or cangrelor 0.3%). Furthermore, only bailout use of GPI was allowed (2.6%).²⁰ The impact of anemia at baseline in such a patient cohort, treated according to contemporary strategies has so far not been adequately assessed. The aim of the current study was therefore to evaluate the prognostic value of baseline anemia in ACS patients enrolled in the VALIDATE-SWEDEHEART trial.

Methods

Study Population

Data, methods and materials used to conduct this research can be made available after clearance by the VALIDATE steering committee. The study design of the VALIDATE-SWEDEHEART trial has already been described in detail.^{20,21} In short, VALIDATE-SWEDEHEART was a registry-based, multicenter, prospective, randomized, controlled, and open-label clinical trial evaluating the efficacy and safety of heparin monotherapy versus bivalirudin in ACS patients undergoing PCI. The trial included patients presenting with STEMI or non-STEMI (NSTEMI) between 2014 and 2016 and for whom PCI was planned at one of the 25 participating Swedish centers. Treatment with ticagrelor, prasugrel, or cangrelor before PCI was mandatory and only bailout use of GPI was allowed. The SWEDEHEART registry was used as case-report form for randomization and for follow-up. A total of 6006 patients underwent randomization in the VALIDATE-SWEDEHEART trial, and all patients without missing values for hemoglobin (n=5482) were included in the present study (Figure 1). Baseline anemia was defined in accordance with the World Health Organization as a hemoglobin value <130 g/L for men and <120 g/L for women.⁹ The hemoglobin values in the cohort represent the first acquired hemoglobin for each patient, meaning that the blood tests were performed at the




Figure 1. Flowchart of study inclusion.

emergency department on arrival or in relation to the PCI procedure in those STEMI patients who went directly to the catheterization lab. The VALIDATE-SWEDEHEART trial and subgroup analyses were approved by the ethics committee at Lund University and the Swedish Medical Products Agency. All patients provided written informed consent to participate in the trial.

End Points

The primary end point was death from any cause at 180 days. Secondary end points were myocardial reinfarction, major bleeding, definite stent thrombosis, and stroke at 180 days. MI was defined according to the third universal definition,²² and major bleeding was defined as type 2, 3, or 5 according to the Bleeding Academic Research Consortium scale.²⁰ Further analyses were done on cause of death, divided into cardiovascular death, fatal bleeding, or other causes.

End point screening and adjudication

Research nurses screened for clinical end point events by contacting the patients or first-degree relatives by telephone 7 and 180 days after PCI. If a patient was suspected to have had a clinical end point event (ie, death, myocardial reinfarction, major bleeding, or stroke), the patient's healthcare records were subjected to central blinded adjudication to determine both the cause of the event and the date of the event according to prespecified criteria. Time to event was calculated using calendar days with date of randomization as time point zero. If the patient or relatives could not be contacted after the nurses had placed repeated telephone calls and mailed a letter, information was collected through review of hospital records.²⁰

Statistical Analyses

Differences between categorical variables were analyzed with the chi-square test and continuous variables were compared using the Mann–Whitney *U* test. Kaplan–Meier event rates were analyzed to compare unadjusted primary and secondary end points between groups. Patients who were lost to follow-up were censored at the last contact with the patient.²⁰ Adjustments for confounders were made using multivariable Cox regression models as well as propensity score matching. We also performed landmark analyses for mortality and major bleeding to distinguish between short- and long-term effects. Statistical significance was defined as a 2-tailed $P < 0.05$. Analyses were performed in IBM SPSS (Armonk, NY: IBM Corp.) (version 25), R (R Foundation for Statistical Computing, Vienna, Austria) (version 3.4.4), Stata (Statacorp, TX) (GraphPad Software, San Diego, California USA) (version 15.1), or GraphPad Prism (version 8.0.2).

Propensity score matching

A propensity score (PS) for baseline anemia was calculated using the variables age, sex, STEMI, medical history of kidney dysfunction, diabetes mellitus, coronary artery disease, and heart failure to achieve a good balancing of baseline differences. PS matching was performed in a 1:1 fashion. End points were subsequently analyzed using univariable Cox regression. In separate sensitivity analyses, downstream adjustments were made for body mass index (BMI).

Multiple imputation and subsequent multivariable Cox regression

Baseline differences that were associated with anemia and with worse clinical outcomes were identified a priori and adjusted for. These variables were age, sex, BMI, STEMI, medical history of kidney dysfunction, hyperlipidemia, hypertension, diabetes mellitus, current smoking, coronary artery disease, stroke, heart failure, Killip class, hours from symptom onset to PCI, radial artery access during PCI, use of potent P2Y12 inhibitors (ticagrelor, prasugrel, or cangrelor) during PCI, and blood transfusion during the hospital stay. In the original data set, 1440 (26.3%) cases had missing values for at least 1 of these variables. Data were assumed to be missing at random, based on Little's Missing Completely at Random test as well as logistic regression modeling. Multiple imputation was performed for missing data by using 10 imputations and 10 iterations for every imputation. In subsequent multivariable Cox regression analysis, adjustments were made for all variables mentioned above. Kidney dysfunction was defined as an estimated glomerular filtration rate < 60 mL/min per 1.73 m². Coronary artery disease was defined as previous MI, previous PCI, or previous coronary artery bypass grafting. The multivariable Cox regression model for mortality was tested for proportionality of hazards using Schoenfeld residuals.

Results

Patient Characteristics

Baseline anemia was present in 792 of 5482 ACS patients (14.4%). Patients with anemia were older ($P < 0.001$), more often women ($P = 0.034$), and had a lower BMI compared with patients without anemia ($P < 0.001$; Table 1). Furthermore, patients with anemia more often had kidney dysfunction, hyperlipidemia, previous hypertension, diabetes mellitus, current smoking, previous coronary artery disease, previous stroke, or a history of heart failure (all $P < 0.001$). Patients with anemia also more frequently presented with STEMI and Killip class 3 or 4 than patients without anemia ($P < 0.001$). The time from symptom onset to PCI did not significantly differ between nonanemic and anemic patients when analyzing

Table 1. Characteristics of Patients With Acute Coronary Syndromes With and Without Anemia

| | No Anemia* (n=4690) | Anemia* (n=792) | P Value | Missing or Unknown, n (%) |
|--|------------------------|--------------------|---------|------------------------------|
| Hemoglobin, median (IQR) | 145 (137–154) | 118 (112–125) | <0.001 | 0 (0.0) |
| Age, y | | | | 0 (0.0) |
| Median (IQR) | 67 (59–74) | 74 (67–81) | <0.001 | |
| ≥65, n (%) | 2695 (57.5) | 642 (81.1) | <0.001 | |
| Female, n (%) | 1217 (25.9) | 234 (29.5) | 0.034 | 0 (0.0) |
| BMI, median (IQR) | 27 (25–30) | 26 (23–28) | <0.001 | 563 (10.3) |
| Medical history, n (%) | | | | |
| Kidney dysfunction | 598 (12.8) | 244 (30.8) | <0.001 | 8 (0.1) |
| Hyperlipidemia | 1419 (30.3) | 299 (37.8) | <0.001 | 43 (0.8) |
| Hypertension | 2351 (50.4) | 475 (60.3) | <0.001 | 27 (0.5) |
| Diabetes mellitus | 702 (15.0) | 198 (25.0) | <0.001 | 14 (0.3) |
| Current smoker | 1174 (25.7) | 146 (19.1) | <0.001 | 142 (2.6) |
| CAD | 847 (18.1) | 221 (27.9) | <0.001 | 56 (1.0) |
| Previous MI | 690 (14.7) | 183 (23.1) | <0.001 | |
| Previous PCI | 639 (13.6) | 156 (19.7) | <0.001 | |
| Previous CABG | 200 (4.3) | 66 (8.3) | <0.001 | |
| Stroke | 170 (3.6) | 51 (6.5) | <0.001 | 26 (0.5) |
| Heart failure | 142 (3.0) | 60 (7.6) | <0.001 | 120 (2.2) |
| STEMI, n (%) | 2365 (50.4) | 457 (57.7) | <0.001 | 0 (0.0) |
| Killip class 3 or 4, n (%) | 38 (0.8) | 13 (1.6) | 0.024 | 0 (0.0) |
| Hours from symptoms to PCI | | | | 592 (10.8) |
| STEMI, median (IQR) | 3 (2–6) | 3 (2–6) | 0.482 | |
| NSTEMI, median (IQR) | 31 (18–57) | 35 (18–64) | 0.177 | |
| Randomized to heparin, n (%) | 2372 (50.6) | 385 (48.6) | 0.306 | 0 (0.0) |
| Potent P2Y12 inhibitors during PCI, n (%) | 4566 (98.9) | 749 (96.9) | <0.001 | 92 (1.7) |
| Radial access, n (%) | 4276 (91.2) | 680 (85.9) | <0.001 | 0 (0.0) |
| Peak high-sensitivity cardiac troponin T, median (IQR) | 628 (146–2490) | 1013 (169–3251) | 0.001 | 1475 (26.9) |
| LVEF during hospital stay, n (%) | | | <0.001 | 700 (12.8) |
| ≥50% | 2672 (65.0) | 371 (55.4) | | |
| 49%–40% | 879 (21.4) | 174 (26.0) | | |
| 39%–30% | 451 (11.0) | 91 (13.6) | | |
| <30% | 110 (2.7) | 34 (5.1) | | |
| Blood transfusion during hospital stay, n (%) | 4 (0.1) | 4 (0.5) | 0.013 | 39 (0.7) |
| Discharge medications, n (%) | | | | |
| Aspirin | 4483 (97.0) | 713 (93.2) | <0.001 | 104 (1.9) |
| Other antiplatelets | | | <0.001 | 97 (1.8) |
| Clopidogrel | 339 (7.3) | 92 (12.0) | | |
| Prasugrel | 46 (1.0) | 9 (1.2) | | |
| Ticagrelor | 4160 (90.0) | 639 (83.5) | | |

Continued

Table 1. Continued

| | No Anemia* (n=4690) | Anemia* (n=792) | P Value | Missing or Unknown, n (%) |
|------------------|------------------------|--------------------|---------|---------------------------|
| Warfarin or NOAC | 291 (6.3) | 69 (9.0) | 0.014 | 102 (1.9) |
| ACEI or ARB | 4106 (88.9) | 651 (85.1) | 0.009 | 98 (1.8) |
| Beta-blockers | 4217 (91.3) | 652 (85.2) | <0.001 | 98 (1.8) |
| Statins | 4522 (97.9) | 732 (95.7) | <0.001 | 97 (1.8) |

ACEI indicates angiotensin-converting enzyme inhibitor; ARB, angiotensin II receptor blocker; BMI, body mass index; CABG, coronary artery bypass grafting; CAD, coronary artery disease; IQR, interquartile range; LVEF, left ventricular ejection fraction; MI, myocardial infarction; NOAC, novel oral anticoagulant; NSTEMI, non-ST-segment-elevation myocardial infarction; PCI, percutaneous coronary intervention; STEMI, ST-segment-elevation myocardial infarction.

*Baseline anemia was defined in accordance with the World Health Organization as a hemoglobin value <130 g/L for men and <120 g/L for women.

STEMI (median time, 3 hours in both groups) and NSTEMI patients separately (31 hours versus 35 hours, respectively). Radial access PCI and potent P2Y12 inhibitors were less frequently used in patients with anemia ($P<0.001$). Patients with anemia had a higher median peak value of high-sensitivity cardiac troponin T and a lower left ventricular ejection fraction during their hospital stay than patients without anemia (both $P=0.001$). Furthermore, patients with anemia more often received blood transfusion ($P=0.013$). At discharge, patients with anemia less often received evidence-based discharge medications such as aspirin and ticagrelor (Table 1). After PS matching, baseline variables were equalized between groups (Table 2). However, differences remained regarding BMI (which was analyzed in separate sensitivity analyses).

Mortality

Patients with ACS and anemia at baseline had a higher unadjusted mortality rate at 180 days compared with patients without anemia (6.9% and 2.1%, respectively; $P<0.001$; Figure 2A). After multiple imputation, and subsequent multivariable adjustments for confounding factors, patients with anemia still had a higher hazard ratio (HR) for 180-day mortality (HR, 1.9; 95% CI, 1.3–2.7; $P<0.001$). Results were consistent in the PS matched cohort (HR, 2.3; 95% CI, 1.4–3.7; $P=0.001$; Table 3; Figure S1A).

Myocardial Reinfarction

Patient with ACS and anemia more often experienced myocardial reinfarction (4.3% versus 1.9%; $P<0.001$; Figure 2B). Results were unchanged after multiple imputation with multivariable adjustments (HR, 1.7; 95% CI, 1.1–2.7; $P=0.013$) as well as in PS matched patients (HR, 2.0; 95% CI, 1.1–3.7; $P=0.022$; Table 3; Figure S1B).

Major Bleeding

ACS patients with anemia experienced higher rates of major bleeding than patients without anemia (13.4% versus 8.2%;

$P<0.001$; Figure 2C). Results were consistent after multiple imputation with multivariable adjustments (HR, 1.3; 95% CI, 1.0–1.6; $P=0.041$) and in PS matched patients (HR, 1.4; 1.0–1.9; $P=0.037$; Table 3; Figure S1C).

Other End Points

No difference was found in event rates of definite stent thrombosis between patients with and without anemia (0.8% and 0.6%, respectively; $P=0.46$; Figure 2D). PS matching showed similar results (HR, 1.5; 95% CI, 0.4–5.4; $P=0.51$; Figure S1D). However, patients with anemia experienced higher rates of stroke (2.0% compared with 0.7%; $P<0.001$; Figure 2E), which was consistent in PS matched patients (HR, 2.9; 95% CI, 1.0–8.0; $P=0.042$; Table 3; Figure S1E).

Sensitivity and Subgroup Analyses

Landmark analyses in patients who survived the first 7 days were performed (Figure 3). Both mortality (4.5% versus 1.2%; $P<0.001$) and major bleeding (6.9% versus 4.9%; $P=0.026$) were significantly increased in the 7-day landmark analyses for ACS patients with anemia. The mortality curve also showed continued separation over time (Figure 3A). Results were also consistent when anemia was stratified according to severity (Figure 4). Patients with anemia with a hemoglobin <100 g/L had a nearly 10 times higher mortality rate, 6 times higher rate of myocardial reinfarction, and 3 times higher rate of major bleeding, compared with patients with normal hemoglobin values. High hemoglobin values (>150 g/L) were not associated with worse outcome compared with normal hemoglobin values. Hemoglobin was also analyzed as a continuous variable for patients with a hemoglobin value between 90 (to exclude outliers, based on visual inspection of a histogram of the hemoglobin level distribution) and 140 (as greater hemoglobin levels have previously been associated with increased risk of adverse

Table 2. Patient Characteristics After Propensity Score Matching

| | No Anemia (n=762) | Anemia* (n=762) | P Value |
|--|-------------------|-----------------|---------|
| Hemoglobin, median (IQR) | 142 (135–151) | 118 (112–125) | <0.001 |
| Age, y | | | |
| Median (IQR) | 74 (67.75–81) | 74 (67–81) | 0.518 |
| ≥65, n (%) | 628 (82.4) | 617 (81.0) | 0.466 |
| Female, n (%) | 231 (30.3) | 225 (29.5) | 0.737 |
| BMI, median (IQR) | 27 (24–30) | 26 (23–28) | <0.001 |
| Medical history, n (%) | | | |
| Kidney dysfunction | 236 (31.0) | 236 (31.0) | >0.999 |
| Hyperlipidemia | 283 (37.4) | 289 (38.4) | 0.690 |
| Hypertension | 478 (63.1) | 451 (59.5) | 0.155 |
| Diabetes mellitus | 165 (21.7) | 189 (24.8) | 0.145 |
| Current smoker | 128 (17.3) | 138 (18.8) | 0.453 |
| CAD | 195 (25.6) | 213 (28.0) | 0.298 |
| Stroke | 38 (5.0) | 49 (6.5) | 0.217 |
| Heart failure | 56 (7.3) | 60 (7.9) | 0.699 |
| STEMI, n (%) | 441 (57.9) | 436 (57.2) | 0.796 |
| Killip class 3 or 4, n (%) | 13 (1.7) | 13 (1.7) | >0.999 |
| Hours from symptoms to PCI | | | |
| STEMI, median (IQR) | 3 (2–6) | 3 (2–6) | 0.743 |
| NSTEMI, median (IQR) | 33 (21–62) | 34 (18–64) | 0.647 |
| Potent P2Y12 inhibitors during PCI, n (%) | 734 (98.1) | 720 (96.8) | 0.097 |
| Radial artery access, n (%) | 677 (88.8) | 655 (86.0) | 0.089 |
| Peak high-sensitivity cardiac troponin T, median (IQR) | 763 (161–3252) | 1017 (184–3217) | 0.388 |
| LVEF during hospital stay, n (%) | | | 0.854 |
| ≥50% | 380 (57.2) | 356 (55.4) | |
| 49%–40% | 161 (24.2) | 169 (26.3) | |
| 39%–30% | 90 (13.6) | 85 (13.2) | |
| <30% | 33 (5.0) | 33 (5.1) | |
| Blood transfusion during hospital stay, n (%) | 2 (0.3) | 4 (0.5) | 0.434 |

BMI indicates body mass index; CAD, coronary artery disease; IQR, interquartile range; LVEF, left ventricular ejection fraction; NSTEMI, non-ST-segment-elevation myocardial infarction; PCI, percutaneous coronary intervention; STEMI, ST-segment-elevation myocardial infarction.

*Baseline anemia was defined in accordance with the World Health Organization as a hemoglobin value <130 g/L for men and <120 g/L for women.

events).¹² This analysis yielded an HR of 0.95 (95% CI, 0.93–0.96; $P<0.001$) for each gram per liter increase of hemoglobin with regard to 180-day mortality. In patients with baseline anemia, randomized treatment (heparin versus bivalirudin) did not affect rates of mortality ($P=0.36$), myocardial reinfarction ($P=0.89$), or major bleeding ($P=0.92$) at 180 days (Table S1). Patients with anemia did not seem to benefit more from radial artery access during PCI than patients without anemia regarding 180-day mortality ($P_{\text{interaction}}=0.437$) or major bleeding ($P_{\text{interaction}}=0.254$). Results after PS matching did not

change when downstream adjustments were made for BMI (Table S2). Moreover, we analyzed causes of death and found that patients with anemia were 3 times as likely to die from cardiovascular causes, 5 times as likely to die from fatal bleeding, and 4 times as likely to die from other causes (noncardiovascular and nonfatal bleeding; Table 4). Among the patients with anemia who died within 180 days ($n=54$), fatal bleeding and other causes (noncardiovascular and nonfatal bleeding) represented a larger proportion of deaths than for patients without anemia who experienced death ($n=96$; Figure 5).




Figure 2. Kaplan–Meier failure functions for mortality (A), myocardial reinfarction (B), major bleeding (C), definite stent thrombosis (D), and stroke (E), at 180 days in acute coronary syndrome patients with and without anemia.

Discussion

The main findings of our study were that baseline anemia in patients with ACS, treated according to contemporary practice with radial artery access, mandatory use of modern P2Y12 inhibitors, and low use of GPI, was

associated with a large excess risk of mortality at 180 days. Patients with anemia also experienced higher risk of myocardial reinfarction as well as major bleeding and stroke. No difference in definite stent thrombosis was seen. The results were consistent over time in landmark analyses. A strong correlation between anemia severity and

Table 3. End Points at 180 Days

| | No Anemia (n=4690) | Anemia* (n=792) | 95% CI | P Value | Missing or Unknown, n (%) |
|---|-----------------------|--------------------|---------|---------|------------------------------|
| Mortality | | | | | |
| Kaplan–Meier event rates, n (%) | 96 (2.1) | 54 (6.9) | | <0.001 | 0 (0.0) |
| Unadjusted HR | 1.0 | 3.4 | 2.5–4.8 | <0.001 | 0 (0.0) |
| Multivariable adjusted HR after multiple imputation | 1.0 | 1.9 | 1.3–2.7 | <0.001 | 0 (0.0) |
| Propensity score matching HR | 1.0 | 2.3 | 1.4–3.7 | 0.001 | 0 (0.0) |
| Reinfarction | | | | | |
| Kaplan–Meier event rates, n (%) | 88 (1.9) | 32 (4.3) | | <0.001 | 0 (0.0) |
| Unadjusted HR | 1.0 | 2.3 | 1.5–3.4 | <0.001 | 0 (0.0) |
| Multivariable adjusted HR after multiple imputation | 1.0 | 1.7 | 1.1–2.7 | 0.013 | 0 (0.0) |
| Propensity score matching HR | 1.0 | 2.0 | 1.1–3.7 | 0.022 | 0 (0.0) |
| Major bleeding[†] | | | | | |
| Kaplan–Meier event rates, n (%) | 377 (8.2) | 102 (13.4) | | <0.001 | 0 (0.0) |
| Unadjusted HR | 1.0 | 1.7 | 1.4–2.1 | <0.001 | 0 (0.0) |
| Multivariable adjusted HR after multiple imputation | 1.0 | 1.3 | 1.0–1.6 | 0.041 | 0 (0.0) |
| Propensity score matching HR | 1.0 | 1.4 | 1.0–1.9 | 0.037 | 0 (0.0) |
| Definite stent thrombosis | | | | | |
| Kaplan–Meier event rates, n (%) | 26 (0.6) | 6 (0.8) | | 0.460 | 0 (0.0) |
| Propensity score matching HR | 1.0 | 1.5 | 0.4–5.4 | 0.510 | 0 (0.0) |
| Stroke | | | | | |
| Kaplan–Meier event rates, n (%) | 32 (0.7) | 15 (2.0) | | <0.001 | 0 (0.0) |
| Propensity score matching HR | 1.0 | 2.9 | 1.0–8.0 | 0.042 | 0 (0.0) |

HR indicates hazard ratio.

*Baseline anemia was defined in accordance with the World Health Organization as a hemoglobin value <130 g/L for men and <120 g/L for women.

[†]Major bleeding was defined as 2, 3, or 5 on the Bleeding Academic Research Consortium scale.

increased mortality, myocardial reinfarction, and major bleeding rates was seen, with especially a hemoglobin level <100 g/L being associated with an overall high

clinical event rate. All causes of death (cardiovascular death, fatal bleedings, and other causes of death) were increased.




Figure 3. Landmark analysis at 7 days, for mortality (A) and major bleeding (B), in acute coronary syndrome patients with and without anemia.




Figure 4. Event rates at 180 days for mortality (green), reinfarction (blue), and major bleeding (red), stratified by hemoglobin level categories in acute coronary syndrome patients: severe anemia (hemoglobin <100 g/L); moderate anemia (hemoglobin 100–109 g/L); mild anemia (hemoglobin 110–129 g/L for men and 110–119 g/L for women); normal hemoglobin (130–150 g/L for men and 120–150 g/L for women); and high hemoglobin (>150 g/L).

Results in Context With Previous Studies

The association between baseline anemia and mortality has been studied and validated in several trials, both in the general population²³ and in coronary artery disease.^{10–19} However, these trials have mostly been performed in an era of extensive usage of GPI, which currently is not recommended as standard practice because of excess bleeding risk.²⁴ Furthermore, these studies were carried out when the radial artery was not the default access route.²⁴ Both reducing the use of GPI and using primarily the radial artery as access route has in multiple randomized trials been associated with less serious bleedings.^{3,6} However, the usage of more potent P2Y12 inhibition could in theory increase bleeding rates, especially in patients with anemia.^{7,8} Therefore, we aimed to evaluate how baseline anemia translates into hard clinical outcomes in an era of >90% radial artery access and only bailout use of GPI (<3%) but with modern P2Y12 inhibition. Our results indicate that baseline anemia still constitutes a highly predictive factor for total mortality in this clinical setting. Regarding myocardial reinfarction, previous studies have reported conflicting data. Whereas some studies report an increase in myocardial reinfarction,^{10,11,14} some studies do not show any association.^{12,13,17,25} Our results indicate an increase in myocardial reinfarction in patients with baseline anemia. We also found that the left ventricular ejection fraction of patients with anemia during their hospital stay was lower compared with that of patients without anemia, which might be indicative of a larger infarct size or area at ischemic risk. Likewise, patients with

anemia had higher peak values of high-sensitivity cardiac troponin T, which is a strong predictor of infarct size and systolic dysfunction at 6 months.²⁶ Moreover, we found an increase in bleeding events in patients with anemia, despite low use of GPI and >85% radial artery access in this patient group. Notably, the bleeding risk was particularly high in patients with a hemoglobin value <100 g/L, which is consistent with the PRECISE-DAPT (Predicting Bleeding Complications in Patients Undergoing Stent Implantation and Subsequent Dual Antiplatelet Therapy) risk score.²⁷ The landmark analyses suggest that there is a continued increased bleeding risk even after the acute phase of MI. However, the majority of bleedings occurred within the first week, despite the default radial access use and low GPI use. Some studies have shown a J-shaped association between hemoglobin levels and outcomes.¹² However, this could not be demonstrated from our data, where only low hemoglobin levels appeared to be linked with worse outcomes. Caution should be applied when adding potent antithrombotic agents to patients with anemia during their hospital stay. However, this must be balanced with the increased risk of MI of this patient group.

Mechanisms Behind Mortality, Myocardial Reinfarction, and Major Bleeding

Baseline hemoglobin level may be a surrogate marker of comorbidities, such as malignancies, that are associated with increased bleeding rate as well as risk of thromboembolic

Table 4. Causes of Death for Acute Coronary Syndrome Patients With and Without Anemia

| | No Anemia (n=4690) | Anemia* (n=792) | 95% CI | P Value |
|-----------------------------------|-----------------------|--------------------|----------|---------|
| Cardiovascular death cause | | | | |
| Kaplan–Meier event rates, n (%) | 75 (1.6) | 40 (5.2) | | <0.001 |
| Unadjusted HR | 1.0 | 3.3 | 2.2–4.8 | <0.001 |
| Fatal bleeding | | | | |
| Kaplan–Meier event rates, n (%) | 5 (0.1) | 4 (0.5) | | 0.009 |
| Unadjusted HR | 1.0 | 4.9 | 1.3–18.1 | 0.018 |
| Other death cause | | | | |
| Kaplan–Meier event rates, n (%) | 16 (0.3) | 10 (1.4) | | <0.001 |
| Unadjusted HR | 1.0 | 3.9 | 1.8–8.6 | 0.001 |

HR indicates hazard ratio.

*Baseline anemia was defined in accordance with the World Health Organization as a hemoglobin value <130 g/L for men and <120 g/L for women.

events.²⁸ Anemia could also be a marker of inflammatory disorders that are in turn associated with increased ischemic risk.^{28,29} Furthermore, anemia could also be suggestive of an ongoing occult gastrointestinal bleeding or an increased propensity for bleeding attributable to coagulopathies.¹¹ In accordance with this, fatal bleeding was in our study 5 times more common in patients with anemia. Moreover, patients with anemia more often received blood transfusion, which

has previously been described as a possible independent factor for mortality in ACS patients.³⁰ Other stipulated mechanisms behind anemia and ischemic outcomes could be a decreased oxygen-carrying capacity of the blood that predisposes for ischemia and arrhythmias.¹² Our data suggest that patients with anemia also receive less optimal guideline-directed discharge medication, including aspirin and potent P2Y12 inhibitors, which could predispose to downstream ischemic events. The factors mentioned above may explain our finding that cardiovascular death causes were 3 times as frequent in patients with anemia. This could also be explained by a higher comorbidity burden of cardiovascular risk factors, or discontinuation of antiplatelet therapy due to bleeding events in patients with anemia, which may generate consequential ischemic complications.³¹ Interestingly, radial access was less often used in patients with anemia, highlighting that patients with anemia are a sicker group, where puncture of the radial artery to a lesser degree was deemed possible. This may be attributable to more extensive arteriosclerosis or hemodynamic instability in this patient group, and therefore a femoral approach might have been preferred.

Limitations

Our study is an observational study, and causality between anemia and worse outcomes is therefore uncertain. Furthermore, a larger sample size including more patients with anemia could have shed further light on the topic of anemia and clinical outcomes. More background knowledge of cause of anemia could have added more information on mechanisms behind anemia and worse clinical outcomes. In addition, we




Figure 5. The causes of death for acute coronary syndrome patients who died within 180 days from their index myocardial infarction (n=150). Patients are divided into those who presented without baseline anemia (n=96) and with baseline anemia (n=54).

lacked data on some potential confounders, including malignancy, coagulopathies, and inflammatory disorders, which could have contributed to the higher event rates in patients with anemia. Finally, the hemoglobin value is subject to natural variation due to variation of the intravascular plasma volume that is subject to dehydration/diuretics as well as volume infusion.

Conclusions

Our results indicate that anemia constitutes a high-risk feature in patients with acute coronary syndromes despite treatment with contemporary therapies, including high use of radial artery access, potent P2Y12 inhibitors, and low use of GPI. Especially anemia with a hemoglobin <100 g/L, constitutes a strong negative predictive sign with a substantially increased risk of death, myocardial reinfarction, and major bleeding. The mechanisms are likely multifactorial: Anemia can both be a marker of a sicker patient cohort and have direct physiological consequences, but it may also lead to a lesser degree of guideline-stipulated discharge medications. Nevertheless, in complex patients with anemia and concomitant coronary disease, a multidisciplinary approach is warranted to maximize benefit and minimize patient risk.

Sources of Funding

This study was supported by the Swedish Heart–Lung Foundation, the Swedish Research Council, unrestricted grants from AstraZeneca and the Medicines Company, the Swedish Foundation for Strategic Research (as part of the TOTAL-AMI [Tailoring of Treatment In All Comers With Acute Myocardial Infarction] project), Märta Winkler's Research Foundation, and Thorsten Westerström's Research Foundation.

Disclosures

Dr Szummer has received a modest lecture fee payment from Vifor. Dr Koul has received modest speaker fees from AstraZeneca, Pfizer, BMS. Dr Erlinge has received modest speaker fees and advisory board fees from the Medicines Company and AstraZeneca. The remaining authors have no disclosures to report.

References

1. Steg PG, Huber K, Andreotti F, Arnesen H, Atar D, Badimon L, Bassand JP, De Caterina R, Eikelboom JA, Gulba D, Hamon M, Helft G, Fox KA, Kristensen SD, Rao SV, Verheugt FW, Widimsky P, Zeymer U, Collet JP. Bleeding in acute coronary syndromes and percutaneous coronary interventions: position paper by the Working Group on Thrombosis of the European Society of Cardiology. *Eur Heart J*. 2011;32:1854–1864.
2. Mehran R, Pocock SJ, Stone GW, Clayton TC, Dangas GD, Feit F, Manoukian SV, Nikolsky E, Lansky AJ, Kirtane A, White HD, Colombo A, Ware JH, Moses JW, Ohman EM. Associations of major bleeding and myocardial infarction with the incidence and timing of mortality in patients presenting with non-ST-elevation acute coronary syndromes: a risk model from the ACUITY trial. *Eur Heart J*. 2009;30:1457–1466.
3. Stone GW, Witzeneichler B, Guagliumi G, Peruga JZ, Brodie BR, Dudek D, Kornowski R, Hartmann F, Gersh BJ, Pocock SJ, Dangas G, Wong SC, Kirtane AJ, Parise H, Mehran R; Investigators H-AT. Bivalirudin during primary PCI in acute myocardial infarction. *N Engl J Med*. 2008;358:2218–2230.
4. Stone GW, Clayton T, Deliargyris EN, Prats J, Mehran R, Pocock SJ. Reduction in cardiac mortality with bivalirudin in patients with and without major bleeding: the HORIZONS-AMI trial (Harmonizing Outcomes With Revascularization and Stents in Acute Myocardial Infarction). *J Am Coll Cardiol*. 2014;63:15–20.
5. Kwok CS, Khan MA, Rao SV, Kinnaird T, Sperrin M, Buchan I, de Belder MA, Ludman PF, Nolan J, Loke YK, Mamas MA. Access and non-access site bleeding after percutaneous coronary intervention and risk of subsequent mortality and major adverse cardiovascular events: systematic review and meta-analysis. *Circ Cardiovasc Interv*. 2015;8:e01645.
6. Valgimigli M, Gagnor A, Calabro P, Frigoli E, Leonardi S, Zaro T, Rubartelli P, Briguori C, Ando G, Repetto A, Limbruno U, Cortese B, Sganzerla P, Lupi A, Galli M, Colangelo S, Ierna S, Ausiello A, Presbitero P, Sardella G, Varbella F, Esposito G, Santarelli A, Tresoldi S, Nazzaro M, Zingarelli A, de Cesare N, Rigattieri S, Tosi P, Palmieri C, Brugaletta S, Rao SV, Heg D, Rothenbuhler M, Vranckx P, Juni P; MATRIX Investigators. Radial versus femoral access in patients with acute coronary syndromes undergoing invasive management: a randomised multicentre trial. *Lancet*. 2015;385:2465–2476.
7. Wallentin L, Becker RC, Budaj A, Cannon CP, Emanuelsson H, Held C, Horrow J, Husted S, James S, Katus H, Mahaffey KW, Scirica BM, Skene A, Steg PG, Storey RF, Harrington RA; Investigators P, Freij A, Thorsen M. Ticagrelor versus clopidogrel in patients with acute coronary syndromes. *N Engl J Med*. 2009;361:1045–1057.
8. Wiviott SD, Braunwald E, McCabe CH, Montalescot G, Ruzyllo W, Gottlieb S, Neumann FJ, Ardissino D, De Servi S, Murphy SA, Riesmeyer J, Weerakkody G, Gibson CM, Antman EM; TRITON-TIMI 38 Investigators. Prasugrel versus clopidogrel in patients with acute coronary syndromes. *N Engl J Med*. 2007;357:2001–2015.
9. Nutritional anaemias. Report of a WHO scientific group. *World Health Organ Tech Rep Ser*. 1968;405:5–37.
10. Lawler PR, Filion KB, Dourian T, Atallah R, Garfinkle M, Eisenberg MJ. Anemia and mortality in acute coronary syndromes: a systematic review and meta-analysis. *Am Heart J*. 2013;165:143–153.e145.
11. Bassand JP, Afzal R, Eikelboom J, Wallentin L, Peters R, Budaj A, Fox KA, Joyner CD, Chrolavicius S, Granger CB, Mehta S, Yusuf S; OASIS 5 and OASIS 6 Investigators. Relationship between baseline haemoglobin and major bleeding complications in acute coronary syndromes. *Eur Heart J*. 2010;31:50–58.
12. Brenner SJ, Mehran R, Dangas GD, Ohman EM, Witzeneichler B, Zhang Y, Parvataneni R, Stone GW. Relation of baseline hemoglobin levels and adverse events in patients with acute coronary syndromes (from the acute catheterization and urgent intervention triage strategy and harmonizing outcomes with revascularization and stents in acute myocardial infarction trials). *Am J Cardiol*. 2017;119:1710–1716.
13. Ducrocq G, Puymirat E, Steg PG, Henry P, Martelet M, Karam C, Schiele F, Simon T, Danchin N. Blood transfusion, bleeding, anemia, and survival in patients with acute myocardial infarction: FAST-MI registry. *Am Heart J*. 2015;170:726–734.e722.
14. Kunadian V, Mehran R, Lincoff AM, Feit F, Manoukian SV, Hamon M, Cox DA, Dangas GD, Stone GW. Effect of anemia on frequency of short- and long-term clinical events in acute coronary syndromes (from the Acute Catheterization and Urgent Intervention Triage Strategy Trial). *Am J Cardiol*. 2014;114:1823–1829.
15. Mamas MA, Kwok CS, Kontopantelis E, Fryer AA, Buchan I, Bachmann MO, Zaman MJ, Myint PK. Relationship between anemia and mortality outcomes in a national acute coronary syndrome cohort: insights from the UK Myocardial Ischemia National Audit Project Registry. *J Am Heart Assoc*. 2016;5:e003348. DOI: 10.1161/JAHA.116.003348.
16. Meneveau N, Schiele F, Seronde MF, Descotes-Genon V, Oettinger J, Chopard R, Ecarnot F, Bassand JP; Reseau de Cardiologie de Franche Comte. Anemia for risk assessment of patients with acute coronary syndromes. *Am J Cardiol*. 2009;103:442–447.
17. Tsujita K, Nikolsky E, Lansky AJ, Dangas G, Fahy M, Brodie BR, Dudek D, Mockel M, Ochala A, Mehran R, Stone GW. Impact of anemia on clinical outcomes of patients with ST-segment elevation myocardial infarction in relation to gender and adjunctive antithrombotic therapy (from the HORIZONS-AMI trial). *Am J Cardiol*. 2010;105:1385–1394.

18. Younger JO, Nauta ST, Akkerhuis KM, Deckers JW, van Domburg RT. Effect of anemia on short- and long-term outcome in patients hospitalized for acute coronary syndromes. *Am J Cardiol.* 2012;109:506–510.
19. Gillespie TW. Effects of cancer-related anemia on clinical and quality-of-life outcomes. *Clin J Oncol Nurs.* 2002;6:206–211.
20. Erlinge D, Omerovic E, Frobert O, Linder R, Danielewicz M, Hamid M, Swahn E, Henareh L, Wagner H, Hardhammar P, Sjogren I, Stewart J, Grimfjard P, Jensen J, Aasa M, Robertsson L, Lindroos P, Haupt J, Wikstrom H, Ulvenstam A, Bhiladvala P, Lindvall B, Lundin A, Todt T, Ioanes D, Ramunddal T, Kellerth T, Zagozdzon L, Gotberg M, Andersson J, Angeras O, Ostlund O, Lagerqvist B, Held C, Wallentin L, Schersten F, Eriksson P, Koul S, James S. Bivalirudin versus heparin monotherapy in myocardial infarction. *N Engl J Med.* 2017;377:1132–1142.
21. Erlinge D, Koul S, Eriksson P, Schersten F, Omerovic E, Linder R, Ostlund OP, Wallentin L, Frobert O, James S. Bivalirudin versus heparin in non-ST and ST-segment elevation myocardial infarction—a registry-based randomized clinical trial in the SWEDEHEART registry (the VALIDATE-SWEDEHEART trial). *Am Heart J.* 2016;175:36–46.
22. Thygesen K, Alpert JS, Jaffe AS, Simoons ML, Chaitman BR, White HD; Writing Group on the Joint ESC/ACCF/AHA/WHF Task Force for the Universal Definition of Myocardial Infarction, Thygesen K, Alpert JS, White HD, Jaffe AS, Katus HA, Apple FS, Lindahl B, Morrow DA, Chaitman BA, Clemmensen PM, Johanson P, Hod H, Underwood R, Bax JJ, Bonow RO, Pinto F, Gibbons RJ, Fox KA, Atar D, Newby LK, Galvani M, Hamm CW, Uretsky BF, Steg PG, Wijns W, Bassand JP, Menasche P, Ravkilde J, Ohman EM, Antman EM, Wallentin LC, Armstrong PW, Simoons ML, Januzzi JL, Niemenen MS, Gheorghide M, Filippatos G, Luepker RV, Fortmann SP, Rosamond WD, Levy D, Wood D, Smith SC, Hu D, Lopez-Sendon JL, Robertson RM, Weaver D, Tendera M, Bove AA, Parkhomenko AN, Vasilieva EJ, Mendis S; ESC Committee for Practice Guidelines. Third universal definition of myocardial infarction. *Eur Heart J.* 2012;33:2551–2567.
23. Martinsson A, Andersson C, Andell P, Koul S, Engstrom G, Smith JG. Anemia in the general population: prevalence, clinical correlates and prognostic impact. *Eur J Epidemiol.* 2014;29:489–498.
24. Neumann FJ, Sousa-Uva M, Ahlsson A, Alfonso F, Banning AP, Benedetto U, Byrne RA, Collet JP, Falk V, Head SJ, Juni P, Kastrati A, Koller A, Kristensen SD, Niebauer J, Richter DJ, Seferovic PM, Sibbing D, Stefanini GG, Windecker S, Yadav R, Zembala MO; Group ESCSD. 2018 ESC/EACTS guidelines on myocardial revascularization. *Eur Heart J.* 2019;40:87–165.
25. Voeltz MD, Patel AD, Feit F, Fazel R, Lincoff AM, Manoukian SV. Effect of anemia on hemorrhagic complications and mortality following percutaneous coronary intervention. *Am J Cardiol.* 2007;99:1513–1517.
26. Mohammad MA, Koul S, Smith JG, Noc M, Lang I, Holzer M, Clemmensen P, Jensen U, Engstrom T, Arheden H, James S, Lindahl B, Metzler B, Erlinge D. Predictive value of high-sensitivity troponin T for systolic dysfunction and infarct size (six months) after ST-elevation myocardial infarction. *Am J Cardiol.* 2018;122:735–743.
27. Valgimigli M, Bueno H, Byrne RA, Collet JP, Costa F, Jeppsson A, Juni P, Kastrati A, Kolh P, Mauri L, Montalescot G, Neumann FJ, Petricevic M, Roffi M, Steg PG, Windecker S, Zamorano JL, Levine GN; ESC Scientific Document Group, ESC Committee for Practice Guidelines, ESC National Cardiac Societies. 2017 ESC focused update on dual antiplatelet therapy in coronary artery disease developed in collaboration with EACTS: the Task Force for dual antiplatelet therapy in coronary artery disease of the European Society of Cardiology (ESC) and of the European Association for Cardio-Thoracic Surgery (EACTS). *Eur Heart J.* 2018;39:213–260.
28. Adamson JW. The anemia of inflammation/malignancy: mechanisms and management. *Hematology Am Soc Hematol Educ Program.* 2008;1:159–165.
29. Sager HB, Nahrendorf M. Inflammation: a trigger for acute coronary syndrome. *Q J Nucl Med Mol Imaging.* 2016;60:185–193.
30. Rao SV, Jollis JG, Harrington RA, Granger CB, Newby LK, Armstrong PW, Moliterno DJ, Lindblad L, Pieper K, Topol EJ, Stamler JS, Califf RM. Relationship of blood transfusion and clinical outcomes in patients with acute coronary syndromes. *JAMA.* 2004;292:1555–1562.
31. Dewilde WJ, Oirbans T, Verheugt FW, Kelder JC, De Smet BJ, Herrman JP, Adriaenssens T, Vrolix M, Heestermans AA, Vis MM, Tijssen JG, van 't Hof AW, ten Berg JM; WOEST Study Investigators. Use of clopidogrel with or without aspirin in patients taking oral anticoagulant therapy and undergoing percutaneous coronary intervention: an open-label, randomised, controlled trial. *Lancet.* 2013;381:1107–1115.

Supplemental Material

Table S1. Kaplan-Meier event rates for patients with and without anemia who received heparin vs bivalirudin.

| | No Anemia n=4690 | | p-value | Anemia n=792 | | p-value |
|---------------------|---------------------|-----------------------|---------|------------------|----------------------|---------|
| | Heparin n=2372 | Bivalirudin n=2318 | | Heparin n=385 | Bivalirudin n=407 | |
| Mortality n(%) | 51(2.2) | 45(2.0) | 0.617 | 23(6.1) | 31(7.8) | 0.364 |
| Reinfarction n(%) | 52(2.2) | 36(1.6) | 0.106 | 16(4.4) | 16(4.2) | 0.889 |
| Major bleeding n(%) | 192(8.2) | 185(8.1) | 0.850 | 50(13.4) | 52(13.3) | 0.917 |

*Baseline anemia was defined in accordance with WHO as a hemoglobin value <130 g/L for men and <120 g/L

for women. †Major bleeding was defined as 2, 3, or 5, on the Bleeding Academic Research Consortium scale.

Table S2. Sensitivity analyses for endpoints at 180 days after propensity score matching with downstream adjustments for body mass index (BMI).



| | No Anemia n=4690 | Anemia n=792 | 95% CI | p-value | Missing or unknown n(%) |
|--|---------------------|-----------------|---------|---------|----------------------------|
| Mortality - hazard ratio | 1.0 | 2.2 | 1.3-3.8 | 0.002 | 156(10.2) |
| Reinfarction – hazard ratio | 1.0 | 2.2 | 1.2-4.2 | 0.013 | 156(10.2) |
| Major bleeding – hazard ratio | 1.0 | 1.4 | 1.0-1.9 | 0.041 | 156(10.2) |
| Definite stent thrombosis – hazard ratio | 1.0 | 1.0 | 0.3-3.9 | 0.979 | 156(10.2) |
| Stroke – hazard ratio | 1.0 | 2.7 | 0.9-7.6 | 0.066 | 156(10.2) |


*Baseline anemia was defined in accordance with WHO as a hemoglobin value <130 g/L for men and <120 g/L

for women. Major bleeding was defined as 2, 3, or 5, on the Bleeding Academic Research Consortium scale.

†Abbreviations: CI=confidence interval

Figure S1. Kaplan-Meier failure functions for mortality (A), myocardial reinfarction (B), major bleeding (C), definite stent thrombosis (D), and stroke (E), at 180 days in acute coronary syndrome patients with and without anemia after propensity score matching.




C

Number at risk

| | | | | |
|-----------|-----|-----|-----|-----|
| Anemia | 762 | 636 | 618 | 606 |
| No Anemia | 762 | 685 | 671 | 658 |

No Anemia — Anemia

D

Number at risk

| | | | | |
|-----------|-----|-----|-----|-----|
| Anemia | 762 | 704 | 694 | 691 |
| No Anemia | 762 | 729 | 726 | 724 |

No Anemia — Anemia

