



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Total sleep deprivation increases pain sensitivity, impairs conditioned pain modulation and facilitates temporal summation of pain in healthy participants

Staffe, Alexander Torp; Bech, Mathias Winther; Clemmensen, Sara Louise Kjær; Nielsen, Henriette Tranberg; Larsen, Dennis Boye; Petersen, Kristian Kjær

Published in:
PLOS ONE

DOI (link to publication from Publisher):
[10.1371/journal.pone.0225849](https://doi.org/10.1371/journal.pone.0225849)

Creative Commons License
CC BY 4.0

Publication date:
2019

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Staffe, A. T., Bech, M. W., Clemmensen, S. L. K., Nielsen, H. T., Larsen, D. B., & Petersen, K. K. (2019). Total sleep deprivation increases pain sensitivity, impairs conditioned pain modulation and facilitates temporal summation of pain in healthy participants. *PLOS ONE*, *14*(12), [e0225849]. <https://doi.org/10.1371/journal.pone.0225849>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- ? Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- ? You may not further distribute the material or use it for any profit-making activity or commercial gain
- ? You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

RESEARCH ARTICLE

Total sleep deprivation increases pain sensitivity, impairs conditioned pain modulation and facilitates temporal summation of pain in healthy participants

Alexander Torp Staffe¹, Mathias Winther Bech¹, Sara Louise Kjær Clemmensen¹, Henriette Tranberg Nielsen¹, Dennis Boye Larsen¹, Kristian Kjær Petersen^{1,2*}

1 SMI, Department of Health Science and Technology, Faculty of Medicine, Aalborg University, Aalborg, Denmark, **2** Department of Health Science and Technology, Center for Neuroplasticity and Pain, Faculty of Medicine, Aalborg University, Aalborg, Denmark

* kkp@hst.aau.dk



OPEN ACCESS

Citation: Staffe AT, Bech MW, Clemmensen SLK, Nielsen HT, Larsen DB, Petersen KK (2019) Total sleep deprivation increases pain sensitivity, impairs conditioned pain modulation and facilitates temporal summation of pain in healthy participants. *PLoS ONE* 14(12): e0225849. <https://doi.org/10.1371/journal.pone.0225849>

Editor: Giuseppe Biagini, University of Modena and Reggio Emilia, ITALY

Received: May 31, 2019

Accepted: November 13, 2019

Published: December 4, 2019

Copyright: © 2019 Staffe et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the manuscript and its Supporting Information files.

Funding: KKP was supported by The Aalborg University Talent Management Programme (j.no. 771126) for this study. KKP is partly hired by Center for Neuroplasticity and Pain (CNAP) to conduct this study and CNAP is supported by the Danish National Research Foundation (DNRF121). The funders had no role in study design, data

Abstract

Chronic pain patients often suffer from insomnia or impaired sleep which has been associated with increased pain sensitivity, but a limited amount of studies have investigated the effects of total sleep deprivation on central pain mechanisms. Therefore, the aim of this study was to determine the effects of total sleep deprivation on temporal summation, conditioned pain modulation, thermal and pressure pain sensitivity in healthy participants. Twenty-four healthy participants took part in this two-session trial. The measurements were conducted after a night of habitual sleep (baseline) and following 24 hours of total sleep deprivation. Detection thresholds for cold and warmth and pain thresholds for cold and heat were assessed. Cuff induced pressure pain detection and tolerance thresholds, temporal summation and conditioned pain modulation were assessed with user-independent, computer-controlled cuff algometry. Conditioned pain modulation was significantly impaired, temporal summation was significantly facilitated and pain sensitivity to pressure and cold pain were significantly increased at follow-up compared with baseline. In conclusion, this study found that one night of total sleep deprivation impaired descending pain pathways, facilitated spinal excitability and sensitized peripheral pathways to cold and pressure pain. Future studies are encouraged to investigate if sleep therapy might normalize pain sensitivity in sleep-deprived chronic pain patients.

Introduction

The sleep quality declines with various chronic pain conditions as shown in, e.g., fibromyalgia [1], burn injuries [2], and back pain [3]. Studies have reported that at least 50% of patients with diverse chronic pain conditions suffer from sleep impairments [4]. Additionally, sleep impairments have been described as valuable predictors for new incidences and worsening of symptoms linked to chronic pain conditions like fibromyalgia [5], rheumatoid arthritis (RA)

collection and analysis, decision to publish, or preparation of the manuscript

Competing interests: NociTech is partly owned by Aalborg University and that I, K. K. Petersen, was partly employed by NociTech from January 2015 to December 2016. The work conducted by me, K. K. Petersen, in this period did not affect any of my research in that period or in future projects. The remaining authors have no conflicts of interest to declare. This does not alter our adherence to PLOS ONE policies on sharing data and materials.

[6], and orofacial and back pain [7] as patients with co-occurring sleep impairments seem to experience worsened pain symptoms [4,8].

Though the underlying mechanisms connecting sleep impairments and decreased descending pain inhibitory control are unclear, brain areas such as the periaqueductal gray are known to modulate both sleep stages and nociception [9]. Furthermore, the raphe nuclei are involved in the descending pain control system [10] and the ascending reticular activating system (ARAS), which is responsible for the transition between sleep and wakefulness [11]. To some degree, overlapping functions in these areas of the brain may account for the link between sleep loss and decreased pain inhibition [4,12–15]. However, this encourages further investigation into the connection between sleep loss and central pain processing.

Mechanistic pain profiling assesses the underlying pain mechanisms in the peripheral and central nervous system and includes pain thresholds, temporal summation of pain (TSP), and conditioned pain modulation (CPM) [16]. TSP and CPM are the human surrogate models for wind-up and descending pain inhibitory control, respectively [17,18]. CPM protocols reflect endogenous pain inhibition by measuring the inhibition of a nociceptive stimulus when interceded by a secondary conditioning stimulus (CS), otherwise referred to as the “pain inhibits pain phenomenon”. [17]. An impaired CPM is therefore reflected by a reduced pain inhibits pain effect. It is well-established that patients suffering from back pain [19], fibromyalgia [20], or severe osteoarthritis [21] exhibit reduced descending pain inhibitory control and that impaired CPM is associated with worsening of pain [22]. In addition, CPM has been shown to be impaired following sleep impairment [4,23]. For instance, decreased sleep quality is associated with impaired CPM in healthy participants after three days of fragmented sleep [24] and in patients suffering from temporomandibular joint disorder [25]. TSP mimics the wind-up process, which indicates central sensitization and is interpreted as the gradual increase in pain sensitivity when continuously exposed to stimuli with a constant intensity. Facilitated TSP is reflected by an intensified gradual response to pain stimuli. TSP is facilitated in multiple chronic pain conditions [22] and emerging evidences suggests that facilitated TSP might be a more reliable predictor for poor response to standard pain therapy [16,26–28]. TSP is facilitated in many chronic pain conditions [29–31]. Yet, limited evidence exists on the effect of total sleep deprivation (TSD) on TSP. For example, some studies have observed increased TSP in participants with sleep impairments due to prolonged REM sleep or osteoarthritis [32,33]. In support, one night of TSD increased the thermal and pressure pain sensitivity in healthy participants [34] and 60 hours of sleep deprivation decreased the pressure pain tolerance [35]. Nonetheless, the evidence supporting thermal hyperalgesia is conflicting as some studies finds increased pain sensitivity in both heat pain thresholds (HPT) and cold pain thresholds (CPT) among healthy participants subjected to 24 hours of TSD [36,37]. Contrary to this, other studies report no significant changes in heat pain thresholds among healthy participants after a night of TSD [35,38]. A better understanding of how sleep impairment influences central and peripheral pain mechanisms is important to further develop its use in clinical practice [39], either by implementing sleep therapy or through pharmacological treatment targeting endogenous pain modulation. Therefore, the current study aimed to investigate changes in peripheral pain by applying pressure and assessing the thermal pain thresholds and central pain mechanisms employing TSP and CPM in healthy participants before and after 24 hours of TSD.

Methods

Participants

Twenty-five healthy participants (nine women) (average age 22.6 ± 2.02 years) were recruited at Aalborg University through advertisements. The participants received detailed written and

oral information and signed an informed consent form prior to enrollment in the experiment. The study was approved by The North Denmark Region Committee on Health Research Ethics (N-20180089) and was conducted in agreement with the Helsinki Declaration.

Experimental design

The experiment consisted of two identical sessions conducted on two consecutive days. In-between sessions, the participants underwent 24 hours of TSD. To ensure that the participants were awake, they were instructed to send an hourly text message. The experimental pain stimuli were applied in the order of heat, cold, pressure, TSP and CPM. In order to control for order effects, test modalities were conducted in the same order for all participants in both sessions.

Questionnaires

Prior to the baseline measurements, the participants were asked to report both sleep duration and sleep quality the night before. The participants also filled out the Pittsburgh Sleep Quality index (PSQI) and the Pain Catastrophizing Scale (PCS). The PSQI is a validated tool for assessment of sleep quality [40] in which participants evaluate their sleep based on a four-point Likert-type scale (ranging from 0 = “not at all” to 3 = “three times or more per week”), with high scores indicating poor sleep quality [41]. The PCS questionnaire is a validated tool for assessment of thoughts regarding pain [42,43] based on a five-point Likert-type scale (ranging from 0 = “not at all” to 4 = “all the time”) with a high score indicating a high level of catastrophizing thoughts concerning pain [44].

Thermal stimuli

Cold and warm detection thresholds (CDT and WDT) and cold and heat pain thresholds (CPT and HPT) were assessed using the Medoc pathway system (Medoc, Israel) in accordance with the protocol of the German Research Network on Neuropathic Pain [45]. Briefly, the 3x3 cm ATS Probe (Medoc, Israel) was placed on the volar forearm, 3 cm below the elbow. Thermal stimuli were applied with ramped stimuli (1°C/s) which were terminated when the participant pressed a button. When assessing CDT or WDT, the participants were instructed to press the stop button as soon as they experienced a change of temperature to either “cool” or “warm”. When assessing CPT or HPT, the participants were instructed to press the stop button immediately following the initial sensation of pain. Cut-off temperatures were 0°C and 50°C and baseline temperature was 32°C. Three consecutive stimuli were applied in each test. The results were averaged to reflect CDT, WDT, CPT, or HPT.

Pressure stimuli

Cuff pressure detection thresholds (cPDTs), tolerance thresholds (cPTT), TSP, and CPM were assessed by a cuff algometer (Cortex Technology, Hadsund, Denmark, and Aalborg University, Denmark). A cuff was placed on the belly of the gastrocnemius muscle of the participants in agreement with Graven-Nielsen et al. [46]. The cuff was inflated at a progression rate of 1 kPa/s and was set to a maximum limit of 100 kPa. The participants were instructed to rate the level of pain as soon as pain was detected on a visual analogue scale (VAS).

Pressure pain thresholds

cPDT was defined as the pain-inducing amount of pressure, equivalent to 1 cm on the VAS. The amount of pressure applied causing unbearable pain was defined as cPTT. cPDT and

cPTT values were determined as the value of the measurements performed on the dominant leg.

Temporal summation of pain

TSP was assessed using a mechanical pressure stimulus. The stimuli were applied 10 times with 1 s interstimulus interval and duration [30]. The applied pressure was equal to the cPTT. The participants were instructed to continuously rate their pain on a VAS. An average of the VAS scores measured during the first three pulses and an average of the VAS scores measured during the last three pulses were used to determine TSP. The difference between these averages was interpreted as TSP [47].

Conditioned pain modulation

On the non-dominant leg, the cuff was promptly inflated to a pressure corresponding to 70% of the cPTT. The cuff on the dominant leg was then inflated at a rate of 1 kPa/s. The participants were instructed to rate the pain on their dominant leg. CPM was calculated as the difference between cPDT with and without the conditioning stimuli. cPDT without the conditioning stimulus was measured as the cuff pressure detection threshold on the dominant leg, when no cuff was inflated on the non-dominant leg.

Statistics

All statistical analyses were conducted using SPSS version 25 (IBM, USA). For all parameters, including thermal detection thresholds (WDT, CDT), pain sensitivity (HPT, CPT, cPDT, cPTT), and central pain mechanisms (TSP, CPM), separate paired sample t-tests were used to compare the differences of means between measurements (pre-TSD, post-TSD). If the data violated the assumptions for parametric data, a Wilcoxon signed-rank test was performed. Bonferroni correction was applied to account for multiple comparisons when assessing the descending pain inhibitory control ($0.05 / 2$ comparisons = 0.025). The level of significance was set to $P \leq 0.05$. Unless otherwise stated, all data are presented as means \pm standard deviation (SD).

Results

Demographics

Twenty-four healthy participants (eight women) with no history of chronic, mental, musculoskeletal, or neurological illness participated in this study (average age 22.6 ± 2.04 years). Twenty-five participants were initially recruited, but one participant was subsequently excluded due to a previously undisclosed history of mental illness. Due to technical issues during the data collection of CDT, WDT, CPT, and HPT from one participant, the thermal stimuli data are based on 23 participants, whereas TSP, CPM, cPDT, and cPTT are based on 24 participants. Demographics of the 24 participants are presented in [Table 1](#).

Thermal stimuli

No significant changes were found when comparing baseline (CDT: 29.5° Celsius ± 1.03 ; WDT: $34.6^\circ \pm 0.77$) and follow-up (CDT: 29° Celsius ± 1.3 ; WDT: $34.7^\circ \pm 1.24$) data for CDT ([Fig 1](#); $t = 1.36$, $p = 0.19$) and WDT ([Fig 1](#); $z = 0.99$, $p = 0.32$). Significantly decreased CPT was observed at follow-up ($15.6^\circ \pm 8.51$) compared with baseline ($12.1^\circ \pm 8.9$, [Fig 1](#); $z = 2.3$, $p = 0.02$). Conversely, HPT showed no difference when comparing follow-up ($44.1^\circ \pm 2.7$) with baseline ($44.6^\circ \pm 2.7$, [Fig 1](#); $t = 0.96$, $p = 0.35$).

Table 1. An overview of demographics, PSQI, sleep duration, and sleep quality at baseline for all participants (mean \pm SD).

Age (years)	22.6 (19–27)
Sex (% females)	33.3%
PSQI	5.04 \pm 1.71
Sleep duration before baseline (hours)	6.85 \pm 1.11
Habitual sleep duration (hours)	7.25 \pm 0.69
Sleep quality before baseline (0–10)	6.92 \pm 1.63
Habitual sleep quality (0–10)	7.48 \pm 1.44

Data are presented as mean \pm SD. PSQI, Pittsburgh Sleep Quality Index. Sleep quality was quantified on a scale from 0 to 10, in which 0 was the worst possible sleep quality and 10 was the best possible sleep quality.

<https://doi.org/10.1371/journal.pone.0225849.t001>

Pressure pain thresholds

cPDT was significantly reduced after 24 hours of TSD (38.8 kPa \pm 11.85) compared with baseline (42.13 kPa \pm 10.45, Fig 2; $t = 2.22$, $p = 0.037$). Similarly, cPTT was significantly reduced at follow-up (83.9 kPa \pm 17.11) compared with baseline (87.83 kPa \pm 14.45, Fig 2; $z = -2.11$, $p = 0.03$).

Temporal summation of pain

TSP was significantly facilitated at follow-up (2.27 VAS \pm 1.66) compared with baseline (1.59 VAS \pm 1.23, Fig 3; $t = -2.68$, $p = 0.01$).

Conditioned pain modulation

cPDT was significantly increased with conditioning stimulus compared with cPDT without conditioning stimulus at baseline (Fig 4; $t = -3.63$, $p = 0.002$, Bonferroni-corrected). Conversely, cPDT did not increase during conditioning after 24 hours of TSD ($t = -1.81$, $p = 0.168$, Bonferroni-corrected).




Fig 1. Thermal QST data before and after total sleep deprivation (mean \pm SD). A significant decrease in cold pain threshold (CPT) was found after total sleep deprivation (Post-TSD) compared to before TSD (Pre-TSD), whereas cold and warm detection thresholds (CDT and WDT), and heat pain threshold (HPT) were unaffected. *, $p < 0.05$.

<https://doi.org/10.1371/journal.pone.0225849.g001>




Fig 2. Cuff induced pain detection and tolerance thresholds before and after total sleep deprivation (mean \pm SD). Cuff induced pain detection (cPDT) and tolerance thresholds (cPTT) significantly decreased after total sleep deprivation (Post-TSD) compared with before TSD (Pre-TSD). *, $p < 0.05$.

<https://doi.org/10.1371/journal.pone.0225849.g002>

Discussion

This study is the first to demonstrate that 24-hours of TSD impairs CPM, facilitates TSP, and increases pain sensitivity to pressure and cold pain stimuli in one experimental setup, indicating that TSD affects both central and peripheral pain pathways.




Fig 3. Temporal summation of pain before and after total sleep deprivation (mean \pm SD). Temporal summation of pain was significantly increased after total sleep deprivation (Post-TSD) compared to before TSD (Pre-TSD). *, $p < 0.05$.

<https://doi.org/10.1371/journal.pone.0225849.g003>




Fig 4. Conditioned pain modulation before and after total sleep deprivation (mean \pm SD). Conditioned pain modulation is assessed using an unconditioned (white bars) and conditioned cuff pain detection threshold (cPDT). Subjects displayed a significant (*, $P < 0.05$) increase in conditioned cPDT compared with unconditioned cPDT at baseline but not following 24 hours of total sleep deprivation (TSD).

<https://doi.org/10.1371/journal.pone.0225849.g004>

Sleep impairment and descending pain inhibitory control

An association between impaired CPM and decreased sleep quality has been shown in several studies investigating sleep impairment in patients with conditions such as RA [6], fibromyalgia [48], and insomnia [49]. The current study sought to investigate whether a similar relationship was present in healthy participants as it would indicate whether the relationship between sleep and pain modulation is limited to chronic pain conditions.

The current study found impaired CPM after 24 hours of TSD, which is in line with a recent study conducted by Eichhorn et al. [50], who demonstrated impaired descending pain inhibitory control following one night of TSD. In contrast, Smith et al. [51] assessed CPM and found reduced descending pain inhibition after partial sleep deprivation (PSD) but found no alterations in CPM following 36 hours of TSD. Finally, Matre et al. [52] found a significantly improved CPM following PSD. These conflicting results concerning the effect of sleep deprivation on CPM may be explained by the heterogeneity in the methodological approaches to the CPM protocols [53]; the current study used cuff algometry whereas earlier studies used a cold-pressor test.

The findings of the current study support the theory that sleep impairments decrease the effect of the descending pain inhibition. The underlying mechanism of the interaction between sleep impairments and impaired descending pain inhibition remains unclear. However, a theory may be that brain areas such as the periaqueductal gray and the raphe nuclei both are involved in the descending pain control system [10], nociception [9], and modulation of sleep [11]. Therefore, neurotransmitter alterations in these areas due to sleep impairments could cause a decrease in the effect of descending pain inhibitory control. This could explain why sleep impairments are often seen in chronic pain patients.

A better understanding of the interaction between sleep and central pain mechanisms could possibly improve the treatment options for chronic pain patients since sleep impairment

is a major problem among chronic pain patients [6,48,49]. Future studies are encouraged to investigate whether the effects of sleep impairments on central pain mechanisms are reversible through sleep therapy.

Sleep impairment and temporal summation of pain

Central sensitization, demonstrated as TSP, represents an essential pathophysiological process that augments the development and maintenance of pain conditions in several clinical contexts [30,54]. TSP has been found to be increased in women suffering from insomnia [32] and in individuals with increased REM-sleep [33]. In contrast, Schuh-Hofer et al. [34] found increased sensitivity to pinprick stimuli and hyperalgesia to cold but found no changes in TSP. Similarly, no association was found between increased TSP and sleep impairments in patients with primary insomnia [49], fibromyalgia [55], and restless legs syndrome [56]. However, previous studies tested TSP using heat pain, and since pressure pain is primarily mediated by A-fiber nociceptor inputs [57] while heat pain depends mainly on C-fiber inputs [58], it was hypothesized that targeting different nociceptive pathways might produce separate results. The current study found that pressure-induced TSP was facilitated following 24 hours of TSD, which is in contrast to previous findings. This could be explained by different methodological setups and by the fact that the aforementioned studies included patients as subjects. Facilitated TSP following 24 hours of TSD indicates increased central sensitization of pain following sleep loss. Sleep therapy may be a possible invention method for pain relief.

Sleep and pain sensitivity

A meta-analysis by Schrimpf et al. [59] demonstrated that decreased sleep quality was associated with increased pain sensitivity across a range of different test modalities such as pressure algometry, laser stimuli, and thermal tests. In this respect, studies have reported that CPT increased [36] and HPT decreased [60] following TSD. Additionally, decreased sleep quality has also been linked to decreased mechanical pain thresholds [34,61,62]. For instance, Chiu et al. [61] found a decrease in PPTs in participants with self-reported poor sleep, while Ağar-gün et al. [63] found a negative correlation between pain and sleep quality as poor sleep led to an increase in pain sensitivity in patients with fibromyalgia.

Increased pain sensitivity has also been reported following experimentally induced sleep deprivation as Azevedo et al. [64] found a significant increase in thermal and mechanical sensitivity after two nights of TSD. Schuh-Hofer et al. [34] found that only one night of TSD was able to promote a state of generalized hyperalgesia among the participants. Like the current study, Schuh-Hofer et al. found no alterations in CDT and WDT. The fact that no changes were demonstrated in non-nociceptive stimulations led Schuh-Hofer et al. [34] to conclude that the sensory alterations following sleep loss were purely nociceptive. The hyperalgesic effects of 24 hours of TSD in healthy participants have also been demonstrated using a cold pressor test [65,66] and radiant heat laser pulses [64]. Together with the results of this study, these previous results support the notion that TSD causes generalized hyperalgesia detectable across a number of QST modalities.

Even reduced sleep duration seems to affect the pain sensitivity as participants with a short sleep duration (< 6 hours) reported increased pain sensitivity the day following the short sleep duration [67]. These results indicate that while TSD does increase the pain sensitivity, even disturbed sleeping patterns may cause the pain sensitivity to increase. This sentiment is supported by Onen et al. [35], who tested the pain sensitivity following TSD as well as PSD. While a significantly decreased cPDT was found after 40 hours of PSD, Onen et al. [35] found the hypersensitivity to be more extensive after TSD compared with PSD, supporting that both sleep loss and impairments increase the pain sensitivity. The connection between sleep restriction and pain

sensitivity has also been applied to clinical trials as Fitzgerald et al. [68] found increased pain sensitivity when examining self-reported pain in RA patients after PSD. Though multiple sleep disturbance paradigms have produced results in the past, the current study applied TSD as it seems to produce hyperalgesia more extensively and consistently than PSD.

In contrast, Arima et al. [69] reported no interaction between partial sleep deprivation and pain sensitivity and Onen et al. [35] found no significant changes in thermal pain after TSD and PSD. However, Onen et al. [35] found hyperalgesia to mechanical stimuli and suggested that these results might have been affected by undetected differences in skin surface temperatures or the fact that thermal tests generally show less reliably when it comes to discriminating small changes in pain thresholds [70].

A generalized hyperalgesic effect from TSD is widely reported [34], with mechanical stimulus tests seemingly producing the most consistent results. The findings presented in the current study support earlier results as the participants exhibited mechanic and thermal hyperalgesia following 24 hours of TSD.

Limitations

This study monitored TSD by hourly text messages and this procedure does not guarantee that the participants did not sleep in between. However, one study observed that partial sleep deprivation has a greater effect than TSD regarding CPM [51]. Therefore, it is unlikely to have impacted the current findings even if some participants did fall asleep. The participants in the current study scored an average of more than 5 on the PSQI and averaged 6.85 hours of sleep the night before the baseline session. The fact that the sample in this study displayed both lowered general sleep quality and average sleep duration the night before baseline suggests that this current sample may be less sensitive to sleep deprivation. Despite this, it is noteworthy that the current study still demonstrated an effect on the central pain mechanisms. A sample with higher average sleep quality and sleep duration closer to recommended amounts may have produced an even more profound hyperalgesic response.

Conclusion

The current study is the first to demonstrate that TSD impairs CPM, facilitates TSP, and increases pain sensitivity to pressure and cold pain stimuli, indicating that TSD affects both central and peripheral pain pathways. Future studies are encouraged to investigate the underlying mechanism of TSP, the effects on both the central and peripheral nervous system, and whether the effects are reversible through sleep therapy.

Supporting information

S1 Table. cPDT and cPTT. Raw data for cuff pressure pain detection & tolerance thresholds before and after total sleep deprivation.
(DOCX)

S2 Table. CPM. Raw pressure detection thresholds (conditioned and unconditioned) measured before and after total sleep deprivation.
(DOCX)

S3 Table. TSP. Difference between the first and last three pain intensity ratings during temporal summation of pain.
(DOCX)

S4 Table. Thermal QST. Averaged cold detection threshold, warm detection threshold, cold pain threshold, and heat pain threshold values before and after total sleep deprivation. (DOCX)

Author Contributions

Conceptualization: Alexander Torp Staffe, Mathias Winther Bech, Sara Louise Kjær Clemmensen, Henriette Tranberg Nielsen, Dennis Boye Larsen.

Data curation: Alexander Torp Staffe, Mathias Winther Bech, Sara Louise Kjær Clemmensen, Henriette Tranberg Nielsen, Kristian Kjær Petersen.

Formal analysis: Alexander Torp Staffe, Mathias Winther Bech, Sara Louise Kjær Clemmensen, Henriette Tranberg Nielsen, Dennis Boye Larsen, Kristian Kjær Petersen.

Funding acquisition: Kristian Kjær Petersen.

Investigation: Alexander Torp Staffe, Mathias Winther Bech, Sara Louise Kjær Clemmensen, Henriette Tranberg Nielsen.

Methodology: Alexander Torp Staffe, Mathias Winther Bech, Sara Louise Kjær Clemmensen, Henriette Tranberg Nielsen, Dennis Boye Larsen, Kristian Kjær Petersen.

Project administration: Alexander Torp Staffe, Mathias Winther Bech, Sara Louise Kjær Clemmensen, Henriette Tranberg Nielsen, Kristian Kjær Petersen.

Resources: Kristian Kjær Petersen.

Supervision: Dennis Boye Larsen, Kristian Kjær Petersen.

Validation: Sara Louise Kjær Clemmensen, Dennis Boye Larsen, Kristian Kjær Petersen.

Writing – original draft: Alexander Torp Staffe, Mathias Winther Bech, Sara Louise Kjær Clemmensen, Henriette Tranberg Nielsen, Dennis Boye Larsen, Kristian Kjær Petersen.

Writing – review & editing: Mathias Winther Bech, Sara Louise Kjær Clemmensen, Henriette Tranberg Nielsen, Dennis Boye Larsen, Kristian Kjær Petersen.

References

1. McCracken LM, Iverson GL. Disrupted sleep patterns and daily functioning in patients with chronic pain. *Pain Res Manag.* 2002; 7(2):75–9. <https://doi.org/10.1155/2002/579425> PMID: 12185371
2. Raymond I, Ancoli-Israel S, Choinière M. Sleep disturbances, pain and analgesia in adults hospitalized for burn injuries. *Sleep Med.* 2004; 5(6):551–9. <https://doi.org/10.1016/j.sleep.2004.07.007> PMID: 15511701
3. Tang NKY, Wright KJ, Salkovskis PM. Prevalence and correlates of clinical insomnia co-occurring with chronic back pain. *J Sleep Res.* 2007; 16(1):85–95. <https://doi.org/10.1111/j.1365-2869.2007.00571.x> PMID: 17309767
4. Smith MT, Haythornthwaite JA. How do sleep disturbance and chronic pain inter-relate? Insights from the longitudinal and cognitive-behavioral clinical trials literature. *Sleep Med Rev.* 2004 Apr; 8(2):119–32. PMID: 15033151
5. Affleck G, Urrows S, Tennen H, Higgins P, Abeles M. Sequential daily relations of sleep, pain intensity, and attention to pain among women with fibromyalgia. Vol. 68, *Pain.* 1996.
6. Lee YC, Lu B, Edwards RR, Wasan AD, Nassikas NJ, Clauw DJ, et al. The role of sleep problems in central pain processing in rheumatoid arthritis. *Arthritis Rheum.* 2013 Jan; 65(1):59–68. <https://doi.org/10.1002/art.37733> PMID: 23124650
7. Alsaadi SM, McAuley JH, Hush JM, Lo S, Bartlett DJ, Grunstein RR, et al. The Bidirectional Relationship Between Pain Intensity and Sleep Disturbance/Quality in Patients With Low Back Pain. *Clin J Pain.* 2014 Sep; 30(9):755–65. <https://doi.org/10.1097/AJP.000000000000055> PMID: 24451630

8. Moldofsky H. Sleep and pain. *Sleep Med Rev.* 2001; 5(5):385–96. <https://doi.org/10.1053/smr.2001.0179> PMID: 12531004
9. Demarco GJ, Baghdoyan HA, Lydic R. Differential cholinergic activation of G proteins in rat and mouse brainstem: Relevance for sleep and nociception. *J Comp Neurol.* 2003 Mar 3; 457(2):175–84. <https://doi.org/10.1002/cne.10548> PMID: 12541317
10. Foo H, Mason P. Brainstem modulation of pain during sleep and waking. *Sleep Med Rev.* 2003 Apr; 7(2):145–54. <https://doi.org/10.1053/smr.2002.0224> PMID: 12628215
11. Lu J, Jhou TC, Saper CB. Identification of Wake-Active Dopaminergic Neurons in the Ventral Periaqueductal Gray Matter. *J Neurosci.* 2006 Jan 4; 26(1):193–202. <https://doi.org/10.1523/JNEUROSCI.2244-05.2006> PMID: 16399687
12. Novati A, Roman V, Cetin T, Hagewoud R, den Boer JA, Luiten PGM, et al. Chronically restricted sleep leads to depression-like changes in neurotransmitter receptor sensitivity and neuroendocrine stress reactivity in rats. *Sleep.* 2008 Nov; 31(11):1579–85. <https://doi.org/10.1093/sleep/31.11.1579> PMID: 19014078
13. Volkow ND, Wang G-J, Telang F, Fowler JS, Logan J, Wong C, et al. Sleep deprivation decreases binding of [¹¹C]raclopride to dopamine D2/D3 receptors in the human brain. *J Neurosci.* 2008 Aug 20; 28(34):8454–61. <https://doi.org/10.1523/JNEUROSCI.1443-08.2008> PMID: 18716203
14. Andersen ML, Martins PJF, D'Almeida V, Bignotto M, Tufik S. Endocrinological and catecholaminergic alterations during sleep deprivation and recovery in male rats. *J Sleep Res.* 2005; 14(1):83–90. <https://doi.org/10.1111/j.1365-2869.2004.00428.x> PMID: 15743338
15. Kim Y, Elmenhorst D, Weisshaupt A, Wedekind F, Kroll T, McCarley RW, et al. Chronic sleep restriction induces long-lasting changes in adenosine and noradrenaline receptor density in the rat brain. *J Sleep Res.* 2015 Oct; 24(5):549–58. <https://doi.org/10.1111/jsr.12300> PMID: 25900125
16. Petersen KK, Olesen AE, Simonsen O, Arendt-Nielsen L. Mechanistic pain profiling as a tool to predict the efficacy of 3-week nonsteroidal anti-inflammatory drugs plus paracetamol in patients with painful knee osteoarthritis. *Pain.* 2019 Feb; 160(2):486–92. <https://doi.org/10.1097/j.pain.0000000000001427> PMID: 30371559
17. Yarnitsky D. Role of endogenous pain modulation in chronic pain mechanisms and treatment. *Pain.* 2015 Apr; 156:S24–31. <https://doi.org/10.1097/01.j.pain.0000460343.46847.58> PMID: 25789433
18. Price D, Hu J, Dubner R, Gracely R. Peripheral suppression of first pain and central summation of second pain evoked by noxious heat pulses. *Pain.* 1977; 3:57–68. [https://doi.org/10.1016/0304-3959\(77\)90035-5](https://doi.org/10.1016/0304-3959(77)90035-5) PMID: 876667
19. Corrêa JB, Costa LOP, de Oliveira NTB, Sluka KA, Liebano RE. Central sensitization and changes in conditioned pain modulation in people with chronic nonspecific low back pain: a case-control study. *Exp Brain Res.* 2015 Aug; 233(8):2391–9. <https://doi.org/10.1007/s00221-015-4309-6> PMID: 25963754
20. van Wijk G, Veldhuijzen DS. Perspective on Diffuse Noxious Inhibitory Controls as a Model of Endogenous Pain Modulation in Clinical Pain Syndromes. *J Pain.* 2010 May 1; 11(5):408–19. <https://doi.org/10.1016/j.jpain.2009.10.009> PMID: 20075013
21. Lewis GN, Rice DA, McNair PJ. Conditioned pain modulation in populations with chronic pain: a systematic review and meta-analysis. *J Pain.* 2012 Oct; 13(10):936–44. <https://doi.org/10.1016/j.jpain.2012.07.005> PMID: 22981090
22. Arendt-Nielsen L, Morlion B, Perrot S, Dahan A, Dickenson A, Kress HG, et al. Assessment and manifestation of central sensitisation across different chronic pain conditions. *Eur J Pain.* 2018 Feb; 22(2):216–41. <https://doi.org/10.1002/ejp.1140> PMID: 29105941
23. Finan PH, Goodin BR, Smith MT. The association of sleep and pain: an update and a path forward. *J Pain.* 2013 Dec; 14(12):1539–52. <https://doi.org/10.1016/j.jpain.2013.08.007> PMID: 24290442
24. Smith Michael T., Edwards Robert R., McCann JAH Una D. The Effects of Sleep Deprivation on Pain Inhibition and Spontaneous Pain in Women. 2007; 30(4):494–505.
25. Edwards RR, Grace E, Peterson S, Klick B, Haythornthwaite JA, Smith MT. Sleep continuity and architecture: Associations with pain-inhibitory processes in patients with temporomandibular joint disorder. *Eur J Pain.* 2009 Nov; 13(10):1043–7. <https://doi.org/10.1016/j.ejpain.2008.12.007> PMID: 19168380
26. Petersen KK, Arendt-Nielsen L, Simonsen O, Wilder-Smith O, Laursen MB. Presurgical assessment of temporal summation of pain predicts the development of chronic postoperative pain 12 months after total knee replacement. *Pain.* 2015; 156(1):55–61. <https://doi.org/10.1016/j.pain.0000000000000022> PMID: 25599301
27. Izumi M, Petersen KK, Laursen MB, Arendt-Nielsen L, Graven-Nielsen T. Facilitated temporal summation of pain correlates with clinical pain intensity after hip arthroplasty. *Pain.* 2017 Feb; 158(2):323–32. <https://doi.org/10.1097/j.pain.0000000000000764> PMID: 27870648
28. Kurien T, Arendt-Nielsen L, Petersen KK, Graven-Nielsen T, Scammell B. Preoperative Neuropathic Pain-like Symptoms and Central Pain Mechanisms in Knee Osteoarthritis Predicts Poor Outcome 6

- Months After Total Knee Replacement Surgery. *J Pain*. 2018; 19(11):1329–41. <https://doi.org/10.1016/j.jpain.2018.05.011> PMID: 29920331
29. Ren K. Wind-up and the NMDA receptor: from animal studies to humans. *Pain*. 1994; 59(2):157–8. [https://doi.org/10.1016/0304-3959\(94\)90067-1](https://doi.org/10.1016/0304-3959(94)90067-1) PMID: 7892012
 30. Graven-Nielsen T, Arendt-Nielsen L. Assessment of mechanisms in localized and widespread musculoskeletal pain. *Nat Rev Rheumatol*. 2010; 6(10):599–606. <https://doi.org/10.1038/nrrheum.2010.107> PMID: 20664523
 31. Price DD, Mao J, Frenk H, Mayer DJ. The N-methyl-D-aspartate receptor antagonist dextromethorphan selectively reduces temporal summation of second pain in man. *Pain*. 1994; 59(2):165–74. [https://doi.org/10.1016/0304-3959\(94\)90069-8](https://doi.org/10.1016/0304-3959(94)90069-8) PMID: 7892014
 32. Petrov ME, Goodin BR, Cruz-Almeida Y, King C, Glover TL, Bulls HW, et al. Disrupted sleep is associated with altered pain processing by sex and ethnicity in knee osteoarthritis. *J Pain*. 2015 May; 16(5):478–90. <https://doi.org/10.1016/j.jpain.2015.02.004> PMID: 25725172
 33. Smith MT, Edwards RR, Stonerock GL, McCann UD. Individual variation in rapid eye movement sleep is associated with pain perception in healthy women: preliminary data. *Sleep*. 2005 Jul; 28(7):809–12. <https://doi.org/10.1093/sleep/28.7.809> PMID: 16124658
 34. Schuh-Hofer S, Wodarski R, Pfau DB, Caspani O, Magerl W, Kennedy JD, et al. One night of total sleep deprivation promotes a state of generalized hyperalgesia: A surrogate pain model to study the relationship of insomnia and pain. *Pain*. 2013 Sep; 154(9):1613–21. <https://doi.org/10.1016/j.pain.2013.04.046> PMID: 23707287
 35. Onen SH, Alloui A, Gross A, Eschallier A, Dubray C. The effects of total sleep deprivation, selective sleep interruption and sleep recovery on pain tolerance thresholds in healthy subjects. *J Sleep Res*. 2001; 10(1):35–42. <https://doi.org/10.1046/j.1365-2869.2001.00240.x> PMID: 11285053
 36. Kundermann B, Sernal J, Huber MT, Krieg J-C, Lautenbacher S. Sleep Deprivation Affects Thermal Pain Thresholds but Not Somatosensory Thresholds in Healthy Volunteers. *Psychosom Med*. 2004 Nov; 66(6):932–7. <https://doi.org/10.1097/01.psy.0000145912.24553.c0> PMID: 15564360
 37. Faraut B, Léger D, Medkour T, Dubois A, Bayon V, Chennaoui M, et al. Napping reverses increased pain sensitivity due to sleep restriction. *PLoS One*. 2015; 10(2):e0117425. <https://doi.org/10.1371/journal.pone.0117425> PMID: 25723495
 38. Drewes AM, Rössel P, Arendt-Nielsen L, Nielsen KD, Hansen LM, Birket-Smith L, et al. Sleepiness does not Modulate Experimental Joint Pain in Healthy Volunteers. *Scand J Rheumatol*. 1997 Jan 12; 26(5):399–400. <https://doi.org/10.3109/03009749709065709> PMID: 9385357
 39. Pud D, Granovsky Y, Yarnitsky D. The methodology of experimentally induced diffuse noxious inhibitory control (DNIC)-like effect in humans. *Pain*. 2009; 144(1–2):16–9. <https://doi.org/10.1016/j.pain.2009.02.015> PMID: 19359095
 40. Beaudreau SA, Spira AP, Stewart A, Kezirian EJ, Lui L-Y, Ensrud K, et al. Validation of the Pittsburgh Sleep Quality Index and the Epworth Sleepiness Scale in older black and white women. *Sleep Med*. 2012 Jan; 13(1):36–42. <https://doi.org/10.1016/j.sleep.2011.04.005> PMID: 22033120
 41. Buysse DJ, Reynolds CF, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res*. 1989 May; 28(2):193–213. [https://doi.org/10.1016/0165-1781\(89\)90047-4](https://doi.org/10.1016/0165-1781(89)90047-4) PMID: 2748771
 42. Osman A, Barrios FX, Kopper BA, Hauptmann W, Jones J, O'neill E1. Factor Structure, Reliability, and Validity of the Pain Catastrophizing Scale. Vol. 20, *Journal of Behavioral Medicine*. 1997.
 43. Cano A, Leonard MT, Franz A. The significant other version of the Pain Catastrophizing Scale (PCS-S): preliminary validation. *Pain*. 2005 Dec 15; 119(1–3):26–37. <https://doi.org/10.1016/j.pain.2005.09.009> PMID: 16298062
 44. Sansone RA, Watts DA, Wiederman MW. The demographics of pain catastrophizing in a primary care sample. *Innov Clin Neurosci*. 2013 Sep; 10(9–10):12–4. PMID: 24307976
 45. Rolke R, Baron R, Maier C, Tolle TR, Treede RD, Beyer A, et al. Quantitative sensory testing in the German Research Network on Neuropathic Pain (DFNS): standardized protocol and reference values. [Erratum appears in *Pain*. 2006 Nov; 125(1–2):197]. *Pain*. 2006; 123(3):231–43. <https://doi.org/10.1016/j.pain.2006.01.041> PMID: 16697110
 46. Graven-Nielsen T, Izumi M, Petersen KK, Arendt-Nielsen L. User-independent assessment of conditioning pain modulation by cuff pressure algometry. *Eur J Pain*. 2017 Mar; 21(3):552–61. <https://doi.org/10.1002/ejp.958> PMID: 27859944
 47. Vaegter HB, Handberg G, Graven-Nielsen T. Isometric exercises reduce temporal summation of pressure pain in humans. *Eur J Pain (United Kingdom)*. 2015; 19(7):973–83.

48. Paul-Savoie E, Marchand S, Morin M, Bourgault P, Brissette N, Rattanavong V, et al. Is the deficit in pain inhibition in fibromyalgia influenced by sleep impairments? *Open Rheumatol J.* 2012; 6:296–302. <https://doi.org/10.2174/1874312901206010296> PMID: 23091577
49. Haack M, Scott-Sutherland J, Santangelo G, Simpson NS, Sethna N, Mullington JM. Pain sensitivity and modulation in primary insomnia. *Eur J Pain.* 2012 Apr; 16(4):522–33. <https://doi.org/10.1016/j.ejpain.2011.07.007> PMID: 22396081
50. Eichhorn N, Treede R-D, Schuh-Hofer S. The Role of Sex in Sleep Deprivation Related Changes of Nociception and Conditioned Pain Modulation. *Neuroscience.* 2018; 1(387):191–200.
51. Smith T. M, Edwards R. R, McCann D. U, Hythornthwaite AJ. The Effects of Sleep Deprivation on Pain Inhibition and Spontaneous Pain in Women. *Sleep.* 2007; 30(4):494–505. <https://doi.org/10.1093/sleep/30.4.494> PMID: 17520794
52. Matre D, Andersen MR, Knardahl S, Nilsen KB. Conditioned pain modulation is not decreased after partial sleep restriction. *Eur J Pain.* 2016 Mar 1; 20(3):408–16. <https://doi.org/10.1002/ejp.741> PMID: 26104968
53. Matre D. Conditioned pain modulation (CPM) is not one single phenomenon—Large intra-individual differences depend on test stimulus (TS) and several other independent factors. *Scand J Pain.* 2013; 4(1):8–9. <https://doi.org/10.1016/j.sjpain.2012.11.003> PMID: 29913888
54. Staud R, Cannon RC, Mauderli AP, Robinson ME, Price DD, Vierck CJ Jr. Temporal summation of pain from mechanical stimulation of muscle tissue in normal controls and subjects with fibromyalgia syndrome. *Pain.* 2003; 102(1–2):87–95. [https://doi.org/10.1016/s0304-3959\(02\)00344-5](https://doi.org/10.1016/s0304-3959(02)00344-5) PMID: 12620600
55. Anderson RJ, McCrae CS, Staud R, Berry RB, Robinson ME. Predictors of clinical pain in fibromyalgia: examining the role of sleep. *J Pain.* 2012 Apr; 13(4):350–8. <https://doi.org/10.1016/j.jpain.2011.12.009> PMID: 22381437
56. Edwards RR, Quartana PJ, Allen RP, Greenbaum S, Earley CJ, Smith MT. Alterations in pain responses in treated and untreated patients with restless legs syndrome: associations with sleep disruption. *Sleep Med.* 2011 Jun; 12(6):603–9. <https://doi.org/10.1016/j.sleep.2010.09.018> PMID: 21570347
57. Ziegler EA, Magerl W, Meyer RA, Treede R-D. Secondary hyperalgesia to punctate mechanical stimuli. *Brain.* 1999 Dec 1; 122(12):2245–57.
58. Carstens E, Klumpp D, Zimmermann M. Differential Inhibitory Effects of Medial and Lateral Midbrain Stimulation on Spinal Neuronal Discharges to Noxious Skin Heating in the Cat. Vol. 43, *JOURNAL OF NEUROPHYSIOLOGY.* 1980.
59. Schrimpf M, Liegl G, Boeckle M, Leitner A, Geisler P, Pieh C. The effect of sleep deprivation on pain perception in healthy subjects: a meta-analysis. *Sleep Med.* 2015 Nov 1; 16(11):1313–20. <https://doi.org/10.1016/j.sleep.2015.07.022> PMID: 26498229
60. Kundermann B, Hemmeter-Spernal J, Huber MT, Krieg J-C, Lautenbacher S. Effects of Total Sleep Deprivation in Major Depression: Overnight Improvement of Mood is Accompanied by Increased Pain Sensitivity and Augmented Pain Complaints. *Psychosom Med.* 2008 Jan; 70(1):92–101. <https://doi.org/10.1097/PSY.0b013e31815c1b5d> PMID: 18158380
61. Chiu YH, Silman AJ, Macfarlane GJ, Ray D, Gupta A, Dickens C, et al. Poor sleep and depression are independently associated with a reduced pain threshold. Results of a population based study. *Pain.* 2005; 115(3):316–21. <https://doi.org/10.1016/j.pain.2005.03.009> PMID: 15911158
62. Ağargün MY, Tekeoğlu I, Güneş A, Adak B, Kara H, Ercan M. Sleep quality and pain threshold in patients with fibromyalgia. *Compr Psychiatry.* 1999 May 1; 40(3):226–8. [https://doi.org/10.1016/s0010-440x\(99\)90008-1](https://doi.org/10.1016/s0010-440x(99)90008-1) PMID: 10360619
63. Ağargün MY, Tekeoğlu I, Güneş A, Adak B, Kara H, Ercan M. Sleep quality and pain threshold in patients with fibromyalgia. *Compr Psychiatry.* 40(3):226–8. [https://doi.org/10.1016/s0010-440x\(99\)90008-1](https://doi.org/10.1016/s0010-440x(99)90008-1) PMID: 10360619
64. Azevedo E, Manzano GM, Silva A, Martins R, Andersen ML, Tufik S. The effects of total and REM sleep deprivation on laser-evoked potential threshold and pain perception. *Pain.* 2011; 152:2052–8. <https://doi.org/10.1016/j.pain.2011.04.032> PMID: 21624774
65. Larson RA, Carter JR. Total Sleep Deprivation and Pain Perception during Cold Noxious Stimuli in Humans. *Scand J pain.* 2016; 13:12–6. <https://doi.org/10.1016/j.sjpain.2016.05.037> PMID: 27867438
66. Sivertsen B, Lallukka T, Petrie KJ, Steingrimsdottir OA, Stubhaug A, Nielsen CS. Sleep and pain sensitivity in adults. *Pain.* 2015; 156(8):1433–9. <https://doi.org/10.1097/j.pain.000000000000131> PMID: 25915149
67. Edwards RR, Almeida DM, Klick B, Haythornthwaite JA, Smith MT. Duration of sleep contributes to next-day pain report in the general population. *Pain.* 2008 Jul; 137(1):202–7. <https://doi.org/10.1016/j.pain.2008.01.025> PMID: 18434020

68. Fitzgerald JD, Carrillo C, Sadeghi N, Olmstead R, Nicassio PM, Irwin MR, et al. Sleep Loss Exacerbates Fatigue, Depression, and Pain in Rheumatoid Arthritis. *Sleep*. 2012; 35(4):537–43. <https://doi.org/10.5665/sleep.1742> PMID: 22467992
69. Arima T, Svensson P, Rasmussen C, Nielsen KD, Drewes AM, Arendt-Nielsen L. The relationship between selective sleep deprivation, nocturnal jaw-muscle activity and pain in healthy men. *J Oral Rehabil*. 2001 Feb 1; 28(2):140–8. <https://doi.org/10.1046/j.1365-2842.2001.00687.x> PMID: 11298262
70. Yarnitsky D, Sprecher E, Zaslansky R, Hemli JA. Multiple session experimental pain measurement. *Pain*. 1996 Oct; 67(2–3):327–33. [https://doi.org/10.1016/0304-3959\(96\)03110-7](https://doi.org/10.1016/0304-3959(96)03110-7) PMID: 8951926