

RESEARCH

Manipulation of Human Primary Endothelial Cell and Osteoblast Coculture Ratios to Augment Vasculogenesis and Mineralization

Amita R. Shah, MD, PhD,*† Joseph C. Wenke, PhD,‡ and Chandra Mauli Agrawal, PhD, PE§

Abstract: Tissue-engineering scaffolds are often seeded with a single type of cell, but there has been more focus on cocultures to improve angiogenesis and bone formation for craniofacial applications. Investigation of bone-derived osteoblasts (OBs) is important because of the use of bone grafts and migration of OBs from native bone into constructs in vivo and therefore, their contribution to bone formation in vivo. The limitation of primary OBs has been their inability to mineralize without osteogenic factors in vitro. Through coculture of OBs and endothelial cells (ECs) and manipulation of the coculture ratio, mineralization can be achieved without osteogenic media or additional growth factors, thus enhancing their utility for tissue-engineering applications. An optimal ratio of EC/OB for vasculogenesis and mineralization has not been determined for human primary cells. Human umbilical vein ECs were cultured with normal human primary OBs in different EC/OB ratios, namely, 10:1, 5:1, 1:1, 1:5, and 1:10 with EC and OB monocultures as controls. The number of vasculogenic networks in a collagen matrix was highest in ratios of 5:1 and 1:1. ECs lined up and formed capillary-like networks by day 10, which was not seen in the other groups. On polystyrene, cells were cocultured with ECs and OBs in direct contact (direct coculture) or separated by a transwell membrane (indirect coculture). At day 21, Alizarin Red staining showed mineralization on the 1:5 and 1:10 direct coculture ratios, with 1:5 having more mineralization nodules present than 1:10. No mineralization was seen in other direct coculture ratios or in any of the indirect coculture ratios. Alkaline phosphatase secretion was highest in the 1:5 direct coculture group. Vascular endothelial growth factor secretion from OBs was present in the 1:5 and 1:10 direct coculture ratios at all time points and inhibited after day 1 in other coculture groups. To improve vasculogenesis, cocultures of primary human ECs and OBs in ratios of 5:1 should be used, but to improve bone formation, the 1:5 direct coculture ratio results in most mineralization.

Key Words: coculture, endothelial cells, osteoblast, human primary cells, bone regeneration, vasculogenesis

(*Ann Plast Surg* 2014;00: 00 00)

There are many types of cells involved with the formation, repair, and remodeling of bone but bone tissue engineering scaffolds have usually been seeded with a single type of cell.¹ There has been more focus on cocultures, the culture of 2 or more types of cells, in in vitro and in vivo studies to more closely model the natural

Received April 16, 2014, and accepted for publication, after revision, June 18, 2014.

From the *Department of Surgery, University of Texas Health Science Center at San Antonio, San Antonio, TX; †Division of Plastic and Reconstructive Surgery, University of North Carolina at Chapel Hill, Chapel Hill, NC; ‡Extremity Trauma and Regenerative Medicine Task Area, United States Army Institute of Surgical Research; and §Department of Biomedical Engineering, University of Texas at San Antonio, San Antonio, TX.

Conflicts of interest and sources of funding: Supported by Department of Army Grant W81XWH-07-1-0717. The US Army Medical Research Acquisition Activity, 820 Chandler Street, Fort Detrick, MD 21702-5014, is the awarding and administering acquisition office.

Reprints: C. Mauli Agrawal, PhD, PE, Department of Biomedical Engineering, University of Texas at San Antonio, One UTSA Blvd, San Antonio, TX 78249. Email: Mauli.Agrawal@utsa.edu.

Copyright © 2014 by Lippincott Williams & Wilkins
ISSN: 0148-7043/14/0000-0000
DOI: 10.1097/SAP.0000000000000318

regeneration of bone and gain more insight into what cell cell interactions can improve on the limited success of bone tissue engineering thus far. The gold standard for regeneration of bone defects remains autologous bone grafts which are limited by donor site morbidity, availability, and decreased efficacy in the elderly.^{2,3} Harvests from bilateral iliac crests have enough tissue to cover 8 cm per person.⁴ Fully exploring the potential of bone derived primary osteoblasts (OBs) may lead to strategies to extend and expand the use of bone grafts for craniofacial applications.

Endothelial cells (ECs) and OBs have multiple interactions during bone formation. Early coculture studies showed that the ECs accelerate bone formation through angiogenesis and support that there is a synergistic relationship between the 2 cell types.⁵⁻⁹ Osteoblasts produce angiogenic factors such as vascular endothelial growth factor (VEGF) and matrix components which in turn stimulate ECs to produce osteogenic factors important in the differentiation of ECs into vessel components.^{8,10-12} Endothelial cells and formed tubules have also showed prolonged survival when in coculture.^{13,14}

Coculture studies of EC and OB have been often been done in 1:1 cell ratios in which the number of ECs is equal to the number of OBs^{8,15,16} but the full potential of the cocultures is not recognized without using the correct cell ratios. In a study which looked at angiogenesis in cocultures ratios of primary cells, the cultures with human dermal microvascular EC with human primary OB (EC/OB) in ratios of 1:1 or less had almost no ECs remaining at 1 week, whereas ratios of 5:1 and 10:1 EC/OB showed presence of both types of cells after 1 week.¹³

The effect of cocultured cells in different ratios has been systematically investigated by Ma et al using human umbilical ECs and human marrow stromal cells. Because the marrow cells in coculture require osteogenic media to induce mineralization, these data cannot be extrapolated to bone derived OBs which do not require the media with osteogenic supplements to mineralize while in coculture.¹⁷ The optimal coculture ratios that should be used to increase vascularization or osteogenesis from these types of OBs still have not been determined. Investigation of bone derived OBs is important because of the use of bone grafts and migration of OBs from native bone into constructs in vivo and therefore, their contribution to bone formation in vivo. Limitations of primary OBs have been their inability to mineralize without osteogenic factors in vitro. Achieving mineralization of these cells without the need for the addition of factors such as bone morphogenic protein would enhance their utility for tissue engineering applications.

It is hypothesized that by manipulating the cell ratio of EC to OB in coculture, vasculogenesis and mineralization can be augmented. The ratios investigated in this study (10:1, 5:1, 1:1, 1:5, 1:10) were chosen through review of current EC and OB literature. The purpose of this study was to determine optimal coculture ratios of human primary ECs and OBs to enhance mineralization and vasculogenesis. In addition, the effects of physical contact between ECs and OBs on mineralization and secretion of alkaline phosphatase and VEGF in coculture were examined.

MATERIALS AND METHODS

Cell Culture

Human umbilical vein ECs (Invitrogen, Carlsbad, CA) were cultured in EC media (Lifeline, Walkersville, MD) and used after

Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

| | | | | | |
|--|------------------------------------|-------------------------------------|---|--|---------------------------------|
| 1. REPORT DATE 01 AUG 2014 | | 2. REPORT TYPE N/A | | 3. DATES COVERED - | |
| 4. TITLE AND SUBTITLE Manipulation of Human Primary Endothelial Cell and Osteoblast Coculture Ratios to Augment Vasculogenesis and Mineralization. | | | | 5a. CONTRACT NUMBER | |
| | | | | 5b. GRANT NUMBER | |
| | | | | 5c. PROGRAM ELEMENT NUMBER | |
| 6. AUTHOR(S) Shah A. R., Wenke J. C., Agrawal C. M., | | | | 5d. PROJECT NUMBER | |
| | | | | 5e. TASK NUMBER | |
| | | | | 5f. WORK UNIT NUMBER | |
| 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) United States Army Institute of Surgical Research, JBSA Fort Sam Houston, TX | | | | 8. PERFORMING ORGANIZATION REPORT NUMBER | |
| 9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) | | | | 10. SPONSOR/MONITOR'S ACRONYM(S) | |
| | | | | 11. SPONSOR/MONITOR'S REPORT NUMBER(S) | |
| 12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited | | | | | |
| 13. SUPPLEMENTARY NOTES | | | | | |
| 14. ABSTRACT | | | | | |
| 15. SUBJECT TERMS | | | | | |
| 16. SECURITY CLASSIFICATION OF: | | | 17. LIMITATION OF ABSTRACT UU | 18. NUMBER OF PAGES 7 | 19a. NAME OF RESPONSIBLE PERSON |
| a. REPORT unclassified | b. ABSTRACT unclassified | c. THIS PAGE unclassified | | | |

passage 2. Normal human primary OBs (Lonza, Allendale, NJ) were cultured in OB media consisting of ascorbic acid and antibiotics (Lonza) and the cells were used after passage 2.

Cocultures

Coculture experiments were performed in different ratios, 10:1, 5:1, 1:1, 1:5, and 1:10, of ECs to OBs denoted EC/OB. Endothelial cell and OB monocultures were used as controls. All cocultures were cultured in media consisting of 2% fetal bovine serum, 15 ng/mL inulin growth factor 1, 5 ng/mL endothelial growth factor, 5 ng/mL basic fibroblast growth factor, 50 μ g/mL ascorbic acid, 1 μ g/mL hydrocortisone, 0.75 U/mL heparin, 10 mM glutamine, and antibiotics in MCDB 131. No VEGF was present in these media and both EC and OB monocultures were able to proliferate in these media.

Direct and Indirect Cocultures on Polystyrene

Polystyrene tissue culture treated 24 well plates (Corning, Edison, NJ) were used to examine mineralization in direct and indirect cocultures. In direct cocultures, the cells were in physical contact with each other. In indirect cocultures, the cells were separated by transwell membranes with 0.4 μ m pores (Corning) so growth factors from the cells could interact with the other cell type without physical cell contact. The direct coculture was performed by mixing the cells in the desired ratios before seeding them on the substrates. For the indirect studies, the OBs were seeded on the well plate surface and ECs seeded on the membrane. The culture medium was changed every other day and 24 hours before each supernatant collection. A 700 μ L of media was used for each medium change. For the indirect cultures, 600 μ L was placed into the well and 100 μ L into the membrane insert. At each time point, phase contrast microscopy (Nikon, Melville, NY) was used to image the OBs on each well at 10 \times (n = 4 constructs).

Alizarin Red S Staining

Staining using a 2% vol/vol Alizarin Red S solution at pH 4.1 was performed after 21 days of culture to detect calcium from mineralization. After staining, the monolayer was examined under phase contrast using an inverted microscope at 10 \times (Nikon). Areas with bright red staining were considered to have calcium present (n = 4 samples).

VEGF Enzyme-Linked Immunosorbent Analysis

At days 1, 3, 7, 12, 14, 18, and 21, the supernatant from each well of the polystyrene cultures was collected and frozen at -80°C . The culture medium was changed 24 hours before each supernatant collection. After all samples were collected, enzyme linked immunosorbent analysis was performed to determine protein concentration of VEGF using commercially prepared plates (R&D Systems, Wiesbaden, Germany) per manufacturer's protocol. Results are expressed as picogram per milliliter (n = 3 samples/time point).

Alkaline Phosphatase Assay

At days 1, 3, 7, 12, 14, 18, and 21, the supernatant from each well was collected and frozen at -80°C . The culture medium was changed 24 hours before each supernatant collection. The AnaSpec Sensolyte FDP Alkaline Phosphatase Assay Kit (Fremont, CA) was used to measure the concentration of alkaline phosphatase in the supernatant per manufacturer's protocol. Results are expressed as microgram per milliliter (n = 3 samples/time point).

Collagen Angiogenesis Assay

Collagen matrix was used to evaluate the ability of the cocultures to form vasculogenic networks in an environment that supports vasculogenesis, but does not contain provasculogenic factors.¹⁵ A 30 μ L of ECM matrix (Cayman Chemical, Denver, CO) was




FIGURE 1. Representative images of polystyrene cocultures in direct contact: (A) separation of ECs and OBs in 1:10 coculture on day 7, (B) nodule formations on day 10 of 1:5 coculture, (D) nodule formation at day 21 in 1:5 coculture, (C) OB monoculture on day 21, (E) EC monocultures at day 14 with apoptotic cells, and (F) 10:1 coculture on day 14 with ECs and OBs growing forming a mesh-like network.




FIGURE 2. Day 7 collagen assay. Circular endothelial formations are seen in the 1:5 coculture.

placed in each well of a 96 well plate and allowed to solidify at 37°C for 1 hour. Cocultures in the following coculture ratios (EC/OB): EC only, 10:1, 5:1, 1:1, 1:5, 1:10, and OB only were seeded in each well at a concentration of 7500 cells/150 μL. Media changes were performed daily with 150 μL. Cells were stained with calcein AM before imaging (n = 3–4 samples).

Fluorescence microscopy images were obtained at days 1, 3, and 7 at 10× using an inverted microscope (Nikon). Five to 10 random images were taken of each sample. Image analysis was performed by pattern recognition as described by Luesch et al.¹⁸ In this method, a number was assigned to the angiogenesis pattern present in each image. Individual cells that were well separated were assigned 0 point. Migrating cells that were aligning were assigned 1 point, capillary tubes without sprouting were assigned 2 points, sprouting of new capillary tubes were assigned 3 points, the formation of closed polygons were assigned 4 points, and if complex mesh like structures were present, 5 points were assigned.¹⁸

Matrigel Angiogenesis Assay

The ability of the different coculture ratios to form vasculogenic networks was also evaluated using a Matrigel, a provasculogenic substrate (Millipore, Billerica, MA). Two sets of experiments were done in Matrigel. The first experiment had the same total cell number

in each well for each group so the seeding density was the same. A 150 μL of the cell solution with a total cell number of 7500 cells per well were added to each well. The second experiment had 7500 ECs per well and OBs were added to achieve the desired coculture ratio. To perform the assay, 50 μL of EC matrix solution was added to pre-cooled 96 well tissue culture plates and incubated at 37°C for 1 hour. human umbilical vein endothelial cells and OBs were trypsinized and resuspended in coculture media in the following coculture ratios (EC/OB): EC only, 5:1, 1:1, 1:5, and OB only. The 1:10 and 10:1 ratios were not done because based on preliminary experiments, the number of cells required to achieve these ratios was not feasible under the restraints of the assay and well sizes. For both experiments, the cells were incubated for 6 hours at 37°C before imaging. Imaging was performed using phase contrast microscopy (Nikon) at 10×. Five random images were taken of each sample. Image analysis was performed by pattern recognition counts as was done with the collagen assay (n = 3 samples).

Statistics

Statistics were performed using 1 way analysis of variance with significance determined at $P < 0.05$. Results are reported as mean (SD).

RESULTS

Cocultures on Polystyrene

The direct coculture 1:5 and 1:10 ratios on polystyrene were the only groups to show evidence of cell organization and mineralization. Endothelial cells clustered together and a separation between the EC and OB could be seen starting at day 3 in these 2 groups (Fig. 1A). This organization of cells was also seen in the 1:5 and 1:10 groups on the collagen assay starting at day 7 (Fig. 2). In the 1:5 and 1:10 polystyrene direct coculture groups at day 10, there was evidence of nodule like formations with OBs situated radially around the circular formation (Fig. 1B) and in subsequent days there was continued nodule formation and mineralization patches present with no evidence of cell death (Fig. 1C). Osteoblast monocultures did not form nodule like formations or patches and the cells remained elongated and parallel to each other through day 21 (Fig. 1D).

Endothelial cell monocultures reached confluence and the cells started to become apoptotic at day 14 and did not survive to day 21 (Fig. 1E). The 10:1 and 5:1 direct coculture groups showed proliferation of both cell types throughout the 21 days but the ECs and OBs grew on top of each other, forming a mesh like network starting at day 14 with no nodule formation (Fig. 1F). The 1:1 group also had ECs and OBs growing on top of each other, but the cell layer partially




FIGURE 3. Indirect polystyrene cocultures: (A) 5:1 indirect coculture with flattened OBs with multiple filopodia on day 3 and (B) 5:1 indirect coculture on day 21.




FIGURE 4. Alizarin Red S staining on day 21: (A) mineralization nodule in 1:5 direct coculture, (B) mineralization patch in 1:5 direct coculture, and (C) no mineralization in OB monoculture.

lifted off the substrate on day 10. From days 10 to 21, mostly OBs remained in the 1:1 culture with a few EC clusters present.

In the indirect coculture group, the OBs were spaced far apart in groups 10:1, 5:1, and 1:1 and the cells took on a flattened appearance with multiple extended filopodia on days 1, 3, and 7 (Fig. 3A). There were no noticeable differences in OB morphology between groups 10:1, 5:1, 1:1, 1:5, 1:10, and OB on and after day 10. From days 10 to 21, the OBs in all groups remained elongated and parallel to each other (Fig. 3B).

Alizarin Red S Staining

Alizarin Red staining was performed at day 21 for the poly styrene direct and indirect cocultures. The staining showed mineralization nodules and patches on the 1:5 and 1:10 direct coculture ratios (Fig. 4A C). No mineralization was seen on the other direct coculture ratios or on any of the indirect coculture ratios.

Alkaline Phosphatase Assay

Although at day 1, there was no difference in alkaline phosphatase secretion between any of the direct coculture groups, by day 7, the 1:5 direct coculture group secreted significantly more alkaline phosphatase than the other groups and remained the highest through day 18. At day 12, 1:10 began to increase its secretion of alkaline phosphatase and remained at levels close to 1:5 for the rest of the culture period. On day 21, the amount of alkaline phosphate dropped back to or below day 1 levels in all groups (Fig. 5A).

In the indirect coculture groups, there were no significant differences in alkaline phosphatase secretion between the groups after day 1. On day 1, there was very little alkaline phosphatase secreted in the 1:5, 1:10, and OB group which does not correlate with the direct culture values and could be due to error; but on day 3, these groups had a levels of alkaline phosphatase to that secreted by the other groups (Fig. 5B).

VEGF Enzyme-Linked Immunosorbent Analysis

Osteoblast monocultures secreted VEGF over 21 days and there was no significant difference between the indirect and direct groups, showing that the presence of the transwell membrane in the indirect group did not alter OB VEGF secretion. The EC monocultures did not secrete any detectable levels of VEGF in direct or indirect cultures. In the direct contact cocultures, VEGF was detected only in the OB, 1:10, and 1:15 groups. After day 1, the amount of VEGF secreted decreased in the 1:5 and 1:10 direct coculture groups, but started increasing at day 7. By day 21, the 1:5 direct coculture ratio VEGF secretion exceeded that of the OB group. Vascular endothelial growth factor was also present in the 1:5 and 1:10 indirect coculture ratios on day 1 but was inhibited afterward and did not reappear in detectable levels until

day 18. For all coculture groups, there is significantly more VEGF secreted than in the indirect groups (Fig. 6).

Collagen Assay

The pattern recognition count defines the visual vasculogenic networks seen in each image of the well. The higher the number, the more vasculogenic progression that was present. At day 1, the vasculogenic response was dependent on EC density. The groups




FIGURE 5. Alkaline phosphatase assay. A, Direct coculture. Increase in secretion for 1:5 at day 7 and at day 12 for 1:10, correlating with formation of mineralization nodules. B, Indirect coculture showed constant secretion of alkaline phosphatase throughout the culture period for all groups (n = 3, P < 0.05).




FIGURE 6. VEGF assay. A, Direct coculture. No VEGF is secreted except in groups OB, 1:5, and 1:10. At day 21, 1:5 had significantly more VEGF than all groups except 1:10 and OB. B, Indirect coculture. OB VEGF secretion is significantly higher than all the coculture and EC groups ($n = 3, P < 0.05$).

with more ECs had more cells lining up and forming capillary like formations. At day 3, EC, 10:1, and 5:1 had significantly more vasculogenic activity than the rest of the groups. At day 7, 5:1 had significantly more vasculogenic formations than 1:10, 1:5, and EC. Endothelial cell and 10:1 groups peaked at day 3 and decreased on days 7 and 10 after they were confluent in the culture showing evidence of contact inhibition. The 1:1, 1:5, and 1:10 continue to show increased vasculogenic network formation up to day 7. Throughout the 7 days, the OB group did not have any vasculogenic formations

and it, along with 1:10 and 1:5, had the lowest values (Fig. 7). On day 7, the cells in the 1:5 and 1:10 groups clustered together into circular formations with a lining of ECs (Fig. 2).

Matrigel Angiogenesis Assay

The Matrigel angiogenesis assay was performed to determine if the presence of OBs was inhibitory to vasculogenic performance because ECs are able to form capillary networks in Matrigel without any additional vasculogenic factors. When the wells were seeded so that each well had the same number of ECs (Equal EC group), there were no significant differences between the groups (Fig. 8). When the wells were seeded so that there was an equal number of total cells seeded (Seeding density group), resulting in a different number of EC seeded in each well, the EC group had significantly more vasculogenic network formation than the 1:1 and 1:5 groups. The EC group also had more network formation than the 5:1 group, but it did not result in a statistically significant difference. There was also a significant difference between the 5:1 and 1:1 groups (Fig. 9). The results show that in Matrigel, the vasculogenic response is dependent on the number of ECs present in the culture and that OBs do not have an inhibitory effect on angiogenesis when in a provasculogenic environment.

DISCUSSION

Successful bone regeneration consists of both bone formation and vascularization, and cocultures of cells have been increasingly explored to achieve this.¹⁹ Various types of cells have been cocultured with ECs, such as mesenchymal stem cells,²⁰ stromal cells,¹⁷ fibro blasts,^{21,22} and human primary cells.^{8,12,13} An advantage of using human primary cells rather than stem cells is that differentiation of the cells through the use of specialized media with osteogenic factors is not required to achieve mineralization, which could translate into eliminating the need for exogenous growth factors such as bone morphogenetic protein 2. Many studies focus on the behavior of the cells in coculture that have an equal amount of ECs and OBs,^{11,16} but it is apparent that the ratio of the cocultured cells is an important factor that should be considered when designing implants.

Vasculogenesis

The vasculogenic response in collagen seems to be dependent on having enough ECs to form the networks, but not too many that the field becomes confluent and contact inhibition occurs, which is what is seen in group 10:1 and EC at days 7 and 10. The 5:1 coculture ratio, followed by 1:1, have a ratio of ECs to OBs which fosters ECs lining up and forming capillary like formation without




FIGURE 7. Collagen assay vasculogenic networks: 5:1 and 1:1 showed the highest vasculogenic networks for day 7; 5:1 had significantly more vasculogenesis than all groups except 1:1 [mean (SE), $n = 3-4, P < 0.05$].




FIGURE 8. Matrigel angiogenesis assay of the Equal EC group. The number of vasculogenic networks was significantly lower for the OB group. There were no differences between the other groups (n = 3, P < 0.05).

the culture overwhelming the OBs and becoming confluent and losing organization. The 1:5 and 1:10 had more vasculogenic formations in collagen than expected at days 7 and 10 because they had the fewest number of ECs in culture. Although the values were not as high as those for EC, 10:1, and 5:1, these cultures demonstrated EC organization which may have become more apparent and significant if the culture period was extended further. Unfortunately, the culture could not be extended past 10 days because the collagen delaminated off of the wells at day 12. The response in these culture groups may be a result of increased VEGF secretion from the OBs which was seen in the polystyrene experiments.

Endothelial cells do not secrete VEGF in nonhypoxic conditions so the VEGF response seen in this set of experiments is only from the OBs.²³ At day 1, measurable amounts of VEGF in both the indirect and direct culture groups were seen only in the OB monoculture and the 1:5 and 1:10 cocultures. The increase in VEGF in the direct cocultures seen at day 7 for 1:5 and 1:10, which coincides with the formation of nodules, can be explained by the EC and OB crosstalk that has been discussed in other studies which occurs with




FIGURE 9. Matrigel angiogenesis assay of the Equal Seeding Density group. EC had the highest vasculogenic pattern counts and the number of ECs present is proportional to the pattern counts seen for the different groups with the exception of 1:5 (n = 3, P < 0.05).

direct contact between the cells.^{10,15} As the ECs interact with the OBs, a release of paracrine factors occurs which has been shown to increase mineralization and, in turn, increase OB secretion and expression of angiogenic factors.

The Matrigel assay provided another way to look at angiogenesis in a gel like matrix. Endothelial cell monocultures are able to form vasculogenic networks due to the angiogenic factors in the Matrigel. The results show that the amount of capillary formation in Matrigel is roughly proportional to the number of ECs seeded. However, the 1:5 group had similar numbers of vasculogenic network formations as the other coculture groups with more ECs, which was not expected. Because the components in the Matrigel speed up the vasculogenesis process, they could also be increasing the effect of the OBs on the ECs in that ratio. In the collagen and polystyrene cultures, the ECs in the 1:5 ratio aggregate which also occurs in the Matrigel. The images of the 1:5 ratio show that there is aggregation of the cells at the junction points of the networks which may be serving as a nidus for the capillary tubules. Additionally, the secretion of VEGF was higher in the 1:5 coculture than in the other coculture ratios which may aid in an increased amount of vasculogenesis. When the OBs were cultured in the Matrigel by themselves, the cells mostly stayed separated with only a few cells migrating together. No filopodia were seen in these cultures so the extended capillary tubules seen in 1:5 were from the ECs and not OB filopodia.

Mineralization

Vascularization is an important part of bone regeneration and healing, not only because it provides a conduit for oxygen and nutrients but also because it helps activate bone morphogenic protein mediated healing processes. In studies in which angiogenesis was inhibited, there was significantly reduced bone healing in vivo.²⁴ Vascular endothelial growth factor is secreted in a differentiation dependent manner during osteogenesis and is an essential coordinator in endochondral bone formation. It also has a role in OB differentiation and mineralization.²⁵⁻²⁷ These factors may also have a part in the increased mineralization present in the 1:5 coculture, but in vivo studies need to be performed to fully explore this phenomenon.

It was hypothesized that a coculture of ECs and OBs where more OBs than ECs are present would result in increased mineralization compared to cultures with an equal amount or more ECs than OBs. In this study, mineralization was seen only in the 1:5 and 1:10 direct cocultures with 1:5 having more mineralization nodules and patches than the 1:10 culture. This is also reflected in the alkaline phosphatase assay where the highest amount of alkaline phosphatase was produced by the 1:5 direct ratio group followed by 1:10 direct ratio group. Because mineralization occurs only in the direct coculture and not in the indirect coculture with the same cell ratios, it suggests that there are more than paracrine effects at work. Multiple groups have demonstrated that in cocultures, the contact between the cells is an important part of the cellular response.^{6,7,28} When the cells do not have the opportunity to interact with each other through physical contact, mineralization is not promoted. When cells are exposed to only cultured media and are not able to form cell to cell contacts, an increase in expression of alkaline phosphatase is also not seen.^{6,7,28} Alkaline phosphatase secretion is an indicator of active calcium deposition and is a byproduct of OB activity. Its secretion has been shown to be dependent on endothelial contact with OBs with Connexin43 as the gap junction involved in the increase in alkaline phosphatase expression in the cocultures. This secretion is not only dependent on the presence of contact but also on how much contact and how long the different cell types have with each other.²⁹⁻³¹ In the groups with more OBs than ECs, the most alkaline phosphatase secretion was seen from days 7 to 14, after which there was a decrease in secretion. The EC monoculture group did not have a significant rise and fall of the alkaline phosphatase levels throughout the culture period for both the indirect and direct

cultures. This reflects a constant secretion of alkaline phosphatase from ECs, whereas the OB secretion of alkaline phosphatase is related to the mineralization activity and OB differentiation.

The choice of culture medium is important because mineralization occurs in OB monocultures in osteogenic media containing β glycerophosphate and dexamethasone. In this study, the media used do not contain dexamethasone or β glycerophosphate and, therefore, did not promote mineralization in OB monocultures after 21 days of culture. Because of this, mineralization seen in the cultures can be attributed to the effects of the coculture and not the culture media. The use of primary differentiated cells in this study is also important. Ma et al¹⁷ have also investigated the effect of coculture ratios on osteogenic and vasculogenic outcomes with human marrow stromal cells; however, osteogenic media were required to differentiate the cells and thereby induce mineralization. In culture medium similar to what was used in this study, no mineralization was seen.

CONCLUSIONS

To improve vasculogenesis in human primary EC and bone derived OB cocultures, ratios of 5:1 or 1:1 should be used; but to increase mineralization, the 1:5 direct coculture ratio, followed by the 1:10 direct coculture ratio, should be used. This information can be used to develop future tissue engineering scaffold systems which use preseeded primary cells or even composite scaffolds for bone segmental defects which have an environment which would support native OB migration into the scaffold. Additionally, this information can be used in systems to augment and expand the use of bone grafts.

For complex tissues, such as bone, a vascular network needs to be established to support cell infiltration and growth into central areas of the scaffold. On the basis of this study, areas needing more vascularization could be implanted with a 5:1 coculture and the areas where bone formation is desired can be implanted with a 1:5 coculture. In these cultures, there is enough of 1 cell type to have enough of an effect on the other cell type, but not sufficient to overwhelm and overtake the culture. The study confirms that contact between the ECs and OBs is important and that the simple addition of growth factors into the milieu may not be enough to induce vasculogenesis and mineralization.

ACKNOWLEDGMENTS

The authors thank Sarita R. Shah for the technical assistance and John Jones for the assistance with statistical analysis.

REFERENCES

- Lieberman JFG, ed. *Bone Regeneration and Repair: Biology and Clinical Applications*. Totowa, NJ: Humana Press; 2005.
- Lee K-B, Taghavi CE, Hsu MS, et al. The efficacy of rhBMP-2 versus autograft for posterolateral lumbar spine fusion in elderly patients. *Eur Spine J*. 2010;19:924-930.
- Mears D. Surgical treatment of acetabular fractures in elderly patients with osteoporotic bone. *J Am Acad Orthop Surg*. 1999;7:128-141.
- Green S. A comparison of bone grafting and bone transport for segmental skeletal defects. *Clin Orthop Relat Res*. 1994;301:111-117.
- Wang D, Miura M, Demura H, et al. Anabolic effects of 1,25-dihydroxyvitamin D3 on osteoblasts are enhanced by vascular endothelial growth factor produced by osteoblasts and by growth factors produced by endothelial cells. *Endocrinology*. 1997;138:2953-2962.
- Stahl A, Wenger A, Weber H, et al. Bi-directional cell contact-dependent regulation of gene expression between endothelial cells and osteoblasts in a three-dimensional spheroidal coculture model. *Biochem Biophys Res Commun*. 2004;322:684-692.
- Villars F, Guillotin B, Amedee T, et al. Effect of HUVEC on human osteoprogenitor cell differentiation needs heterotypic gap junction communication. *Am J Physiol Cell Physiol*. 2002;282:775-785.
- Wenger A, Stahl A, Weber H, et al. Modulation of in vitro angiogenesis in a three-dimensional spheroidal coculture model for bone tissue engineering. *Tissue Eng*. 2004;10:1536-1547.
- Finkenzeller G, Torio-Padron N, Momeni A, et al. In vitro angiogenesis properties of endothelial progenitor cells: a promising tool for vascularization of ex vivo engineered tissues. *Tissue Eng*. 2007;13:1413-1420.
- Clarkin CE, Emery RJ, Pitsillides AA, et al. Evaluation of VEGF-mediated signaling in primary human cells reveals a paracrine action for VEGF in osteoblast-mediated crosstalk to endothelial cells. *J Cell Physiol*. 2008;214:537-544.
- Fuchs S, Hofmann A, Kirkpatrick CJ. Microvessel-like structures from outgrowth endothelial cells from human peripheral blood in 2-dimensional and 3-dimensional co-cultures with osteoblastic lineage cells. *Tissue Eng*. 2007;13:2577-2588.
- Wenger A, Stahl A, Kneser U, et al. Introduction of a three-dimensional coculture module for in vitro angiogenesis in bone tissue engineering. *Eur Cell Mater*. 2003; 5.
- Unger RE, Sartoris A, Peters K, et al. Tissue-like self-assembly in cocultures of endothelial cells and osteoblasts and the formation of microcapillary-like structures on three-dimensional porous biomaterials. *Biomaterials*. 2007;28:3965-3976.
- Koike N, Fukumura D, Gralla O, et al. Tissue engineering: creation of long-lasting blood vessels. *Nature*. 2004;428:138-139.
- Santos MI, Unger RE, Sousa RA, et al. Crosstalk between osteoblasts and endothelial cells co-cultured on a polycaprolactone-starch scaffold and the in vitro development of vascularization. *Biomaterials*. 2009;30:4407-4415.
- Hofmann A, Ritz U, Verrier S, et al. The effect of human osteoblasts on proliferation and neo-vessel formation of human umbilical vein endothelial cells in a long-term 3D co-culture on polyurethane scaffold. *Biomaterials*. 2008;29:4217-4226.
- Ma J, Beucken JJ, Yang F, et al. Coculture of osteoblasts and endothelial cells: optimization of culture medium and cell ratio. *Tissue Eng*. 2011;17:349-357.
- Luesch H, Chanda S, Raya R, et al. A functional genomics approach to the mode of action of apratoxin A. *Nat Chem Biol*. 2006;2:158-167.
- Grellier M, Bordenave L, Amedee J. Cell-to-cell communication between osteogenic and endothelial lineages: implications for tissue engineering. *Trends Biotechnol*. 2009;27:562-571.
- Zhou J, Lin H, Fang T, et al. The repair of large segmental bone defects in the rabbit with vascularized tissue engineered bone. *Biomaterials*. 2010;31:1171-1179.
- Choong CS, Huttmacher DW, Triffitt JT. Co-culture of bone marrow fibroblasts and endothelial cells on modified polycaprolactone substrates for enhanced potentials in bone tissue engineering. *Tissue Eng*. 2006;12:2521-2531.
- Sieminski A, Padera R, Blunk T, et al. Systemic delivery of human growth hormone using genetically modified tissue-engineered microvascular networks: prolonged delivery and endothelial survival with inclusion of non-endothelial cells. *Tissue Eng*. 2002;8:1057-1069.
- Liu Y, Cox S, Morita T, et al. Hypoxia regulates vascular endothelial growth factor gene expression in endothelial cells. *Circ Res*. 1995;77:638-643.
- Hausman MR, Schaffler MB, Majeska RJ. Prevention of fracture healing in rats by an inhibitor of angiogenesis. *Bone*. 2001;29:560-564.
- Gerber H, Vu T, Ryan A, et al. VEGF couples hypertrophic cartilage remodeling, ossification and angiogenesis during endochondral bone formation. *Nat Med*. 1999;5:623-628.
- Keramaris NC, Calori GM, Nikolau VS, et al. Fracture vascularity and bone healing: a systematic review of the role of VEGF. *Injury*. 2008;39:S45-S57.
- Mayer H, Bertram H, Lindenmaier W, et al. Vascular endothelial growth factor (VEGF-A) expression in human mesenchymal stem cells: autocrine and paracrine role on osteoblastic and endothelial differentiation. *J Cell Biochem*. 2005;95:827-839.
- Guillot B, Bourcet R, Remy-Zolghadri M, et al. Human primary endothelial cells stimulate human osteoprogenitor cell differentiation. *Cell Physiol Biochem*. 2004;14:325-332.
- Association AM. Pathophysiology of osteoporosis. *Osteoporosis Management [serial online]*. 2004.
- Hager S, Lampert F, Orimo H, et al. Up-regulation of alkaline phosphatase expression in human primary osteoblasts by cocultivation with primary endothelial cells is mediated by p38 mitogen-activated protein kinase-dependent mRNA stabilization. *Tissue Eng Part A*. 2009;15:3437-3447.
- Xue Y, Xing Z, Hellem S, et al. Endothelial cells influence the osteogenic potential of bone marrow stromal cells. *Biomed Eng Online*. 2009;8:34.