



PHS PUBLIC ACCESS

Author manuscript

Cancer Epidemiol Biomarkers Prev. Author manuscript; available in PMC 2017 November 01.

Published in final edited form as:

Cancer Epidemiol Biomarkers Prev. 2017 May ; 26(5): 787–794. doi:10.1158/1055-9965.EPI-16-0792.

Alcohol Intake and Breast Cancer Risk in African American Women from the AMBER Consortium

Lindsay A Williams¹, Andrew F Olshan¹, Chi-Chen Hong², Elisa V Bandera³, Lynn Rosenberg⁴, Ting-Yuan David Cheng², Kathryn L. Lunetta⁵, Susan E McCann², Charles Poole¹, Laurence N Kolonel⁶, Julie R Palmer⁴, Christine B Ambrosone², and Melissa A Troester^{1,*}

¹Department of Epidemiology, University of North Carolina at Chapel Hill, 135 Dauer Drive, Chapel Hill, NC 27599, USA

²Department of Cancer Prevention and Control, Roswell Park Cancer Institute, Elm & Carlton Streets, Buffalo, NY 14263, USA

³Cancer Prevention and Control Program, Rutgers Cancer Institute of New Jersey, Rutgers, The State University of New Jersey, 195 Little Albany Street, New Brunswick, NJ 08903, USA

⁴Slone Epidemiology Center, Boston University, 1010 Commonwealth Avenue, Boston, MA, 02215-1201, USA

⁵Department of Biostatistics, Boston University School of Public Health, Crosstown Center, 801 Massachusetts Avenue, 3rd Floor, Boston, MA 02118, USA

⁶Cancer Epidemiology Program, University of Hawai'i Cancer Center, 701 Ilalo Street, Honolulu, HI 96813, USA

Abstract

Background—Alcohol is a recognized risk factor for invasive breast cancer, but few studies involve African American women.

Methods—The present analysis included 22,338 women (5,108 cases of invasive breast cancer) from the African American Breast Cancer Epidemiology and Risk (AMBER) Consortium. The association between number of alcoholic drinks per week (dpw) and breast cancer was estimated using logistic regression, adjusting for potential confounders, and stratifying by breast cancer subtype.

Results—Approximately 35% of controls were current drinkers at interview. Women who reported current drinking of 14 dpw had an elevated risk of breast cancer compared to light drinkers (>0–<4 dpw) [adjusted odds ratio (adjOR), 95% confidence interval (CI): 1.33 (1.07–1.64)]. We observed elevated risk among women drinking 7 dpw for ER– [adjOR, 95% CI: 1.31 (1.00–1.72)], PR– [adjOR, 95% CI: 1.28 (1.00–1.63)], HER2– [adjOR, 95% CI: 1.36 (1.09–1.70)], and triple negative [adjOR, 95% CI: 1.39 (0.98–2.00)] molecular subtype. Among receptor-

*Corresponding Author: Melissa A Troester, (919) 966-7408, troester@email.unc.edu, Department of Epidemiology, University of North Carolina at Chapel Hill, 135 Dauer Drive, Chapel Hill, NC 27599, USA.

Conflicts of Interest: The authors declare no potential conflicts of interest.

positive cases, ORs remained elevated but attenuated relative to receptor-negative cases. Sensitivity analysis of age-defined windows of exposure (<30 years, 30–49, 50+ years of age) did not reveal variation in patterns of association. Risk associated with alcohol intake did not vary significantly by oral contraceptive use, smoking status, or menopausal status.

Conclusions—Among African American women, similar to women of European descent, drinking 7 alcoholic dpw was associated with an increased risk of breast cancer regardless of subtype.

Impact—Alcohol intake is a modifiable risk factor for breast cancer and reduced intake among African American women should be encouraged.

INTRODUCTION

Although alcohol is an established risk factor for breast cancer, most studies have been conducted in predominantly white populations (1–9). African Americans report less alcohol intake than whites for a variety of reasons including religious beliefs and prevalence of comorbid conditions, such as type 2 diabetes or hypertension (10–18). Furthermore, African American women have different patterns of exposure to other breast cancer risk factors, including parity, oral contraceptive use, age at menarche, breastfeeding, and age at first birth (11,13,19–21). Many risk factors, including alcohol exposure, contribute to breast cancer development by altering duration of exposure or activity of hormones (22,23). Thus, it is important to understand the relationship between alcohol and breast cancer risk in context of other exposures.

Many breast cancer risk factors have distinct effects on risk of ER-positive versus ER-negative breast cancers (21,24,25), and several studies have shown distinct risk factor profiles for triple-negative breast cancers (26–31). Despite suggestions that alcohol exposure may modulate estrogen metabolism pathways (32), there has been limited evidence that alcohol exposure produces distinct effects on risk of ER-positive vs. ER-negative breast cancer. Moreover, very few studies have evaluated alcohol-associated risk of triple-negative breast cancers (1,14). Alcohol is hypothesized to be a ‘complete carcinogen’ acting as both an initiator and promoter of the disease through inhibition of DNA synthesis and repair (33–36). The current study used data from the African American Breast Cancer Epidemiology and Risk (AMBER) consortium to examine alcohol drinking among African American women as a risk factor for invasive breast cancer overall, by hormone receptor status, and triple negative subtype. Both recent and age-defined periods of alcohol exposure were evaluated in association with risk.

MATERIALS AND METHODS

Study population

This analysis included 22,338 African American women from the African American Breast Cancer Epidemiology and Risk (AMBER) consortium of four large epidemiologic studies of breast cancer. The parent studies include the Carolina Breast Cancer Study (CBCS) (37), the Black Women’s Health Study (BWHS) (38), the Multiethnic Cohort Study (MEC) (39) and the Women’s Circle of Health Study (WCHS) (40). The study details for the AMBER

consortium have been previously described (41). Briefly, CBCS Phases 1 and 2 is a population-based case-control study conducted in North Carolina from 1993–2001. Breast cancer cases were identified through rapid case ascertainment in cooperation with the NC Central Cancer Registry. Controls were selected from North Carolina Department of Motor Vehicles (women aged 20 to 64) and Health Care Administration Financing lists (women aged 65 to 74). BWHS is a prospective cohort study of 59,000 African American women from around the United States enrolled by mailed questionnaire starting in 1995 with follow-up questionnaires administered every other year. Cases were identified by self-report and confirmed by medical record review or linkage with state cancer registries. MEC is a prospective cohort study based in Hawaii and Los Angeles, California, consisting of women from five different racial-ethnic groups with over 16,000 African American women enrolled from 1993–1996. Cases were identified via linkage to the Los Angeles County Cancer Surveillance Program, the State of California Cancer Registry, and the Hawaii State Cancer Registry. WCHS is a case-control study started in 2002 in New York City hospitals and expanded into ten counties in New Jersey, with cases identified by rapid case ascertainment by the New Jersey State Cancer Registry. Controls were identified by Random Digit Dialing in both sites, complemented in NJ with community-based recruitment (42). MEC and BWHS are prospective cohort studies sampled as nested case-control studies with cases and controls frequency matched by 5-year age categories, geographic location, and most recent questionnaire completed (43). Research protocols for each study were approved by the Institutional Review Board at the respective institutions. All subjects provided informed consent for study enrollment.

Eligible cases were women with a first diagnosis of invasive breast cancer (N=5,108). Tumor subtype for Estrogen Receptor (ER), Progesterone Receptor (PR) and Human Epidermal Growth Factor Receptor 2 (HER2) classification was based on pathology data from hospital records or cancer registry records. Cases were classified as ER+ or PR+ if marker expression was recorded as positive or borderline in the clinical record. ER and PR status was missing for approximately 25% of cases in the AMBER consortium (24). HER2 negative status was defined as immunohistochemistry (IHC) reported as 0 or 1+ staining intensity or a combination of negative by *fluorescence in situ hybridization* and 2+ by IHC. HER2 testing was not routine until the mid-2000's and is missing for approximately 50% of AMBER cases (24). Triple negative cases were classified as negative for ER, PR and HER2.

Exposure Assessment

Each study ascertained alcohol intake via study questionnaire. For CBCS, type of alcohol (beer, wine, liquor) and amount (drinks per day, week, month) was queried for the following age ranges for each participant: <25, 25 to 49, and 50. WCHS collected alcoholic drinks/day for each decade of life. For BWHS, type of alcohol (beer, wine, liquor) and amount (drinks per day, week, month) was asked at baseline and amount was reported on the follow-up survey administered every other year. For MEC, alcohol intake data was gathered at baseline in 1993, and on the follow-up questionnaire in 2003. Participants reported the number of days per week they drank alcohol and how many glasses/cans/bottles/drinks of beer/wine/liquor they consumed. Level of recent alcohol intake was determined by the self-reported drinking in the age category that included diagnosis age (for cases) or enrollment

age (for non-cases) for all studies. Forty-four cases (0.8%) and 256 controls (1.5%) were excluded from all analyses due to missing information on alcohol drinking.

Statistical Analyses

Participants were coded as never, past, or current drinkers based on recent use. To be classified as a never drinker, participants had to report never drinker for each of the surveys preceding their index date. To be a past drinker, participants had to report drinking during a time period before the period most proximal to the index date and report no alcohol intake in the period most proximal to diagnosis. Exposure categories (drinks/week) for analyses stratified by subtype and modifiers were: never drinkers, past drinkers, >0 to <4 (referent), 4 to <7, and ≥7. The highest exposure category for stratified analyses was chosen because seven or more drinks exceeds the Dietary Guidelines for Americans as determined by Health and Human Services recommending no more than 1 drink a day for women. Further, a relatively small number of women in the study exceeded 7 drinks/week, so stratified analyses resulted in too few women in the highest category to produce reliable estimates. For the main analysis (overall invasive breast cancer risk), the highest category was divided into two: ≥7–<14 and ≥14 drinks/week. The referent category, light drinking (>0 to <4 drinks/week), was selected using flexible modeling techniques with drinks/week modeled as a squared-term and graphed against the log odds of breast cancer with the corresponding 95% confidence intervals (Supplemental Figure 1).

To examine critical exposure windows for alcohol intake, we created 3 age categories to describe intake during the following time periods: young adult (age <25 for CBCS, <30 for WCHS), middle adult (age 25–49 for CBCS, and 30–49 WCHS), and older adult (age 50 and older). Since WCHS asked about intake for each decade, the number of drinks/week for <20 and 20–29 was averaged to obtain the number of drinks/week for <30 years of age. This same rule was applied for 30–49 years of age (averaging the thirties and forties), and 50 years of age or greater (averaging the fifties and subsequent decades prior to diagnosis). The following alcohol intake categories (drinks/week) were used: 0, >0 to <4 (referent), 4 to <7, and ≥7.

Logistic regression was used to calculate the odds ratio (OR) and 95% confidence interval (95% CI) for alcohol drinking and breast cancer risk. Two-sided p-values and a significance level of 0.05 were used for all tests of statistical significance. To evaluate modification by study, duration of oral contraceptive use, smoking status, and menopausal status, likelihood ratio tests were conducted comparing the model with the interaction term to a reduced model. Covariates selected as confounders in the multivariable (MV) logistic regression models were selected based on subject matter knowledge and were used in the construction of a directed acyclic graph (44). Covariates were coded as follows: study time period (1993–1998, 1999–2005, 2006–2013), United States geographic location of the participant (Northeast, South, Midwest, and West), parent study, age at diagnosis for cases or age at index date for controls (less than 40, 40–49, 50–59, 60–69, 70 or older), level of education (<12 years, 12 years, 13–15 years, 16 years, >16 years), age at menarche (<11 years, 11–12, 13–14, 15–16, 17 or older), parity (nulliparous, 1, 2, 3, or 4 or more live births), postmenopausal hormone therapy (HT) use defined as duration of combined estrogen and

progesterone use (never used, ever used), recent body mass index (BMI). Smoking status (never, former, current smoker), duration of oral contraceptive use (never, 1–9 years, 10 or more years) and menopausal status (pre-, post-) were considered as possible effect modifiers, but ultimately added as adjustment variables in unstratified MV models. A complete case analysis was used for the basic and MV logistic models (resulting in exclusion of approximately 15% percent of cases and controls in the MV model, largely due to missing menopausal status). Analyses among excluded participants produced similar estimates to those from MV models presented. Tests for trend for drinks/week were conducted using >0 to <4 (referent), 4 to <7, 7–<14 and 14 where noted, and excluded never/past drinkers. Alcohol categories were treated as an ordinal variable with the beta coefficient p-value serving as the measure of significance for the test of linear trend. All analyses were done in SAS version 9.3 (SAS Institute, Cary, NC).

RESULTS

The present analysis includes 5,108 cases of invasive breast cancer and 17,230 controls. Forty-five percent of participants were never drinkers and 20.8% were past drinkers. Recent drinking patterns differed by study (Table 1). Approximately 11% of cases and controls reported drinking 4 drinks/week. CBCS, BWHS and MEC had higher proportions of current drinkers (CBCS: 40.6% of cases, 42.1% of controls; BWHS: 35.4% of cases, 37.8% of controls; MEC: 35.1% of cases, 34.5% of controls) than WCHS (16.9% of cases, 17.5% of controls). Additional participant characteristics overall and by study can be found in Supplemental Table 1.

Case-control odds ratios (OR) and 95% confidence intervals (95% CI) for the overall association between alcohol intake and breast cancer are presented in Table 2. Odds ratios for a minimally adjusted basic model and a multivariable (MV) model (adjusted ORs) are reported in the tables. We observed a J-shaped curve between recent alcohol intake categories (excluding past drinking) and risk of invasive breast cancer. Compared to light drinkers, overall, never drinking was associated with elevated risk of breast cancer [adjusted OR 1.12, 95% CI (1.02–1.24), Table 2]. Drinking 14 drinks/week was associated with significantly increased risk of invasive breast cancer [adjusted OR 1.33, 95% CI (1.07–1.64), Table 2]. When analyses were stratified by hormone receptor (HR) status, elevated risk was observed in the highest intake category for all subtypes. Table 2 shows slightly stronger risk associated with 7 drinks/week among ER negative [adjusted OR: 1.31, 95% CI (1.00–1.72)], PR negative [adjusted OR: 1.28, 95% CI (1.00–1.63)], HER2 negative [adjusted OR: 1.36, 95% CI (1.09–1.70)], and triple negative cases [adjusted OR: 1.39, 95% CI (0.98–2.00)]. Associations were in the same direction for ER+, PR+, and HER2+ breast cancers, but were slightly attenuated and non-significant.

Among the four studies, CBCS had the strongest J-shaped pattern of risk: compared to drinking >0–<4 drinks/week, the adjusted ORs were [1.23, 95% CI (0.92–1.65)], [1.50, 95% CI (1.11–2.03)], and [2.03, 95% CI (1.29–3.18)], respectively, for never drinking, past drinking, and drinking 7 drinks/week (Table 3). There were no associations between risk of breast cancer and any alcohol intake in WCHS. In BWHS and MEC, associations between alcohol intake and risk were similar in direction to those in CBCS, but lower in magnitude

(Table 3). BWHS never drinkers had a significantly elevated risk of breast cancer [adjusted OR 1.17, 95% CI (1.02–1.34)]. The multivariable p-value for heterogeneity was 0.10.

We evaluated effect modification by smoking status, duration of oral contraceptive use, and menopausal status (Supplemental Table 2, Supplemental Table 3, and Supplemental Table 4, respectively). We did not observe evidence for statistical interaction between alcohol intake and smoking status in the multivariable model (MV $p=0.58$, Supplemental Table 2). Risk was elevated in the highest alcohol intake category across all strata of smoking status. Similarly, we did not observe evidence of statistical interaction for alcohol intake and duration of oral contraceptive use (MV $p=0.17$, Supplemental Table 3) or menopausal status (MV $p=0.38$, Supplemental Table 4).

We evaluated drinking during specific age-defined periods. Associations between recent drinking and risk were attenuated when alcohol drinking was assessed by age-at-exposure, but only two of the four AMBER studies (CBCS and WCHS) had data available for drinking at different time periods so sample size is limited. ORs for early, middle, and later life were close to the null and did not show the dose-response trend observed for recent drinking (Table 4).

DISCUSSION

In this study among African American women in the AMBER consortium, we found evidence of a J-shaped curve for alcohol drinking and invasive breast cancer risk. Women who reported drinking 14 or more drinks/week as compared to those drinking $>0 - <4$ drinks/week experienced the highest risk of invasive breast cancer. Never drinkers also experienced a significantly elevated risk of breast cancer. The J-shaped curve was not substantially altered when stratifying by hormone receptor status or triple negative subtype. In subtype-stratified analyses, alcohol intake of 7 or more drinks/week was associated with increased risk of breast cancer across all subtypes studied. There was no evidence for statistical interaction by smoking, oral contraceptive use, or menopausal status. Interestingly, past drinkers were most often found to have a lower risk of breast cancer than women reporting recent use, suggesting that decreasing alcohol consumption may reduce risk. We conclude that alcohol is a risk factor for invasive breast cancer among African American women, as has been consistently shown among studies primarily involving white women.

Previous studies of alcohol and cancer have found J-shaped curves similar to those observed in this study (6,45), but few breast cancer studies have assessed or reported this dose-response pattern (3–8,11,46–48). We hypothesize that J-shaped curve morphology may be attributable in part to women with unmeasured comorbidities that preclude alcohol drinking, yet contribute to increased risk of breast cancer among never drinkers, such as type 2 diabetes (6,49). A behavioral study found that as self-reported health status moves from poor to excellent among African American women, there is a 10% increase in the odds of alcohol drinking (16). In our study, women who were never drinkers were more frequently of obese BMI, had lower education (with the exception of BWHS), and were more likely report never having a mammogram. Future studies that explicitly capture comorbidities and reasons for

abstaining from alcohol would help to elucidate the reasons for our observed elevated risk in never drinkers.

Our findings of elevated risk among women drinking 14 or more drinks/week are consistent with two large meta-analyses and other epidemiologic studies conducted among primarily white women (3–8,11,46,47). Our findings are also consistent with the literature showing an increased risk of both HR+ and HR- subtypes with increasing alcohol intake (3,8,47,50–54). Considering other co-exposures, similar to our findings, smoking did not modify alcohol-associated risk in a meta-analysis of 53 epidemiologic studies and a separate pooled analysis where elevated risk was seen in the highest intake category of alcohol intake regardless of smoking status (5,53). The patterns of risk for duration of oral contraceptive use mirrored those of our overall risk estimates, with a J-shaped curve and no evidence of statistical interaction, consistent with previous findings (2,14).

The most important difference between our findings and previous literature was with regard to menopausal status and age-related patterns. We observed no differences in associations by menopausal status; previous studies among white women have suggested that postmenopausal women who consumed alcohol were at an elevated risk of breast cancer compared to premenopausal women (3,8,48,51,53). However, our menopause findings are echoed in our analysis of risk by age-period. Previous studies have suggested that drinking prior to 40–50 years of age has a greater impact on breast cancer risk relative to drinking at older ages (8,51). These previous studies have been conducted predominantly in white women, and different patterns of co-exposures such as reproductive behavior may underlie inconsistencies.

Our study has some limitations. First, we did not have dietary intake or complete physical activity information for all the studies in the AMBER consortium and did not account for these covariates in our analyses. Additionally, we did not have information on reasons for alcohol abstinence, which could shed light on the elevated risk observed among never drinkers in our study. Our pooled analysis approach may be affected by differences between studies. Differences in categorical data collection between studies prevented evaluation of intake as a continuous variable, by grams of alcohol or by alcohol type, but previous studies suggest that cancer risk is most strongly associated with the type of alcoholic beverage most frequently consumed in a population (3,6). Furthermore, while we did not find statistical evidence of heterogeneity by study, we did observe some study-specific qualitative differences in the alcohol-risk association. Finally, case-control studies in the consortium may have been subject to recall bias, but work from the Nurses' Health Study did not find significant differences in recall of alcohol intake comparing pre- and post-diagnosis questionnaires among women with breast cancer (55).

To conclude, in a large consortium of African American women from around the United States, risk of breast cancer was elevated among never drinkers as well as those who reported drinking 7 or more drinks/week, irrespective of HR status or triple negative subtype. Our findings support alcohol as a risk factor for breast cancer in African American women, where risk associations have been less well-studied. Future work should seek to understand why African American women who abstain from alcohol have an increased risk

of breast cancer. Finally, our results suggest that drinking cessation may be associated with a reduced risk of invasive breast cancer and may be a targetable public health intervention strategy among African American women.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding Sources: This work was supported by the National Institutes of Health: P01 CA151135 (C.B. Ambrosone, J.R. Palmer, and A.F. Olshan), R01 CA058420 (L. Rosenberg), UM1 CA164974 (L. Rosenberg and J.R. Palmer), R01 CA098663 (J.R. Palmer), R01 CA100598 (C.B. Ambrosone), UM1 CA164973 (L.N. Kolonel), R01 CA54281 (L.N. Kolonel), P50 CA58223 (M.A. Troester and A.F. Olshan), U01 CA179715 (M.A. Troester and A.F. Olshan); and by the Komen for the Cure Foundation, the Breast Cancer Research Foundation (C.B. Ambrosone); and the University Cancer Research Fund of North Carolina. The results do not necessarily reflect the views of the NIH or the sponsors, who had no role in study design; data collection, analysis, or interpretation; or writing and submission of the manuscript.

References

1. Kabat GC, Kim M, Phipps AI, Li CI, Catherine R, Wactawski-wende J, et al. Smoking and alcohol consumption in relation to risk of triple-negative breast cancer in a cohort of postmenopausal women. *Cancer causes & control: CCC*. 2012; 22:775–83.
2. Dumeaux V, Lund E, Hjarta A. Use of Oral Contraceptives, Alcohol, and Risk for Invasive Breast Cancer. *Cancer Epidemiology, Biomarkers & Prevention*. 2004; 13:1302–7.
3. Fagherazzi G, Vilier A, Boutron-Ruault M-C, Mesrine S, Clavel-Chapelon F. Alcohol consumption and breast cancer risk subtypes in the E3N-EPIC cohort. *European journal of cancer prevention: the official journal of the European Cancer Prevention Organisation (ECP)*. 2014:1–6. [PubMed: 23492954]
4. Tjønneland A, Christensen J, Olsen A, Stripp C, Thomsen BL, Overvad K, et al. Alcohol intake and breast cancer risk: the European Prospective Investigation into Cancer and Nutrition (EPIC). *Cancer causes & control: CCC*. 2007; 18:361–73. [PubMed: 17364225]
5. Hamajima N, Hirose K, Tajima K, Rohan T, Calle EE, Heath CW, et al. Alcohol, tobacco and breast cancer--collaborative reanalysis of individual data from 53 epidemiological studies, including 58,515 women with breast cancer and 95,067 women without the disease. *British journal of cancer*. 2002; 87:1234–45. [PubMed: 12439712]
6. Allen NE, Beral V, Casabonne D, Kan SW, Reeves GK, Brown A, et al. Moderate alcohol intake and cancer incidence in women. *Journal of the National Cancer Institute*. 2009; 101:296–305. [PubMed: 19244173]
7. Longnecker MP, Berlin JA, Orza MJ, Chalmers TC. A Meta-analysis of Alcohol Consumption in Relation to Risk of Breast Cancer. *JAMA: the journal of the American Medical Association*. 1988; 260:652–6. [PubMed: 3392790]
8. Terry MB, Zhang FF, Kabat G, Britton Ja, Teitelbaum SL, Neugut AI, et al. Lifetime alcohol intake and breast cancer risk. *Annals of epidemiology*. 2006; 16:230–40. [PubMed: 16230024]
9. Park SY, Kolonel LN, Lim U, White KK, Henderson BE, Wilkens LR. Alcohol consumption and breast cancer risk among women from five ethnic groups with light to moderate intakes: The Multiethnic Cohort Study. *International Journal of Cancer*. 2014; 134:1504–10. [PubMed: 24037751]
10. Volcik, Ka, Ballantyne, CM., Fuchs, FD., Sharrett, aR, Boerwinkle, E. Relationship of alcohol consumption and type of alcoholic beverage consumed with plasma lipid levels: differences between Whites and African Americans of the ARIC study. *Annals of epidemiology*. 2008; 18:101–7. [PubMed: 17855114]

11. Kinney, aY, Millikan, RC., Lin, YH., Moorman, PG., Newman, B. Alcohol consumption and breast cancer among black and white women in North Carolina (United States). *Cancer causes & control: CCC*. 2000; 11:345–57. [PubMed: 10843445]
12. Jackson CL, Hu FB, Kawachi I, Williams DR, Mukamal KJ, Rimm EB. Black–White Differences in the Relationship Between Alcohol Drinking Patterns and Mortality Among US Men and Women. *American Journal of Public Health*. 2015; 105:S534–43. [PubMed: 25905819]
13. Darrow SL, Russell M, Cooper ML, Mudar PJ, Frone M. Sociodemographic Correlates of Alcohol Consumption Among African-American and White Women. *Women & Health*. 1992; 18:1–15.
14. Williams, LA., Olshan, AF, Tse, CK., Bell, ME., Troester, MA. *Cancer Causes & Control*. Vol. 27. Springer International Publishing; Alcohol intake and invasive breast cancer risk by molecular subtype and race in the Carolina Breast Cancer Study; p. 259-69.
15. Jones-Webb R. Drinking patterns and problems among African-Americans: recent findings. *Alcohol health and research world*. 1998; 22:260–4. [PubMed: 15706752]
16. Holt CL, Roth DL, Huang J, Clark EM. Gender Differences in the Roles of Religion and Locus of Control on Alcohol Use and Smoking Among African Americans. *Journal of Studies on Alcohol and Drugs*. 2015; 76:482–92. [PubMed: 25978836]
17. Angelica MD, Fong Y. Ethnicity and Alcohol Consumption among U.S. Adults with Diabetes. *Oct. 2008* 141:520–9.
18. Zhao G, Ford ES, Mokdad AH. Racial/ethnic variation in hypertension-related lifestyle behaviours among US women with self-reported hypertension. *Journal of human hypertension*. 2008; 22:608–16. [PubMed: 18496555]
19. Bernstein L, Teal CR, Joslyn S, Wilson J. Ethnicity-related variation in breast cancer risk factors. *Cancer*. 2003; 97:222–9. [PubMed: 12491485]
20. Hall IJ, Moorman PG, Millikan RC, Newman B. Comparative analysis of breast cancer risk factors among African-American women and white women. *American Journal of Epidemiology*. 2005; 161:40–51. [PubMed: 15615914]
21. Warner ET, Tamimi RM, Boggs Da, Rosner B, Rosenberg L, Colditz Ga, et al. Estrogen receptor positive tumors: do reproductive factors explain differences in incidence between black and white women? *Cancer causes & control: CCC*. 2013; 24:731–9. [PubMed: 23380944]
22. Singletary KW, Gapstur SM. Alcohol and Breast Cancer: Review of Epidemiologic and Experimental Evidence and Potential Mechanisms. *JAMA: the journal of the American Medical Association*. 2001; 286:2143–51. [PubMed: 11694156]
23. Seitz HK, Stickel F. Molecular mechanisms of alcohol-mediated carcinogenesis. *Nature reviews Cancer*. 2007; 7:599–612. [PubMed: 17646865]
24. Ambrosone CB, Zirpoli G, Hong C-C, Yao S, Troester Ma, Bandera EV, et al. Important Role of Menarche in Development of Estrogen Receptor-Negative Breast Cancer in African American Women. *J Natl Cancer Inst*. 2015; 107:djv172–djv172. [PubMed: 26085483]
25. Palmer JR, Boggs DA, Wise LA, Ambrosone CB, Adams-Campbell LL, Rosenberg L. Parity and Lactation in Relation to Estrogen Receptor Negative Breast Cancer in African American Women. *Cancer Epidemiology Biomarkers & Prevention*. 2011; 20:1883–91.
26. Phipps AI, Chlebowski RT, Prentice R, McTiernan A, Wactawski-Wende J, Kuller LH, et al. Reproductive history and oral contraceptive use in relation to risk of triple-negative breast cancer. *Journal of the National Cancer Institute*. 2011; 103:470–7. [PubMed: 21346227]
27. Li CI, Beaber EF, Tang M-TC, Porter PL, Daling JR, Malone KE. Reproductive factors and risk of estrogen receptor positive, triple-negative, and HER2-neu overexpressing breast cancer among women 20–44 years of age. *Breast cancer research and treatment*. 2013; 137:579–87. [PubMed: 23224237]
28. Millikan RC, Newman B, Tse C-K, Moorman PG, Conway K, Dressler LG, et al. Epidemiology of basal-like breast cancer. *Breast cancer research and treatment*. 2008; 109:123–39. [PubMed: 17578664]
29. Dolle JM, Daling JR, White E, Brinton La, Doody DR, Porter PL, et al. Risk factors for triple-negative breast cancer in women under the age of 45 years. *Cancer Epidemiology, Biomarkers & Prevention*. 2009; 18:1157–66.

30. Yang XR, Sherman ME, Rimm DL, Lissowska J, Brinton LA, Peplonska B, et al. Differences in risk factors for breast cancer molecular subtypes in a population-based study. *Cancer Epidemiology, Biomarkers & Prevention*. 2007; 16:439–43.
31. Bandera EV, Chandran U, Hong C-C, Troester Ma, Bethea TN, Adams-Campbell LL, et al. Obesity, body fat distribution, and risk of breast cancer subtypes in African American women participating in the AMBER Consortium. *Breast Cancer Research and Treatment*. 2015; 150:655–66. [PubMed: 25809092]
32. Eriksson CJ, Fukunaga T, Sarkola T, Lindholm HAL. Estrogen-related acetaldehyde elevation in women during alcohol intoxication. *Alcoholism, clinical and experimental research*. 1996; 20:1192–5.
33. Beland FA, Benson RW, Mellick PW, Kovatch RM, Roberts DW, Fang JLDD. Effect of ethanol on the tumorigenicity of urethane (ethyl carbamate) in B6C3F1 mice. *Food and chemical toxicology*. 2005; 43:1–19. [PubMed: 15582191]
34. Soffritti M, Belpoggi F, Cevolani D, Guarino M, Padovani MMC. Results of long-term experimental studies on the carcinogenicity of methyl alcohol and ethyl alcohol in rats. *Annals of the New York Academy of Sciences*. 2002; 982:46–69. [PubMed: 12562628]
35. Watabiki T, Okii Y, Tokiyasu T, Yoshimura S, Yoshida M, Akane A, Shikata NTA. Long-term ethanol consumption in ICR mice causes mammary tumor in females and liver fibrosis in males. *Alcoholism, clinical and experimental research*. 2000; 24:117S–122S.
36. Wang M, McIntee EJ, Cheng G, Shi Y, Villalta PW, Hecht SS. Identification of DNA Adducts of Acetaldehyde. *Chemical Research in Toxicology*. 2000; 13:1149–57. [PubMed: 11087437]
37. Newman B, Moorman PG, Millikan R, Qaqish BF, Gerads J, Aldrich TE, et al. The Carolina Breast Cancer Study: integrating population-based epidemiology and molecular biology. *Breast cancer research and treatment*. 1995; 35:51–60. [PubMed: 7612904]
38. Rosenberg L, Adams-Campbell L, Palmer JR. The Black Women’s Health Study: a follow-up study for causes and preventions of illness. *Journal of the American Medical Women’s Association*. 1972; (50):56–8.
39. Kolonel LN, Henderson BE, Hankin JH, Nomura AM, Wilkens LR, Pike MC, et al. A multiethnic cohort in Hawaii and Los Angeles: baseline characteristics. *American journal of epidemiology*. 2000; 151:346–57. [PubMed: 10695593]
40. Ambrosone CB, Ciupak GL, Bandera EV, Jandorf L, Bovbjerg DH, Zirpoli G, et al. Conducting Molecular Epidemiological Research in the Age of HIPAA: A Multi-Institutional Case-Control Study of Breast Cancer in African-American and European-American Women. *Journal of oncology*. 2009; 2009:871250. [PubMed: 19865486]
41. Palmer JR, Ambrosone CB, Olshan AF. A collaborative study of the etiology of breast cancer subtypes in African American women: the AMBER consortium. *Cancer causes & control: CCC*. 2014; 25:309–19. [PubMed: 24343304]
42. Bandera EV, Chandran U, Zirpoli G, McCann SE, Ciupak G, Ambrosone CB. Rethinking sources of representative controls for the conduct of case-control studies in minority populations. *BMC medical research methodology*. 2013; 13:71. [PubMed: 23721229]
43. Bethea TN, Rosenberg L, Castro-Webb N, Lunetta KL, Sucheston-Campbell LE, Ruiz-Narvaez EA, et al. Family History of Cancer in Relation to Breast Cancer Subtypes in African American Women. *Cancer Epidemiology Biomarkers & Prevention*. 2016; 25:366–73.
44. Greenland S, Pearl J, Robins JM. Causal diagrams for epidemiologic research. *Epidemiology*. 1999; 10:37–48. [PubMed: 9888278]
45. Bagnardi V, Randi G, Lubin J, Consonni D, Lam TK, Subar AF, et al. Alcohol consumption and lung cancer risk in the environment and genetics in lung cancer etiology (EAGLE) study. *American Journal of Epidemiology*. 2010; 171:36–44. [PubMed: 19933698]
46. Li CI, Chlebowski RT, Freiberg M, Johnson KC, Kuller L, Lane D, et al. Alcohol consumption and risk of postmenopausal breast cancer by subtype: the women’s health initiative observational study. *Journal of the National Cancer Institute*. 2010; 102:1422–31. [PubMed: 20733117]
47. Falk RT, Maas P, Schairer C, Chatterjee N, Mabie JE, Cunningham C, et al. Alcohol and risk of breast cancer in postmenopausal women: an analysis of etiological heterogeneity by multiple tumor characteristics. *American journal of epidemiology*. 2014; 180:705–17. [PubMed: 25150269]

48. Li CI. Relationship between Established Breast Cancer Risk Factors and Risk of Seven Different Histologic Types of Invasive Breast Cancer. *Cancer Epidemiology Biomarkers & Prevention*. 2006; 15:946–54.
49. Larsson SC, Mantzoros CS, Wolk A. Diabetes mellitus and risk of breast cancer: A meta-analysis. *International Journal of Cancer*. 2007; 121:856–62. [PubMed: 17397032]
50. Gapstur SM, Potter JD, Drinkard C, Folsom R. Synergistic effect between alcohol and estrogen replacement therapy on risk of breast cancer differs by estrogen / progesterone receptor status in the Iowa Women's Health Study. *Cancer Epidemiology, Biomarkers & Prevention*. 1995; 4:313–8.
51. Chen W, Rosner B, Hankinson SE, Colditz GA, Willett WC. Moderate Alcohol Consumption During Adult Life, Drinking Patterns, and Breast Cancer Risk. *Breast Diseases: A Year Book Quarterly*. 2011; 23:231–2.
52. Suzuki R, Orsini N, Mignone L, Saji S, Wolk A. Alcohol intake and risk of breast cancer defined by estrogen and progesterone receptor status--a meta-analysis of epidemiological studies. *International journal of cancer Journal international du cancer*. 2008; 122:1832–41. [PubMed: 18067133]
53. Jung S, Wang M, Anderson K, Baglietto L, Bergkvist L, Bernstein L, et al. Alcohol consumption and breast cancer risk by estrogen receptor status: in a pooled analysis of 20 studies. *International journal of epidemiology*. 2015:1–13. [PubMed: 25859586]
54. Schonberg, Ma, Marcantonio, ER., Li, D., Silliman, Ra, Ngo, L., McCarthy, EP. Breast cancer among the oldest old: tumor characteristics, treatment choices, and survival. *Journal of clinical oncology: official journal of the American Society of Clinical Oncology*. 2010; 28:2038–45. [PubMed: 20308658]
55. Giovannucci E, Stampfer MJ, Colditz Ga, Manson JE, Rosner Ba, Longnecker MP, et al. Recall and selection bias in reporting past alcohol consumption among breast cancer cases. *Cancer causes & control: CCC*. 1993; 4:441–8. [PubMed: 8218876]

Table 1

Exposure characteristics of cases and controls in the AMBER Consortium stratified by study: recent alcohol intake, smoking status, oral contraceptive use, and menopausal status

	CBCSN (%)		WCHS N (%)		MEC N (%)		BWHS N (%)		Total N (%)	
	Cases	Controls	Cases	Controls	Cases	Controls	Cases	Controls	Cases	Controls
Number of alcoholic drinks per week										
Never drinkers	293 (37.5)	309 (39.4)	798 (65.4)	791 (65.1)	629 (55.5)	2,360 (53.3)	797 (40.4)	4,063 (37.6)	2,517 (49.3)	7,523 (43.4)
>0-<4	193 (24.7)	232 (29.6)	113 (9.3)	111 (9.1)	226 (19.9)	906 (20.5)	492 (24.9)	2,868 (26.6)	1,024 (20.1)	4,117 (23.9)
4-<7	32 (4.1)	44 (5.6)	40 (3.3)	43 (3.5)	43 (3.8)	174 (3.9)	109 (5.5)	615 (5.7)	224 (4.4)	876 (5.1)
7-<14	46 (5.9)	24 (3.1)	30 (2.5)	38 (3.1)	46 (4.1)	192 (4.3)	68 (3.5)	351 (3.3)	190 (3.7)	605 (3.5)
14	42 (5.4)	30 (3.8)	23 (1.9)	22 (1.8)	83 (7.3)	259 (5.9)	30 (1.5)	135 (1.3)	178 (3.5)	446 (2.6)
Past drinkers	175 (22.4)	145 (18.5)	216 (17.7)	211 (17.4)	107 (9.4)	540 (12.2)	477 (24.2)	2,767 (25.6)	975 (19.1)	3,663 (21.3)
Smoking Status										
Never Smoker	435 (55.7)	467 (59.6)	745 (61.1)	702 (57.7)	508 (45.7)	2,050 (46.9)	1,168 (59.3)	6,288 (58.3)	2,856 (56.2)	9,507 (55.4)
Current Smoker	134 (17.2)	150 (19.1)	199 (16.3)	244 (20.1)	216 (19.4)	807 (18.5)	343 (17.4)	1,978 (18.3)	892 (17.6)	3,179 (18.5)
Former Smoker	212 (27.1)	167 (21.3)	276 (22.6)	270 (22.2)	388 (34.9)	1,515 (34.7)	460 (23.3)	2,525 (23.4)	1,336 (26.3)	4,477 (26.1)
Missing	0	0	0	0	22	59	2	8	24	67
Oral Contraceptive Use										
Never, <1 year	380 (48.7)	386 (49.4)	589 (48.6)	593 (49.1)	688 (63.1)	2,780 (66.1)	825 (41.8)	4,831 (44.7)	2,482 (49.1)	8,590 (55.6)
1-9 years	276 (35.3)	295 (37.8)	422 (34.8)	441 (36.5)	279 (25.6)	994 (23.7)	820 (41.6)	4,398 (40.7)	1,797 (35.5)	6,128 (36.1)
10+ years	125 (16.0)	100 (12.8)	202 (16.7)	175 (14.5)	123 (11.3)	429 (10.2)	328 (16.6)	1,570 (14.5)	778 (15.4)	2,274 (13.4)
Missing	0	3	7	7	44	228	0	0	51	238
Menopausal Status										
Pre-	286 (40.9)	298 (43.1)	574 (49.6)	565 (49.1)	140 (12.7)	388 (9.0)	717 (42.7)	4,179 (43.6)	1,717 (37.0)	5,430 (34.5)
Post-	413 (59.1)	393 (56.9)	583 (50.4)	587 (50.9)	964 (87.3)	3,905 (90.0)	964 (57.4)	5,412 (56.4)	2,924 (63.0)	10,297 (65.5)
Missing	82	93	63	64	30	138	292	1,208	467	1,503

Table 2
Recent alcohol intake in relation to Hormone Receptor status among invasive breast cancer cases in the AMBER Consortium

Drinks per week	Basic model ^a α-cr1>OR (95% CI)						Cases N (%)	MV model ^b OR (95% CI)	Cases N (%)	Basic model ^c OR (95% CI)	Cases N (%)	MV model ^b OR (95% CI)
	Controls N (%)	Cases N (%)	Controls N (%)	OR	Cases N (%)	MV model ^b OR (95% CI)						
All Invasive Breast Cancer												
Never drinkers	7,516 (43.7)	2,517 (49.3)	1.12 (1.03–1.22)	6,476 (43.4)	2,134 (49.1)	1.12 (1.02–1.24)						
>0–<4	4,114 (23.9)	1,023 (20.0)	Ref.	3,539 (23.7)	862 (19.8)	Ref.						
4–<7	876 (5.1)	223 (4.4)	0.99 (0.84–1.17)	750 (5.0)	192 (4.4)	1.04 (0.86–1.25)						
7–<14	605 (3.5)	190 (3.7)	1.13 (0.94–1.36)	534 (3.6)	159 (3.7)	1.07 (0.88–1.32)						
14	446 (2.6)	178 (3.5)	1.34 (1.10–1.63)	394 (2.6)	149 (3.4)	1.33 (1.07–1.64)						
Past drinkers	3,661 (21.3)	973 (19.1)	1.01 (0.91–1.12)	3,224 (21.6)	853 (19.6)	1.04 (0.93–1.17)						
p-trend *			0.01			0.09						
ER+												
Never drinkers	7,516 (43.7)	1,259 (49.4)	1.10 (0.98–1.23)	6,476 (43.4)	1,072 (48.9)	1.13 (0.99–1.29)	638 (46.3)	1.12 (0.96–1.30)	536 (45.7)	1.07 (0.90–1.27)		
>0–<4	4,114 (23.9)	508 (19.9)	Ref.	3,539 (23.7)	428 (19.5)	Ref.	297 (21.5)	Ref.	254 (21.7)	Ref.		
4–<7	876 (5.1)	109 (4.3)	0.98 (0.78–1.23)	750 (5.0)	98 (4.5)	1.10 (0.86–1.40)	61 (4.4)	0.97 (0.72–1.31)	53 (4.5)	1.02 (0.74–1.41)		
7	1,051 (6.1)	189 (7.4)	1.20 (1.00–1.45)	928 (6.2)	161 (7.3)	1.20 (0.97–1.47)	111 (8.1)	1.35 (1.06–1.72)	91 (7.8)	1.31 (1.00–1.72)		
Past drinkers	3,661 (21.3)	484 (19.0)	0.99 (1.00–1.45)	3,224 (21.6)	434 (19.8)	1.07 (0.92–1.24)	272 (19.7)	1.11 (0.92–1.33)	238 (20.3)	1.11 (0.91–1.35)		
p-trend *			0.12			0.18		0.03		0.08		
ER–												
Never drinkers	7,516 (43.7)	1,001 (49.5)	1.12 (0.99–1.28)	6,476 (43.4)	852 (48.8)	1.14 (0.98–1.32)	825 (47.0)	1.10 (0.96–1.26)	688 (46.6)	1.07 (0.92–1.25)		
>0–<4	4,114 (23.9)	395 (19.5)	Ref.	3,539 (23.7)	336 (19.2)	Ref.	377 (21.5)	Ref.	318 (21.5)	Ref.		
4–<7	876 (5.1)	77 (3.8)	0.87 (0.67–1.13)	750 (5.0)	72 (4.1)	1.02 (0.77–1.35)	82 (4.7)	1.04 (0.80–1.35)	70 (4.7)	1.08 (0.81–1.43)		
7	1,051 (6.1)	142 (7.0)	1.18 (0.95–1.45)	928 (6.2)	121 (6.9)	1.19 (0.95–1.50)	138 (7.9)	1.32 (1.06–1.64)	112 (7.6)	1.28 (1.00–1.63)		
Past drinkers	3,661 (21.3)	406 (20.1)	1.07 (0.95–1.46)	3,224 (21.6)	366 (21.0)	1.14 (0.97–1.35)	333 (19.0)	1.02 (0.87–1.20)	290 (19.6)	1.04 (0.88–1.25)		
p-trend *			0.30			0.29		0.03		0.08		
PR+												
Never drinkers	7,516 (43.7)	1,001 (49.5)	1.12 (0.99–1.28)	6,476 (43.4)	852 (48.8)	1.14 (0.98–1.32)	825 (47.0)	1.10 (0.96–1.26)	688 (46.6)	1.07 (0.92–1.25)		
>0–<4	4,114 (23.9)	395 (19.5)	Ref.	3,539 (23.7)	336 (19.2)	Ref.	377 (21.5)	Ref.	318 (21.5)	Ref.		
4–<7	876 (5.1)	77 (3.8)	0.87 (0.67–1.13)	750 (5.0)	72 (4.1)	1.02 (0.77–1.35)	82 (4.7)	1.04 (0.80–1.35)	70 (4.7)	1.08 (0.81–1.43)		
7	1,051 (6.1)	142 (7.0)	1.18 (0.95–1.45)	928 (6.2)	121 (6.9)	1.19 (0.95–1.50)	138 (7.9)	1.32 (1.06–1.64)	112 (7.6)	1.28 (1.00–1.63)		
Past drinkers	3,661 (21.3)	406 (20.1)	1.07 (0.95–1.46)	3,224 (21.6)	366 (21.0)	1.14 (0.97–1.35)	333 (19.0)	1.02 (0.87–1.20)	290 (19.6)	1.04 (0.88–1.25)		
p-trend *			0.30			0.29		0.03		0.08		
HER2+												
HER2–												

	Basic model ^a model ^a _{cr}I>OR (95% CI)						MV model ^b OR (95% CI)	Cases N (%)	Basic model ^a OR (95% CI)	Cases N (%)	MV model ^b OR (95% CI)	Cases N (%)	MV model ^b OR (95% CI)
	Controls N (%)	Cases N (%)	model ^a _{cr}I>OR (95% CI)	Controls N (%)	Cases N (%)	MV model ^b OR (95% CI)							
Never drinkers	7,516 (43.7)	254 (46.8)	1.01 (0.80–1.28)	6,476 (43.4)	212 (45.8)	0.93 (0.72–1.21)	857 (46.4)	1.01 (0.88–1.15)	1,007 (46.9)	0.93 (0.72–1.21)	857 (46.4)	1.04 (0.90–1.21)	
>0–4	4,114 (23.9)	115 (21.2)	Ref.	3,539 (23.7)	101 (21.8)	Ref.	367 (19.9)	Ref.	436 (20.3)	Ref.	367 (19.9)	Ref.	
4–7	876 (5.1)	31 (5.7)	1.21 (0.80–1.83)	750 (5.0)	28 (6.0)	1.34 (0.87–2.09)	79 (4.3)	0.91 (0.70–1.18)	87 (4.1)	1.04 (0.79–1.37)	79 (4.3)	1.04 (0.79–1.37)	
7	1,051 (6.1)	44 (8.1)	1.33 (0.93–1.92)	928 (6.2)	33 (7.1)	1.17 (0.77–1.77)	149 (8.0)	1.27 (1.03–1.56)	169 (7.9)	1.36 (1.09–1.70)	149 (8.0)	1.36 (1.09–1.70)	
Past drinkers	3,661 (21.3)	99 (18.2)	0.94 (0.72–1.25)	3,224 (21.6)	89 (19.2)	0.99 (0.73–1.73)	398 (21.5)	1.08 (0.93–1.26)	450 (20.9)	1.15 (0.99–1.37)	398 (21.5)	1.15 (0.99–1.37)	
p-trend*			0.06			0.38		0.05				0.01	
Non Triple Negative													
Triple Negative													
Never drinkers	7,516 (43.7)	2,195 (50.3)	1.14 (1.04–1.25)	6,476 (43.4)	1,867 (50.1)	1.14 (1.03–1.26)	267 (42.7)	0.97 (0.77–1.20)	322 (43.4)	0.97 (0.77–1.20)	267 (42.7)	0.96 (0.76–1.22)	
>0–4	4,114 (23.9)	859 (19.7)	Ref.	3,539 (23.7)	725 (19.5)	Ref.	137 (21.9)	Ref.	164 (22.1)	Ref.	137 (21.9)	Ref.	
4–7	876 (5.1)	188 (4.3)	0.99 (0.82–1.18)	750 (5.0)	161 (4.3)	1.02 (0.84–1.24)	31 (5.0)	1.01 (0.68–1.51)	35 (4.7)	1.14 (0.74–1.74)	31 (5.0)	1.14 (0.74–1.74)	
7	1,051 (6.1)	305 (7.0)	1.21 (1.04–1.40)	928 (6.2)	255 (6.9)	1.16 (0.98–1.37)	53 (8.5)	1.39 (1.01–1.92)	63 (8.5)	1.39 (0.98–2.00)	53 (8.5)	1.39 (0.98–2.00)	
Past drinkers	3,661 (21.3)	815 (18.7)	0.99 (0.89–1.11)	3,224 (21.6)	716 (19.2)	1.02 (0.91–1.15)	137 (21.9)	1.19 (0.93–1.50)	158 (21.3)	1.21 (0.93–1.58)	137 (21.9)	1.21 (0.93–1.58)	
p-trend*			0.04			0.18		0.06				0.06	

^aBasic model adjusted for age, study, time period and geographic region.

^bMultivariable model adjusted for age, study, time period, geographic region, education, age at menarche, BMI, menopausal status (pre-/post-), HT use (ever/never), parity, smoking status, and duration of oral contraceptive use.

* P-trend excludes never and past drinkers.

Table 3

Recent alcohol intake and invasive breast cancer by study from the AMBER consortium

	Controls N (%)	Cases N (%)	Basic model ^a OR (95% CI)	Controls N (%)	Cases N (%)	MV model ^b OR (95% CI)
CBCS						
Number of alcoholic drinks per week						
Never drinkers	309 (39.4)	293 (37.5)	1.16 (0.90–1.50)	258 (39.0)	253 (37.5)	1.23 (0.92–1.65)
>0–<4	232 (29.6)	193 (24.7)	Ref.	199 (30.0)	166 (24.6)	Ref.
4–<7	44 (5.6)	32 (4.1)	0.88 (0.53–1.43)	36 (5.4)	28 (4.2)	0.95 (0.55–1.64)
7	54 (6.9)	88 (11.3)	2.02 (1.37–2.99)	46 (6.9)	72 (10.7)	2.03 (1.29–3.18)
Past drinkers	145 (18.5)	75 (22.4)	1.50 (1.11–2.03)	123 (18.6)	155 (23.0)	1.53 (1.09–2.14)
p-trend*			<0.01			0.01
WCHS						
Number of alcoholic drinks per week						
Never drinkers	791 (65.1)	798 (65.4)	0.96 (0.72–1.27)	742 (66.0)	727 (65.1)	0.89 (0.66–1.21)
>0–<4	111 (9.1)	113 (9.3)	Ref.	101 (8.9)	104 (9.3)	Ref.
4–<7	43 (3.5)	40 (3.3)	0.91 (0.55–1.51)	37 (3.3)	38 (3.4)	1.01 (0.59–1.73)
7	60 (4.9)	53 (4.3)	0.84 (0.53–1.32)	57 (5.1)	50 (4.5)	0.81 (0.51–1.31)
Past drinkers	211 (17.4)	216 (17.7)	0.96 (0.70–1.34)	187 (16.6)	198 (17.7)	0.98 (0.69–1.39)
p-trend*			0.42			0.40
MEC						
Number of alcoholic drinks per week						
Never drinkers	2,360 (53.3)	629 (55.5)	1.07 (0.90–1.27)	1,908 (51.4)	478 (53.0)	1.10 (0.91–1.34)
>0–<4	906 (20.5)	226 (19.9)	Ref.	759 (20.5)	183 (20.3)	Ref.
4–<7	174 (3.9)	43 (3.8)	0.99 (0.69–1.43)	150 (4.0)	32 (3.6)	0.92 (0.60–1.39)
7	451 (10.2)	129 (11.4)	1.15 (0.90–1.47)	411 (11.8)	108 (12.0)	1.06 (0.81–1.40)
Past drinkers	540 (12.2)	107 (9.4)	0.79 (0.62–1.03)	481 (13.0)	101 (11.2)	0.87 (0.66–1.14)
p-trend*			0.29			0.63

	Controls N (%)	Cases N (%)	Basic model ^a OR (95% CI)	Controls N (%)	Cases N (%)	MV model ^b OR (95% CI)
BWHS						
Number of alcoholic drinks per week						
Never drinkers	4,056 (37.6)	797 (40.5)	1.15 (1.02–1.30)	3,568 (37.9)	676 (40.8)	1.17 (1.02–1.34)
>0–4	2,865 (26.6)	491 (24.9)	Ref.	2,480 (26.3)	409 (24.7)	Ref.
4–7	615 (5.7)	108 (5.5)	1.02 (0.82–1.28)	527 (5.6)	94 (5.7)	1.08 (0.85–1.38)
7	486 (4.5)	98 (5.0)	1.17 (0.92–1.49)	414 (4.4)	78 (4.7)	1.13 (0.87–1.48)
Past drinkers	2,765 (25.6)	475 (24.1)	0.99 (0.87–1.14)	2,433 (25.8)	399 (24.1)	1.02 (0.88–1.19)
p-trend *			0.23			0.38

^aBasic model adjusted for age, study, time period and geographic region.

^bMultivariable model adjusted for age, study, time period, geographic region, education, age at menarche, BMI, menopausal status (pre-/post-), HT use (ever/never), parity, smoking status, and duration of oral contraceptive use

* p-trend excludes never and past drinkers

Sensitivity analysis to examine alcohol intake (drinks per week) among ever drinkers during different age periods prior to diagnosis in relation to invasive breast cancer for CBCS and WCHS

Table 4

	Controls N (%)	Cases N (%)	Basic model ^a OR (95% CI)	Controls N (%)	Cases N (%)	MV Model ^b OR (95% CI)
Young Age Drinks per week (<30 years old)*						
Number of alcoholic drinks per week						
0	1,287 (64.3)	1,295 (64.7)	1.05 (0.90–1.23)	1,156 (64.7)	1,159 (64.6)	1.04 (0.88–1.23)
>0-<4	463 (23.1)	439 (21.9)	Ref.	410 (22.9)	396 (22.1)	Ref.
4-<7	95 (4.8)	96 (4.8)	1.06 (0.78–1.45)	82 (4.6)	88 (4.9)	1.09 (0.78–1.52)
7	157 (7.8)	173 (8.6)	1.16 (0.90–1.49)	139 (7.8)	150 (8.4)	1.07 (0.81–1.41)
p-trend			0.25			0.64
Middle age Drinks per week (30–<50 years old)[#]						
Number of alcoholic drinks per week						
0	1,265 (63.3)	1,227 (62.1)	1.01 (0.86–1.19)	1,140 (64.4)	1,102 (62.4)	0.97 (0.81–1.16)
>0-<4	442 (22.3)	420 (21.3)	Ref.	383 (21.6)	369 (20.9)	Ref.
4-<7	109 (5.5)	132 (6.7)	1.26 (0.95–1.68)	92 (5.2)	124 (7.0)	1.38 (1.01–1.88)
7	176 (8.9)	197 (10.0)	1.17 (0.92–1.49)	155 (8.7)	171 (9.7)	1.14 (0.87–1.49)
p-trend			0.12			0.25
Older Age Drinks per week (50+ years old)						
Number of alcoholic drinks per week						
0	879 (78.8)	906 (80.3)	1.04 (0.81–1.34)	770 (78.7)	786 (79.9)	1.00 (0.75–1.32)
>0-<4	145 (13.0)	141 (12.5)	Ref.	132 (13.4)	125 (12.6)	Ref.
4-<7	45 (4.0)	30 (2.7)	0.70 (0.41–1.17)	36 (3.7)	27 (2.7)	0.76 (0.43–1.34)
7	46 (4.1)	51 (4.5)	1.08 (0.68–1.72)	42 (4.3)	47 (4.7)	1.07 (0.65–1.76)
p-trend			0.98			0.94
Restricted to Women 50+ with data for all age periods**						
Number of alcoholic drinks per week						

	Controls N (%)	Cases N (%)	Basic model ^a OR (95% CI)	Controls N (%)	Cases N (%)	MV Model ^b OR (95% CI)
0	875 (78.9)	893 (80.2)	1.25 (0.92–1.69)	767 (78.7)	774 (79.6)	1.22 (0.87–1.71)
>0–<4	145 (13.1)	140 (12.6)	Ref.	132 (13.5)	124 (12.8)	Ref.
4–<7	44 (4.0)	30 (2.7)	0.63 (0.36–1.11)	35 (3.6)	27 (2.8)	0.68 (0.37–1.26)
7	45 (4.1)	51 (4.6)	1.14 (0.68–1.89)	41 (4.2)	47 (4.8)	1.14 (0.66–1.97)
p-trend			0.69			0.65

^aBasic model adjusted for age, study, time period and geographic region.

^bMultivariable model adjusted for age, study, time period, geographic region, education, age at menarche, BMI, menopausal status (pre-/post-), HT use (ever/never), parity, smoking status, and duration of oral contraceptive use

* <25 years old for CBCS

25–<50 years old for CBCS