



HHS PUBLIC ACCESS

Author manuscript

Curr Opin Neurobiol. Author manuscript; available in PMC 2017 August 01.

Published in final edited form as:

Curr Opin Neurobiol. 2016 August ; 39: 108–115. doi:10.1016/j.conb.2016.05.002.

Axon degeneration: context defines distinct pathways

Matthew J Geden^{1,2} and Mohanish Deshmukh^{1,2}

¹Department of Cell Biology and Physiology, University of North Carolina, Chapel Hill, NC 27599, USA

²Neuroscience Center, University of North Carolina, Chapel Hill, NC 27599, USA

Abstract

Axon degeneration is an essential part of development, plasticity, and injury response and has been primarily studied in mammalian models in three contexts: 1) Axotomy-induced Wallerian degeneration, 2) Apoptosis-induced axon degeneration (axon apoptosis), and 3) Axon pruning. These three contexts dictate engagement of distinct pathways for axon degeneration. Recent advances have identified the importance of SARM1, NMNATs, NAD⁺ depletion, and MAPK signaling in axotomy-induced Wallerian degeneration. Interestingly, apoptosis-induced axon degeneration and axon pruning have many shared mechanisms both in signaling (*e.g.* DLK, JNKs, GSK3 α/β) and execution (*e.g.* Puma, Bax, caspase-9, caspase-3). However, the specific mechanisms by which caspases are activated during apoptosis *versus* pruning appear distinct, with apoptosis requiring Apaf-1 but not caspase-6 while pruning requires caspase-6 but not Apaf-1.

Introduction

Neuronal development and target innervation by axons and dendrites are essential processes required for the establishment of the nervous system. Simultaneously however, the selective destruction of neurons or only their projections is also important for the refinement of the wired brain [1]. While the mechanisms regulating the survival and elimination of neurons *via* apoptosis are well studied, the pathways governing the selective degeneration of the axons are less understood. Axon degeneration has been observed to occur in many contexts (*e.g.* pruning, axotomy, apoptosis, dying back) and models (*e.g.* *D. melanogaster*, *C. elegans*, mice), which has been discussed in several recent reviews [2,3,4,5,6].

In the mammalian model, mechanistic details of axon degeneration has been predominantly studied *in vitro* in three contexts: 1) **Axotomy** (also known as Wallerian degeneration), where the severing of axons results in the degeneration of axons distal to the cut site; 2) **Apoptosis-induced Axon Degeneration**, which we define here as ‘**Axon Apoptosis**’, where the entire neuron is exposed to apoptotic stimuli (*e.g.* global deprivation of trophic factors) resulting in the degeneration of both axons and soma; and 3) **Pruning-induced Axon Degeneration**, which we refer to here as ‘**Axon Pruning**’, where a subset of axons are

Corresponding author: Deshmukh, Mohanish (mohanish@med.unc.edu).

Conflict of interest statement

Nothing declared.

selectively exposed to a pruning stimuli (*e.g.* axon-only or 'local' deprivation of trophic factors) which results in the selective degeneration of only the axons exposed to the stimulus. Axon degeneration within these three contexts appears to engage overlapping but also distinct pathways. While axon degeneration is ultimately the convergent outcome of these pathways, the molecular mechanisms engaged are defined by the context of degeneration (just as cell death can be executed by distinct mechanisms, dependent on context). In this article, we primarily focus on the recent progress in mammalian models and define the shared and distinct mechanisms of axon degeneration that occur in these three contexts. These *in vitro* studies have been instrumental in providing significant insight into the distinct mechanisms of axon degeneration, which are now poised for future *in vivo* investigations.

Axotomy (Wallerian degeneration)

Wallerian degeneration is a unique and structured form of axon degeneration that occurs when a neuronal axon is crushed or severed (axotomized) from its cell body (illustrated in Figure 1). Wallerian degeneration has been well examined in the *Wld^s* mouse, where axotomy-induced axon degeneration is strikingly inhibited [7]. *Wld^s* mice express a unique gain-of-function gene product responsible for this protection, which was identified as the fusion of the N-terminal fragment of ubiquitination factor E4B (UBE4b) with NMNAT1 (nicotinamide mononucleotide adenylyltransferase 1) [8]. These findings implicated the role of NAD⁺ (nicotinamide adenine dinucleotide) metabolism in Wallerian degeneration, as NMNATs are key enzymes in the NAD⁺ salvage pathway [9]. NAM (Nicotinamide), which is a byproduct of NAD⁺ metabolism, is recycled into NMN (nicotinamide mononucleotide). NAD⁺ can then be regenerated from NMN by the NMNAT enzymes to maintain steady-state pools of intracellular NAD⁺. The rapid depletion of NAD⁺ has been identified to be a central event in axons undergoing Wallerian degeneration and addition of exogenous NAD⁺ (albeit at high concentrations) is sufficient to inhibit this degeneration [9,10,11]. However, recent evidence suggest that rather than NAD⁺ depletion, it could be the accumulation of NMN instead that triggers axon degeneration [12].

Recently, SARM1 (Sterile Alpha and TIR Motif 1) was identified as the first loss-of-function mutation to provide protection against Wallerian degeneration [13^{*},14]. SARM1-deficiency robustly protects axons following axotomy *in vivo* and *in vitro* [13^{*},14] and maintains NAD⁺ levels *in vitro* [15^{**}]. Conversely, direct activation of SARM1 rapidly depletes intracellular NAD⁺ and is sufficient to induce axon degeneration [14,15^{**}]. A key event that occurs after axotomy is the depletion of NMNAT2, which is sufficient to trigger axon degeneration [16]. SARM1 appears to be the effector of NMNAT2-depletion induced axon degeneration, as the degeneration triggered by NMNAT2 depletion can be fully rescued by the co-deletion of SARM1 [17]. The exact role of SARM1 in degeneration is not fully understood, but these data and a recent review suggest that NMNAT2 depletion causes the reduction of NAD⁺ levels which leads to subsequent SARM1 activation and axon degeneration [4,15^{**},17]. Downstream of SARM1, a MAP Kinase signaling pathway is activated to propagate the degenerative signal [18^{**},19]. This cascade includes DLK (Dual Leucine Zipper Kinase), MEKK4, and MLK2 which leads to the activation of c-Jun-N-terminal kinases (JNKs) *via* MKK4 and MKK7 [18^{**}]. Exactly how these signaling

pathways induce axon degeneration is not fully understood, but this involves additional mediators of Wallerian degeneration which include the ubiquitin-proteasome system [20], the ubiquitin ligase Phr1/Highwire [21,22], and SCG10 [23,24]. Furthermore, Ca^{2+} signaling and calpain activation are also important for mediating late events in axon degeneration [25,26,27].

Axon apoptosis (apoptosis-induced axon degeneration)

Neuronal apoptosis, which results in the degeneration of both the soma and axons, occurs extensively during development and in response to various neuronal insults [28,29]. This pathway has been well studied in the developing peripheral nervous system where sympathetic and sensory neurons, which are produced in excess, are acutely dependent on the limiting amounts of target-derived nerve growth factor (NGF). During target innervation, only those neurons that obtain robust NGF signaling survive whereas those deprived of NGF undergo apoptosis, a phenomenon that serves to match the number of innervating neurons with the size of target [30,31]. While neurons can undergo developmental or pathological apoptosis in multiple contexts, the model of NGF deprivation-induced apoptosis is a useful and well-recognized model for studying the molecular mechanisms of neuronal apoptosis. This phenomenon can also be recapitulated in cell culture where dissociated sympathetic or sensory neurons, or ganglia explants, are maintained in the presence of NGF. It should be noted that in these models both the soma and axons are exposed to direct NGF signaling, while *in vivo* NGF signaling primarily occurs at the axon terminals and the signal is retrogradely transported to the cell body. Importantly, elimination of NGF from the culture media, to deprive the entire neuron of NGF (often referred to as 'global deprivation'), results in the apoptotic degeneration of both the soma and axons (Illustrated in Figure 2) [30,32]. Since axon degeneration in this context is a consequence of the activation of the global apoptotic program, we refer to this apoptosis-induced axon degeneration here as 'axon apoptosis'.

Global deprivation of NGF initiates a well-defined apoptotic pathway that we briefly outline here. NGF deprivation induces the dephosphorylation of its receptor, Tropomyosin receptor kinase A (TrkA), which leads to the activation of a DLK-mediated apoptotic signal. DLK signaling activates downstream mediators (including JNKs, MAPKs, and GSK α/β) and leads to the transcriptional upregulation of proapoptotic BH3-only family proteins (including Bim, Puma, Bmf, Hrk/DP5) by transcription factors such as c-Jun, NF- κ B, and FOXOs [30,33**]. Recently, Puma was shown to be a key mediator of axon apoptosis. Not only is Puma induced in the cell body after global NGF deprivation, but Puma deficiency markedly reduces axon apoptosis [33**]. A major function of the BH3-only proteins such as Puma is to activate Bax, a pro-apoptotic member of the Bcl-2 family, which is required for neuronal apoptosis [30]. Some BH3-only proteins directly interact with Bax to induce its conformation activation, while others inhibit anti-apoptotic proteins of the Bcl-2 protein family (Bcl-2, Bcl-w, Bcl-XL, Mcl-1) [34]. In particular, Bcl-w and Bcl-XL are known to be localized to axons where they are important for maintaining axonal survival [33**,35].

Once Bax is activated, it translocates to mitochondria and inserts into the outer mitochondrial membrane, leading to the release of cytochrome *c* (cyt *c*) into the cytoplasm

[36]. Released cyt *c* can then bind to Apaf-1 (Apoptotic protease activating factor 1) to induce a conformational change in Apaf-1 that results in the recruitment of and activation of procaspase-9 (Casp9). Activated Casp9 functions to activate Caspase-3 (Casp3), and subsequently calpains, which together execute the degeneration of both soma and axons by targeting many cellular proteins for proteolysis [26]. Interestingly, while there are numerous mammalian caspases with diverse cellular functions, only Casp9 and Casp3 appear to be strictly required for global NGF-deprivation induced sympathetic neuronal apoptosis [37]. Lastly, X-linked Inhibitor of Apoptosis Protein (XIAP) inhibits caspase activity in both soma and axons, but is degraded during global NGF deprivation [38,39*]. Consistent with this, XIAP-deficient neurons exhibit enhanced axon degradation when globally deprived of NGF [39*].

This apoptotic pathway activated upon global deprivation of NGF has been recognized to be important for the degeneration of both the cell bodies and axons. Indeed, deletion or depletion of the key effectors of this apoptotic program (*e.g.* DLK, JNKs, c-Jun, FOXO3a, Puma, Bax, Apaf-1, Casp9, Casp3) blocks the death of neurons in response to global NGF deprivation [30,33**,37,40*,41]. Overall, these data suggest that in the context of global apoptotic stimulation, axonal degeneration is a consequence of the apoptotic program activated in the cell body.

Axon pruning (pruning-induced axon degeneration)

In addition to neuronal apoptosis, the nervous system undergoes substantial refinement by the pruning of axons and dendritic connections, which are selectively dismantled and rewired. Pruning is critical not only for establishing specific and appropriate neuronal circuitry during development but also for neuronal plasticity in the adult nervous system [1,42]. Precise spatial and temporal control of the degenerative machinery is important for pruning to ensure that only the targeted region of the axons are degraded. This spatial regulation is in contrast to apoptosis, where the entire neuron undergoes degeneration.

Similarly to axon apoptosis, mechanistic insights into the axon pruning pathway have primarily come from studies in peripheral neurons (sensory and sympathetic) subjected to NGF deprivation. An important distinction however is that while axonal apoptosis is triggered by ‘global’ deprivation of NGF (where neurons are completely deprived of NGF), axon pruning is observed in response to the selective deprivation of NGF from the axon while maintaining the neuronal cell body in NGF (illustrated in Figure 3). To study this pathway *in vitro*, neurons are cultured in compartmentalized chambers (*e.g.* Campenot [43] or microfluidic chambers [44]) where the cell bodies and the axons are maintained in different compartments, separated by a physical barrier, to keep them spatially and fluidically isolated. Selective deprivation of NGF from only the axon compartment (known as ‘local’ NGF deprivation), with cell bodies maintained in NGF, triggers the selective degeneration (pruning) of only the axons in the NGF-deprived axon compartment [43]. Although axon apoptosis and axon pruning are both induced by NGF deprivation, the distinction between global and local NGF deprivation is important as it defines the context, and thus, the specific pathway activated.

While multiple factors are known to mediate axon pruning *in vivo* [1], the molecular pathway of axon pruning has been best studied *in vitro* in the context of local (axon-only) NGF deprivation. Amyloid Precursor Protein (APP) and Death Receptor 6 (DR6) have been shown to regulate pruning *in vivo* [45*,46]. However, loss of either APP or DR6 was not protective against apoptosis (global NGF deprivation) *in vitro* [46], suggesting that APP and DR6 may play a role in axon pruning but are dispensable for apoptosis. During local NGF deprivation-mediated pruning, the loss of TrkA activity leads to the activation of DLK to induce a retrograde signal to the soma [33**]. This signal is important, as DLK-deficient neurons are protected from axon pruning [33**,40*]. DLK activation leads to the local phosphorylation of JNKs to propagate the pruning signal [40*,47*]. Interestingly, inhibition of JNK (and also p38 MAPK) in only the axons, and not the cell bodies, is sufficient to protect against axon pruning, confirming a function of JNK in axons during pruning [47*]. One of the functions of JNK in axons could be to mediate the transmission of the retrograde signal to soma. Interestingly, GSK3 (both α and β) could be a potential JNK-regulated mediator of this retrograde signal as inhibition of GSK3 activity in the soma, but not the axons, protects against axon pruning [47*]. While GSK3 could be activated locally in axons, its kinase activity may be required only in the soma where its substrates important for pruning could be spatially localized. c-Jun has been identified as another soma-dependent transcriptional regulator that is activated during pruning [40*,48]. Whether c-Jun is required for axon pruning is undetermined however, as previous results examined the requirement of c-Jun in the context of axon apoptosis (global NGF deprivation) [33**,40*].

Recently, the BH3-only protein Puma was identified as an essential mediator for axon pruning where its deficiency protects axons against local NGF deprivation [33**]. As Bax is also required for axon pruning [45*,49**,50*], the role of Puma is likely to directly activate Bax. Consistent with Bax activation, cyt *c* is released from mitochondria in axons undergoing pruning [49**], but whether cyt *c* is required for pruning is not known. This is particularly relevant as cyt *c* is known to activate Casp9 *via* Apaf-1 on the apoptosome. However, Apaf-1 is not required for axon pruning [49**]. While Apaf-1-deficient neurons are protected from undergoing apoptosis after global NGF deprivation [33**,37], they still undergo pruning after local NGF deprivation [49**]. Interestingly, despite not requiring Apaf-1, pruning requires both Casp9 [49**] and Casp3 [49**,51]. Thus, Casp9 appears to be activated *via* an Apaf-1-independent mechanism during pruning. This is particularly relevant in post-developmental mature neurons, which are known to shut down the apoptotic pathway by turning off Apaf-1 expression [52], yet maintain plasticity and remain competent to undergo pruning [49**]. This finding not only supports but highlights the Apaf-1-independent nature of the pruning pathway and illustrates how neurons are able to exert precise and differential control over the apoptosis and pruning pathways. Additionally, pruning is dependent on Casp6, a caspase which is not essential for apoptosis [45*,49**,51]. However, exactly how caspases are activated during pruning, or even their precise order of activation, remains unknown. As pruning allows for the selective degeneration only the targeted axon, without degeneration outside of that target region, it is likely that caspase activation during pruning is spatially restricted [53]. One such mediator that restricts caspase activity during pruning is XIAP (dIAP in *Drosophila*) [49**,54], which could serve to spatially constrain degeneration to the desired region. Indeed, XIAP-deficient neurons are

unable to spatially restrict caspase activity and inappropriately accumulate active Casp3 in their cell bodies during pruning [49**].

Mechanistic overlap in the three distinct pathways of axon degeneration

Despite the fact that the three pathways of axon degeneration that we highlight in this review are engaged in different contexts, there are intriguing overlaps in both the signaling as well as the execution of these pathways. We highlight here some of these similarities and differences, although in many instances the function of individual proteins has not been examined in all three contexts of axon degeneration (Table 1). While it is too early to tell whether the shared proteins have exactly the same functional role in all contexts of axon degeneration, future research will help clarify these roles.

An obvious difference between these axon degeneration pathways is that during axotomy-induced Wallerian degeneration, the cell body is not involved as it is physically separated from the axon undergoing degeneration. In contrast, the cell body is required, and in fact orchestrates, the degeneration programs during both apoptosis and pruning. Despite this fundamental contextual difference, key proteins essential for Wallerian degeneration (*e.g.* Wld^s, NMNATs, SARM1) appear to be involved in axon apoptosis as well. Demonstrating this overlap, neurons isolated from Wld^s mice degenerate their cell bodies but surprisingly not their axons upon global NGF deprivation [55]. Similarly, overexpression of cytosolic NMNAT1 also provides protection against axon apoptosis during global NGF deprivation [56]. Interestingly, this block appears to be in parallel to or after the point of caspase activation [56] but exactly how this protection is mediated remains to be determined. The exact importance of SARM1 for axon apoptosis is unclear as one study found SARM1 deficiency to protect axons [14] while another failed to see protection [13*] after global NGF deprivation. The ability of these proteins to protect axons but not the cell bodies in the context of apoptosis suggests that there could be mechanistically distinct pathways mediating the degeneration of axons *versus* cell bodies during neuronal apoptosis. Interestingly, all three contexts of axon degeneration mediate degeneration by engaging many of the same signaling pathways and proteins (*e.g.* various MAPKs and JNKs). While during apoptosis and pruning these signaling proteins are required for the transcriptional activation of the axon degeneration [33**,40*,47*], their function in Wallerian degeneration is downstream of SARM1 and independent of transcription as the degenerating axons are separate from the cell body [18**].

Significant mechanistic similarities exist between axon apoptosis and axon pruning. Both pathways are initiated by a common signal (NGF deprivation) and in both contexts (global or local NGF deprivation), the initiating signals appear to originate from axons [47*,48]. Also, most of the upstream signals up to the point of Bax activation appear to be identical in both contexts. This includes DLK, JNKs, p38 MAPK, GSK3, the BH3-only protein Puma, and Bax itself [33**,40*,45*,47*,49**,51]. However, downstream of Bax, these two pathways diverge. Axon apoptosis strictly requires the Apaf-1 and Casp9-dependent apoptosome pathway to activate Casp3 [33**,37]. In this context, Casp6 is cleaved but its deficiency does not block axon apoptosis [49**]. In contrast, axon pruning requires Casp6 as well as Casp9 and Casp3 [49**,51], but importantly not Apaf-1[49**]. This is surprising as there is very

limited precedence of a Casp9-dependent but Apaf-1-independent pathway for activating Casp3. Thus, many exciting details of how these two distinct pathways use similar signaling machinery yet diverge after the point of Bax activation remain to be discovered. Undoubtedly, future research will help uncover the mechanisms by which neurons exert both spatial and temporal control of these axon degeneration pathways.

Acknowledgments

Thank you to members of the Deshmukh lab for comments on the manuscript. This work was supported by grants from the National Institutes of Health (F31CA186654 to MJG; NS083384 and GM105612 to MD).

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Riccomagno MM, Kolodkin AL. Sculpting neural circuits by axon and dendrite pruning. *Annu Rev Cell Dev Biol.* 2015; 31:779–805. [PubMed: 26436703]
2. Pease SE, Segal RA. Preserve and protect: maintaining axons within functional circuits. *Trends Neurosci.* 2014; 37:572–582. [PubMed: 25167775]
3. Neukomm LJ, Freeman MR. Diverse cellular and molecular modes of axon degeneration. *Trends Cell Biol.* 2014; 24:515–523. [PubMed: 24780172]
4. Gerdts J, Summers Daniel W, Milbrandt J, DiAntonio A. Axon self-destruction: new links among SARM1, MAPKs, and NAD⁺ metabolism. *Neuron.* 2016; 89:449–460. [PubMed: 26844829]
5. Wang JT, Medress ZA, Barres BA. Axon degeneration: molecular mechanisms of a self-destruction pathway. *J Cell Biol.* 2012; 196:7–18. [PubMed: 22232700]
6. Maor-Nof M, Yaron A. Neurite pruning and neuronal cell death: spatial regulation of shared destruction programs. *Curr Opin Neurobiol.* 2013; 23:990–996. [PubMed: 23871216]
7. Conforti L, Gilley J, Coleman MP. Wallerian degeneration: an emerging axon death pathway linking injury and disease. *Nat Rev Neurosci.* 2014; 15:394–409. [PubMed: 24840802]
8. Mack TGA, Reiner M, Beirowski B, Mi W, Emanuelli M, Wagner D, Thomson D, Gillingwater T, Court F, Conforti L, et al. Wallerian degeneration of injured axons and synapses is delayed by a Ube4b/Nmnat chimeric gene. *Nat Neurosci.* 2001; 4:1199–1206. [PubMed: 11770485]
9. Wang J, Zhai Q, Chen Y, Lin E, Gu W, McBurney MW, He Z. A local mechanism mediates NAD-dependent protection of axon degeneration. *J Cell Biol.* 2005; 170:349–355. [PubMed: 16043516]
10. Araki T, Sasaki Y, Milbrandt J. Increased nuclear NAD biosynthesis and SIRT1 activation prevent axonal degeneration. *Science.* 2004; 305:1010–1013. [PubMed: 15310905]
11. Wang JT, Medress ZA, Vargas ME, Barres BA. Local axonal protection by WldS as revealed by conditional regulation of protein stability. *Proc Natl Acad Sci.* 2015; 112:10093–10100. [PubMed: 26209654]
12. Di Stefano M, Nascimento-Ferreira I, Orsomando G, Mori V, Gilley J, Brown R, Janeckova L, Vargas ME, Worrell LA, Loreto A, et al. A rise in NAD precursor nicotinamide mononucleotide (NMN) after injury promotes axon degeneration. *Cell Death Differ.* 2015; 22:731–742. [PubMed: 25323584]
- 13•. Osterloh JM, Yang J, Rooney TM, Fox AN, Adalbert R, Powell EH, Sheehan AE, Avery MA, Hackett R, Logan MA, et al. dSarm/Sarm1 is required for activation of an injury-induced axon death pathway. *Science.* 2012; 337:481–484. Identified SARM1 in *Drosophila* as a key regulator of Wallerian degeneration and demonstrated that SARM1-deficient mouse neurons are protected from axotomy-induced axon degeneration. [PubMed: 22678360]

14. Gerdt J, Summers DW, Sasaki Y, DiAntonio A, Milbrandt J. Sarm1-mediated axon degeneration requires both SAM and TIR interactions. *J Neurosci*. 2013; 33:13569–13580. [PubMed: 23946415]
- 15••. Gerdt J, Brace EJ, Sasaki Y, DiAntonio A, Milbrandt J. SARM1 activation triggers axon degeneration locally via NAD⁺ destruction. *Science*. 2015; 348:453–457. Showed that SARM1 activation, via dimerization, is sufficient to trigger axon degeneration. This degeneration is signaled by NAD⁺ depletion as increased NAD synthesis protected against SARM1-initiated axon destruction. [PubMed: 25908823]
16. Gilley J, Coleman MP. Endogenous Nmnat2 Is an essential survival factor for maintenance of healthy axons. *PLoS Biol*. 2010; 8:e1000300. [PubMed: 20126265]
17. Gilley J, Orsomando G, Nascimento-Ferreira I, Coleman Michael P. Absence of SARM1 rescues development and survival of NMNAT2-deficient axons. *Cell Rep*. 2015; 10:1974–1981. [PubMed: 25818290]
- 18••. Yang J, Wu Z, Renier N, David DJ, Uryu K, Park DS, Greer PA, Tournier C, Davis RJ, Tessier-Lavigne M. Pathological axonal death through a MAPK cascade that triggers a local energy deficit. *Cell*. 2015; 160:161–176. Shows that SARM1-mediated NAD⁺ depletion leads to the activation of a MAPK signaling cascade that regulates axon degeneration during axotomy. The depletion of MEKK4, MLK2, and DLK as well as their downstream regulators MKK4/MKK7 or JNKs all are protective against Wallerian degeneration. [PubMed: 25594179]
19. Miller BR, Press C, Daniels RW, Sasaki Y, Milbrandt J, DiAntonio A. A dual leucine kinase-dependent axon self-destruction program promotes Wallerian degeneration. *Nat Neurosci*. 2009; 12:387–389. [PubMed: 19287387]
20. Zhai Q, Wang J, Kim A, Liu Q, Watts R, Hoopfer E, Mitchison T, Luo L, He Z. Involvement of the ubiquitin-proteasome system in the early stages of wallerian degeneration. *Neuron*. 2003; 39:217–225. [PubMed: 12873380]
21. Babetto E, Beirowski B, Russler Emilie V, Milbrandt J, DiAntonio A. The Phr1 ubiquitin ligase promotes injury-induced axon self-destruction. *Cell Rep*. 2013; 3:1422–1429. [PubMed: 23665224]
22. Xiong X, Hao Y, Sun K, Li J, Li X, Mishra B, Soppina P, Wu C, Hume RI, Collins CA. The highwire ubiquitin ligase promotes axonal degeneration by tuning levels of nmnat protein. *PLoS Biol*. 2012; 10:e1001440. [PubMed: 23226106]
23. Liu Y, Wang Y, Chen Y, Li X, Yang J, Liu Y, Shen A. Spy1 protein mediates phosphorylation and degradation of SCG10 protein in axonal degeneration. *J Biol Chem*. 2015; 290:13888–13894. [PubMed: 25869138]
24. Shin JE, Miller BR, Babetto E, Cho Y, Sasaki Y, Qayum S, Russler EV, Cavalli V, Milbrandt J, DiAntonio A. SCG10 is a JNK target in the axonal degeneration pathway. *Proc Natl Acad Sci*. 2012; 109:E3696–E3705. [PubMed: 23188802]
25. Vargas ME, Yamagishi Y, Tessier-Lavigne M, Sagasti A. Live imaging of calcium dynamics during axon degeneration reveals two functionally distinct phases of calcium influx. *J Neurosci*. 2015; 35:15026–15038. [PubMed: 26558774]
26. Yang J, Weimer Robby M, Kallop D, Olsen O, Wu Z, Renier N, Uryu K, Tessier-Lavigne M. Regulation of axon degeneration after injury and in development by the endogenous calpain inhibitor calpastatin. *Neuron*. 2013; 80:1175–1189. [PubMed: 24210906]
27. Loreto A, Di Stefano M, Gering M, Conforti L. Wallerian degeneration is executed by an NMN-SARM1-dependent late Ca²⁺ influx but only modestly influenced by mitochondria. *Cell Rep*. 2015; 13:2539–2552. [PubMed: 26686637]
28. Buss RR, Sun W, Oppenheim RW. Adaptive roles of programmed cell death during nervous system development. *Annu Rev Neurosci*. 2006; 29:1–35. [PubMed: 16776578]
29. Yuan J, Yankner BA. Apoptosis in the nervous system. *Nature*. 2000; 407:802–809. [PubMed: 11048732]
30. Kristiansen M, Ham J. Programmed cell death during neuronal development: the sympathetic neuron model. *Cell Death Differ*. 2014; 21:1025–1035. [PubMed: 24769728]
31. Glebova NO, Ginty DD. Growth and survival signals controlling sympathetic nervous system development. *Annu Rev Neurosci*. 2005; 28:191–222. [PubMed: 16022594]

32. Eichler ME, Rich KM. Death of sensory ganglion neurons after acute withdrawal of nerve growth factor in dissociated cell cultures. *Brain Res.* 1989; 482:340–346. [PubMed: 2706491]
- 33•. Simon DJ, Pitts J, Hertz NT, Yang J, Yamagishi Y, Olsen O, Teši Mark M, Molina H, Tessier-Lavigne M. Axon degeneration gated by retrograde activation of somatic pro-apoptotic signaling. *Cell.* 2016; 164:1031–1045. Highlight the importance of the cell body in promoting apoptosis-induced axon degeneration in the context of NGF deprivation. Utilise a variety of genetic mouse models and inhibitors to define key elements of this pathway including TrkA and Akt signaling, JNKs, FOXO3a, Puma, Bcl-XL, Bcl-w, and others. [PubMed: 26898330]
34. Walensky LD, Gavathiotis E. BAX unleashed: the biochemical transformation of an inactive cytosolic monomer into a toxic mitochondrial pore. *Trends Biochem Sci.* 2011; 36:642–652. [PubMed: 21978892]
35. Courchesne SL, Karch C, Pazyra-Murphy MF, Segal RA. Sensory neuropathy attributable to loss of Bcl-w. *J Neurosci.* 2011; 31:1624–1634. [PubMed: 21289171]
36. Wang X. The expanding role of mitochondria in apoptosis. *Genes Dev.* 2001; 15:2922–2933. [PubMed: 11711427]
37. Wright KM, Vaughn AE, Deshmukh M. Apoptosome dependent caspase-3 activation pathway is non-redundant and necessary for apoptosis in sympathetic neurons. *Cell Death Differ.* 2007; 14:625–633. [PubMed: 16932756]
38. Potts PR, Singh S, Knezek M, Thompson CB, Deshmukh M. Critical function of endogenous XIAP in regulating caspase activation during sympathetic neuronal apoptosis. *J Cell Biol.* 2003; 163:789–799. [PubMed: 14623868]
- 39•. Unsain N, Higgins Julia M, Parker Kristen N, Johnstone Aaron D, Barker Philip A. XIAP regulates caspase activity in degenerating axons. *Cell Rep.* 2013; 4:751–763. Focuses on the importance of caspase-3 and the regulation of its activity by XIAP during apoptotic axon degeneration. Shows that NGF deprivation leads to the degradation of XIAP to permit caspase activation in axons and that XIAP-deficient neurons exhibit enhanced axon degeneration after NGF deprivation. [PubMed: 23954782]
- 40•. Ghosh AS, Wang B, Pozniak CD, Chen M, Watts RJ, Lewcock JW. DLK induces developmental neuronal degeneration via selective regulation of proapoptotic JNK activity. *J Cell Biol.* 2011; 194:751–764. Demonstrates the importance of DLK as an essential factor for axon degeneration in the context of both global and local deprivation of NGF. Also reports that inhibition of JNK in the axons, and not the soma, during local deprivation prevents nuclear accumulation of phosphorylated c-Jun. [PubMed: 21893599]
41. Wyttenbach A, Tolkovsky AM. The BH3-only protein Puma is both necessary and sufficient for neuronal apoptosis induced by DNA damage in sympathetic neurons. *J Neurochem.* 2006; 96:1213–1226. [PubMed: 16478523]
42. Luo L, O’Leary DD. Axon retraction and degeneration in development and disease. *Annu Rev Neurosci.* 2005; 28:127–156. [PubMed: 16022592]
43. Campenot RB. Local control of neurite development by nerve growth factor. *Proc Natl Acad Sci USA.* 1977; 74:4516–4519. [PubMed: 270699]
44. Taylor AM, Blurton-Jones M, Rhee SW, Cribbs DH, Cotman CW, Jeon NL. A microfluidic culture platform for CNS axonal injury, regeneration and transport. *Nat Methods.* 2005; 2:599–605. [PubMed: 16094385]
- 45•. Nikolaev A, McLaughlin T, O’Leary DD, Tessier-Lavigne M. APP binds DR6 to trigger axon pruning and neuron death via distinct caspases. *Nature.* 2009; 457:981–989. Initial finding demonstrating the importance of DR6 for axon pruning both *in vitro* and *in vivo*. Also report that axon pruning is mediated by Bax and caspase-6. [PubMed: 19225519]
46. Olsen O, Kallop DY, McLaughlin T, Huntwork-Rodriguez S, Wu Z, Duggan CD, Simon DJ, Lu Y, Easley-Neal C, Takeda K, et al. Genetic analysis reveals that amyloid precursor protein and death receptor 6 function in the same pathway to control axonal pruning independent of β -secretase. *J Neurosci.* 2014; 34:6438–6447. [PubMed: 24806670]
- 47•. Chen M, Maloney JA, Kallop DY, Atwal JK, Tam SJ, Baer K, Kissel H, Kaminker JS, Lewcock JW, Weimer RM, et al. Spatially coordinated kinase signaling regulates local axon degeneration. *J Neurosci.* 2012; 32:13439–13453. Identifies GSK3 α/β as regulators of axon pruning, both *in vitro* and *in vivo*. Inhibition of GSK3 α/β in the cell bodies alone protects axons from pruning.

Also identify *dleu2* and *tbx6* as two GSK3 α/β -regulated genes upregulated during pruning. [PubMed: 23015435]

48. Mok SA, Lund K, Campenot RB. A retrograde apoptotic signal originating in NGF-deprived distal axons of rat sympathetic neurons in compartmented cultures. *Cell Res.* 2009; 19:546–560. [PubMed: 19188931]
- 49••. Cusack CL, Swahari V, Hampton Henley W, Michael Ramsey J, Deshmukh M. Distinct pathways mediate axon degeneration during apoptosis and axon-specific pruning. *Nat Commun.* 2013; 4:1876. Defines that axon degeneration in the context of apoptosis and pruning are mediated by distinct pathways While Bax, caspase-9, and caspase-3 are common to both pathways, caspase-6 is essential for pruning but not apoptosis. Also demonstrates that Apaf-1, which is required for apoptosis, is not essential for pruning. Shows that XIAP is important for the spatial restriction of caspase activation to axons during pruning. Also demonstrates that mature neurons are able to restrict apoptosis but remain permissive for pruning. [PubMed: 23695670]
- 50•. Schoenmann Z, Assa-Kunik E, Tiomny S, Minis A, Haklai-Topper L, Arama E, Yaron A. Axonal degeneration is regulated by the apoptotic machinery or a NAD⁺-sensitive pathway in insects and mammals. *J Neurosci.* 2010; 30:6375–6386. Finds that both Caspase 6 and Caspase 3 are activated in axons and that XIAP is degraded in axons in response to global NGF deprivation. Found that combined treatment of neurons with NAD⁺ and caspase inhibitors is significantly more protective than either alone during axon apoptosis induced by global NGF deprivation. [PubMed: 20445064]
51. Simon DJ, Weimer RM, McLaughlin T, Kallop D, Stanger K, Yang J, O’Leary DD, Hannoush RN, Tessier-Lavigne M. A caspase cascade regulating developmental axon degeneration. *J Neurosci.* 2012; 32:17540–17553. [PubMed: 23223278]
52. Wright KM, Smith MI, Farrag L, Deshmukh M. Chromatin modification of Apaf-1 restricts the apoptotic pathway in mature neurons. *J Cell Biol.* 2007; 179:825–832. [PubMed: 18056406]
53. Unsain N, Barker Philip A. New views on the misconstrued: executioner caspases and their diverse non-apoptotic roles. *Neuron.* 2015; 88:461–474. [PubMed: 26539888]
54. Kuo CT, Zhu S, Younger S, Jan LY, Jan YN. Identification of E2/ E3 ubiquitinating enzymes and caspase activity regulating *Drosophila* sensory neuron dendrite pruning. *Neuron.* 2006; 51:283–290. [PubMed: 16880123]
55. Deckwerth TL, Johnson EM Jr. Neurites can remain viable after destruction of the neuronal soma by programmed cell death (apoptosis). *Dev Biol.* 1994; 165:63–72. [PubMed: 8088451]
56. Vohra BPS, Sasaki Y, Miller BR, Chang J, DiAntonio A, Milbrandt J. Amyloid precursor protein cleavage-dependent and -independent axonal degeneration programs share a common nicotinamide mononucleotide adenylyltransferase 1-sensitive pathway. *J Neurosci.* 2010; 30:13729–13738. [PubMed: 20943913]
57. Glass JD, Brushart TM, George EB, Griffin JW. Prolonged survival of transected nerve fibres in C57BL/Ola mice is an intrinsic characteristic of the axon. *J Neurocytol.* 1993; 22:311–321. [PubMed: 8315413]
58. Sasaki Y, Araki T, Milbrandt J. Stimulation of nicotinamide adenine dinucleotide biosynthetic pathways delays axonal degeneration after axotomy. *J Neurosci.* 2006; 26:8484–8491. [PubMed: 16914673]
59. Watkins TA, Wang B, Huntwork-Rodriguez S, Yang J, Jiang Z, Eastham-Anderson J, Modrusan Z, Kaminker JS, Tessier-Lavigne M, Lewcock JW. DLK initiates a transcriptional program that couples apoptotic and regenerative responses to axonal injury. *Proc Natl Acad Sci.* 2013; 110:4039–4044. [PubMed: 23431164]
60. Xu Z, Maroney AC, Dobrzanski P, Kukekov NV, Greene LA. The MLK family mediates c-Jun N-terminal kinase activation in neuronal apoptosis. *Mol Cell Biol.* 2001; 21:4713–4724. [PubMed: 11416147]
61. Ma M, Ferguson TA, Schoch KM, Li J, Qian Y, Shofer FS, Saatman KE, Neumar RW. Calpains mediate axonal cytoskeleton disintegration during Wallerian degeneration. *Neurobiol Dis.* 2013; 56:34–46. [PubMed: 23542511]
62. Whitmore AV, Lindsten T, Raff MC, Thompson CB. The proapoptotic proteins Bax and Bak are not involved in Wallerian degeneration. *Cell Death Differ.* 2003; 10:260–261. [PubMed: 12700655]


63. Deckwerth TL, Elliott JL, Knudson CM, Johnson EM Jr, Snider WD, Korsmeyer SJ. Bax is required for neuronal death after trophic factor deprivation and during development. *Neuron*. 1996; 17:401–411. [PubMed: 8816704]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**Figure 1.**

Axotomy activates the Wallerian degeneration pathway for the selective elimination and degeneration of severed axons. This mechanism of axon degeneration occurs in response to the physical severing of axons from their cell bodies. Wallerian degeneration occurs independently of NGF signaling and the presence of NGF is illustrated in this context only for contrast with the axon apoptosis and axon pruning pathways.




Figure 2.

Global NGF deprivation activates the apoptotic pathway to degenerate both soma and axons, in the model of sympathetic neurons. We refer to the axon degeneration that occurs during apoptosis as ‘axon apoptosis’ for clarity and to distinguish this axon degeneration from that observed during axon pruning. Here, major components of the apoptotic pathway engaged during axon apoptosis are highlighted for clarity.




Figure 3.

Pruning is the selective degeneration of only the axon exposed to the pruning stimulus, leaving the cell body intact. Pruning can be studied in compartmentalized (illustrated by the gray barrier) cultures where NGF can be selectively deprived from the axons, while cell bodies are maintained in NGF. The axon pruning pathway shares many of the same machinery as the axon apoptosis pathway, but several significant differences exist which are discussed in the text and are highlighted in Table 1.

Table 1

Shared and distinct components of the axon degeneration pathways:

	Axotomy	Apoptosis	Pruning
Wld ^s	+++ [57]	Axons ++ Soma – [55]	?
NMNAT overexpression	+++ [58]	Axons ++ Soma – [56]	?
Sarm1 –/–	+++ [13*,14]	+/- [13*,14]	?
MLK-depletion (DLK, MLK, MEKK4)	+++ [18**,19,59]	+++ [40*,59,60]	+++ [40*]
Inhibition or loss of JNKs	+++ [19,33**]	+++ [19,40*]	+++ [47*]
Inhibition of Calpains	+ [26,61]	+ [26]	?
NGF/TrkA stimulation	– [33**]	+++ [33**]	+++ [33**]
Bax –/–	– [62]	+++ [63]	+++ [45,49**]
Apaf-1 –/–	?	+++ [33**,37]	– [49**]
Caspase 9 –/–	?	+++ [37]	+++ [49**]
Caspase 6 –/–	?	– [49**]	+++ [49**,51]
Caspase 3 –/–	– [51]	+++ [37]	+++ [45,49**]

Summary of the functional importance of select key proteins in mediating axon degeneration in the contexts of axotomy, apoptosis, and pruning. The degree of protection reported is represented as: +++ excellent or complete protection, ++ significant protection, + some protection, – no protection, ? not yet known. The relevant publications (not an exhaustive list) are also indicated.