



# DEVOTE 3: temporal relationships between severe hypoglycaemia, cardiovascular outcomes and mortality

Thomas R. Pieber<sup>1</sup> · Steven P. Marso<sup>2</sup> · Darren K. McGuire<sup>3</sup> · Bernard Zinman<sup>4</sup> · Neil R. Poulter<sup>5</sup> · Scott S. Emerson<sup>6</sup> · Richard E. Pratley<sup>7,8</sup> · Vincent Woo<sup>9</sup> · Simon Heller<sup>10</sup> · Martin Lange<sup>11</sup> · Kirstine Brown-Frandsen<sup>11</sup> · Alan Moses<sup>11</sup> · Jesper Barner Lekdorf<sup>11</sup> · Lucine Lehmann<sup>11</sup> · Kajsa Kvist<sup>11</sup> · John B. Buse<sup>12</sup> · on behalf of the DEVOTE Study Group

Received: 14 July 2017 / Accepted: 25 August 2017 / Published online: 15 September 2017

© The Author(s) 2017. This article is an open access publication

## Abstract

**Aims/hypothesis** The double-blind Trial Comparing Cardiovascular Safety of Insulin Degludec vs Insulin Glargine in Patients with Type 2 Diabetes at High Risk of Cardiovascular Events (DEVOTE) assessed the cardiovascular safety of insulin degludec. The incidence and rates of adjudicated severe hypoglycaemia, and all-cause mortality were also determined. This paper reports a secondary analysis investigating associations of severe hypoglycaemia with cardiovascular outcomes and mortality.

**Methods** In DEVOTE, patients with type 2 diabetes were randomised to receive either insulin degludec or insulin glargine U100 (100 units/ml) once daily (between dinner and bedtime) in an event-driven, double-blind, treat-to-target cardiovascular outcomes trial. The primary outcome was the first occurrence of an adjudicated major adverse cardiovascular event (MACE; cardiovascular death, non-fatal myocardial infarction or non-fatal stroke). Adjudicated severe

hypoglycaemia was the pre-specified secondary outcome. In the present analysis, the associations of severe hypoglycaemia with both MACE and all-cause mortality was evaluated in the pooled trial population using time-to-event analyses, with severe hypoglycaemia as a time-dependent variable and randomised treatment as a fixed factor. An investigation with interaction terms indicated that the effect of severe hypoglycaemia on the risk of MACE and all-cause mortality were the same for both treatment arms, and so the temporal association for severe hypoglycaemia with subsequent MACE and all-cause mortality is reported for the pooled population. **Results** There was a non-significant difference in the risk of MACE for individuals who had vs those who had not experienced severe hypoglycaemia during the trial (HR 1.38, 95% CI 0.96, 1.96;  $p = 0.080$ ) and therefore there was no temporal relationship between severe hypoglycaemia and MACE. There was a significantly higher risk of all-cause mortality for patients who had vs those who had not experienced severe

**Electronic supplementary material** The online version of this article (<https://doi.org/10.1007/s00125-017-4422-0>) contains peer-reviewed but unedited supplementary material, which is available to authorised users.

✉ Thomas R. Pieber  
thomas.pieber@medunigraz.at

<sup>1</sup> Department of Internal Medicine, Medical University of Graz, Auenbruggerplatz 15, A-8036 Graz, Austria

<sup>2</sup> Research Medical Center, Kansas City, MO, USA

<sup>3</sup> University of Texas Southwestern Medical Center, Dallas, TX, USA

<sup>4</sup> Lunenfeld-Tanenbaum Research Institute, Mount Sinai Hospital, University of Toronto, Toronto, ON, Canada

<sup>5</sup> Imperial Clinical Trials Unit, Imperial College London, London, UK

<sup>6</sup> University of Washington, Seattle, WA, USA

<sup>7</sup> Florida Hospital Translational Research Institute for Metabolism and Diabetes, Orlando, FL, USA

<sup>8</sup> Sanford Burnham Prebys Medical Discovery Institute, Orlando, FL, USA

<sup>9</sup> University of Manitoba, Winnipeg, MB, Canada

<sup>10</sup> Academic Unit of Diabetes, Endocrinology and Metabolism, University of Sheffield, Sheffield, UK

<sup>11</sup> Novo Nordisk A/S, Søborg, Denmark

<sup>12</sup> University of North Carolina School of Medicine, Chapel Hill, NC, USA

hypoglycaemia during the trial (HR 2.51, 95% CI 1.79, 3.50;  $p < 0.001$ ). There was a higher risk of all-cause mortality 15, 30, 60, 90, 180 and 365 days after experiencing severe hypoglycaemia compared with not experiencing severe hypoglycaemia in the same time interval. The association between severe hypoglycaemia and all-cause mortality was maintained after adjustment for the following baseline characteristics: age, sex, HbA<sub>1c</sub>, BMI, diabetes duration, insulin regimen, hepatic impairment, renal status and cardiovascular risk group.

**Conclusions/interpretation** The results from these analyses demonstrate an association between severe hypoglycaemia and all-cause mortality. Furthermore, they indicate that patients who experienced severe hypoglycaemia were particularly at greater risk of death in the short term after the hypoglycaemic episode. These findings indicate that severe hypoglycaemia is associated with higher subsequent mortality; however, they cannot answer the question as to whether severe hypoglycaemia serves as a risk marker for adverse outcomes or whether there is a direct causal effect.

**Trial registration** [ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT01959529) NCT01959529

**Keywords** Hypoglycaemia · Insulin therapy · Macrovascular disease

### Abbreviations

ACCORD	Action to Control Cardiovascular Risk in Diabetes
DEVOTE	Trial Comparing Cardiovascular Safety of Insulin Degludec vs Insulin Glargine in Patients with Type 2 Diabetes at High Risk of Cardiovascular Events
LEADER	Liraglutide Effect and Action in Diabetes: Evaluation of Cardiovascular Outcome Results
MACE	Major adverse cardiovascular event

### Introduction

People with diabetes are at an increased risk of cardiovascular disease and cardiovascular-related death compared with those without diabetes [1]. Hypoglycaemic events, particularly when severe, have been linked to subsequent adverse cardiovascular outcomes and mortality in individuals with diabetes, although it is currently unknown whether this link is causal, predictive of greater vulnerability or both [2–4].

The Action to Control Cardiovascular Risk in Diabetes (ACCORD) trial was the first large-scale diabetes trial to report that intensive blood glucose management to normalise blood glucose levels (HbA<sub>1c</sub> target < 6% [42 mmol/mol]) was associated with a significant increase in the risk of cardiovascular-specific mortality, a factor that led to the early termination of the trial [5]. Hypoglycaemia was suggested as a

possible mechanism for the increased number of fatal events in the intensive treatment arm of ACCORD, although this association was not clearly demonstrated [6]. However, the ACCORD trial did describe a significantly increased risk of a fatal event after a severe hypoglycaemic event, in both the standard and the intensive treatment arms [6]. Furthermore, a subsequent meta-analysis of several clinical trials and observational studies suggested that severe hypoglycaemia was associated with a higher risk of cardiovascular events [7, 8]. Conclusive evidence of a direct causal relationship between hypoglycaemia and cardiovascular events and mortality is lacking, but experimental evidence in adults without diabetes suggests that hypoglycaemia-induced abnormalities of cardiac repolarisation may contribute to the risk of sudden death [9].

Insulin degludec is a once-daily basal insulin with an ultra-long duration of action, approved for use in adults, adolescents and children with either type 1 or type 2 diabetes [10, 11]. A major clinical benefit of insulin degludec is that it significantly lowers the risk of hypoglycaemia compared with insulin glargine U100 (100 units/ml) [12–15].

The double-blind Trial Comparing Cardiovascular Safety of Insulin Degludec vs Insulin Glargine in Patients with Type 2 Diabetes at High Risk of Cardiovascular Events (DEVOTE) was initiated to assess the cardiovascular safety of insulin degludec compared with insulin glargine U100. DEVOTE demonstrated that in a treat-to-target trial design, insulin degludec was non-inferior to insulin glargine in terms of cardiovascular events and superior with regard to hypoglycaemia risk, with lower rates of both severe and nocturnal severe hypoglycaemia at equivalent glycaemic control [15]. Because of the size and design of the trial and the relatively large number of episodes of severe hypoglycaemia, DEVOTE provides a valuable opportunity to explore the associations of severe hypoglycaemia with cardiovascular outcomes and mortality.

### Methods

The detailed methods of the trial, the trial protocol, the statistical analysis plan and the list of members of the trial teams and committees have been published previously [15, 16]. DEVOTE is registered at [ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT01959529) (number NCT01959529). The trial was conducted in accordance with the Declaration of Helsinki and the ICH Good Clinical Practice Guideline [17, 18]. The protocol was approved by independent ethics committees or institutional review boards for each centre; written informed consent was obtained from each participant before any trial-related activities.

In brief, DEVOTE was a multicentre, prospective, treat-to-target, randomised, double-blind, active comparator cardiovascular outcomes trial, designed to continue until at least 633 major adverse cardiovascular events (MACEs), confirmed by a central, blinded, independent Event Adjudication Committee

(EAC), had accrued [15, 16]. All participants had type 2 diabetes treated with at least one oral or injectable glucose-lowering agent with  $\text{HbA}_{1c} \geq 7.0\%$  (53 mmol/mol), or with  $\geq 20$  units/day of basal insulin. Patients were eligible for the trial if they either had at least one co-existing cardiovascular or renal condition and were aged  $\geq 50$  years or had at least one of a list of pre-specified cardiovascular risk factors and were aged  $\geq 60$  years. Patients were not excluded if they had experienced severe hypoglycaemia prior to randomisation.

Patients with type 2 diabetes at high risk of cardiovascular events were randomised 1:1 to receive either insulin degludec (Novo Nordisk, Bagsværd, Denmark) or insulin glargine (Sanofi, Paris, France), in a blinded fashion, both in identical 100 U/ml, 10 ml vials, administered once daily between dinner and bedtime, in addition to standard care. All patients were allowed to continue their pre-trial glucose-lowering therapy, with the exception of basal and premix insulins, which were discontinued.

The primary adjudicated composite endpoint of DEVOTE was the time from randomisation to the first occurrence of death from cardiovascular causes, non-fatal myocardial infarction or non-fatal stroke. Secondary outcomes included an expanded composite cardiovascular outcome (the primary composite endpoint plus adjudicated unstable angina leading to hospitalisation) and time from randomisation to death from any cause. Adjudication-confirmed severe hypoglycaemia was a pre-specified, multiplicity-adjusted secondary outcome, as defined by the ADA as an episode requiring the assistance of another person to actively administer carbohydrate or glucagon, or to take other corrective actions. Plasma glucose levels may not be available during an event, but neurological recovery after the return of plasma glucose to a normal level is considered sufficient evidence that the event was induced by a low plasma glucose level [19].

In the present analysis, the association between severe hypoglycaemia and either MACE or all-cause mortality was investigated by comparing the risk of an event with or without having experienced severe hypoglycaemia in different time periods (15, 30, 60, 90, 180 and 365 days) prior to the event. Cox regression models were used to analyse these associations for each time period. The indicator of whether a severe hypoglycaemic event had occurred was included in the model as a time-dependent variable. All episodes of severe hypoglycaemia prior to first MACE or all-cause mortality were included in the analysis. Randomised treatment was also included in the model as a fixed factor. For sensitivity analyses, additional baseline information (age, sex,  $\text{HbA}_{1c}$ , BMI, diabetes duration, insulin regimen, hepatic impairment, renal status and cardiovascular risk group inclusion criteria) was accounted for. An investigation with interaction terms indicated that the effect of severe hypoglycaemia on the risk of MACE and all-cause mortality were the same for both treatment arms (insulin degludec and insulin glargine), and so the

temporal association for severe hypoglycaemia with subsequent MACE and all-cause mortality is reported for the pooled population. For MACEs occurring on the same day as a severe hypoglycaemic event, 0.5 days was added to the day of the MACE. All analyses were conducted using SAS, version 9.4 ([https://www.sas.com/en\\_ca/software/sas9.html](https://www.sas.com/en_ca/software/sas9.html)). A  $p$  value of  $< 0.05$  was considered statistically significant.

## Results

**Overall results from DEVOTE** Detailed results from DEVOTE have been published previously [15]. To summarise, a total of 7637 patients were randomised to either insulin degludec ( $n = 3818$ ) or insulin glargine ( $n = 3819$ ). Of these, 98% completed the final follow-up visit or died during the trial. Vital status was known for 99.9% of participants. The median observation time was 1.99 years in both treatment arms.

The pre-specified analysis demonstrated that insulin degludec was non-inferior to insulin glargine in terms of cardiovascular events (HR 0.91, 95% CI 0.78, 1.06;  $p < 0.001$  for non-inferiority), and superior with regard to hypoglycaemia risk, with a lower rate of both severe and nocturnal severe hypoglycaemia (by 40% and 53%, respectively; both  $p < 0.001$ ) [15].

### Severe hypoglycaemia and its association with cardiovascular outcomes and all-cause mortality (secondary analysis)

Of the 681 patients who experienced a MACE and the 439 patients who experienced a severe hypoglycaemic event during the trial, 32 patients had a severe hypoglycaemic event prior to a MACE (14 patients treated with insulin degludec and 18 patients treated with insulin glargine) and 16 patients experienced a MACE prior to severe hypoglycaemia (Table 1 and Electronic supplementary material [ESM] Fig. 1). Of the 423 patients who died from any cause, 38 patients died after experiencing a severe hypoglycaemic event (Table 1). The baseline characteristics of participants who experienced severe hypoglycaemia were not different from those who did not experience severe hypoglycaemia during the trial (ESM Table 1). There was no between-treatment difference in terms of risk of MACE ( $p = 0.679$ ) or all-cause mortality ( $p = 0.209$ ) following severe hypoglycaemia. On this basis, the association between severe hypoglycaemia and time to first MACE or all-cause mortality is reported for the pooled population.

In the pooled population, there was a non-significant difference in the risk of MACE between participants who had and those who had not experienced severe hypoglycaemia during the trial (HR 1.38, 95% CI 0.96, 1.96;  $p = 0.080$ ) (Fig. 1). A similar result was observed for the expanded four-point MACE definition (HR 1.37, 95% CI 0.99, 1.91;  $p = 0.060$ ) (Fig. 1). When the individual components of the three-point and four-point MACE were investigated, there was a significantly higher risk of cardiovascular death at any

**Table 1** Overview of outcomes (pooled data)

Outcome	Number of patients	Rate (events/100 PYO)
MACE	681	4.50
Non-fatal myocardial infarction	313	2.07
Non-fatal stroke	150	0.99
Cardiovascular death	278	1.84
Unstable angina requiring hospitalisation	145	0.96
Severe hypoglycaemia	439	4.97
≥ 1 severe hypoglycaemic events prior to MACE	32	6.34
≥ 2 severe hypoglycaemic events prior to MACE	6	4.35
MACE prior to severe hypoglycaemia	16	–
All-cause mortality	423	2.80
Severe hypoglycaemia prior to all-cause mortality	38	7.32

PYO, patient-years of observation


time following a severe hypoglycaemic event vs not experiencing a severe hypoglycaemic event (HR 2.14, 95% CI 1.37, 3.35;  $p < 0.001$ ), whereas there was not a significantly higher risk of non-fatal myocardial infarction (HR 0.74, 95% CI 0.36, 1.49;  $p = 0.395$ ), non-fatal stroke (HR 1.81, 95% CI 0.92, 3.57;  $p = 0.085$ ) or unstable angina requiring hospitalisation (HR 1.34, 95% CI 0.59, 3.04,  $p = 0.490$ ) (Fig. 1). When dividing the time period following a severe hypoglycaemic event into time intervals of different durations (15, 30, 60, 90, 180 and 365 days), there was no temporal relationship between severe hypoglycaemia and MACE (Fig. 2). The non-significant relationship between severe hypoglycaemia and MACE was maintained after adjustment for the following baseline characteristics: age, sex, HbA<sub>1c</sub>, BMI, diabetes duration, insulin regimen, hepatic impairment, renal status and cardiovascular risk group inclusion criteria (ESM Table 2). Only a small number of participants

experienced a MACE prior to severe hypoglycaemia ( $n = 16$ ) (Table 1), and therefore a possible opposite temporal association could not be analysed. Only three nocturnal severe hypoglycaemic events occurred prior to a MACE and therefore these could also not be analysed separately.


In the pooled population, experiencing severe hypoglycaemia at any time was a significant predictor of all-cause mortality, as the risk for these participants was 2.5-fold that for the participants who did not experience an event (HR 2.51, 95% CI 1.79, 3.50;  $p < 0.001$ ) (Fig. 3). When dividing the time period following severe hypoglycaemia into time intervals of different durations, there was a higher risk of all-cause mortality 15, 30, 60, 90, 180 and 365 days after experiencing severe hypoglycaemia compared with not experiencing severe hypoglycaemia in the same time interval (Fig. 3). The risk appeared to be highest in the shorter-term windows and decreased with the longer-term windows, but remained significant for all ( $p < 0.05$  for all). The relationship between severe hypoglycaemia and all-cause mortality was maintained after adjustment for the following baseline characteristics: age, sex, HbA<sub>1c</sub>, BMI, diabetes duration, insulin regimen, hepatic impairment, renal status and cardiovascular risk group inclusion criteria (ESM Table 2). The cause of each death occurring after a severe hypoglycaemic event is listed in ESM Table 3, including the days the severe hypoglycaemic events occurred and the time between the last event and the fatal event. The majority of these fatal events were ascribed to non-cardiovascular reasons ( $n = 17$ ) and the remainder to cardiovascular ( $n = 14$ ) or undetermined causes ( $n = 7$ ).

**Discussion**

The results of DEVOTE demonstrated that insulin degludec was superior with regard to severe hypoglycaemia risk at equivalent glycaemic control compared with insulin glargine, thereby confirming observations from earlier studies including the open-label Phase 3a programme and the double-blind,



**Fig. 1** Risk of MACE following a severe hypoglycaemic event. Cardiovascular death includes patients with an unknown cause of death. MI, myocardial infarction;  $n$ , number of patients; rate, events per 100 patient-years of observation




**Fig. 2** Risk of MACE following a severe hypoglycaemic event by time period. *n*, number of patients; rate, events per 100 patient-years of observation

crossover SWITCH trial in patients with type 2 diabetes [12, 13, 15]. The results from the new analyses reported here demonstrate an association between severe hypoglycaemic events and a higher risk of all-cause mortality ( $p < 0.001$ ) in the overall DEVOTE population. In addition, the DEVOTE data suggest that an elevated risk of a fatal event might persist for many weeks and months after a severe hypoglycaemic event, although the highest risk appears to be in the shorter time periods, albeit with small numbers of events. The null hypothesis for the temporal analyses was that there would be an increased risk of an event immediately following severe hypoglycaemia. On this basis we would therefore expect that the ‘any time’ hazard ratio would be lower than those for the shorter time periods, which is what was observed for all-cause mortality. These results are similar to those observed in the Liraglutide Effect and Action in Diabetes: Evaluation of Cardiovascular Outcome Results (LEADER) trial, where a similar temporal analysis was conducted [20].

Hypoglycaemia can have a number of adverse cardiovascular pathological effects in addition to the symptomatic effects

experienced by patients. In healthy people, hypoglycaemia triggers a counter-regulatory response with a subsequent increase in catecholamine levels [21–24]. This response induces an increase in myocardial contractility and cardiac output, with a corresponding increase in elasticity of blood vessels and a reduction in central arterial pressure [25–27]. People with a long duration of diabetes may have arterial stiffness or underlying cardiac disease that can compromise the benefits of this counter-regulatory response and lead to adverse outcomes. In an observational study investigating the risk of arrhythmias during spontaneous hypoglycaemia in people with type 2 diabetes at high risk of cardiovascular events, clinically important hypoglycaemia ( $< 3.5$  mmol/l) was common, occurring 6% of the time [28]. These hypoglycaemic events were associated with ECG changes consistent with ischaemia, prolonged QT intervals, repolarisation defects and various cardiac arrhythmias, suggesting that these events could be interconnected. Bradycardia and atrial and ventricular ectopic counts were also significantly higher during episodes of nocturnal vs daytime hypoglycaemia [28]. Furthermore, animal studies have shown that the counter-



**Fig. 3** Risk of all-cause death following a severe hypoglycaemic event by time period. *n*, number of patients; rate, events per 100 patient-years of observation

regulatory sympathoadrenal response is capable of inducing fatal cardiac arrhythmias during severe events [29].

Hypoglycaemia has also been linked to both prothrombotic and proinflammatory effects, which could alter blood flow, increasing the risk of cardiovascular events [21, 30]. The catecholamines released during the counter-regulatory response, as well as the release of coagulation factors and inflammatory cytokines into the circulation, can all increase blood viscosity and promote platelet aggregation and activation, thereby affecting vascular flow [22, 29, 31, 32]. These proinflammatory and procoagulant factors may remain elevated for several days after the hypoglycaemia has resolved, leaving the person vulnerable for some time after the hypoglycaemic event, and potentially contributing to the occurrence of major vascular events [33, 34]. In a clinical setting, analyses from the Veterans Affairs Diabetes Trial (VADT) also demonstrated that a severe hypoglycaemic event was an independent predictor of death at 90 days [19, 35].

There remains considerable controversy over the potential causal relationship between severe hypoglycaemia and adverse cardiovascular events. Several clinical outcome trials and observational studies have shown such an association [23]. In ACCORD, patients who had one or more severe hypoglycaemic episodes had higher rates of death than those who did not experience such episodes [6]. In the Outcome Reduction with Initial Glargine Intervention (ORIGIN) trial, severe hypoglycaemia increased the risk of arrhythmic death (by 77%), all-cause death (by 74%) and cardiovascular death (by 71%) [36]. In addition, a post hoc analysis of the Action in Diabetes and Vascular Disease: PreterAx and Diamicron MR Controlled Evaluation (ADVANCE) trial suggested that severe hypoglycaemia was associated with significantly higher risks for major macrovascular events, cardiovascular death and all-cause death [37]. The major limitation, however, is the inability to definitively attribute the cause of death or cardiovascular event to hypoglycaemia and to determine the exact temporal relationship between hypoglycaemia and a subsequent vascular outcome. A recent analysis of The Examination of Cardiovascular Outcomes with Alogliptin Versus Standard of Care (EXAMINE) trial demonstrated that the relationship between severe hypoglycaemia and MACE was less strong for events occurring after a severe hypoglycaemic event compared with all events and severe hypoglycaemia, suggesting that confounding by comorbidities and hypoglycaemia is important [38]. Indeed, in DEVOTE, the trial population was at particularly high risk of MACE and fatal events, with a long duration of diabetes (> 16 years) and previous insulin use (85%), and the majority of participants with established cardiovascular or chronic kidney disease (85%) [15]. This is in line with other studies that have reported that these patient characteristics were more common amongst those who died following a severe hypoglycaemic event [39]. Similarly, results from a recent post hoc analysis of the LEADER trial data have demonstrated a significantly higher risk

of MACE, cardiovascular death and non-cardiovascular death following a severe hypoglycaemic event, particularly in the short term. Those patients who experienced severe hypoglycaemia had a longer duration of diabetes, greater insulin use and a higher incidence of heart failure and kidney disease at baseline [20]. However, confounding by comorbidities does not appear to account for the association alone. A meta-analysis of several clinical trials and observational studies, including over 900,000 patients with type 2 diabetes, observed that severe hypoglycaemia was associated with a higher risk of cardiovascular disease, but that this was unlikely to be entirely a consequence of confounding by comorbid severe illness [7]. In addition, as described above, there is also considerable experimental evidence that hypoglycaemia can lead to arrhythmias and other adverse cardiovascular pathological effects. Overall, it is most likely that hypoglycaemia is just a single contributory factor to cardiovascular events in a much larger multifactorial landscape.

The analyses reported here have several limitations. DEVOTE was designed to collect only severe hypoglycaemic events and therefore the contribution of non-severe events could not be assessed. There are several studies that show that both non-severe and severe hypoglycaemic events are associated with a higher risk of cardiovascular events, hospitalisation and all-cause mortality [40, 41]. Although DEVOTE did not collect non-severe events, the severe events collected were independently adjudicated and provide an accurate view of these events in an at-risk patient population. In addition, while the overall DEVOTE trial population was large, only a small proportion of patients experienced severe hypoglycaemia prior to a MACE or a fatal event, particularly during the shorter time periods (15–60 days), which limits the statistical power of our time-to-event analysis. Furthermore, history of a patient's previous experience of severe hypoglycaemia prior to the trial was not collected, therefore the contribution of these events to the risk of MACE or all-cause mortality could not be assessed. In addition, it was observed from our analyses and those of other trials that the association between a severe hypoglycaemic event and a higher risk of a fatal event lasts for at least 1 year [20]. It is therefore possible that with a population that mostly used insulin prior to randomisation, confounding from severe hypoglycaemic events prior to trial initiation could have occurred. However, it is also important to note that approximately 30% of the DEVOTE population were on basal insulin alone and 15% were insulin-naïve at baseline, and therefore the relative risk of severe hypoglycaemia-induced cardiovascular events in these populations is unlikely to be very high compared with the risk for patients treated with a basal–bolus regimen. Finally, the incorporation of a continuous glucose monitoring element in future trials may also be warranted to provide further information on blood glucose levels at the time of severe hypoglycaemic events as well as up to the time of a MACE.

These analyses also have several strengths. DEVOTE is the first cardiovascular outcomes trial to compare the cardiovascular

safety of two basal insulins in a double-blind fashion. Many previous cardiovascular outcome trials have been limited in their potential to explore the relationship between hypoglycaemia and MACE because the therapies used in the two treatment arms were different and hence might have had differing pharmacological influences on the risk of MACE. DEVOTE, a trial comparing two basal insulins, allowed the effect of hypoglycaemia alone to be explored more precisely. Furthermore, the independent adjudication of both severe hypoglycaemic events and MACE provided additional strength to the analyses.

The results from these analyses add to the evidence for an association between severe hypoglycaemia and mortality. However, they do not establish whether hypoglycaemia serves as a risk marker for these events or directly contributes to their occurrence.

**Acknowledgements** Open access funding provided by Medical University of Graz. We thank the trial investigators, trial staff and trial participants for their participation, P.-M. Haahr (Novo Nordisk A/S, Denmark) for insights that assisted in the development of this article, and F. Hemingway and R. McDonald from Watermeadow Medical (UK; sponsored by Novo Nordisk) for providing medical writing and editorial support. DEVOTE research activities were supported at numerous US centres by Clinical and Translational Science Awards from the National Institutes of Health's National Center for Advancing Translational Science.

**Data availability** The data generated during and/or analysed during the current trial are available from the corresponding author on reasonable request.

**Funding** This trial and secondary analysis was sponsored and funded by Novo Nordisk (Bagsvaerd, Denmark). JBB received support from The National Institutes of Health (UL1TR001111). The trial sponsor was involved in the design of the trial; the collection, and analysis of data; and writing the clinical report.

**Duality of interest** TRP has received research support from Novo Nordisk and AstraZeneca (paid directly to the Medical University of Graz); personal fees as a consultant from AstraZeneca, Bristol-Myers Squibb, Eli Lilly, Novo Nordisk and Roche Diabetes Care. TRP is also the Chief Scientific Officer of CBmed (Center for Biomarker Research in Medicine), a public-funded biomarker research company.

SPM has received personal fees from Abbott Vascular, Novo Nordisk, University of Oxford, AstraZeneca and Bristol-Myers Squibb; and research support from Novo Nordisk, The Medicines Company and Terumo Medical.

DKM has received personal fees from Boehringer Ingelheim, Janssen Research and Development LLC, Sanofi US, Merck Sharp and Dohme Corp., Eli Lilly USA, Novo Nordisk, GlaxoSmithKline, AstraZeneca, Lexicon Pharmaceuticals, Eisai and Esperion.

BZ has received grant support from Boehringer Ingelheim, AstraZeneca and Novo Nordisk; and consulting fees from AstraZeneca, Boehringer Ingelheim, Eli Lilly, Janssen, Merck, Novo Nordisk and Sanofi.

NRP has received personal fees from Servier, Takeda, Novo Nordisk and AstraZeneca in relation to speakers' fees and advisory board activities (concerning diabetes mellitus); and research grants for his research group (relating to type 2 diabetes mellitus) from Diabetes UK, National Institute for Health Research Efficacy and Mechanism Evaluation (NIHR EME), Julius Clinical and the British Heart Foundation.

SSE has received personal fees related to Data Monitoring Committees from CTI BioPharma, Arena Pharmaceuticals, SFJ Pharmaceuticals, BioMarin, Medivation, Biom'up, Dynavax, Genentech, GlaxoSmithKline, Janssen Research, Novartis, Novo Nordisk, Pfizer, Roche, Sarepta Therapeutics and Xoma; personal fees related to other statistical consulting from AstraZeneca, Celltrion, Sprout Pharmaceuticals, Sanofi, Collegium Pharmaceutical, Intercept, Coherus BioMedical and Emmaus Life Sciences; and research grant support from National Heart, Lung, and Blood Institute (NHLBI).

REP's services were paid directly to Florida Hospital, a non-profit organisation. Consultancy and speaker fees from AstraZeneca, Takeda and Novo Nordisk; consultancy fees from Boehringer Ingelheim, GlaxoSmithKline, Hanmi Pharmaceutical Co. Ltd., Janssen Scientific Affairs LLC, Ligand Pharmaceuticals, Inc., Eli Lilly, Merck, Pfizer, Eisai, Inc.; research grant from Gilead Sciences, Lexicon Pharmaceuticals, Ligand Pharmaceuticals, Inc., Eli Lilly, Merck, Sanofi US LLC and Takeda.

VW has received personal fees in relation to speakers' fees and advisory board activities from Novo Nordisk, Eli Lilly, Sanofi, AstraZeneca, Boehringer Ingelheim and Merck; he also reports serving as an investigator for clinical trials sponsored by Merck, Novo Nordisk, Eli Lilly, Sanofi, AstraZeneca, Boehringer Ingelheim, GlaxoSmithKline and Locemia Solutions.

SH has served on speaker panels for Sanofi, Eli Lilly, Takeda, Novo Nordisk and AstraZeneca, for which he has received remuneration. He has served on advisory panels or as a consultant for Boehringer Ingelheim, Novo Nordisk, Eli Lilly and Takeda for which his institution has received remuneration.

ML, KBF, AM, JBL, LL and KK are full-time employees of, and hold stock in, Novo Nordisk A/S.

JBB reports receiving contracted consulting fees, paid to his institution, and travel support from Novo Nordisk, Eli Lilly, GI Dynamics, Elcylex, Merck, Metavention, vTv Pharma, PhaseBio, AstraZeneca, Dance Biopharm, Sanofi, Lexicon Pharmaceuticals, Orexigen, Takeda, Adocia, Roche, NovaTarg, Shenzen HighTide, Fractyl and Dexcom; grant support from Eli Lilly, Bristol-Myers Squibb, GI Dynamics, Merck, PhaseBio, AstraZeneca, Medtronic Minimed, Sanofi, Johnson & Johnson, Andromeda, Boehringer Ingelheim, GlaxoSmithKline, MacroGenics, Intarcia Therapeutics, Lexicon Pharmaceuticals, Scion NeuroStim, Orexigen, Takeda, Theracos and Bayer; he also reports receiving fees and holding stock options in PhaseBio and Insulin Algorithms; he also reports serving on the board of the AstraZeneca Healthcare Foundation.

**Contribution statement** All authors confirm that they meet the International Committee of Medical Journal Editors uniform requirements for authorship. Specifically, all authors made substantial contributions to the interpretation of data for the manuscript, drafted and critically revised the manuscript, provided final approval of the version to be published and agreed to be accountable for all aspects of the manuscript. All authors had access to the final results and vouch for the fidelity of the trial to the protocol. Medical writing and editorial support, under the guidance of the authors, was provided by Watermeadow Medical, an Ashfield company, part of UDG Healthcare plc, funded by Novo Nordisk. All authors are responsible for the integrity of the work as a whole.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## References

1. Dailey G (2011) Overall mortality in diabetes mellitus: where do we stand today? *Diabetes Technol Ther* 13(Suppl. 1):S65–S74

2. Lung TW, Petrie D, Herman WH et al (2014) Severe hypoglycemia and mortality after cardiovascular events for type 1 diabetic patients in Sweden. *Diabetes Care* 37:2974–2981
3. Paty BW (2015) The role of hypoglycemia in cardiovascular outcomes in diabetes. *Can J Diabetes* 39(Suppl. 5):S155–S159
4. Snell-Bergeon JK, Wadwa RP (2012) Hypoglycemia, diabetes, and cardiovascular disease. *Diabetes Technol Ther* 14(Suppl. 1):S51–S58
5. Gerstein HC et al (2008) Effects of intensive glucose lowering in type 2 diabetes. *N Engl J Med* 358:2545–2559
6. Bonds DE, Miller ME, Bergenstal RM et al (2010) The association between symptomatic, severe hypoglycaemia and mortality in type 2 diabetes: retrospective epidemiological analysis of the ACCORD study. *BMJ* 340:b4909
7. Goto A, Arah OA, Goto M et al (2013) Severe hypoglycaemia and cardiovascular disease: systematic review and meta-analysis with bias analysis. *BMJ* 347:f4533
8. Skyler JS, Bergenstal R, Bonow RO et al (2009) Intensive glycemic control and the prevention of cardiovascular events: implications of the ACCORD, ADVANCE, and VA diabetes trials: a position statement of the American Diabetes Association and a scientific statement of the American College of Cardiology Foundation and the American Heart Association. *Circulation* 119:351–357
9. Robinson RT, Harris ND, Ireland RH et al (2003) Mechanisms of abnormal cardiac repolarization during insulin-induced hypoglycemia. *Diabetes* 52:1469–1474
10. Tresiba® [prescribing information] (2015) Available from [http://www.accessdata.fda.gov/drugsatfda\\_docs/label/2015/203314lbl.pdf](http://www.accessdata.fda.gov/drugsatfda_docs/label/2015/203314lbl.pdf). Accessed 5 July 2017
11. Tresiba® Summary of product characteristics (2015) Available from [http://www.ema.europa.eu/docs/en\\_GB/document\\_library/EPAR\\_-\\_Product\\_Information/human/002498/WC500138940.pdf](http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Product_Information/human/002498/WC500138940.pdf). Accessed 5 July 2017
12. Ratner RE, Gough SC, Mathieu C et al (2013) Hypoglycaemia risk with insulin degludec compared with insulin glargine in type 2 and type 1 diabetes: a pre-planned meta-analysis of phase 3 trials. *Diabetes Obes Metab* 15:175–184
13. Wysham C, Bhargava A, Chaykin L et al (2017) Effect of insulin degludec vs insulin glargine U100 on hypoglycemia in patients with type 2 diabetes; the SWITCH 2 randomized clinical trial. *JAMA* 318:45–56
14. Lane W, Bailey TS, Gerety G et al (2017) Effect of insulin degludec vs insulin glargine U100 on hypoglycemia in patients with type 1 diabetes; the SWITCH 1 randomized clinical trial. *JAMA* 318:33–44
15. Marso S, McGuire DK, Zinman B et al (2017) Efficacy and safety of degludec versus glargine in type 2 diabetes. *N Engl J Med* 377:723–732
16. Marso SP, McGuire DK, Zinman B et al (2016) Design of DEVOTE (trial comparing cardiovascular safety of insulin degludec vs insulin glargine in patients with type 2 diabetes at high risk of cardiovascular events)—DEVOTE 1. *Am Heart J* 179:175–183
17. World Medical Association (2013) Declaration of Helsinki. Ethical principles for medical research involving human subjects. *JAMA* 310:2191–2194
18. ICH (2001) ICH harmonised tripartite guideline: guideline for good clinical practice. *J Postgrad Med* 47:199–203
19. Seaquist ER, Anderson J, Childs B et al (2013) Hypoglycemia and diabetes: a report of a workgroup of the American Diabetes Association and the Endocrine Society. *Diabetes Care* 36:1384–1395
20. Zinman B, Marso SP, Christiansen E, Calanna S, Rasmussen S, Buse JB (2017) Severe hypoglycemia, cardiovascular outcomes and death—the LEADER experience. *Diabetes* 66(Suppl. 1):A95 Abstract
21. Hanefeld M, Frier BM, Pistrosch F (2016) Hypoglycemia and cardiovascular risk: is there a major link? *Diabetes Care* 39(Suppl. 2):S205–S209
22. Wright RJ, Frier BM (2008) Vascular disease and diabetes: is hypoglycaemia an aggravating factor? *Diabetes Metab Res Rev* 24:353–363
23. Frier BM, Schernthaner G, Heller SR (2011) Hypoglycemia and cardiovascular risks. *Diabetes Care* 34(Suppl. 2):S132–S137
24. Desouza CV, Bolli GB, Fonseca V (2010) Hypoglycemia, diabetes, and cardiovascular events. *Diabetes Care* 33:1389–1394
25. Fisher BM, Gillen G, Hepburn DA, Dargie HJ, Frier BM (1990) Cardiac responses to acute insulin-induced hypoglycemia in humans. *Am J Phys* 258:H1775–H1779
26. Hilsted J, Bonde-Petersen F, Nørgaard MB et al (1984) Haemodynamic changes in insulin-induced hypoglycaemia in normal man. *Diabetologia* 26:328–332
27. Sommerfield AJ, Wilkinson IB, Webb DJ, Frier BM (2007) Vessel wall stiffness in type 1 diabetes and the central hemodynamic effects of acute hypoglycemia. *Am J Physiol Endocrinol Metab* 293:E1274–E1279
28. Chow E, Bernjak A, Williams S et al (2014) Risk of cardiac arrhythmias during hypoglycemia in patients with type 2 diabetes and cardiovascular risk. *Diabetes* 63:1738–1747
29. Reno CM, Daphna-Iken D, Chen YS, VanderWeele J, Jethi K, Fisher SJ (2013) Severe hypoglycemia-induced lethal cardiac arrhythmias are mediated by sympathoadrenal activation. *Diabetes* 62:3570–3581
30. Dandona P, Chaudhuri A, Dhindsa S (2010) Proinflammatory and prothrombotic effects of hypoglycemia. *Diabetes Care* 33:1686–1687
31. Wright RJ, Newby DE, Stirling D, Ludlam CA, Macdonald IA, Frier BM (2010) Effects of acute insulin-induced hypoglycemia on indices of inflammation: putative mechanism for aggravating vascular disease in diabetes. *Diabetes Care* 33:1591–1597
32. Gogitidze Joy N, Hedrington MS, Briscoe VJ, Tate DB, Ertl AC, Davis SN (2010) Effects of acute hypoglycemia on inflammatory and pro-atherothrombotic biomarkers in individuals with type 1 diabetes and healthy individuals. *Diabetes Care* 33:1529–1535
33. Hutton RA, Mikhailidis D, Dormandy KM, Ginsburg J (1979) Platelet aggregation studies during transient hypoglycaemia: a potential method for evaluating platelet function. *J Clin Pathol* 32:434–438
34. Pistrosch F, Hanefeld M (2015) Hypoglycemia and cardiovascular disease: lessons from outcome studies. *Curr Diab Rep* 15:117
35. Duckworth W, Abraira C, Moritz T et al (2009) Glucose control and vascular complications in veterans with type 2 diabetes. *N Engl J Med* 360:129–139
36. Mellbin LG, Rydén L, Riddle MC et al (2013) Does hypoglycaemia increase the risk of cardiovascular events? A report from the ORIGIN trial. *Eur Heart J* 34:3137–3144
37. Zoungas S, Patel A, Chalmers J et al (2010) Severe hypoglycemia and risks of vascular events and death. *N Engl J Med* 363:1410–1418
38. Heller SR, Bergenstal RM, White WB et al (2017) Relationship of glycated haemoglobin and reported hypoglycaemia to cardiovascular outcomes in patients with type 2 diabetes and recent acute coronary syndrome events: the EXAMINE trial. *Diabetes Obes Metab* 19:664–671
39. Cha SA, Yun JS, Lim TS et al (2016) Severe hypoglycemia and cardiovascular or all-cause mortality in patients with type 2 diabetes. *Diabetes Metab J* 40:202–210
40. Hsu PF, Sung SH, Cheng HM et al (2013) Association of clinical symptomatic hypoglycemia with cardiovascular events and total mortality in type 2 diabetes: a nationwide population-based study. *Diabetes Care* 36:894–900
41. Finfer S, Liu B, Myburgh JA et al (2012) Hypoglycemia and risk of death in critically ill patients. *N Engl J Med* 367:1108–1118