

Newborn Sequencing in Genomic Medicine and Public Health

Jonathan S. Berg, MD, PhD,^a Pankaj B. Agrawal, MD, MMSc,^{b,c} Donald B. Bailey Jr., PhD,^d Alan H. Beggs, PhD,^e Steven E. Brenner, PhD,^e Amy M. Brower, PhD,^f Julie A. Cakici, BA, BSN,^g Ozge Ceyhan-Birsoy, PhD,^h Kee Chan, PhD,ⁱ Flavia Chen, MPH,^j Robert J. Currier, PhD,^k Dmitry Dukhovny, MD, MPH,^l Robert C. Green, MD, MPH,^m Julie Harris-Wai, MPH, PhD,^{j,n} Ingrid A. Holm, MD, MPH,^o Brenda Iglesias,^o Galen Joseph, PhD,^p Stephen F. Kingsmore, MD, DSc,^g Barbara A. Koenig, PhD,^o Pui-Yan Kwok, MD, PhD,^{j,q} John Lantos, MD,^r Steven J. Leeder, PharmD, PhD,^r Megan A. Lewis, PhD,^d Amy L. McGuire, JD, PhD,^s Laura V. Milko, PhD,^a Sean D. Mooney, PhD,^t Richard B. Parad, MD, MPH,^u Stacey Pereira, PhD,^s Joshua Petrikin, MD,^r Bradford C. Powell, MD, PhD,^a Cynthia M. Powell, MD,^v Jennifer M. Puck, MD,^w Heidi L. Rehm, PhD,^h Neil Risch, PhD,^j Myra Roche, MS,^y Joseph T. Shieh, MD, PhD,^{j,x} Narayanan Veeraraghavan, PhD,^g Michael S. Watson, MS, PhD,^f Laurel Willig, MD, MS,^r Timothy W. Yu, MD, PhD,^o Tiina Urv, PhD,^y Anastasia L. Wise, PhD^o

The rapid development of genomic sequencing technologies has decreased the cost of genetic analysis to the extent that it seems plausible that genome-scale sequencing could have widespread availability in pediatric care. Genomic sequencing provides a powerful diagnostic modality for patients who manifest symptoms of monogenic disease and an opportunity to detect health conditions before their development. However, many technical, clinical, ethical, and societal challenges should be addressed before such technology is widely deployed in pediatric practice. This article provides an overview of the Newborn Sequencing in Genomic Medicine and Public Health Consortium, which is investigating the application of genome-scale sequencing in newborns for both diagnosis and screening.

Universal newborn screening (NBS) is an extraordinarily successful public health program, preventing morbidity and mortality through early diagnosis and management of conditions including rare inborn errors of metabolism.¹ Conditions such as phenylketonuria are not clinically evident at birth but lead to significant irreversible harm or death if not treated promptly.² NBS has saved countless lives and vastly improved the quality of children's lives by allowing timely therapeutic interventions, and technological advances such as the use of tandem mass spectrometry (MS/MS) have played a significant role in expansion of NBS.^{3,4}

The ability to analyze many or all genes in the genome simultaneously provides new opportunities for genomic medicine. The capacity of genome-scale sequencing for disease

gene discovery is well documented,⁵ and it is increasingly being applied as a diagnostic test in children with suspected monogenic disorders.^{6,7} The tangible potential of genomic sequencing more broadly in medicine, as part of "personalized" or "precision" medicine,⁸ was foreshadowed in 1990 by Walter Gilbert, who extrapolated from the exponential growth of DNA sequencing that all newborns would have their genomes sequenced by 2030 or 2040. The idea that genomic sequencing will someday become part of the standard care of newborns is carried forward into today's dialogue:

As we learn more about effective interventions for genetic risk factors, and recognize that interventions early in life provide significant advantages, it will become more and more compelling to determine this information at birth.

—F. S. Collins, *The Language of Life: DNA and the Revolution in Personalized Medicine*. New York: Harper Perennial (2010)

abstract



Departments of ^aGenetics and ^vPediatrics, University of North Carolina at Chapel Hill School of Medicine, Chapel Hill, North Carolina; Divisions of ^bNewborn Medicine and ^cGenetics and Genomics, The Manton Center for Orphan Disease Research, Boston Children's Hospital, Harvard Medical School, Boston, Massachusetts; ^hLaboratory for Molecular Medicine, ^mDivision of Genetics, and ^uDepartment of Pediatric Newborn Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts; ^dRTI International, Research Triangle Park, North Carolina; ^eUniversity of California, Berkeley, California; ^fAmerican College of Medical Genetics and Genomics, Bethesda, Maryland; ^gRady Children's Institute for Genomic Medicine, San Diego, California; ⁱChicago School of Public Health, University of Illinois, Chicago, Illinois; ^jInstitute for Human Genetics, ^kInstitute for Health and Aging, ^lDepartment of Anthropology, History, and Social Medicine, ⁿCardiovascular Research Institute, and ^oDepartment of Dermatology, ^pDepartment of Pediatrics, ^qBenioff Children's Hospital, and ^rDepartment of Pediatrics, University of California, San Francisco, California; ^sGenetic Disease Screening Program, California Department of Public Health, Sacramento, California; ^tDepartment of Pediatrics and Division of Neonatology, Oregon Health & Science University, Portland, Oregon; ^uNational Human Genome Research Institute and ^vNational Center for Advancing Translational Sciences, National Institutes of Health, Bethesda, Maryland; ^wDepartment of Pediatrics, Children's Mercy Hospital, Kansas City, Missouri; ^xCenter for Medical Ethics and Health Policy, Baylor College of Medicine, Houston, Texas; and ^yUniversity of Washington, Seattle, Washington

Dr Berg conceptualized and designed the study, drafted the initial manuscript, participated in data analysis, and reviewed and revised the manuscript; Drs Agrawal and Parad drafted the initial manuscript, supervised data collection at 1 of the sites, participated in data analysis, and reviewed

To cite: Berg JS, Agrawal PB, Bailey DB, et al. Newborn Sequencing in Genomic Medicine and Public Health. *Pediatrics*. 2017;139(2):e20162252

Such predictions raise important questions: Could genomic sequencing become part of the universal standard NBS performed by each state, replacing conventional biochemical testing, or could it be offered as an optional supplement? How would sequencing be paid for, and how would parental informed consent be obtained? What impact would genomic sequencing in newborns have on children's health? What implications would it have for issues such as protecting an autonomous person's right not to know?

In the diagnostic setting, the goal of genome-scale sequencing is to identify genetic variants that provide a molecular etiology for the patient's symptoms. All other variants are considered incidental findings (or secondary findings when discovered through intentional analysis).⁹ These additional findings differ widely with regard to predictive capacity and clinical actionability. Not surprisingly, there is disagreement about how much genomic information should be routinely returned in a pediatric setting,¹⁰⁻¹⁴ and children and their parents are likely to express unique preferences.^{15,16} Central challenges for clinical implementation thus revolve around the boundaries of professional responsibility and individual or parental choice, best practices for informed consent and determining parental preferences, standards regarding the types of findings that should be reported, long-term storage of genomic information so that it can be acted on at the appropriate time, and whether and how genomic data should be reanalyzed and reinterpreted.

The application of genomic sequencing in asymptomatic newborns intensifies many of these challenges and exposes deep societal questions about nonmaleficence, beneficence, autonomy, and the preservation of each child's open future.¹⁷⁻¹⁹ Additionally, the

economics of such screening must be considered before implementation on a population level. Although the technical features needed for rapid genomic sequencing appear feasible, interpretation of each asymptomatic person's variant profile will remain labor intensive for the foreseeable future, and clinically useful prediction of future disease may prove elusive. Recognizing these trends, the *Eunice Kennedy Shriver* National Institute of Child Health and Human Development (NICHD), National Human Genome Research Institute, and the Office of Rare Disease Research held a workshop in December 2010 to identify elements of a trans-National Institutes of Health research agenda that could inform the possible application of new genomic technologies to NBS and child health (https://www.nichd.nih.gov/about/meetings/2010-retired/Documents/Newborn_Research_Agenda.pdf). A consensus finding was the need to examine the technical, clinical, social, and ethical issues related to sequencing in the newborn period in unison. Subsequently, the NICHD and the National Human Genome Research Institute issued a funding opportunity to develop a consortium to explore, in a limited but deliberate manner, opportunities to use genomic information for broadening our understanding of diseases identified in the newborn period, in the context of public health NBS or clinical sequencing of newborns. The consortium, called Newborn Sequencing in Genomic Medicine and Public Health (NSIGHT), is addressing 3 key research questions:

- For disorders currently screened in newborns, how can genomic sequencing replicate or augment known NBS results? Can sequencing replace current screening modalities?
- What knowledge could genomic sequencing provide about

conditions not currently screened for in newborns?

- What additional clinical information could be learned from genomic sequencing relevant to the clinical care of newborns?

In 2014, 4 groups were funded to explore newborn sequencing in different clinical contexts by using unique study designs (Table 1 and Fig 1). Each study also included an aim examining ethical, legal, and social considerations. This article examines some of the challenges of newborn sequencing in 3 distinct clinical settings, describes the 4 projects, and puts this research in the context of current and future strategies for NBS and sequencing of newborns in clinical care settings.

CLINICAL SETTINGS FOR NEWBORN SEQUENCING RESEARCH: DIAGNOSTIC, PREVENTIVE, AND PREDICTIVE

The setting in which genomic sequencing is performed affects its technical performance, yield, and potential benefits and harms. In a diagnostic context, newborn sequencing is much like any other genetic diagnostic modality, and the ability to provide a molecular diagnosis depends on a number of factors including genetic architecture, phenotypic expressivity, locus heterogeneity and the fraction of cases accounted for by known disorders, mutational spectrum, the types of variants that are detectable by sequencing, and the quality of the assay performed in the patient. In contrast, the application of genomic sequencing in an asymptomatic newborn substantially depends on the power of genotype to predict disease status in the present or the future. Thus, a critical question is whether genomic variant data can be used to accurately predict disease when the pretest probability of disease is low, especially in cases where there may be no secondary

TABLE 1 NSIGHT Project Overviews

	BWH, BCH, Baylor College of Medicine	Rady, Children's Mercy Hospital	UCSF	University of North Carolina
Patient cohorts	Sick newborns (ICUs). Healthy newborns (well nursery).	Sick newborns (NICU)	Deidentified DBS samples from California NBS program Newborn DBS from consenting individuals with primary immunodeficiency	Children affected with known NBS findings Healthy newborns (prenatal recruitment)
Biospecimens	Whole blood (newborns) and saliva (newborns and parents).	Whole blood (parent–infant trios)	DBS retrieved from NBS program biobank	Cheek swabs
Sequencing	Exome sequencing. Illumina Content Exome.	WGS Illumina HiSeq 2500, rapid run mode	Exome sequencing Nimblegen v3 capture	Exome sequencing Agilent V6.0 capture
Informatics	Illumina HiSeq. ≥100 × mean coverage. All newborns: analysis of variants in genes responsible for childhood-onset disorders (<18 y). Sick newborns/Later-onset healthy and sick newborns: indication-based analysis of all variants in genes relevant to the phenotype (if applicable).	Clinical features translated into phenotype terms and differential diagnosis by Phenomizer (and custom lists)	Illumina HiSeq 40–80 × average coverage Metabolic disease NBS samples: blinded assessment as an initial screen; second-tier analysis combining genomic, clinical, and MS/MS data Immunodeficiency cohort: analysis of genes relevant to patient's phenotype	Illumina HiSeq 40–80 × average coverage All participants: primary analysis blinded to phenotype • Known diagnosis cohort: diagnostic or indication-based analysis based on phenotypic findings
Primary results returned	Diagnostic findings (ICU patients and others with clinical indications). Highly penetrant childhood-onset or childhood treatable conditions (all participants).	Diagnostic or likely diagnostic findings	No contact or return of results for metabolic NBS program cohort Offer clinical confirmatory testing to follow up likely pathogenic variants for immunodeficiency cohort	Childhood-onset medically actionable conditions (all participants) Diagnostic findings (affected cohort)
Secondary results returned	Carrier status for childhood-onset conditions and selected pharmacogenomics (all participants).	Incidental genetic disease diagnosis if life-threatening in childhood	None	Parents randomly assigned to decision group can select from 3 categories: • Childhood-onset non–medically actionable conditions • Carrier status for recessive conditions • Adult-onset medically actionable conditions

TABLE 1 Continued

	BWH, BCH, Baylor College of Medicine	Rady, Children's Mercy Hospital	UCSF	University of North Carolina
Psychosocial and medical outcomes research	<p>Surveys of parents and physicians assess impact of genomic sequencing across key domains:</p> <ul style="list-style-type: none"> • Attitudes and preferences. • Health care utilization. • Health behaviors and intentions. • Decisional satisfaction. • Psychological impact. <p>Psychosocial impact on the family.</p>	<p>Surveys of parents and physicians assess impact of genomic sequencing across key domains:</p> <ul style="list-style-type: none"> • Attitudes and preferences • Health care utilization • Health behaviors and intentions • Decisional satisfaction • Psychological impact <p>• Psychosocial impact on the family</p> <p>Correlation of parents' attitudes about test results with their health literacy, genomic literacy, anxiety, depression, and religiosity</p>	<p>Focus groups with:</p> <ul style="list-style-type: none"> • Parents of immunodeficiency patients • Healthy pregnant women • Obstetric and pediatric clinicians 	<p>Surveys with parents assess decision-making about genomic sequencing and its effects on individual and dyadic parent outcomes:</p> <ul style="list-style-type: none"> • Parents' collaborative decision-making and conflict • Prenatal anxiety (parents of healthy newborns only) • Parental bonding with child • Attitudes and beliefs about genomic sequencing • Decision conflict and regret • Test-related and general distress • Beliefs and concerns about the child's future health
Other project aims	<p>Although Sanger confirmation is used for the main project, the study is exploring orthogonal sequencing by 2 NGS methods as an alternative for variant confirmation.</p>	<p>Comparison of rate of diagnosis and time to diagnosis in WGS and no-WGS groups</p> <p>Rates and types of actionability measured</p>	<p>Assess suitability of stored DBS as a source of DNA for deep sequencing</p>	<p>Semiquantitative metric to determine medical actionability</p> <p>Electronic decision aid for parental preference setting</p>

gold standard clinical evaluation by which to validate the genomic prediction.

Sequencing in a NICU population of sick infants extends current molecular diagnostic strategies to a genomic scale, whereas sequencing otherwise healthy infants targets the prevention of future disease and is more akin to current NBS performed in a public health setting. Sequence information could also be used to guide a patient's care throughout life by predicting disease and directing management strategies for clinical scenarios that emerge, blending the predictive, reproductive, diagnostic, therapeutic, and prognostic value of genomic information. The NSIGHT projects differ in the clinical scenarios in which the use of genome-scale sequencing is being studied, spanning the areas described below.

Diagnostic Sequencing in the NICU

Level III and IV NICUs care for many neonates with genetic disorders, including metabolic disorders²⁰ and congenital malformations.^{21,22} Indeed, genetic disorders and congenital anomalies are the leading cause of death in the NICU.²³ Current approaches for diagnosis of suspected genetic disorders in NICU patients include karyotyping, chromosomal microarrays, single-gene testing, and gene panels. In many cases, the etiologic diagnosis remains elusive despite several rounds of genetic testing, and clinical genome-scale sequencing is used for only a small fraction of cases. If genomic sequencing were used earlier in the diagnostic process, it could provide more timely definitive diagnoses and thereby increase the precision of treatments, whether

therapeutic or palliative, and provide answers about prognosis and family recurrence risk. In addition, sequence data could aid in interpretation of the false-positive NBS results commonly reported in premature infants because of their immature organ systems, liver enzymatic activity, long-term parenteral nutrition requirement, and other comorbid conditions.²⁴

Preliminary studies have shown potential cost reductions resulting from genomic sequencing in neonates with genetic disorders.^{7,25,26} However, uptake has been impeded by concerns about clinical interpretation of ambiguous results, secondary findings and potential parental anxieties, economic factors including resistance on the part of third-party payers in the absence of definitive data on clinical utility




FIGURE 1

A, Diagram of the protocol in progress at BWH and BCH, Boston, Massachusetts. Infants are recruited from the well baby nursery at BWH and from the ICUs at BCH and BWH. After a pre-enrollment session with a study genetic counselor and completion of baseline outcomes, enrolled infants are randomly assigned to receive NBS and family history or NBS, family history, and exome sequencing. Results are disclosed to the family by a study genetic counselor and physician, and postdisclosure outcomes are collected. Follow-up is performed at 3 and 10 months after disclosure. Medical, behavioral, and economic outcomes are collected throughout the study from surveys, medical record reviews, and consultation with the families. B, Diagram of the protocol in progress at NICUs at Children’s Mercy Hospital, Kansas City (CMH) and Rady Children’s Hospital, San Diego (RCHSD). Eligible patients are infants <4 months old in whom a clinical genetic test or genetic consult was ordered, or those with 1 major anomaly or 3 minor structural anomalies, or an abnormal laboratory test suggestive of a genetic disease, or an abnormal response to standard therapy for a major underlying condition. Enrolled infants are randomly assigned to receive standard diagnostic testing or standard tests and rapid WGS of parent–infant trios. Diagnostic results are returned. Primary outcome measures are rate of molecular diagnosis in 28 days, time to diagnosis, and whether the diagnosis provided a change in clinical management. C, Diagram of the NBSex protocol in progress at UCSF. NBSex uses archived residual DBS from a very large and diverse population to examine the diagnostic utility of exome sequencing in 2 parallel projects. For metabolic disorders (left), DBS samples from deidentified true positives (TP), false positives (FP), and false negatives (FN) are subjected to exome sequencing, variant calling, and interpretation under blinded models for evaluation of first-tier screening and in conjunction with MS/MS and clinical data for evaluation of second-tier screening. A predetermined list of 93 genes relevant to the metabolic disorders is the basis for the exome assessment. Consented, identified patients with immune disorders with no gene identified (right) also undergo NBS DBS retrieval and exome sequencing and analysis, in this case with a pipeline restricted to immune system genes. Analysis protocols for both

and cost-effectiveness, and difficulty ordering tests due to institutional concerns over high costs. The process is also complicated by a need for detailed phenotypic information for optimal analysis by diagnostic laboratories. Provisional genomic diagnosis of genetic disease is technically feasible in as little as 26 hours.^{27,28} However, cost is inversely related to the turnaround time, and the optimal balance of cost and clinical utility is unknown. Although some aspects of rapid turnaround time can be improved through technology, limits imposed by variant review and clinical interpretation standards will remain, including the need for communication between the clinical team and the laboratory for clinical genomic assessment. Parental acceptance of testing, given the unclear future implications for insurability and privacy, is not yet known. Finally, the responses of neonatologists and parents to genomic information warrant more study, and rigorous frameworks for translating such information into precision care plans in NICUs remain to be developed.

Preventive Sequencing in a Public Health Setting

Current NBS yields few false-negative results but does incur substantial numbers of false positives, with attendant emotional and financial costs.²⁹ For example, in a study of 176 186 specimens screened by MS/MS, there were 51 true positives, 2 false negatives,

and 454 false positives that were ultimately resolved as nondisease after referral to a metabolic center.³⁰ Genomic sequencing could function as a multiplexed second-tier screen, increasing the specificity of current MS/MS screening tests by distinguishing false-positive results from true disease and aiding in the differential diagnosis of nonspecific biochemical profiles. Sequencing could also confirm conditions identified through other screening methods,^{31–33} aid in providing prognosis and appropriate treatment,^{34–37} determine the etiology of conditions identified through point-of-care testing,³⁸ and provide families with information about pathogenic variants that could be used for family testing. Such analyses could also increase knowledge of correlations between genotypes and phenotypes and might reveal possible genetic contributions to false positives, such as abnormal MS/MS screening data due to carrier status or hypomorphic variants. These goals will require longitudinal follow-up to obtain clinical data and examine genotype–phenotype correlations to ultimately determine the predictive capacity and clinical impact of genetic variants.

Genomic sequencing could also be deployed as a first-tier screen, particularly for rare disorders that currently lack methods for conventional biochemical NBS. In the public health context,

selecting exactly which disorders to screen for requires careful consideration of factors such as age of onset, severity, penetrance, treatability, confirmatory testing, and opportunities for surveillance. Some genetic conditions will fulfill the original Wilson and Jungner criteria for screening,^{39–41} but many will not. Thus, there is a need for scalable methods to determine which conditions would be appropriate for inclusion in a public health screening setting. In addition, the current practice of returning findings consistent with carrier status might be unsustainable given that nearly every person is likely to be a carrier for a handful of rare recessive conditions.

A significant challenge in the use of sequencing for NBS is the lack of data regarding the analytic and clinical performance of sequencing as a predictive test. Little is known about the positive or negative predictive value of genomic sequencing in asymptomatic people whose previous probability of disease is very small. Although it may seem self-evident that Mendelian diseases would be best identified from genetic sequence, this assumption remains to be demonstrated and may not be true. Monogenic illnesses are often influenced by additional and as-yet-unidentified genetic or environmental factors, whereas MS/MS measures analytes that are typically closer

FIGURE 1 Continued

applications undergo refinement based on integration of observed sensitivity and specificity with genome analysis tools. Enrollees with immune disorders can obtain confirmational clinical gene testing, and their assessment of risks, benefits, and uncertainties of exome sequencing for NBS are solicited. D, Diagram of the North Carolina Newborn Exome Sequencing for Universal Screening protocol in progress at the University of North Carolina at Chapel Hill. This study is enrolling healthy newborns, identified prenatally, and children affected with known conditions identified through standard NBS (metabolic disorders, hearing loss, pulmonary disorders). Parents use an electronic decision aid in addition to an in-person consultation with a genetic counselor to determine whether to have their child undergo sequencing. Exome sequencing is performed, with analysis of a panel of genes associated with childhood-onset medically actionable conditions (NGS-NBS) for all participants and indication-based analysis for patients from the diagnosed cohort. Participants are randomly assigned at the time of return of results. Parents in the control arm receive only the primary diagnostic findings and NGS-NBS results, whereas parents in the decision arm will use the electronic decision aid to choose between 3 additional categories of optional genomic information (adult-onset medically actionable conditions, childhood-onset non–medically actionable conditions, and carrier status for recessive conditions). Parents will also participate in longitudinal surveys to assess their responses to the genomic information.

to relevant phenotypes. Even for the best-studied diseases, there are substantial challenges in interpreting rare variants. Variant selection algorithms that maximize sensitivity necessarily sacrifice specificity, leading to increased false-positive results and potential downstream harms due to unnecessary medical interventions. A strategy of excluding “variants of uncertain significance” (which by definition have poor predictive value) from genomic screening results may be necessary. Historically, the focus of NBS on preventable disorders has led to low tolerance for false-negative screening results. With genomic sequencing it might be necessary to shift the screening paradigm from finding all affected individuals to finding an optimal proportion of cases for a larger number of potentially treatable conditions.

Predictive Sequencing of Newborns in Genomic Medicine

The broadest vision of genomic medicine, and potentially the most challenging for societal and practical reasons, involves the use of sequence data to guide a patient’s care throughout life. Genomic sequencing reveals information well beyond the scope of conventional NBS. Some of this information could result in medical action, but most will not, raising questions of exactly what information should be reported and when.^{42,43} Variant data could conceivably be held for future diagnostic analysis in the event that the patient develops symptoms of a genetic condition. Currently, our ability to interpret genetic variants is largely confined to simple monogenic and oligogenic conditions. As we learn to use multifactorial models for risk stratification or management in a clinically useful way, additional information will

increasingly be available. Within this spectrum is the potential for genomic information to alert clinicians to reconsider the family history or interpret physical examination findings in a new light, and potentially the ability to benefit other family members before they develop a disease. Pharmacogenomic variants could guide the real-time selection and dosing of medications, yielding safer and more effective treatments. Recessive carrier traits detected in newborns could alert parents to genetic risks that provide information valuable to reproductive planning. Common variation for complex conditions may motivate families to be more vigilant about diet and other lifestyle choices. Finally, there is the potential for voluntary personal exploration of one’s own genomic data.

If genomic sequence data are available, parents will need to make decisions about whether to learn about additional categories of information that may predict future events about their child with differing levels of certainty and ability to intervene, ranging from childhood-onset conditions that may not have direct interventions or preventive measures, adult-onset medically actionable conditions, or carrier status for recessive disorders. Studies suggest that parents are interested in their child’s genetic variants, even when that information has no defined clinical utility,⁴⁴ although these preferences have largely been elicited in hypothetical scenarios and may not reflect real-life choices.

Genomic information may enable families to become aware of otherwise unsuspected familial risks, including potentially actionable adult-onset conditions in the infant that a parent is unknowingly

carrying. However, some findings may be at odds with professional guidance that genetic testing in asymptomatic minors should generally be done only when identification before adulthood is needed to prevent harm and directly benefit the child.^{11,13,17} The potential to query genome-scale data in children for secondary findings has elicited vigorous debate over the ethical boundaries of return of results. The argument has been made that benefit to family (eg, a parent who is unknowingly at risk) may be a valid consideration in decisions regarding return of results for actionable adult-onset conditions in children.^{45,46} Genome-scale sequencing and analysis in newborns would probably require modification of current informed consent procedures^{41,47} compared with a targeted screen focusing on a restricted number of conditions. If implemented across the entire population, genomic sequencing would fundamentally alter contemporary public health NBS procedures, necessitating innovative approaches to facilitate parental decision-making. Alternatively, such testing could be implemented through voluntary, out-of-pocket testing.

Expanding genomic sequencing in newborns to include a broad range of conditions demands a close partnership between clinical providers and parents, similar to other areas of medicine in which shared decision-making is becoming the norm.^{48,49} Determining criteria for disclosure of information will be a challenge for clinicians and policymakers and will require development of decisional supports to help parents determine the information they want to learn.⁵⁰ The clinical interactions needed for support of parental decision-making would move this activity beyond the realm of current public health service

provision and into the clinical domain. Workforce shortages may present significant obstacles if genome-scale sequencing were to become widely available in the public health setting,⁵¹ suggesting that new and more scalable materials and procedures for communicating the potential benefits and risks of learning such information would need to be developed, validated, and deployed. Systems may need to be established that enable parents to request certain results from their child's genomic information over time, allowing them to decide iteratively as their values and perceptions of risk change or as their child attains an age to assent or consent.

The complexities of genomic result interpretation currently demand trained geneticists and genetic counselors to provide guidance and follow-up management. As genomic medicine becomes more mainstream in health care, a broader range of health care providers will need to interact with genetic information, which is likely to be increasingly viewed as 1 of many risk factors influencing future conditions; reports will have to be constructed with clarity that makes them useful for pediatricians and primary care providers. The potential use of this type of genomic information over time would necessitate development of new infrastructure to manage reporting, reanalysis, storage, and integration with electronic health records. Such data could be used for iterative phenotyping of the individual to define the clinical relevance of genetic variants. However, thresholds for reporting or acting on potentially relevant variants would have to be calibrated against the possible harms of false-positive results or overdiagnosis, which would lead to unnecessary, dangerous, or costly

medical treatments.⁵² The benefits of detecting true positives must therefore be balanced against the magnitude of harms.

NSIGHT RESEARCH GROUPS, STUDY DESIGNS, AND KEY QUESTIONS BEING ADDRESSED

Each of the 4 members of the NSIGHT Consortium independently designed and implemented study designs that focus on somewhat different populations and research questions (Fig 1 and Table 1).

Genome Sequence-Based Screening for Childhood Risk and Newborn Illness project (Fig 1A), led by Brigham and Women's Hospital (BWH) and Boston Children's Hospital (BCH) at Harvard Medical School and by Baylor College of Medicine, is a randomized controlled trial assessing the impact of providing genomic sequencing information to parents and physicians of newborns (clinicaltrials.gov identifier NCT02422511). The study is enrolling a cohort of healthy newborns approached in the postpartum period from the BWH Well Baby Nursery and a cohort of sick newborns from the BWH NICU and the ICUs at BCH. Within each cohort newborns are randomly assigned to either the control arm (conventional NBS results and a detailed family history) or the experimental arm (genomic sequencing in addition to conventional NBS and a detailed family history). Parents and physicians are surveyed to assess the impact of the genomic information across several key domains including attitudes and preferences, health care utilization, health behaviors and intentions, decisional satisfaction, and psychosocial impact on the family.⁵³

The Clinical and Social Implications of 2-Day Genome Results in Acutely III Newborns project, led by Rady Children's Institute for Genomic Medicine and Children's Mercy Hospital, studies the use of rapid whole genome sequencing (WGS) at 2 large level-IV NICUs (~1000 admissions per year) in children's hospitals (Fig 1B). WGS was adapted for diagnosis of rare genetic diseases in NICUs, including shortening the minimum time to provisional diagnosis to 26 hours, increasing the analytic sensitivity and specificity of variant detection to >99.5%, and gaining US Food and Drug Administration approval to report a provisional diagnosis verbally to an attending neonatologist if death was imminent and the diagnosis would inform implementation of a treatment that could change the outcome.^{27,28} The current study is a prospective, randomized controlled trial of the diagnostic utility, cost-effectiveness, and psychosocial implications of rapid WGS at the 2 NICUs (clinicaltrials.gov identifier NCT02225522). As part of this study, doctors and parents are surveyed about their perceptions of the risks and benefits of rapidly obtaining WGS results.

The goal of the Sequencing of Newborn Blood Spot DNA to Improve and Expand Newborn Screening (NBSseq) project, led by the University of California, San Francisco (UCSF) and collaborating institutions, is to evaluate the potential application of exome sequencing to public health NBS by using dried blood spots (DBS) (Fig 1C). The project explores the feasibility of exome sequencing to augment or replace current MS/MS technologies in NBS. Exome sequencing is performed at UCSF on deidentified archival DBS from all California newborns found to have metabolic disorders in the

TABLE 2 Cross-Consortium Working Groups

Ethical, Legal, Economic, and Social Issues	Common Data Elements	Outcomes and Measure
<p>Key questions</p> <ul style="list-style-type: none"> • Differences in perceptions of benefits and risks of sequencing between symptomatic and asymptomatic populations • Parent willingness to accept sequencing and factors associated with parents' decisions • Extent to which parents are willing to accept uncertainties inherent in test interpretation • How key stakeholders make decisions about whom to test, how to share results, under what circumstances, and with what goals • Public policy regarding use of genome sequencing as part of mandated screening programs 	<p>Key questions</p> <ul style="list-style-type: none"> • Identify common data elements in the NSIGHT projects to be collected systematically across the consortium • Collaboration with NBSTRN to use LPDR for individual cohort and combined cohort analysis where applicable • Define data elements to be shared more broadly, in a deidentified fashion, with other researchers in the NBSTRN 	<p>Key questions</p> <ul style="list-style-type: none"> • Identify common outcome measures in the NSIGHT projects to be collected systematically across the consortium • Considerations of the overall cost/benefit ratio of newborn sequencing

LPDR, Longitudinal Pediatric Data Resource (<https://www.nbstrn.org/research-tools/longitudinal-pediatric-data-resource>).

past decade, as well as samples that were false positives on the MS/MS screening. Sequence data from newborn DBS⁵⁴ are also being interrogated in a cohort of 50 patients who have been clinically diagnosed with immunodeficiency disorders to determine whether sequencing DBS as part of NBS could facilitate early diagnosis and optimal management of non-severe combined immunodeficiency immune defects. Stakeholder views, perspectives, and value preferences about the potential expansion of NBS are being evaluated through focus groups.⁵⁵ In addition, legal and constitutional issues surrounding the potential use of genome-scale analysis in NBS are being examined.⁵⁶

The North Carolina Newborn Exome Sequencing for Universal Screening study at the University of North Carolina at Chapel Hill evaluates exome sequencing from saliva samples in 2 groups: children and

infants affected with conditions identified through standard NBS and a cohort of healthy newborns whose parents are approached for participation prenatally (Fig 1D). Parents will use an electronic decision aid to assist in decisions about exome sequencing.⁵⁷ After providing informed consent at an in-person study visit, those who accept sequencing will receive results from a “next-generation sequencing newborn screening (NGS-NBS)” panel of genes implicated in childhood-onset medically actionable conditions.⁵⁸ Parents will also be enrolled in a randomized trial of decision-making regarding whether to learn about 3 types of additional genomic findings in their child by using the electronic decision aid (clinicaltrials.gov identifier NCT02826694). The study seeks to understand how parents think about and consider different categories of information,⁵⁹ and, combined with longitudinal quantitative surveys,

the study will reveal the spectrum of results parents decide to learn, issues surrounding returning these findings, and consequences of decision-making and results disclosure.

CROSS-CUTTING CONSORTIUM ACTIVITIES

Although the research projects address distinct research questions and have unique study designs, the 4 NSIGHT groups participate in consortium activities that build on the strengths at each site and harmonize data collection to improve cross-cutting analysis (Table 2). An ethical, legal, and social implications workgroup brings together the perspectives of each project regarding the responses of clinicians, families, payers, and other stakeholders; parental informed consent and decision-making; and implications for public health NBS programs.^{19,60} The Common Data Elements workgroup is charged with creating a common set of data elements collected across the 4 research groups to enable data sharing and combined data analysis. The Outcomes and Measures workgroup examines the standardized instruments being used by each group to assess stakeholder responses to newborn sequencing to facilitate cross-consortium analyses. The NSIGHT Consortium is working closely with the Newborn Screening Translational Research Network (NBSTRN) Coordinating Center, housed at the American College of Medical Genetics and Genomics. The NBSTRN, created as part of the NICHD's Hunter Kelly Newborn Screening Research Program to create a shared research infrastructure to support NBS researchers, provides a mechanism to increase understanding of conditions that

are currently part of routine NBS or may be future candidates for screening. The NBSTRN resources, tools, and network of experts are used in population-based pilots of new screening technologies and natural history studies of screened conditions. NSIGHT Consortium investigators also maintain close ties with other National Institutes of Health-funded consortia including the Clinical Sequencing Exploratory Research consortium⁶¹ and Clinical Genome Resource consortium.⁶²

CONCLUSIONS

Data gathered from the projects in the NSIGHT Consortium will address technical, clinical, and ethical questions that are fundamental to the future consideration of sequencing in newborns. The projects will determine the feasibility and utility of this technology in critical care and public health settings.

Design and implementation of rapid high-throughput methods will be essential to maximize the benefit of sequencing for certain conditions in the neonatal population. Longitudinal follow-up of parents will allow the study of parental decision-making, measure parental preferences in real-world settings, and assess test-related stress or anxiety. Medical outcomes of the children who undergo sequencing will need to be monitored over many years. Ultimately, these data will aid in the development of best clinical practices and provide guidance on the implementation of sequencing in newborns. Although genomic sequencing will expand our ability to diagnose conditions and offer personalized treatments, health care providers and public health entities must be good stewards of this technology, ensuring careful attention to ethical standards and evidence-based outcomes in making recommendations about its use.

ABBREVIATIONS

BCH: Boston Children's Hospital
BWH: Brigham and Women's Hospital
DBS: dried blood spots
MS/MS: tandem mass spectrometry
NBS: newborn screening
NBSeq: Sequencing of Newborn Blood Spot DNA to Improve and Expand Newborn Screening
NBSTRN: Newborn Screening Translational Research Network
NGS-NBS: next-generation sequencing newborn screening
NICHD: Eunice Kennedy Shriver National Institute of Child Health and Human Development
NSIGHT: Newborn Sequencing in Genomic Medicine and Public Health
UCSF: the University of California, San Francisco
WGS: whole genome sequencing

and revised the manuscript; Dr Bailey conceptualized and designed the study and reviewed and revised the manuscript; Dr Beggs conceptualized and designed the study, drafted the initial manuscript, coordinated and supervised data collection at 1 of the sites, participated in data analysis, and reviewed and revised the manuscript; Dr Brenner drafted the initial manuscript, conceptualized and designed the study, designed the data collection instruments, participated in data analysis, and reviewed and revised the manuscript; Dr Brower drafted the initial manuscript, designed data collection instruments, and reviewed and revised the manuscript; Ms Cakici and Dr Petrikin participated in data analysis, designed data collection instruments, coordinated and supervised data collection at 1 of the sites, and reviewed and revised the manuscript; Drs Ceyhan-Birsoy and Chan, Ms Chen, and Drs Leeder, Shieh, and Yu drafted the initial manuscript, participated in data analysis, and reviewed and revised the manuscript; Dr Currier drafted the initial manuscript, conceptualized and designed the study, designed the dried blood spot data collection instruments, coordinated and supervised data collection at 1 of the sites, participated in data analysis, and reviewed and revised the manuscript; Drs Dukhovny and Pereira drafted the initial manuscript, designed data collection instruments, participated in data analysis, and reviewed and revised the manuscript; Dr Green conceptualized and designed the study, drafted the initial manuscript, supervised data collection at one of the sites, participated in data analysis, and reviewed and revised the manuscript; Drs Harris-Wai, B. Powell, and Veeraraghavan participated in data analysis and reviewed and revised the manuscript; Dr Holm drafted the initial manuscript, coordinated and supervised data collection at one of the sites, participated in data analysis, and reviewed and revised the manuscript; Ms Iglesias participated in programmatic oversight and reviewed and revised the manuscript; Dr Joseph participated in data analysis, designed data collection instruments, and reviewed and revised the manuscript; Dr Kingsmore conceptualized and designed the study, drafted the initial manuscript, participated in data analysis, designed data collection instruments, coordinated and supervised data collection at 1 of the sites, and reviewed and revised the manuscript; Dr Koenig conceptualized and designed the study, drafted the initial manuscript, participated in data analysis, designed data collection instruments, and reviewed and revised the manuscript; Drs Kwok and Puck conceptualized and designed the study, participated in data analysis, coordinated and supervised data collection at 1 of the sites, and reviewed and revised the manuscript; Dr Lantos drafted the initial manuscript, participated in data analysis, coordinated and supervised data collection at 1 of the sites, and reviewed and revised the manuscript; Dr Lewis drafted the initial manuscript, designed the electronic decision aid, and reviewed and revised the manuscript; Drs McGuire and Rehm drafted the initial manuscript, designed data collection instruments, coordinated and supervised data collection at 1 of the sites, participated in data analysis, and reviewed and revised the manuscript; Dr Milko and Mr Watson supervised data collection at 1 of the sites and reviewed and revised the manuscript; Dr Mooney participated in data analysis, coordinated and supervised data collection at 1 of the sites, and reviewed and revised the manuscript; Dr C. Powell conceptualized and designed the study, drafted the initial manuscript, coordinated and supervised data collection at 1 of the sites, and reviewed and revised the manuscript; Dr Risch conceptualized and designed the study, participated in data analysis, and reviewed and revised the manuscript; Ms Roche coordinated and supervised data collection at 1 of the sites and reviewed and revised the manuscript; Dr Willig drafted the initial manuscript, participated in data analysis, designed data collection instruments, coordinated and supervised data collection at one of the sites, and reviewed and revised the manuscript; Drs Urv and Wise conceptualized and designed the study, drafted

the initial manuscript, participated in programmatic oversight, and reviewed and revised the manuscript; and all authors approved the final manuscript as submitted.

DOI: 10.1542/peds.2016-2252

Accepted for publication Nov 15, 2016

Address correspondence to Jonathan S. Berg, MD, PhD, Department of Genetics, The University of North Carolina at Chapel Hill, 120 Mason Farm Rd, Chapel Hill, NC 27599-7264. E-mail: jonathan_berg@med.unc.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2017 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: Dr Bailey receives funding from the John Merck Fund and Ovid Therapeutics; Dr Beggs received a 1-time ad hoc consultancy in early 2016 from Human Longevity Inc; Dr Brenner receives research support from Tata Consultancy Services; Dr Green is compensated for speaking or advisory services by AIA, Genome Medical, Helix, Illumina, Invitae, Prudential, and Roche; Dr McGuire is an unpaid consultant to Human Longevity Inc; Dr Parad is an unpaid scientific advisor to Parabase Genomics, and he has stock valued at <\$50 on a next-generation sequencing NICU-based gene panel diagnostic test; Dr Puck's spouse is employed by Invitae, a clinical gene sequencing company; Dr Willig has a patent for a bioinformatics pipeline and data analysis tool for genome sequencing; Dr Yu consults for Claritas Genomics, a commercial genomic medicine company that offers DNA diagnostic services including single-gene, panel, and exome sequencing; and the remaining authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: Supported by National Institute of Health grants 1U19HD077693, 1U19HD077632, 1U19HD077627, and 1U19HD077671. Newborn Screening Translational Research Network is funded through a contract from the *Eunice Kennedy Shriver* National Institute of Child Health and Human Development (HHSN275201300011C). Funded by the National Institutes of Health (NIH).

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

REFERENCES

1. American Academy of Pediatrics, Newborn Screening Task Force. Newborn screening: a blueprint for the future. *Pediatrics*. 2000;106(suppl 2):383–427
2. Blau N. Genetics of phenylketonuria: then and now. *Hum Mutat*. 2016;37(6):508–515
3. Wilcken B, Wiley V, Hammond J, Carpenter K. Screening newborns for inborn errors of metabolism by tandem mass spectrometry. *N Engl J Med*. 2003;348(23):2304–2312
4. Centers for Disease Control and Prevention (CDC). Impact of expanded newborn screening: United States, 2006. *MMWR Morb Mortal Wkly Rep*. 2008;57(37):1012–1015
5. Chong JX, Buckingham KJ, Jhangiani SN, et al; Centers for Mendelian Genomics. The genetic basis of Mendelian phenotypes: discoveries, challenges, and opportunities. *Am J Hum Genet*. 2015;97(2):199–215
6. Biesecker LG, Green RC. Diagnostic clinical genome and exome sequencing. *N Engl J Med*. 2014;370(25):2418–2425
7. Willig LK, Petrikon JE, Smith LD, et al. Whole-genome sequencing for identification of Mendelian disorders in critically ill infants: a retrospective analysis of diagnostic and clinical findings. *Lancet Respir Med*. 2015;3(5):377–387
8. Collins FS, Varmus H. A new initiative on precision medicine. *N Engl J Med*. 2015;372(9):793–795
9. Presidential Commission for the Study of Bioethical Issues. Anticipate and communicate: ethical management of incidental and secondary findings in the clinical, research, and direct-to-consumer contexts. 2013. Available at: http://bioethics.gov/sites/default/files/FINALAnticipateCommunicate_PCSBI_0.pdf. Accessed March 2, 2016
10. Green RC, Berg JS, Grody WW, et al; American College of Medical Genetics and Genomics. ACMG recommendations for reporting of incidental findings in clinical exome and genome sequencing. *Genet Med*. 2013;15(7):565–574
11. Borry P, Evers-Kiebooms G, Cornel MC, Clarke A, Dierickx K; Public and Professional Policy Committee (PPPC) of the European Society of Human Genetics (ESHG). Genetic testing in asymptomatic minors: background considerations towards ESHG recommendations. *Eur J Hum Genet*. 2009;17(6):711–719
12. Committee on Bioethics, Committee on Genetics, and the American College of Medical Genetics and Genomics Social, Ethical, and Legal Issues Committee. Ethical and policy issues in genetic testing and screening of children. *Pediatrics*. 2013;131(3):620–622
13. Botkin JR, Belmont JW, Berg JS, et al. Points to consider: ethical, legal, and psychosocial implications of genetic testing in children and adolescents. *Am J Hum Genet*. 2015;97(1):6–21
14. McCullough LB, Brothers KB, Chung WK, et al; Clinical Sequencing Exploratory Research (CSER) Consortium Pediatrics Working Group. Professionally responsible disclosure of genomic sequencing results in pediatric practice. *Pediatrics*. 2015;136(4). Available at: www.pediatrics.org/cgi/content/full/136/4/e974
15. Goldenberg AJ, Dodson DS, Davis MM, Tarini BA. Parents' interest in whole-genome sequencing of newborns. *Genet Med*. 2014;16(1):78–84
16. Roche MI, Berg JS. Incidental findings with genomic testing: implications for genetic counseling practice. *Curr Genet Med Rep*. 2015;3(4):166–176
17. Ross LF, Saal HM, David KL, Anderson RR; American Academy of Pediatrics; American College of Medical Genetics and Genomics. Technical report: ethical and policy issues in genetic testing and screening of children [published correction appears in *Genet Med*. 2013;15(4):321]. *Genet Med*. 2013;15(3):234–245

18. Bredenoord AL, de Vries MC, van Delden H. The right to an open future concerning genetic information. *Am J Bioeth.* 2014;14(3):21–23
19. Frankel LA, Pereira S, McGuire AL. Potential psychosocial risks of sequencing newborns. *Pediatrics.* 2016;137(suppl 1):S24–S29
20. Meng M, Zhang YP. Impact of inborn errors of metabolism on admission in a neonatal intensive care unit: a 4-year report. *J Pediatr Endocrinol Metab.* 2013;26(7–8):689–693
21. Lu XY, Phung MT, Shaw CA, et al. Genomic imbalances in neonates with birth defects: high detection rates by using chromosomal microarray analysis. *Pediatrics.* 2008;122(6):1310–1318
22. Yoon PW, Rasmussen SA, Lynberg MC, et al. The National Birth Defects Prevention Study. *Public Health Rep.* 2001;116(suppl 1):32–40
23. Weiner J, Sharma J, Lantos J, Kilbride H. How infants die in the neonatal intensive care unit: trends from 1999 through 2008. *Arch Pediatr Adolesc Med.* 2011;165(7):630–634
24. Kamrath C, Hartmann MF, Boettcher C, Wudy SA. Reduced activity of 11 β -hydroxylase accounts for elevated 17 α -hydroxyprogesterone in preterms. *J Pediatr.* 2014;165(2):280–284
25. Soden SE, Saunders CJ, Willig LK, et al. Effectiveness of exome and genome sequencing guided by acuity of illness for diagnosis of neurodevelopmental disorders. *Sci Transl Med.* 2014;6(265):265ra168
26. Stark Z, Tan TY, Chong B, et al; Melbourne Genomics Health Alliance. A prospective evaluation of whole-exome sequencing as a first-tier molecular test in infants with suspected monogenic disorders. *Genet Med.* 2016;18(11):1090–1096
27. Saunders CJ, Miller NA, Soden SE, et al. Rapid whole-genome sequencing for genetic disease diagnosis in neonatal intensive care units. *Sci Transl Med.* 2012;4(154):154ra135
28. Miller NA, Farrow EG, Gibson M, et al. A 26-hour system of highly sensitive whole genome sequencing for emergency management of genetic diseases. *Genome Med.* 2015;7(1):100
29. Tarini BA, Christakis DA, Welch HG. State newborn screening in the tandem mass spectrometry era: more tests, more false-positive results. *Pediatrics.* 2006;118(2):448–456
30. Hall PL, Marquardt G, McHugh DM, et al. Postanalytical tools improve performance of newborn screening by tandem mass spectrometry. *Genet Med.* 2014;16(12):889–895
31. Hoppe CC. Prenatal and newborn screening for hemoglobinopathies. *Int J Lab Hematol.* 2013;35(3):297–305
32. Bhattacharjee A, Sokolsky T, Wyman SK, et al. Development of DNA confirmatory and high-risk diagnostic testing for newborns using targeted next-generation DNA sequencing. *Genet Med.* 2015;17(5):337–347
33. Kösel S, Burggraf S, Fingerhut R, Dörr HG, Roscher AA, Olgemöller B. Rapid second-tier molecular genetic analysis for congenital adrenal hyperplasia attributable to steroid 21-hydroxylase deficiency. *Clin Chem.* 2005;51(2):298–304
34. Rampersad A, Hampton K, Duncan N, et al. Noninvasive buccal swab antigen sample and molecular testing provides extended antigen typing for patients with hemoglobinopathies. *J Pediatr.* 2014;165(5):1003–1007
35. Barry PJ, Ronan N, Plant BJ. Cystic fibrosis transmembrane conductance regulator modulators: the end of the beginning. *Semin Respir Crit Care Med.* 2015;36(2):287–298
36. Rivers L, Gaspar HB. Severe combined immunodeficiency: recent developments and guidance on clinical management. *Arch Dis Child.* 2015;100(7):667–672
37. Matern D, Rinaldo P. Medium-chain acyl-coenzyme A dehydrogenase deficiency. In: Paçon RA, Adam MP, Ardinger HH, et al, eds. *GeneReviews*. Seattle, WA: University of Washington, Seattle:1993–2016). Available at: www.ncbi.nlm.nih.gov/books/NBK1424/. Accessed May 1, 2016
38. Lammens F, Verhaert N, Devriendt K, Debruyne F, Desloovere C. Aetiology of congenital hearing loss: a cohort review of 569 subjects. *Int J Pediatr Otorhinolaryngol.* 2013;77(9):1385–1391
39. American College of Medical Genetics Newborn Screening Expert Group. Newborn screening: toward a uniform screening panel and system executive summary. *Pediatrics.* 2006;117(5 pt 2):S296–S307
40. Wilson JMG, Jungner G. *Principles and Practice of Screening for Disease*. Geneva, Switzerland: World Health Organization; 1968
41. Ross LF. Mandatory versus voluntary consent for newborn screening? *Kennedy Inst Ethics J.* 2010;20(4):299–328
42. Green RC, Lupski JR, Biesecker LG. Reporting genomic sequencing results to ordering clinicians: incidental, but not exceptional. *JAMA.* 2013;310(4):365–366
43. Berg JS, Powell CM. Potential uses and inherent challenges of using genome-scale sequencing to augment current newborn screening. *Cold Spring Harb Perspect Med.* 2015;5(12):a023150
44. Waisbren SE, Bäck DK, Liu C, et al. Parents are interested in newborn genomic testing during the early postpartum period. *Genet Med.* 2015;17(6):501–504
45. McGuire AL, Joffe S, Koenig BA, et al. Point-counterpoint. Ethics and genomic incidental findings. *Science.* 2013;340(6136):1047–1048
46. Wilfond BS, Fernandez CV, Green RC. Disclosing secondary findings from pediatric sequencing to families: considering the “benefit to families.” *J Law Med Ethics.* 2015;43(3):552–558
47. Goldenberg AJ, Sharp RR. The ethical hazards and programmatic challenges of genomic newborn screening. *JAMA.* 2012;307(5):461–462
48. Elwyn G, Frosch D, Thomson R, et al. Shared decision making: a model for clinical practice. *J Gen Intern Med.* 2012;27(10):1361–1367
49. Légaré F, Witteman HO. Shared decision making: examining key elements and barriers to adoption into routine clinical practice. *Health Aff (Millwood).* 2013;32(2):276–284
50. Holm IA, Iles BR, Ziniel SI, et al. Participant satisfaction with a preference-setting tool for the return of individual research results in pediatric genomic research.

J Empir Res Hum Res Ethics.
2015;10(4):414–426

51. Bailey DB Jr, Skinner D, Davis AM, Whitmarsh I, Powell C. Ethical, legal, and social concerns about expanded newborn screening: fragile X syndrome as a prototype for emerging issues. *Pediatrics.* 2008;121(3). Available at: www.pediatrics.org/cgi/content/full/121/3/e693
52. Adams MC, Evans JP, Henderson GE, Berg JS. The promise and peril of genomic screening in the general population. *Genet Med.* 2016;18(6):593–599
53. Waisbren SE, Weipert CM, Walsh RC, Petty CR, Green RC. Psychosocial factors influencing parental interest in genomic sequencing of newborns. *Pediatrics.* 2016;137(suppl 1):S30–S35
54. Hollegaard MV, Grauholm J, Nielsen R, Grove J, Mandrup S, Hougaard DM. Archived neonatal dried blood spot samples can be used for accurate whole genome and exome-targeted next-generation sequencing. *Mol Genet Metab.* 2013;110(1–2):65–72
55. Joseph G, Chen F, Harris-Wai J, Puck JM, Young C, Koenig BA. Parental views on expanded newborn screening using whole-genome sequencing. *Pediatrics.* 2016;137(suppl 1):S36–S46
56. King JS, Smith ME. Whole-genome screening of newborns? The constitutional boundaries of state newborn screening programs. *Pediatrics.* 2016;137(suppl 1):S8–S15
57. Lewis MA, Paquin RS, Roche MI, et al. Supporting parental decisions about genomic sequencing for newborn screening: the NC NEXUS decision aid. *Pediatrics.* 2016;137(suppl 1):S16–S23
58. Berg JS, Foreman AKM, O'Daniel JM, et al. A semiquantitative metric for evaluating clinical actionability of incidental or secondary findings from genome-scale sequencing. *Genet Med.* 2016;18(5):467–475
59. Lewis MA, McBride CM, Pollak KI, Puleo E, Butterfield RM, Emmons KM. Understanding health behavior change among couples: an interdependence and communal coping approach. *Soc Sci Med.* 2006;62(6):1369–1380
60. Thiffault I, Lantos J. The challenge of analyzing the results of next-generation sequencing in children. *Pediatrics.* 2016;137(suppl 1):S3–S7
61. Green RC, Goddard KA, Jarvik GP, et al; CSER Consortium. Clinical Sequencing Exploratory Research Consortium: accelerating evidence-based practice of genomic medicine. *Am J Hum Genet.* 2016;98(6):1051–1066
62. Rehm HL, Berg JS, Brooks LD, et al; ClinGen. ClinGen: the clinical genome resource. *N Engl J Med.* 2015;372(23):2235–2242