



Connecting Mentally Ill Detainees in Large Urban Jails with Community Care

Sean K. Sayers¹ · Marisa E. Domino^{1,2} · Gary S. Cuddeback¹ · Nadine J. Barrett^{3,4,5} · Joseph P. Morrissey^{1,2}

© Springer Science+Business Media New York 2016

Abstract Large urban jails have become a collection point for many persons with severe mental illness. Connections between jail and community mental health services are needed to assure in-jail care and to promote successful community living following release. This paper addresses this issue for 2855 individuals with severe mental illness who received community mental health services prior to jail detention in King County (Seattle), Washington over a 5-year time period using a unique linked administrative data source. Logistic regression was used to determine the probability that a detainee with severe mental illness received mental health services while in jail as a function of demographic and clinical characteristics. Overall, 70 % of persons with severe mental illness did receive in-jail mental health treatment. Small, but statistically significant sex and race differences were observed in who received treatment in the jail psychiatric unit or from the jail infirmary. Findings confirm the jail's central role in mental health treatment and emphasize the need for greater information sharing and collaboration with community mental health agencies to minimize jail use and to facilitate successful community reentry for detainees with severe mental illness.

Keywords Jail · Mental illness · Community connections · Health disparities

✉ Marisa E. Domino
domino@unc.edu

¹ Cecil G. Sheps Center for Health Services Research, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

² Department of Health Policy and Management, The Gillings School of Global Public Health, University of North Carolina at Chapel Hill, 1104G McGavran-Greenberg Hall, CB#7411, 135 Dauer Dr., Chapel Hill, NC 27599-7411, USA

³ Office of Health Equity and Disparities, Duke Cancer Institute, Durham, NC, USA

⁴ Duke Community Connections and Collaborations Core, Duke Center for Community and Population Health Improvement and Clinical Translational Science Award, Durham, NC, USA

⁵ Department of Community and Family Medicine, School of Medicine, Duke University, Durham, NC, USA

Introduction

Over the past few decades, people with severe mental illness have become a growing part of jail caseloads. Today, the prevalence of severe mental illness in large urban jails is at least three to four times higher than found in the general population. Specifically, about 4 % of the general population has a severe mental illness [1–3] whereas more than 17 % of people admitted to jail have a severe mental illness [4]. Moreover, jail detainees with severe mental illness are known to have disproportionately high rates of substance abuse [5–8], poverty [9–11], and chronic medical conditions [5, 12], which make them complex and costly to treat.

Although men (87 %) make up the large majority of the jail population, the prevalence of severe mental illness among jail admissions is twice as high among women (31 %) than among men (15 %). As applied to fiscal year 2011 [13], these percentages indicate that there were about 2 million jail admissions with severe mental illness including about a half-million women. At midyear 2012, nearly 6 in 10 detainees in local jails were racial or ethnic minorities with African Americans constituting about 37 % and Hispanics 15 % of the total population [14].

Jails have become the default mental health system in communities throughout the US for many people with severe mental illness [15]. Some estimates suggest that approximately three-fourths of detainees with severe mental illness in large urban jails receive acute psychiatric inpatient treatment in the criminal justice system rather than in the mental health system [16]. This trend is of particular concern because there is a profound lack of adequate mental health treatment within all levels of the criminal justice system [17–20].

While much is known about racial, ethnic and gender disparities in receipt of community-based outpatient mental health care [21–25], little is known about whether these same disparities carry-over into jail settings. Some recent studies suggest that racial disparities in mortality rates are smaller in prison populations than in the general community population [26–28], for example, but research has yet to establish whether jails mirror similar racial and gender disparities that exist in community treatment for people with severe mental illness.

In this paper, we address two questions to advance research in this area. First, how many people with severe mental illness who received community-based treatment prior to jail admission also receive treatment in jail? Second, do rates of in-jail treatment vary by race and gender? These questions are particularly important given that jail stays are relatively brief and that comprehensive mental health services in jails are often lacking.

Methods

Study Setting

This study used a unique administrative database that linked detailed information from Medicaid, the county jail, state psychiatric hospitals, and community mental health agency records in King County (Seattle), Washington. The King County jail ranked 34th on the list of largest jail jurisdictions in the United States [29], with an average daily jail census in 2000 of 2400 and a total number of jail bookings of 60,992. The jail incarceration rate was 138 per 100,000, which was below the US average (226 per 100,000) (see [30] and [31] for more details). Further, the community population in King County was largely white (75.7 %); among minorities, there was a larger percentage of Asians (10.8 %) as compared to African Americans (5.4 %) and Hispanics or Latinos (5.5 %).

Sample

The linked data combined random samples drawn from administrative data sources from the community mental health, Medicaid, and jail service sectors over a 5-year period (1993–1998). This data source is unique in its ability to track individuals in and out of jail during this time period, and identify the use of several types of mental health services received in the jail. For our study we included adults between 18 and 64 years of age who were detained at least once during our study period and had previously used the community mental health or state hospital system. We then restricted our sample to those participants who had a diagnosis of severe mental illness (schizophrenia or bipolar disorder) as recorded in community mental health or state hospital records prior to jail entry. Data were collapsed at the person-level to indicate service use over the 5-year period.

Study participants ($n = 2855$) were predominantly male (68 %), White (72 %) and, on average, 38 years of age ($SD = 9.5$) (Table 1). Non-whites included African Americans (23 %), 3 % Asians (3 %), and Native Americans (2 %). Although Latinos constituted 5 % of the overall administrative data files, no Latinos appeared in the estimation sample for this analysis. Participants spent an average of 52.6 days ($SD = 102.9$) in jail. With regard to prior service contacts, 97 % of the sample had at least one community-based outpatient mental health encounter and 10 % had at least one state psychiatric hospital admission. The mean number of outpatient mental health encounter visits prior to jail detention was 146 ($SD = 210$), ranging from 0 to 1372 visits.

Measures

Dependent Variables

We identified persons who received mental health services in jail in two ways. First, we created a dichotomous variable measuring whether each person ever was in the jail psychiatric unit. Second, we created another dichotomous variable measuring whether individuals had received any mental health services on an outpatient basis at the jail infirmary staffed by county health department employees or in the general population. Infirmary contacts were recorded in a Public Health Information System (PHIS). Contacts associated with mental health diagnoses or procedure codes were used to identify receipt of an in-jail mental health service. We examined the two components separately and also combined them into a composite measure of jail mental health service receipt.

Demographics

The available data allowed us to examine differences in the use of jail-based mental health services by sex as well as by four race/ethnic groups: White, African-American, Asian, and Native American. We controlled for age as a continuous variable in quadratic form.

Covariates and Independent Variables

Because treatment of mental illness is likely an increasing function of the length of stay in jail, we controlled for the overall number of days in jail. We also controlled for the number of outpatient mental health encounters in the community mental health system prior to jail admission and expenditures on state psychiatric hospital care, as crude measures of severity of illness.

Table 1 Sample characteristics of king county jail detainees with severe mental illness ($n = 2855$)

Variable	<i>n</i>	%
Gender		
Male	1926	67.5
Female	929	32.5
Race/ethnicity		
White	2058	72.1
Male	1395	67.8
Female	663	32.2
African American	649	22.7
Male	432	66.6
Female	217	33.4
Asian	82	2.9
Male	64	78.0
Female	18	22.2
Native American	66	2.3
Male	35	53.0
Female	31	47.0
State hospital use prior to jail detention	284	9.9
Outpatient mental health encounters prior to jail detention	2766	96.9
Average of outpatient mental health encounter visits prior to jail detention [mean (SD)]	146 (210)	
Jail days [mean (SD)]	52.6 (102.9)	
Indicator of jail mental health service use		
Stay in jail psychiatric unit (JPU)	1741	61.0
Jail infirmary contact (JIC)	1752	61.4
Either JPU or JIC	2008	70.3
Treatment from both JPU and JIC	1485	52.0
No mental health service use	847	29.7

JIC refers to an out-patient mental health service contact with the jail infirmary

Analysis

We used logistic regression models to examine the association between our covariates and the probability that a detainee received mental health services while in jail. All continuous explanatory variables were entered in quadratic form, which improved model fit. We report the logit coefficients along with the average marginal effects (AMEs) of each covariate on the predicted probability of mental health treatment. Statistical significance was attributed to p values of .05 or less.

Results

Overall, 70 % of study participants received mental health treatment while in jail (Table 1) and this percentage was roughly the same across racial groups (Table 2). Women made up 32 % of the total sample; however, women accounted for 41 % of those who did not

Table 2 Mental health service receipt by gender and race/ethnicity while in jail ($n = 2855$)

Variable	Mental health service either JPU or JIC		Stay in jail psychiatric unit (JPU)		Jail infirmary contact for mental health (JIC)		No receipt of mental health services in jail	
	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%
Gender								
Male	1429	71.7	1275	73.2	1262	72.0	497	58.7
Female	579	28.8	466	26.8	490	28.0	350	41.3
Mental health service by race/ethnicity								
White	1439	71.7	1251	71.9	1252	71.5	619	73.1
Male	1026	71.3	914	73.1	903	72.1	369	59.6
Female	413	28.7	337	26.9	349	27.9	250	40.4
African American	471	23.5	402	23.1	423	24.1	178	21.0
Male	330	70.1	294	73.1	300	70.9	102	57.3
Female	141	29.1	108	26.9	123	29.1	76	42.7
Asian	52	2.6	51	2.9	36	2.1	30	3.5
Male	43	82.7	42	82.4	30	83.3	21	70.0
Female	9	17.3	9	17.7	6	16.7	9	30.0
Native American	46	2.3	37	2.1	41	2.3	20	2.4
Male	30	65.2	25	67.6	29	70.7	5	25.0
Female	16	34.8	12	32.4	12	29.3	15	75.0

receive mental health services while in jail. Service recipients were just as likely to receive treatment from the jail mental health unit (61 %, $n = 1741$) as from the jail infirmary (61 %, $n = 1752$) and 52 % ($n = 1485$) received services from both sources. Men (74 %, $n = 1492$) were more likely to receive mental health treatment than women (62 %, $n = 579$) and this unadjusted difference was consistent for Whites (+11 %), African Americans (+11 %), Asians (+16 %) and Native Americans (+34 %).

Logistic regression models examining sex and race differences in receipt of mental health treatment from the jail psychiatric unit, the jail infirmary, or the composite of the two, while controlling for covariates, are presented in Table 3. Across all models, individuals were more likely to receive services if they were older ($p < .01$), had longer jail stays ($p < .01$), had received more community-based outpatient visits prior to jail entry ($p < .05$), and had more state psychiatric hospital costs prior to jail entry ($p < .01$).

Controlling for these differences, the average marginal effects revealed a number of small, but statistically significant differences by race and gender. Across models, women were 4–8 % points less likely to receive mental health services as compared to men ($p < .01$). We also found that African Americans were 3–5 % points less likely to receive services across models than Whites ($p < .01$). Asians were 23 % points less likely than Whites to receive mental health services from the jail infirmary ($p < .01$) and 10 % points less likely than Whites to receive services on the composite service receipt measure ($p < .05$). Finally, we ran additional models interacting African-American race with sex gender (not presented in tables) and found that African-American men had a lower rate of mental health service receipt than White men both overall and in the jail psychiatric unit, but no differences were detected for women by race in any of the models.

Table 3 Logistic regression model results for probability of mental health service receipt in jail ($n = 2855$)

Variables	Stay in jail psychiatric unit (JPU)		Jail infirmary contact for mental health (JIC)		Service receipt from either JPU or JIC	
	β (SE)	AME	β (SE)	AME	β (SE)	AME
Female	-.43** (.09)	-.08** (.0)	-.30** (.09)	-.06** (.02)	-.26** (.09)	-.04** (.02)
Race (White = referent)						
African American	-.27* (.11)	-.05* (.02)	-.14 (.11)	-.03 (.02)	-.22* (.11)	-.04* (.02)
Native American	.13 (.27)	.02 (.05)	.31 (.27)	.06 (.06)	.29 (.29)	.05 (.05)
Asian	-.15 (.26)	-.03 (.05)	-1.13** (.27)	-.23** (.053)	-.58 (.27)	-.10* (.05)
Age	.12** (.030)	.002** (.0009)	.09** (.03)	.003** (.0009)	.11** (.031)	.004** (.0008)
Age squared	-.001** (.0004)		-.001** (.0004)		-.001** (.0004)	
Length of jail stay	.01** (.001)	.002** (.0002)	.02** (.001)	.003** (.0002)	.02** (.002)*	.003** (.0003)
Jail days squared	-9.31e-06* (1.56e-06)		-1.63e-05** (2.68e-06)		-1.53e-05** (1.87e-06)	
Outpatient mental health encounter days	.004* (.0006)	.0007** (.00008)	.004* (.0005)	5.38e-04** (7.69e-05)	.004** (.0006)*	.0006** (.00008)
Outpatient mental health encounter days squared	-3.07e-06** (6.60e-07)		-3.11e-06** (6.16e-07)		-3.17e-06** (7.02e-07)	
State psychiatric hospital costs	.03** (.005)	.006** (.0009)	.008** (.002)	.002** (.0004)	.03** (.006)	.005** (.001)
State psychiatric hospital costs squared	-4.45e-05** (8.09e-06)		-1.07e-05* (4.35e-06)		-4.37e-05** (9.63e-06)	

* $p \leq .05$; ** $p \leq .01$

Discussion

In the past, state mental hospitals functioned as the institutions of last resort for the care and confinement of persons with severe mental illness, but increasingly over the past three decades, this role has shifted throughout the US to local jails. Our results indicate that 70 % of jail detainees who had severe mental illness and a prior history of mental health service use in the community received mental health services in jail. This statistic underscores the many reports in the literature about jails, by default, having to assume mental health provider roles for their many detainees who have severe mental illness. It also suggests that this jail identified the majority, but not all, inmates with severe mental illness.

We also found small, but statistically significant sex and racial differences after adjusting for covariates. Women with severe mental illness were less likely than their male counterparts to receive mental health services while detained in jail. Further, this difference between males and females was consistent across racial groups. Women are a minority in

male-dominated jail populations and their particular needs are often not addressed [32–35]. We also observed racial differences in the locus of services receipt. African Americans differed from Whites in a lower likelihood of receiving jail psychiatric unit services whereas Asians differed from Whites in their greater receipt of infirmary versus psychiatric unit services. Most of these differences are smaller than the sex-race disparities observed in community settings suggesting that jail mental health services such as those in King County can mitigate some of the social inequities that occur in less controlled community settings.

Several limitations of this study need to be acknowledged. Our administrative dataset does not provide clinical information on the levels of functional disability associated with psychiatric diagnoses. Consequently, we were unable to determine whether detainees most in need of treatment actually received it. Also, we are unable to say whether the 30 % of the sample who did not receive treatment were in need of mental health treatment during their jail stay.

These limitations are partially offset by several strengths. It is rare in this research area to have studies based upon a large, multi-year, population-based sample of jail detainees with severe mental illness. Further, study participants' severe mental illness was established by chart diagnoses at community agencies prior to jail entry and thus was independent of any detection bias that might exist in identification by jail authorities [36].

Ideally, a large epidemiological study on a cohort of detainees using a measure of mental health status based upon psychiatric rating scales rather than chart diagnoses would be most useful in determining the extent of racial and gender disparities in who is identified and treated in jail or other criminal justice settings. Clearly, further research along these lines would provide a firmer foundation for intervention efforts to enhance necessary mental health care for detainees with severe mental illness while in jail as well as efforts to connect them to community services upon release.

It has long been recognized that the jail is a public health outpost where high-risk individuals are concentrated for brief periods of time and thus amenable to interventions that otherwise are difficult to implement given their mobile and chaotic lives in the community [37]. The findings from this study suggest that we need to think of the jail as a mental health outpost as well.

Despite a range of moral arguments that jail detention amounts to criminalization of the mentally ill [38, 39], jails today are de facto mental health way-stations for significant numbers of people with severe mental illness. As a result, more attention has to focus on efforts to prevent detention and foster successful community reentry after detention. Our data did not include the reentry phase for this study cohort but it is clear that community transition requires further collaboration and information sharing between community and correctional authorities. Continuity of care and treatment for very disabled individuals with severe mental illness has to occur both at the front door of the jail prior to entry and at its back door upon release to the community to interrupt the revolving door experience that many of these individuals encounter.

Numerous efforts have been directed at diversion of mentally ill detainees from jail with mixed success [38, 40, 41]. Part of the problem lies with the low quality and intensity of the community services to which diverted individuals are referred. Although virtually all (97 %) participants in this study received community mental health services prior to arrest and jail entry (Table 1), those treatment experiences did not deter their jail stays. The implication is that much more than simply connecting these individuals to routine community mental health services is needed to change their criminal justice trajectories. New approaches are needed to figure out how to keep people with severe mental illness out of

the criminal justice system and to provide appropriate services to those who end-up detained.

The expansion of Medicaid in many states as part of the Affordable Care Act holds great promise for improving these circumstances [42–45]. Large numbers of those released from the criminal justice system will be newly eligible for expanded Medicaid services [46]. With an influx of dollars from new Medicaid revenue streams both community and criminal justice authorities will have the opportunity to implement more targeted interventions aimed at improving the lives of the many thousands of individuals with severe mental illness who now repeatedly cycle through jails and other parts of the criminal justice system.

Conclusion

Urban jails end up serving large numbers of detainees with severe mental illness. Findings from this study indicate that small but statistically significant differences in who received in-jail mental health services occurred along sex and race lines but these differences are smaller than those often reported for receipt of community-based services. New efforts are needed to link-up these jails with community-based services both at the point of detention and release to promote continuity of care for the many thousands of persons with severe mental illness who are detained in these settings across the US.

Funding This research was supported by National Institute of Mental Health Grants MH063883, a grant from the John D. Catherine T. MacArthur Foundation Mental Health Policy Research Network, and by the Unc-Duke Postdoctoral Training Program in Mental Health Services Research under a National Research Services Award from the National Institute of Mental Health (T32 MH019117).

Compliance with Ethical Standards

Conflict of interest All authors declare that they have no conflicts of interest.

Ethical Approval All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. This study uses only secondary sources of data without an individual identifiers. A waiver of consent was obtained for this research.

References

1. Kessler RC, Berglund P, Demler O, Jin R, Merikangas KR, Walters EE: Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Archives of General Psychiatry*, 2005; doi:[10.1001/archpsyc.62.6.593](https://doi.org/10.1001/archpsyc.62.6.593).
2. Swanson J, Swartz M, Van Dorn RA, Monahan J, McGuire TG, Steadman HJ, et al.: Racial disparities in involuntary outpatient commitment: are they real? *Health Affairs (Project Hope)*, 2009; doi:[10.1377/hlthaff.28.3.816](https://doi.org/10.1377/hlthaff.28.3.816).
3. Whitley R, Lawson WB: The psychiatric rehabilitation of African Americans with severe mental illness. *Psychiatric Services*, 2010; doi:[10.1176/appi.ps.61.5.508](https://doi.org/10.1176/appi.ps.61.5.508).
4. Steadman HJ, Osher FC, Robbins PC, Case B, Samuels S: Prevalence of serious mental illness among jail inmates. *Psychiatric Services*, 2009; doi:[10.1176/ps.2009.60.6.761](https://doi.org/10.1176/ps.2009.60.6.761).
5. Binswanger IA, Merrill JO, Krueger PM, White MC, Booth RE, Elmore JG: Gender differences in chronic medical, psychiatric, and substance-dependence disorders among jail inmates. *American Journal of Public Health*, 2010; doi:[10.2105/AJPH.2008.149591](https://doi.org/10.2105/AJPH.2008.149591).
6. Cropsey KL, Binswanger IA, Clark CB, Taxman FS: The unmet medical needs of correctional populations in the United States. *Journal of the National Medical Association* 104(11–12):487–92, 2012.

7. James D, Glaze, L: Mental health problems of prison and jail inmates. In: Bureau of Justice Statistics Special Report, 2006. <http://www.bjs.ojp.usdoj.gov/content/pub/pdf/mhppji.pdf>.
8. Karberg J, James, D: Substance dependence, abuse, and treatment of jail inmates, 2002. In: Bureau of Justice Statistics Special Report, 2005. <http://www.bjs.gov/content/pub/pdf/sdatji02.pdf>.
9. Hatcher SS, Toldson IA, Godette DC, Richardson JB, Jr: Mental health, substance abuse, and HIV disparities in correctional settings: practice and policy implications for African Americans. *Journal of Health care for the Poor and Underserved* 20(2 Suppl):6–16, 2009.
10. Lurigio A: Examining prevailing beliefs about people with serious mental illness in the criminal justice system. *Federal Probation* 75(1), 2011.
11. Pettit B, Western B: Mass imprisonment and the life course: race and class inequality in US incarceration. *American Sociological Review*, 2004; doi:10.1177/000312240406900201.
12. Binswanger IA, Krueger PM, Steiner JF: Prevalence of chronic medical conditions among jail and prison inmates in the USA compared with the general population. *Journal of Epidemiology and Community Health*, 2009; doi:10.1136/jech.2009.090662.
13. Minton TD: Jail inmates at midyear 2011—statistical tables. Washington, DC, US Department of Justice, 2012. <http://www.bjs.gov/content/pub/pdf/jim11st.pdf>.
14. Minton TD: Jail inmates at midyear 2012—statistical tables. Washington, DC, US Department of Justice, 2013. <http://www.bjs.gov/content/pub/pdf/jim12st.pdf>.
15. Safran MA, Mays RA, Jr, Huang LN, McCuan R, Pham PK, Fisher SK, et al.: Mental health disparities. *American Journal of Public Health*, 2009; doi:10.2105/AJPH.2009.167346.
16. Lamb HR, Weinberger LE, Marsh JS, Gross BH: Treatment prospects for persons with severe mental illness in an urban county jail. *Psychiatric Services*, 2007; doi:10.1176/ps.2007.58.6.782.
17. Freudenberg N: Jails, prisons, and the health of urban populations: a review of the impact of the correctional system on community health. *Journal of Urban Health*, 2001; doi:10.1093/jurban/78.2.214.
18. Lamb HR, Weinberger LE: Persons with severe mental illness in jails and prisons: a review. *Psychiatric Services*, 1998; doi:10.1176/ps.49.4.483.
19. Primm AB, Osher FC, Gomez MB: Race and ethnicity, mental health services and cultural competence in the criminal justice system: are we ready to change? *Community Mental Health Journal*, 2005; doi:10.1007/s10597-005-6361-3.
20. Watson R, Stimpson A, Hostick T: Prison health care: a review of the literature. *International Journal of Nursing Studies*, 2004; doi:10.1016/S0020-7489(03)00128-7.
21. Cook BL, McGuire TG, Lock K, Zaslavsky AM: Comparing methods of racial and ethnic disparities measurement across different settings of mental health care. *Health Services Research*, 2010; doi:10.1111/j.1475-6773.2010.01100.x.
22. Cook BL, Doksum T, Chen CN, Carle A, Alegria M: The role of provider supply and organization in reducing racial/ethnic disparities in mental health care in the US. *Social Science and Medicine*, 2013; doi:10.1016/j.socscimed.2013.02.006.
23. Hankerson SH, Fenton MC, Geier TJ, Keyes KM, Weissman MM, Hasin DS: Racial differences in symptoms, comorbidity, and treatment for major depressive disorder among black and white adults. *Journal of the National Medical Association* 103(7):576–84, 2011.
24. Samnaliev M, McGovern MP, Clark RE: Racial/ethnic disparities in mental health treatment in six Medicaid programs. *Journal of Health Care for the Poor and Underserved*, 2009; doi:10.1353/hpu.0.0125.
25. Stockdale SE, Lagomasino IT, Siddique J, McGuire T, Miranda J: Racial and ethnic disparities in detection and treatment of depression and anxiety among psychiatric and primary health care visits, 1995–2005. *Medical Care*, 2008; doi:10.1097/MLR.0b013e3181789496.
26. Patterson EJ: Incarcerating death: mortality in US state correctional facilities, 1985–1998. *Demography* 47(3):587–607, 2010.
27. Rosen DL, Wohl DA, Schoenbach VJ: All-cause and cause-specific mortality among black and white North Carolina state prisoners, 1995–2005. *Annals of Epidemiology*, 2011; doi:10.1016/j.annepidem.2011.04.007.
28. Spaulding AC, Seals RM, McCallum VA, Perez SD, Brzozowski AK, Steenland NK: Prisoner survival inside and outside of the institution: implications for health-care planning. *American Journal of Epidemiology*, 2011; doi:10.1093/aje/kwq422.
29. Beck A, Karberg, J: Prison and jail inmates at midyear 2000. Washington, DC, US Department of Justice, Bureau of Justice Statistics, 2001.
30. Morrissey JP, Steadman HJ, Dalton KM, Cuellar A, Stiles P, Cuddeback GS: Medicaid enrollment and mental health service use following release of jail detainees with severe mental illness. *Psychiatric Services*, 2006; doi:10.1176/appi.ps.57.6.809.

31. Morrissey JP, Cuddeback GS, Cuellar AE, Steadman HJ: The role of Medicaid enrollment and outpatient service use in jail recidivism among persons with severe mental illness. *Psychiatric Services*, 2007; doi:[10.1176/ps.2007.58.6.794](https://doi.org/10.1176/ps.2007.58.6.794).
32. Bloom BE, Covington S: Addressing the mental health needs of women offenders. In: Gido RL, Dalley LP (Eds): *Women's mental health issues across the criminal justice system*. Upper Saddle River, NJ, Prentice Hall, pp. 160–76, 2008.
33. Pimlott Kubiak S, Beeble M: Assessing the mental health of female jail detainees [Abstract]. Annual meeting of the SCRA Biennial Meeting; June 18, 2009; Montclair State University, Montclair, NJ2009.
34. Steadman HJ, Scott JE, Osher F, Agnese TK, Robbins PC: Validation of the brief jail mental health screen. *Psychiatric Services*, 2005; doi:[10.1176/appi.ps.56.7.816](https://doi.org/10.1176/appi.ps.56.7.816).
35. van den Bergh BJ, Moller LF, Hayton P: Women's health in prisons: it is time to correct gender insensitivity and social injustice. *Public Health*, 2010; doi:[10.1016/j.puhe.2010.08.006](https://doi.org/10.1016/j.puhe.2010.08.006).
36. Steadman HJ, Robbins PC, Islam T, Osher FC: Revalidating the brief jail mental health screen to increase accuracy for women. *Psychiatric Services*, 2007; doi:[10.1176/ps.2007.58.12.1598](https://doi.org/10.1176/ps.2007.58.12.1598).
37. Glaser JB, Greifinger RB: Correctional health care: a public health opportunity. *Annals of Internal Medicine* 118(2):139–45, 1993.
38. Steadman HJ: Reducing the involvement of persons with mental illness and co-occurring disorder in the criminal justice system through jail diversion programs. *Criminalization of mental illness*; 2013; USC Gould School of Law, University of Southern California, Los Angeles, CA. Los Angeles, CA, Saks Institute for Mental Health Law, Policy and Ethics, 2013.
39. Torrey EF, Kennard AD, Eslinger D, Lamb R, Pavle J: More mentally ill persons are in jails and prisons than hospitals: a survey of the states. Arlington, VA, Treatment Advocacy Center, 2010.
40. Sirotych F: The criminal justice outcomes of jail diversion programs for persons with mental illness: a review of the evidence. *Journal of the American Academy of Psychiatry and the Law*, 37(4):461–472, 2009.
41. Steadman HJ, Naples M: Assessing the effectiveness of jail diversion programs for persons with serious mental illness and co-occurring substance use disorders. *Behavioral Sciences and the Law*, 2005; doi:[10.1002/bsl.640](https://doi.org/10.1002/bsl.640).
42. States implement health reform: Medicaid and the Affordable Care Act. National Conference of State Legislatures, 2011; Washington, DC. Washington, DC, National Conference of State Legislatures, 2011.
43. Dey J, Rosenoff, E, West, K, Ali, M, Lynch, S, McClellan, C, Mutter, R, Patton, L, Teich, J, Woodward, A: Benefits of Medicaid expansion for behavioral health. ASPE Issue Brief. Washington, DC, Department of Health and Human Services, Office of Assistant Secretary for Planning and Evaluation, 2016.
44. Mechanic D: Seizing opportunities under the Affordable Care Act for transforming the mental and behavioral health system. *Health Affair (Project Hope)*, 2012; doi:[10.1377/hlthaff.2011.0623](https://doi.org/10.1377/hlthaff.2011.0623).
45. Mechanic D, Olsson M: The relevance of the Affordable Care Act for improving mental health care. *Annual Review of Clinical Psychology*, 2016; doi:[10.1146/annurev-clinpsy-021815-092936](https://doi.org/10.1146/annurev-clinpsy-021815-092936).
46. Cuellar AE, Cheema J: As roughly 700,000 prisoners are released annually, about half will gain health coverage and care under federal laws. *Health Affair (Project Hope)*, 2012; doi:[10.1377/hlthaff.2011.0501](https://doi.org/10.1377/hlthaff.2011.0501).

Sean K. Sayers is a Senior Psychologist I at the NCDHHS—Disability Determination Services and an Adjunct Professor with the Ph.D. Clinical Psychology Program at Carlos Albizu University, San Juan Campus, Puerto Rico. He was a Postdoctoral Fellow at UNC when this work was conducted. He received his Ph.D. in Clinical Psychology from Carlos Albizu University, San Juan Campus, Puerto Rico.

Marisa E. Domino is a Professor in the Department of Health Policy and Management at The Gillings School of Global Public Health at the University of North Carolina at Chapel Hill and the Director of the Program on Mental Health and Substance Abuse Systems and Services Research at the Cecil G. Sheps Center for Health Services Research. She received her PhD from the Johns Hopkins University.

Gary S. Cuddeback is an Associate Professor in School of Social Work at the University of North Carolina at Chapel Hill and a Research Fellow at the Cecil G. Sheps Center for Health Services Research. He received his PhD from the University of Tennessee.

Nadine J. Barrett is the Director of the Office of Health Equity and Disparities at DCI and the Director of the Duke CTSA Community Connections and Collaborations Core at Duke University. She received her PhD from Texas Women's University and is a former Postdoctoral Fellow at UNC.

Joseph P. Morrissey is a Professor in the Department of Health Policy and Management at The Gillings School of Global Public Health at the University of North Carolina at Chapel Hill and a Research Fellow at the Cecil G. Sheps Center for Health Services Research. He received his PhD in Sociology from the University of North Carolina.