



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

23rd International Grassland Congress

Economics of BN Hybrid Grass Production Bundelkhand Region of India: A Case Study

Satendra Kumar

Indian Grassland and Fodder Research Institute, India

Ravi Pratap Singh

Indian Grassland and Fodder Research Institute, India

Rajiv K. Agrawal

Indian Grassland and Fodder Research Institute, India

J. B. Singh

Indian Grassland and Fodder Research Institute, India

M. M. Das

Indian Grassland and Fodder Research Institute, India

See next page for additional authors

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/23/2-8-1/7>

The 23rd International Grassland Congress (Sustainable use of Grassland Resources for Forage Production, Biodiversity and Environmental Protection) took place in New Delhi, India from November 20 through November 24, 2015.

Proceedings Editors: M. M. Roy, D. R. Malaviya, V. K. Yadav, Tejveer Singh, R. P. Sah, D. Vijay, and A. Radhakrishna

Published by Range Management Society of India

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Presenter Information

Satendra Kumar, Ravi Pratap Singh, Rajiv K. Agrawal, J. B. Singh, M. M. Das, and Sunil Kumar

Paper ID: 1369

Theme 2. Grassland production and utilization

Sub-theme 2.8. Grassland-market linkage

Economics of BN Hybrid grass production Bundelkhand region of India: A case study.

Satendra Kumar*, Ravi Pratap Singh, R. K. Agrawal, J. B. Singh, M. M. Das, Sunil Kumar

ICAR-Indian Grassland and Fodder Research Institute, Jhansi-284003, India

*Corresponding author e-mail: stndkumar543@gmail.com

Keywords: Drought, Harvesting, Napier Bajra Hybrid, Net return, Rooted slip

Introduction

Napier grass is also called as Elephant Grass due to its tallness and vigorous vegetative growth. The plants tiller freely and a single clump may produce more than 60 tillers under favorable climatic and soil conditions. Unfortunately, the grass coarse-textured, the leaf blade sheaths as well as sharply serrated, leaf margins. Therefore, cross was made between Bajra which is more succulent, leafy, fine-textured, palatable, fast growing and drought resistant and Napier to combine these qualities with its high yielding potential. The outcome Hybrid Napier is a perennial grass which can be retained on field for 2-3 years. Hybrid napier grass are mostly grown under assured water supply but cultivating under varying agro-climatic condition is also possible. Low grass production in dry land area is mainly due to the limited availability of soil moisture and plant nutrients. Agrawal *et al.*, (2001) reported NB hybrid is superior to guinea grass or *setaria* grass. However the economy of production plays an important role in introduction/ adoption of a crop in an area, village or farm. Many earlier studies have been conducted on economics of BN Hybrid production at research farm, or other government farms. But the information on economics of BN hybrid production at farmer's field and its integration livestock feeding as well rooted slips are limited. Therefore, the present study was conducted with the object to study the economics of BN hybrid at farmer's field in Bundelkhand.

Materials and Methods

Field study was conducted at 4 farmers field during July 2013 to June 2015, at Kadesara Kalan village of, Talbehat, Lalitpur district (24° 11' to 25° 14' N and 78° 10' to 79° 00') of Bundelkhand region. The grass was sole planted in 0.1 to 0.2 hectare areas at 75 cms apart rows. The intra row spacing was kept at 60 cm. Planting was done manually after commencement of rains in the month of July. The basal dose of 60 kg N and 40 kg P₂O₅/ha was given besides 4000 kg FYM/ha (20-25 days prior to planting). First cut was taken at 65 days after planting. One manual weeding was done in the month of August and July in respective years. The standard irrigation schedules and other management practices were followed as per recommendation (Kumar *et al.*, 2012). In first year seven and in second year eight cuts were taken manually with the sickle. The need for renovation or dressing of tussock was not realized as tussocks were pruned regularly to uproot and sale the rooted slips. Rooted slips were sold twice in a year.

Table 1: Average cost of production (four farmers)

Agricultural operations	Cost (Rs.)
Field Preparation	3,300
Plantation material and plantation	24,000
Fertilizer cost + FYM	5,910
Weed management	6,000
Irrigation	16,000
Other cost	5,000
Total Input	60,210
Output Income	
Return from green fodder	1,63,540
Return from rooted slip sale	18,937
Total output	1,82,477
Net return	1,22,268

Table 2: Economic returns

Farmer's Name	Area (ha)	Green fodder yield (q/ha)			Rooted slips sold	Return (Rs./ha)				B: C Ratio
		2013-14	2014-15	Total		GFY	Rooted slips	Gross	Net	
Sh. Hariram	0.20	1011	821	1832	20000	146560	15000	161560	101350	1.68
Sh. Jasrath Ahirwar	0.20	1402	1125	2527	31000	202160	23250	225410	165200	2.74
Sh. Dhaniram	0.10	1024	877	1901	28000	152080	21000	173080	112870	1.87
Md. Siddiqui	0.10	1055	862	1917	22000	153360	16500	169860	109650	1.82
Average	0.15	1123	921	2044	25250	163540	18938	182478	122268	




Fig. 1: Farmer's Gross and Net return.

Results and Discussion

On the basis of study Sh. Jasrath Ahirwar got maximum total return (Rs. 1, 65,200) followed by Sh. Dhaniram (Rs. 1,12,870), Md. Siddiqui, (Rs.1,09,650) and Sh. Hariram (Rs.1,01,350). The average cost of cultivation was recorded Rs 60,210 per hectare (Table 1) and net return was Rs. 1, 22,268 per hectare (Table 2). The highest B: C ratio was recorded at the farm of Sh. Jasrath Ahirwar.

Conclusion

The cultivation of BN Hybrid for green fodder and rooted slips is highly profitable at farmers' field in Kadesara Kalan of Bundelkhand region of India.

References

- Agrawal, R. K., K. K. Singh and K. C. Sharma. 2001. Effect of split application of nitrogen on yield and quality of perennial grasses. *Indian Journal of Animal Nutrition* 18 (3): 267-270.
- Kumar, S., R. K. Agrawal, A. K. Dixit, A. K. Rai, J. B. Singh and S. K. Rai. 2012. *Forage Production Technology for Arable Lands*. Technical Bulletin. IGFR, Jhansi