

University of Louisville

ThinkIR: The University of Louisville's Institutional Repository

Undergraduate Arts and Research Showcase

Undergraduate Research

Spring 2020

GLYCINE INHIBITION OF ON-OFF DIRECTIONALLY SELECTIVE RETINAL GANGLION CELLS

Madeline C Mitchell
madeline.mitchell@louisville.edu

Maureen McCall Ph.D
maureen.mccall@louisville.edu

Follow this and additional works at: <https://ir.library.louisville.edu/uars>



Part of the [Neuroscience and Neurobiology Commons](#)

Recommended Citation

Mitchell, Madeline C and McCall, Maureen Ph.D, "GLYCINE INHIBITION OF ON-OFF DIRECTIONALLY SELECTIVE RETINAL GANGLION CELLS" (2020). *Undergraduate Arts and Research Showcase*. 24. <https://ir.library.louisville.edu/uars/24>

This Book is brought to you for free and open access by the Undergraduate Research at ThinkIR: The University of Louisville's Institutional Repository. It has been accepted for inclusion in Undergraduate Arts and Research Showcase by an authorized administrator of ThinkIR: The University of Louisville's Institutional Repository. For more information, please contact thinkir@louisville.edu.

GLYCINE INHIBITION OF ON-OFF DIRECTIONALLY SELECTIVE RETINAL GANGLION CELLS

Madeline Mitchell and Maureen McCall Ph.D

Department of Anatomical Sciences and Neurobiology, University of Louisville

Abstract

Retinal ganglion cells (RGCs) are a group of over 40 different types that use both GABAergic and Glycine inhibition. There are four different types of Glycine receptors (GlyRs) known as GlyR α 1, GlyR α 2, GlyR α 3, and GlyR α 4 (Sanes & Masland, 2015; C. Zhang & McCall, 2012). Electrophysiological data has supported the theory that the RGC cell type On-Off directionally selective (ooDS) cells express both GlyR α 4 and GlyR α 2. If ooDS cells do express only GlyR α 4 and GlyR α 2, an immunohistochemical analysis of these cells should support this theory. By comparing the expression of GlyR α 4 and GlyR α 2 in Glra4 $^{-/-}$, Glra2 $^{-/-}$, Glra4 $^{-/-}$, and Glra2 $^{-/-}$ the expression we can support or weaken the findings made by previous electrophysiological data. Comparing cell images of colocalized dendrites and GlyR α 4 or GlyR α 2 in an original and randomized orientation can determine if an immunohistochemical analysis confirms the presence of GlyR α 4 and GlyR α 2 on ooDS RGCs. Our findings confirmed that both GlyR α 4 and GlyR α 2 are expressed on ooDS RGCs.

Introduction

Retinal Ganglion cells (RGCs) are responsible for receiving information from the bipolar and amacrine cells and transporting this information to the brain (Sanes & Masland, 2015; C. Zhang & McCall, 2012). Ganglion cells have been grouped morphologically by their dendritic ramification patterns in the inner plexiform layer ON and OFF sublaminae, soma diameter, dendritic field size, and branch patterning (C. Zhang & McCall, 2012). There are around 40 RGC types and each utilizes both GABA and glycine receptors. (Sanes & Masland, 2015; C. Zhang, Nobles, & McCall, 2015). Glycine receptors are pentameric ligand-gated chloride ion channels. A typical glycine receptor contains one type of GlyR β subunit and one of four different types of GlyR α -subunits. These are known as GlyR α 1, GlyR α 2, GlyR α 3, and GlyR α 4 (Weiss et al., 2008). As a whole, glycinergic inhibition modulates the light responses of retinal neurons and plays a role in creating specificity, preferentially by crossover inhibition (Wassle et al., 2009). The pattern of expression of each of the GlyRs differs across the inner plexiform layer (IPL) and there is a significant amount of GlyR α subunit diversity across currently identified RGCs (Nobles et al., 2012; Wassle et al., 2009). WT ON/OFF direction selective (ooDS) cells respond preferentially to movement in a preferred direction. There are four types of ooDS cells, each encoding for motion in one of the four cardinal directions (Im & Shelley, 2016). OoDS cells are one of the most common RGCs and represent about 20% of all RGCs in the mouse retina (Sanes & Masland, 2015). Current electrophysiological data has indicated that ooDS RGCs should express GlyR α 2 & GlyR α 4 (C. Zhang et al., 2014; Haverkamp et al., 2003; Haverkamp et al., 2004). By staining for specific GlyRs, we should find WT ON/OFF direction selective (DS) RGCs express GlyR α 2 & GlyR α 4 as indicated by electrophysiological data. By comparing the immunohistochemical analysis of GlyRs ON/OFF directional selective (DS) we can strengthen or weaken findings made previously using electrophysiological data.

Table 1: Primary and Secondary Antibodies used for Immunohistochemistry

Primary Antibodies	Concentration	Host	Manufacturer
anti-Lucifer Yellow	1:1000	rabbit polyclonal	Life Technologies
anti-GlyR α 2	1:50	goat polyclonal	Santa Cruz Biotech
anti-GlyR α 4	1:100	rabbit polyclonal	Chemicon
Secondary Antibodies	Concentration	Host	Manufacturer
anti-goat IgG Cy3	1:200	donkey polyclonal	Jackson ImmunoResearch
anti-rabbit IgG 488	1:200	donkey polyclonal	Jackson ImmunoResearch
streptavidin Cy2 or 633	1:200	conjugate	Life Technologies
Hoechst nuclear stain	1:1000	conjugate	Life Technologies

Methods and Materials

Animals

For this experiment mice of the TRHR and PVCre/Thy1STP mouse lines were used. In the TRHR and PVCre/Thy1STP mouse lines, some ON/OFF DS ganglion cells express yellow fluorescent protein (YFP), which glows green under the right fluorescent light. This makes the cells easier to find and target. TRHR and PVCre/Thy1STP mice were crossed and backcrossed to Glra2 $^{-/-}$, Glra4 $^{-/-}$ and Glra2 $^{-/-}$ 4 $^{-/-}$ to obtain GlyR global knockout models along with labeled RGCs. All experiments were carried out in accordance with the Association for Research in Vision and Ophthalmology (ARVO) Statement for the Use of Animals in Ophthalmic and Visual Research and with the approval of the Institutional Animal Care and Use Committee (IACUC) of the University of Louisville.

Immunohistochemistry

RGCs from pieces of mouse retina were filled with Lucifer Yellow and neurobiotin, fixed in 4% paraformaldehyde for 12 minutes, washed three times with 0.01 M phosphate buffer saline (PBS), and placed into a 24-well plate. For 60 minutes the retina pieces were incubated in blocking solution consisting of PBX (0.5% Triton-X 100 in PBS) and 10% normal donkey or goat serum. Each retina piece was then reacted with a combination of primary antibodies to stain for the either GlyR α 2, GlyR α 4, or a combination of the two GlyRs, and Lucifer yellow overnight at 4°C (Table 1). The primary antibodies were washed off by three washes with PBX. A combination of secondary antibodies and Hoechst stain in normal serum was added to label the primary antibodies and it was left on the tissue overnight at 4°C. The Hoechst stain was used as a label for DNA in the somas and distinguish between the retina nuclear and plexiform layers. The tissue was washed with PBS one last time and mounted onto a slide and covered using VECTASHIELD (Vector Labs, Burlingame, CA) clear mounting medium and a coverslip.

Confocal Imagery and Colocalization Analysis

The filled RGCs were imaged, using an Olympus Fluoview FV1000 confocal microscope with Fluoview software (Olympus, Tokyo, Japan) with a 60x oil immersion objective (NA 1.4). Depending on the size of the dendritic field 3-5 images were taken of each cell, taking care to capture as much of the dendritic field as possible without capturing the soma. Images of ooDS RGC dendrites and GlyR puncta were deconvolved, using constrained iterative deconvolution in cellSens (Olympus, Tokyo, Japan). Using the Imaris (Bitplane, Zurich, Switzerland) surfaces and distance transformation tools, coincident GlyR puncta (color channel 1) on or within the dendritic processes (color channel 2) were counted. The length of the dendritic processes within the image was measured using the filaments tool. The coincident puncta divided by the dendritic length estimated puncta density.

To determine which puncta were colocalized onto the dendrite, the puncta channel (channel 1) was flipped along the x axis and on the y axis and coincident GlyR puncta were recounted. Then, the flipped or “random” (RAN) puncta density was subtracted from the “original” (ORI) puncta density and the result estimated the “corrected” puncta density.

The ORI and RAN puncta density counts were compared using a paired one-tailed t-test.

Results

OoDS RGCs Express GlyR α 4-Positive Puncta

To assess the expression of GlyR α 4 positive puncta on the ooDS RGC dendrites we compared the original puncta density to an estimate of random puncta coincidence. The average ooDS GlyR α 4-positive puncta density was 1.05 ± 0.31 puncta $\cdot\mu\text{m}^{-1}$, which was significantly more than the random GlyR α 4-positive puncta density, which was 0.82 ± 0.24 puncta $\cdot\mu\text{m}^{-1}$ (Figure 1) $p < 0.05$ (paired one-tailed T-test). When corrected for our estimate of randomly associated GlyR α 4-positive puncta, the corrected density of GlyR α 4 puncta expression on ooDS RGCs was 0.23 ± 0.04 puncta $\cdot\mu\text{m}^{-1}$. This corrected GlyR α 4-positive puncta value suggest positive expression of GlyR α 4 on ooDS RGCs.




Figure 1: OoDS RGCs Express GlyR α 4: A) Dendrites on ooDS RGCs have a significantly higher GlyR α 4 puncta density co-localization than coincidence co-localization in the flipped (RAN) orientation. B) GlyR α 4 puncta on ooDS RGC dendrites in original orientation. C) GlyR α 4 puncta on ooDS RGC dendrites in randomized orientation.

OoDS RGCs Express GlyR α 2-Positive Puncta

We also assessed GlyR α 2 positive puncta on the ooDS RGC dendrites and compared the original density to an estimate of random puncta coincidence. The average ooDS RGC GlyR α 2-positive puncta density was 0.92 ± 0.73 puncta $\cdot\mu\text{m}^{-1}$. This was significantly more than the random average GlyR α 2-positive puncta density, which was 0.72 ± 0.69 puncta $\cdot\mu\text{m}^{-1}$ (Figure 2) $p < 0.05$ (paired one-tailed T-test). When corrected for our estimate of randomly associated GlyR α 2-positive puncta, the corrected density of GlyR α 2 puncta expression on ooDS RGCs was 0.21 ± 0.05 puncta $\cdot\mu\text{m}^{-1}$. The similarity between the GlyR α 2-positive puncta and chance puncta coincidence suggests GlyR α 2 expression is present ooDS RGCs.




Figure 2: OoDS RGCs Express GlyR α 2: A) Dendrites on ooDS RGCs have a significantly higher GlyR α 2 puncta density co-localization than coincidence co-localization in the flipped (RAN) orientation. Min and max lines not available due to small sample size B) GlyR α 2 puncta on ooDS RGC dendrites in original orientation. C) GlyR α 2 puncta on ooDS RGC dendrites in randomized orientation.

Discussion

The established immunohistochemical approach we used to identify GlyR puncta expression was able to positively identify GlyR α 2 and GlyR α 4 on ooDS RGCs. This supports the conclusion that GlyR α 2 and GlyR α 4 are both expressed on ooDS RGCs. Our immunohistochemical data combined with previous electrophysiological data provides strong support of the theory that ooDS RGCs that these cells express GlyR α 4 and GlyR α 2.

The expression of GlyR α 4 and GlyR α 2 on ooDS RGCs leads us to believe that it is likely that GlyR α 4 and GlyR α 2 play an important role in modulated inhibition in ooDS RGCs.

Only 3 cell images were used to determine GlyR α 2 expression and 5 cell images were used to determine GlyR α 4 expression. While the number of cell images we were able to incorporate into our experiment is not ideal, we were limited in our ability to gather more data by the COVID-19 pandemic. Repeating the experiment with a larger sample size would increase our confidence in the results of the experiment.

References

- Haverkamp, S. (1995). Glycine Receptor Diversity in the Mammalian Retina. In H. Kolb, E. Fernandez, & R. Nelson (Eds.), *Webvision: The Organization of the Retina and Visual System*. Salt Lake City (UT).
- Haverkamp, S., Muller, U., Harvey, K., Harvey, R. J., Betz, H., & Wassle, H. (2003). Diversity of glycine receptors in the mouse retina: localization of the alpha3 subunit. *J Comp Neurol*, 465(4), 524-539. doi:10.1002/cne.10852
- Haverkamp, S., Muller, U., Zeilhofer, H. U., Harvey, R. J., & Wassle, H. (2004). Diversity of glycine receptors in the mouse retina: localization of the alpha2 subunit. *J Comp Neurol*, 477(4), 399-411. doi:10.1002/cne.20267
- Im, M., Fried, S. Directionally selective retinal ganglion cells suppress luminance responses during natural viewing. *Sci Rep* 6, 35708 (2016). <https://doi.org/10.1038/srep35708>
- Masland, R. H. (2012). The neuronal organization of the retina. *Neuron*, 76(2), 266-280. doi:10.1016/j.neuron.2012.10.002
- Nobles, R. D., Zhang, C., Muller, U., Betz, H., & McCall, M. A. (2012). Selective glycine receptor alpha2 subunit control of crossover inhibition between the on and off retinal pathways. *J Neurosci*, 32(10), 3321-3332. doi:10.1523/JNEUROSCI.5341-11.2012
- Sanes, J. R., & Masland, R. H. (2015). The types of retinal ganglion cells: current status and implications for neuronal classification. *Annu Rev Neurosci*, 38, 221-246. doi:10.1146/annurev-neuro-071714-034120
- Wassle, H., Heinze, L., Ivanova, E., Majumdar, S., Weiss, J., Harvey, R. J., & Haverkamp, S. (2009). Glycinergic transmission in the Mammalian retina. *Front Mol Neurosci*, 2, 6. doi:10.3389/neuro.02.006.2009
- Weiss, J., O'Sullivan, G. A., Heinze, L., Chen, H. X., Betz, H., & Wassle, H. (2008). Glycinergic input of small-field amacrine cells in the retinas of wildtype and glycine receptor deficient mice. *Mol Cell Neurosci*, 37(1), 40-55. doi:10.1016/j.mcn.2007.08.012
- Zhang, C., & McCall, M. A. (2012). Receptor targets of amacrine cells. *Vis Neurosci*, 29(1), 11-29. doi:10.1017/S0952523812000028
- Zhang, C., Nobles, R. D., & McCall, M. A. (2015). GlyR α 2, not GlyR α 3, Modulates the Receptive Field Surround of OFF Retinal Ganglion Cells. *Vis Neurosci*. doi:10.1017/S0952523815000280
- Zhang, C., Rompani, S. B., Roska, B., & McCall, M. A. (2014). Adeno-associated virus-RNAi of GlyR α 1 and characterization of its synapse-specific inhibition in OFF alpha transient retinal ganglion cells. *J Neurophysiol*, 112(12), 3125-3137. doi:10.1152/jn.00505.2014