


On: 28 April 2015, At: 10:36

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Critical Reviews in Food Science and Nutrition

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/bfsn20>

The glycaemic index of rice and rice products: a review, and table of GI values

Bhupinder Kaur^a, Viren Ranawana^a & C. Jeya. K. Henry^a

^a Clinical Nutrition Research Centre, Singapore Institute for Clinical Sciences, 30 Medical Drive, Singapore 117609, Singapore

Accepted author version posted online: 15 Jan 2015.



[Click for updates](#)

To cite this article: Bhupinder Kaur, Viren Ranawana & C. Jeya. K. Henry (2015): The glycaemic index of rice and rice products: a review, and table of GI values, *Critical Reviews in Food Science and Nutrition*, DOI: [10.1080/10408398.2012.717976](https://doi.org/10.1080/10408398.2012.717976)

To link to this article: <http://dx.doi.org/10.1080/10408398.2012.717976>

Disclaimer: This is a version of an unedited manuscript that has been accepted for publication. As a service to authors and researchers we are providing this version of the accepted manuscript (AM). Copyediting, typesetting, and review of the resulting proof will be undertaken on this manuscript before final publication of the Version of Record (VoR). During production and pre-press, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal relate to this version also.

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

ACCEPTED MANUSCRIPT

The glycaemic index of rice and rice products: a review, and table of GI values

BHUPINDER KAUR, VIREN RANAWANA and C. JEYA. K. HENRY

Clinical Nutrition Research Centre, Singapore Institute for Clinical Sciences, 30 Medical Drive, Singapore 117609, Singapore.

**Address correspondence to Viren Ranawana, Clinical Nutrition Research Centre, Singapore Institute for Clinical Sciences, 30 Medical Drive, Singapore 117609, Singapore.
Tel: (+65) 6407-0741, Fax: (+65) 6778-4193, E-mail: viren_ranawana@sics.a-star.edu.sg**

ABSTRACT

Rice is the principle staple and energy source for nearly half the world's population and therefore has significant nutrition and health implications. Rice is generally considered a high glycaemic index (GI) food, however this depends on varietal, compositional, processing and accompaniment factors. Being a major contributor to the glycaemic load in rice eating populations, there is increasing concern that the rising prevalence of insulin resistance is as a result of the consumption of large amounts of rice. Devising ways and means of reducing the glycaemic impact of rice is therefore imperative. This review gathers studies examining the GI of rice and rice products and provides a critical overview of the current state of the art. A table collating published GI values for rice and rice products is also included.

Running head: GI of rice and rice products

Keywords: rice; rice products; glycaemic index; glycaemic response; diabetes

INTRODUCTION

The increasing prevalence of diabetes and related chronic diseases in recent times has prompted greater research attention into ways and means of curbing their further escalation. Dietary intervention methods which are the cornerstone of diabetes prevention and management are primarily aimed at maintaining a low and stable postprandial blood glucose concentration (Jenkins et al., 1984). The glycaemic index (GI) which is based on the glycaemic response is a well established indicator of the blood glucose raising potential of a carbohydrate food. Studies show that regular consumption of high-GI diets is associated with an increased risk for type 2 diabetes mellitus (Salmeron et al., 1997a; 1997b). There is evidence to suggest that low-GI diets reduce the incidence of diabetes (Bjorck et al., 1994; Augustin et al., 2002), hyperlipidaemia (Jenkins et al., 1987b) and cardiovascular disease (Murakami et al., 2006; Larsen et al., 2010; Ma et al., 2005; Ludwig et al., 1999; Ludwig et al., 2002). Although simple carbohydrates such as glucose and maltose, have been historically regarded as the greater inducers of hyperglycaemia, recent data conclusively shows that complex carbohydrates, including starches, are able to produce equal or even larger blood glucose excursions (Kalergis et al., 1998).

Rice (*Oryza sativa* L.) is the most widely eaten staple in the world with global consumption levels more than tripling from 156 million tons to 456 million tons between 1960 and 2010 (Figure 1) (IRRI, 2012). It is the main energy source for a large segment of the population; around 3 billion depend on rice for 35–59% of their caloric intake (Meng et al., 2005). Asia has the highest consumption of rice with intake exceeding 100 kg per capita in many countries

(IRRI, 2010). Whilst polished white rice is the most widely consumed form, unmilled brown rice is regularly eaten in some cultures.

White rice elicits a relatively large glycaemic response and is thus associated with exacerbating impaired glucose tolerance (Sun et al., 2010). It contributes a large glycaemic load (GL) to the diets of those in countries where it is the main staple due to both the large quantity eaten and its greater GI. These countries incidentally are also where diabetes rates are markedly increasing (Chan et al., 2009; Shaw et al., 2010). Therefore, devising ways and means of reducing the glycaemic response of rice is imperative to reduce the risk of developing the metabolic syndrome. A good understanding of all the factors affecting the GI of rice is essential to achieve this. Although a considerable number of studies have focused on the GI of rice and factors affecting it a systematic review of the state-of-the-art has not been carried out. The objective of this article therefore was to review the literature surrounding the GI of rice and factors affecting it.

REVIEW

Composition of rice

China, India and Indonesia are believed to be where rice was first cultivated, and thus the origin of the three varieties of rice – *japonica*, *javanica* and *indica* (Juliano, 1992). *Japonica* grains are short, roundish grains which do not shatter easily and contain 0-20% amylose. *Javanica* grains

(long grain), which are long, broad, and thick, contains 0-25% amylose and tend to remain intact after cooking. *Indica* grains may be short or long and are slender and somewhat flat, and contains a higher amylose content (23-31%).

White rice is produced from the polishing of brown rice. Polishing removes the pericarp, seed-coat, testa, aleurone layer and embryo which results in the loss of fat, protein, crude and neutral detergent fibre, ash, vitamins and polyphenols whilst increasing starch content (Shobana et al., 2011). Due to the removal of the bran layer, white rice has a lower total dietary fibre content (0.7-2%) than brown rice (3-4%) (Juliano, 1985). The energy content of brown rice is slightly higher than white rice due to lipids in the bran (Juliano, 2003).

Starch is the predominant macronutrient in milled rice. The rice starch granule is composed of 99% α -glucans which consists of two distinct glucose polymers, amylose and amylopectin (Takeda et al., 1993). Amylose is an essentially linear molecule of α -(1 \rightarrow 4)-linked D-glucopyranosyl units with a few branches. Amylopectin has a larger molecular weight and a highly branched structure consisting of straight chains of D-glucopyranosyl units linked by α -(1 \rightarrow 4) bonds to which branched chains attach via α -(1 \rightarrow 6) bonds (Hizukuri, 1986). Rice amylose has 2-4 chains with an average DP_n of 900 and a β -amylolysis limit of 73-87% whilst rice amylopectins have β -amylolysis limits of 56 to 59%, chain lengths of 19 to 22 glucose units and a DP_n of 5 000 to 15 000 glucose units (Hizukuri et al., 1989).

The health impact of rice

Recent studies suggest that the high consumption of rice accelerates the development of diabetes (Hu et al, 2012). A high dietary glycemic load (predominantly from rice) has been associated with increased risk of type 2 diabetes in Chinese and Japanese (Villegas et al., 2007; Murakami et al., 2006; Nanri et al., 2008) and Indian (Mohan et al., 2009) populations. White rice is considered a high GL food (of about 23) (Foster-Powell et al., 2002). However, this depends on the quantity of rice consumed, and it is not a high GL food when consumed at levels typically found in Western diets compared to Asian diets.

Frequent consumption of high GI and GL meals over time is associated with insulin resistance, β -cell dysfunction, dyslipidemia, obesity, breast cancer, endothelial dysfunction, cardiovascular disease and a reduction in cognitive functions (Ludwig, 2002; Venn & Green, 2007; Liu et al., 2000; Sieri et al., 2007; Nakshima et al., 2010; Ingwerseb et al., 2007). The relationship between rice consumption and diabetes is equivocal (Nanri et al., 2010; Villegas et al., 2007; Sun et al., 2010; Hodge et al., 2004). A recent meta analysis by Hu et al. (2012) found a significant relationship between white rice consumption and the risk of type 2 diabetes. The authors concluded that the association was stronger for Asian (Chinese and Japanese) populations than their Western counterparts which is probably due to the larger quantities eaten by the former group. Intervention studies also show a relationship between rice consumption and diabetes. A study in Japan found that the provision of white bread compared to boiled white rice to children at breakfast caused less glucose fluctuations and better cognitive function (Taki et al., 2010). This was attributed to white bread having a lower GI than boiled white rice. Another study

showed that substituting white rice with the same amount of low GI brown rice significantly lowered the risk of developing diabetes in U.S. men and women (Sun et al., 2010).

In summary observational studies suggest that rice increases the risk of diabetes and other chronic diseases. However, the evidence is equivocal, and associations in epidemiological studies do not always prove causality. Randomised controlled and intervention studies show that rice has a large impact both on glycaemia and diabetes indicators. They also suggest that brown rice may be a better alternative to white rice. However, the acceptability of brown rice is lower than of white rice especially in communities where it is not traditionally consumed (Kumar et al., 2011) which often becomes a stumbling block to its integration into diets.

The GI of rice and rice products

The impact of rice on health appears to be mainly due to the greater GL it exerts in rice eating populations. Traditionally rice constitutes the major proportion of the plate content in these populations and advocacy to eat less of it would not be a sustainable public health strategy. Therefore, lowering the GI of rice may be a better strategy to achieve a low GL. A large number of studies have examined the GI of different rices and these have demonstrated it to be a food with very diverse values. A recent systematic tabulation of GI values showed that rice GI ranged between 24-160 when white bread was used as the reference (Atkinson et al., 2008). This diversity highlights the importance of considering rice on an individual type and variety basis.

Factors affecting the GI of rice and rice products

Rice variety and starch content (amylose and amylopectin)

The GI of rice depends on the variety. Varietal effects on GI appear to be mainly mediated by its amylose content. Based on the amylose content, rice can be classified as waxy (0-2%), very low amylose (5-12 %), low amylose (12 – 20%), medium amylose (20-25%) and high amylose (25-33 %) rice (Juliano, 1992). Apparent amylose content correlates positively with water absorption and volume expansion during cooking and with the hardness of boiled rice (Juliano, 2003). The amylose content therefore influences the textural qualities of cooked rice and is an important consideration when selecting rice for specific applications (Table 1).

The amylose:amylopectin ratio effects processing properties and eating quality. Amylose is directly related to the hardness, whiteness and dullness of cooked rice, and volume expansion and water absorption during cooking. The texture of cooked rice has been closely correlated with the relative abundance of long chains in its amylopectin (Rani & Bhattacharya,1995). The molecular structure of amylose is tighter and more compact, thus less susceptible to breakdown than amylopectin whose structure is more vulnerable to digestion. Therefore, the amylose:amylopectin ratio determines starch digestion rates and is thus used to predict blood glucose and insulin responses to rice (Goddard et al., 1984; Juliano & Goddard, 1986; Miller et al., 1992; Frei et al., 2003; Hu et al., 2004).

Chung et al. (2011) recently studied the in vitro digestibility of rice starches differing in amylose content (Long-grain, Arborio, Calrose, Glutinous). Long-grain rice starch showed the highest amylose content (27%) while glutinous was the lowest (4%). Long-grain rice starch had the highest gelatinization temperature (GT), and the lowest rapidly digestible starch (RDS) content (39% compared to 71% in glutinous rice). High RDS containing food have greater GIs and cause more rapid changes in blood glucose (Thondre et al., 2010).

Goddard et al. (1984) investigated the effect of amylose and amylopectin content on glucose and insulin responses to long grain (Labelle with 25% amylose and 75-77% amylopectin), medium grain (Pecos with 14-17% amylose and 83-86% amylopectin) and sweet rice (Mochi gome with 100% amylopectin). Although the nutritional composition (protein, total lipid content, fibre) of the rices were similar the high-amylose rice showed lower initial responses and slower declines in both glucose and insulin levels. The greater levels of complexed starch lipids and lower levels of non-starch lipids in high amylose rices were suggested to contribute to its lower impact on glycaemia. Similarly Juliano & Goddard (1986) observed the lowest glucose and insulin response curves when high amylose (>24%) rice was consumed. Miller et al. (1992) determined the GI of white and brown variants of three commercial rice varieties (Doongara, Calrose and Pelde). Doongara, a high amylose rice (28% amylose) produced significantly lower GIs for both the white and brown types (Doongara white rice-64; Doongara brown rice-66). The other two medium amylose (20%) varieties, Calrose and Pelde had higher GI values (Calrose white rice-83; Calrose brown rice-87 and Pelde white rice-93; Pelde brown rice-76). But in contrast, Holt and Miller (1995) found that Doongara rice had a lower GI (54). Both studies on Doongara rice

had a similar cooking time (14 min) but the lower GI in Holt and Miller's study was probably as a result of storing the rice overnight at 4°C. Cooling has been shown to increase resistant starch (RS) content and thereby reduce the GI, and this is discussed later in the article. Panlasigui et al. (1991) examined three varieties of rice containing the same amount of amylose (27%) and found that they all had similar GI values (75, 78, 81).

Studies conducted on local rice varieties showed similar trends. Bangladeshi white rice varieties containing 27% and 28% amylose had GIs of 37 and 39 respectively (Larsen et al., 1996; 2000) whilst traditional and improved medium to high amylose Sri Lankan rice varieties showed GI values between 57 and 73 (Pathiraje et al., 2010). White basmati rice, a popular Indian rice with a medium amylose content (20-25%) (Bhattacharjee et al., 2002) has a GI around 50-58 (Holt & Brand Miller, 1995; Ranawana et al., 2009).

Glutinous rice generally has a high GI due to its high amylopectin content. Hu et al. (2004) and Frei et al. (2003) found that starch hydrolysis for waxy cultivars was more rapid and more complete than for high amylose rice cultivars. Hu et al. (2004) reported that Yunuo No. 1 waxy *indica* rice containing 1% and 0.8% amylose content was high GI (106 and 101 respectively) whilst ZF201 high amylose rice containing 27% and 26% amylose content was low GI (63 and 60 respectively). Several studies show that Thai glutinous rice has a GI of over 90 (Chan et al., 2001; Ranawana et al., 2009). Japanese Koshikari rice similarly showed high GI values (89- Sato et al., 2010; 80- Sugiyama et al., 2003). Frei et al. (2003) reported that waxy rice cultivars in the Philippines had GIs above 90 compared to non-waxy cultivars. The amylose content of

rice also affects the GI of processed products. Puffed rice cakes made with high amylose rice showed comparatively lower GI values in one study (Holt & Brand-Miller, 1995).

In summary the literature suggests that high amylose rice has notably lower GI values than the high amylopectin variants. They also produce more attenuated insulin responses. However, amylose content alone is not a good predictor of the glycaemic response and other factors (discussed later) also affect starch digestion rates and glycaemia.

Cooking

Rice is predominantly eaten in the boiled form (conventional boiling or pressure cooking) and is steamed on some occasion. The method and degree of cooking, and the volume of cooking water determines the extent of starch gelatinisation, and this in turn directly correlates with digestibility and the glycaemic response (Collings et al., 1981). One study showed that boiling rice for only 5 minutes produced less swelled and intact grains (producing a lower GI) compared to rice boiled for 15 minutes where the grains were swollen and split (producing a higher GI) (Wolever et al., 1986) (Table 3).

Ranawana et al (2009) showed that basmati rice cooked for longer times elicited greater glycaemic responses. White basmati rice cooked for 10 min had a GI of 50 (low) but basmati and wild rice, brown and white basmati rice cooked for 25 min was high GI. Similarly Al Mssallem et al (2011) found that rice cooked for longer had a higher GI when they compared the glycaemic

response of Hassawi and long grain white rice. Parastouei et al. (2011) showed that the GI of Iranian rice differed after cooking. Rice soaked for 35 min and then boiled for 10 min elicited a GI of 55 and rice that was boiled for 5-8 min and then simmered for 30 min had a GI of 66. As might be expected this shows that the boiling time has a greater effect on starch gelatinisation than soaking. Daomukda et al. (2011) studied the degree of gelatinization of brown rice cooked by various cooking methods (electric cooking, microwaving, steaming and conventional boiling) and found that steaming produced the least degree of gelatinisation and the lowest GI. However, steamed rice cakes (Pinto) commonly consumed in the Philippines showed high GI values (80-90) (Trinidad et al., 2010) as did steamed Sri Lankan rice noodles (103) (Hettiarachchi et al, 2003). The use of milled rice flour combined with steaming may have caused a greater gelatinisation and produced a higher GI in these products.

Some processed rice products requiring longer cooking times have higher GIs than those requiring shorter times presumably as a result of gelatinization and a weakened structure. Instant white rice boiled for 1 min had a GI of 65 (Wolever et al., 1986) compared to instant rice cooked for 6 min which showed a higher GI of 87 (Brand et al.,1985). Holt & Brand Miller (1995) reported a GI of 102 for instant Doongara white rice. Instantization involves pre-cooking and dehydrating the rice which enables consumers to shorten cooking time to 2-3 min.

The water-to-rice ratio during cooking affects the degree of gelatinization and thereby digestibility (Daomukda et al. 2011). The degree of gelatinization of cooked rice tended to decrease with decreasing ratios of water to rice. The ratios studied were 2:1, 3:1 and 4:1 (w/v).

Brown rice cooked at a ratio of 2:1 yielded a hard texture which may be due to the cooking water being insufficient for complete starch gelatinization, whereas a ratio of 3:1 and 4:1 produced a rice with a more desirable texture. A water to rice ratio of 2:1 was also found to significantly increase RS content in white rice (Kim et al., 2006). There are marked differences in the way rice is cooked around the world. Whilst in predominantly rice eating countries it is cooked such that all the cooking liquid has evaporated at the end of cooking, rice in Western countries is cooked in excess water and drained before consumption. A notable amount of starch may be lost along with the cooking liquor in the latter method.

In summary the cooking method, cooking time, cooking conditions and cooking liquid volume all impact on rice GI. Lesser boiled and steamed rice grains have a smaller GI. However steamed products made with rice flour shows a high GI. The literature highlights the importance of standardising the cooking method in GI studies.

Processing

Processing methods such as explosion puffing, extrusion and instantisation increase starch digestibility. Instantisation has been shown to increase digestibility even in high amylose rices (Brand et al., 1985; Holt & Brand Miller, 1995). Puffing increases the digestibility and the GI of rice presumably due to gelatinisation, and a weakened and porous structure. Puffed white rice cooked for 5 min generated a GI of 74 in an unpublished observation (Human Nutrition Unit; Sydney, Australia). Similarly rice bubbles (puffed rice) and puffed rice cakes showed GIs of 116

and 95, and 117 and 128 respectively (Brand et al., 1985; Holt et al., 1992; Miller et al., 1992; Holt & Brand Miller, 1995). However in contrast to instantization, puffing still produced a lower glycaemic response if a high amylose rice was used.

Extrusion uses high pressure and temperature conditions to gelatinise starch and is sometimes followed by drying. The principle extruded rice-based product is noodles. Fresh rice noodles are sometimes cooked again in boiling water at the last stage to further gelatinize the starch (Hou et al., 2010). Extrusion reduces the molecular weight of both amylose and amylopectin (Politz et al., 1994). These shorter branches could form novel indigestible cross-links and lower the GI (Theander & Westerlund, 1987). Further, rice vermicelli is predominantly made with high amylose rice which inherently has a lower GI (Juliano & Goddard, 1986). Panlasigui et al. (1992) showed that extruded high amylose rice noodles lowered the in vitro starch digestion and the in vivo blood glucose response in healthy and diabetic volunteers in Toronto and Philippines. A study by Juliano et al. (1990) showed that rice noodles elicited a lower starch digestibility and glycaemic index in healthy volunteers. Dry (Vietnamese) and fresh (Australian) rice vermicelli showed GI values of 61 and 40 respectively in one study (Chan et al., 2001). Differences in GI values between these two types may have been due to the type of rice used, processing and storage conditions. Two rice noodles, Jianxi rice vermicelli and Taiwan vermicelli, consumed in Hong Kong, was studied by Lok et al. (2009). Whilst Taiwan vermicelli had a GI of 68, Jianxi vermicelli had a GI of 56. Chinese Jianxi and Guilin vermicellis however showed lower GI values (37 and 40 respectively) in another study (Ranawana et al, 2009). Philippino bihon was similarly found to be low GI (49) (Trinidad et al., 2010). The higher GI values for Taiwan and

Vietnamese vermicelli (compared to the Chinese types) may be due to different amylopectin contents and processing conditions. *Idiyappam* (steamed fresh rice vermicelli or string hoppers) is made from red or white rice flour and is eaten in India and Sri Lanka. Hettiaratchi et al. (2009) reported a GI of 103 for red rice *idiyappam*. Sato et al (2010) investigated the GI of Thai long grain *indica* rice and rice vermicelli variants in healthy subjects and found that the GI of rice vermicelli (55) and rice flat noodles (kway teow) (50) made from high amylose rice was lower than that of long grain rice (60) (Sato et al., 2010).

Processes that cause amylose retrogradation such as parboiling may reduce the GI through the formation RS. Parboiling is a hydrothermal treatment where the starch in the grain undergoes gelatinisation followed by retrogradation. Niba (2003) showed that moist heat processing and the post-process storage temperature both significantly affect starch digestibility. Parboiled rice due to its harder texture is more resistant to milling losses and leaching during cooking (Bhattacharya, 2004). It is the preferred form in some communities due to its greater yield, better storability and greater grain volume. Some studies show that parboiling reduces the GI of rice (Casiraghi et al, 1993; Wolever, 1986; Ranawana et al, 2009). A reduced glycaemic response was observed in healthy subjects who consumed parboiled Sri Lankan rice compared to an unparboiled version (Pathiraje et al., 2010). The authors suggest that besides the retrogradation of rice starch during the parboiling process, the higher protein content in the parboiled version and the resulting protein-starch interactions may have contributed to the lower starch digestibility compared to raw rice. However, other studies show that parboiling has little effect on the GI (Larsen et al, 1996). They compared the GI of high and low amylose parboiled rice and found

that although the former produced a significantly lower GI than the latter the values were not different from the un-parboiled controls. Aston et al (2008) reported that easy-cook basmati rice (instant rice or minute rice) which is a pre-gelatinised rice, had a significantly higher GI than standard basmati rice. This was confirmed by Ranawana et al (2009) who compared the GI of raw and easy-cook Basmati rice. The authors suggested that the higher GI of the latter was possibly due to a longer cooking time which resulted in a greater degree of starch gelatinisation which may have undermined the GI lowering effects of RS formed during parboiling. Although an increased RS content due to parboiling might be expected to bring down the GI, traditional parboiling may not be producing adequate amounts of RS to affect the glycaemic response. Larsen et al (2000) suggested that the effect of parboiling on the GI depends on the severity of processing. They compared the effect of non-parboiled, mildly traditional parboiled and severely pressure parboiled rices on the GI in diabetics and found that the latter produced the lowest value. The formation of RS seemed to have little effect on the GI reduction as amounts were quite small in both traditionally (0.2%) and pressure parboiled (1.6%) rice. Differential scanning calorimetry (DSC) showed that the pressure parboiled variant had retrograded amylopectin and this may have reduced available carbohydrate content. Further, amylopectin crystallites retain some of the associating forces during reheating and could be partly responsible for the low GI of severely pressured parboiled rice (Faisant et al., 1993). The authors observed no amylose retrogradation in any of the rice samples and this concurs with other reports (Ong & Blanshard, 1995). However, Lamberts et al. (2009) using DSC and wide-angle X-ray diffraction (WAXD) showed amylose crystallites in parboiled rice, and found that amounts depended both on parboiling conditions and cultivar. Mildly parboiled rice showed lower amylose crystallites than

intermediate and severely parboiled rice. Although parboiled rice may be a better alternative to white rice due to its low GI properties there is limited acceptance of it in communities not traditionally consuming it. Parboiling alters the physical and sensory properties of rice and renders it less palatable to some (Heinemann et al., 2006).

In summary the impact of processing on rice GI depends on the method and intensity. Processes that disrupt structure and promote gelatinisation increase the GI whilst those promoting RS formation and amylose/amylopectin complexes reduce it. The relative impact also depends on the chemical composition of the starch in the rice. Parboiling has the potential to reduce the GI of rice but this may depend on the parboiling technique used and cooking conditions. Extruded rice products generally have a lower GI than whole rice. However this may again depend on rice type, finishing method (fresh or dried) and cooking procedure.

Cooling

Cooling reduces the GI of cooked rice through retrogradation and resultant increase in RS levels (Englyst et al., 1992; Sievert & Pomeranz, 1989). Highest crystallisation rates can be observed around 4°C (Baik et al., 1997). In vitro studies showed storing cooked rice at refrigerated temperatures (4°C for 24 hours) led to a reduction in their digestibility and estimated GI for both brown (Frei et al., 2003) and milled rice (Hu et al., 2004). Goñi et al (1997) estimated the expected GI of cooked and cooked refrigerated rice and found substantial differences between rice cultivars. Starch hydrolysis tended to be more rapid and more complete for waxy cultivars

than for high amylose cultivars. Storing rice in the refrigerator led to a reduction in the estimated GI for all cultivars. The highest decrease in starch hydrolysis after cool storing was seen for the waxy cultivars which suggest that high amylopectin rice may be more susceptible to cold induced retrogradation. Aging of noodles (as part of processing) in a cold location increased RS formation in rice noodles and lowered the GI (Sato et al., 2010). Therefore cooling seems to reduce the digestibility of both rice grains and processed products. Cooling appears to be a simple and effective intervention to reduce the GI of rice.

Soaking

Rice is sometimes soaked prior to cooking for various reasons. Punjabi Basmati rice is commonly pre-soaked to elongate the grains with cooking and produce a more aesthetically pleasing product (Juliano, 2003). White and brown rice is often soaked prior to milling to soften the grain, and also is a processing step in parboiling. Brown rice is sometimes soaked in water before cooking to reduce hardness and aid in cooking. Soaking allows starch granule expansion and better gelatinization leading to improved digestibility and a higher GI. The soaking temperature and moisture content of rice kernels affects the digestive properties of the cooked rice. Han & Lim, (2009) studied the digestibility of brown rice kernels (*japonica* type) soaked in water at different temperatures (25 or 50°C) before cooking to a moisture content of 20 or 30%. The cooked brown rice that had 30% moisture before cooking was digested to a greater extent than rice with 20% moisture. Rice soaked at 50°C (equal moisture content) was digested more readily. It was suggested that the amount of soluble material leached during soaking differed

according to the soaking temperature and moisture content, which subsequently affected the texture and digestive properties of the cooked rice. The rice cooked in its own soaking water was harder, more adhesive, had higher levels of RS, and exhibited smaller GI values than its distilled water cooked counterpart. Soluble material leached during soaking makes the cooked rice harder and less digestible, which the authors attributed to the interactions between these molecules and the gelatinized rice during cooking. In a study of Chinese starchy foods, Lin et al. (2010) pre-soaked brown rice overnight in water at a ratio of 1.5:1 before cooking and found that it elicited a high GI (82; white bread as reference), the highest in comparison to other starchy foods (taro, adlay, yam and mung bean noodles). The authors suggested that the process of soaking, besides reducing hardness and chewy mouthfeel, allows for starch granule expansion and better gelatinization, leading to improved digestibility and a high GI. The literature thus suggests that soaking may increase the GI of rice but it is dependent on soaking and subsequent cooking conditions. Omitting this step may be beneficial if a low GI rice is required.

Germination

Pre-germinated brown rice (PGBR) is a type of Japanese rice that is slightly germinated following soaking in water. Pre-germinated brown rice was better than white rice in preventing rapid postprandial blood glucose and insulin increases (Ito et al., 2005). In a study using rats Seki et al. (2005) found that insoluble dietary fibre in PGBR bran was mainly responsible for lowering postprandial blood glucose levels. Daniel (2008) reported that PGBR helps lower blood glucose level in diabetics due to the action of acylated sterol glucoside (ASG) dysfunctional

enzymes. Acylated steryl glucoside is a growth factor found in brown rice after germination. Pre-germinating rice appears to be an effective method of reducing the GI. However, pre-germinated rice consumption is limited to a few countries and making it a culturally acceptable elsewhere would be the greatest challenge to its more widespread use. The observed beneficial effects and the dearth of literature on this product justifies further research.

Fibre

The evidence on the effect of fibre on rice GI remains equivocal. While some studies have shown that brown rice has a lower GI than its white counterpart (O'Dea et al., 1981; Brand-Miller et al., 1992) other studies have shown it to be higher (Lin et al, 2010; Ranawana et al, 2009). No correlation has been observed between fibre content in rice and its GI (Jenkins et al., 1981). Therefore it may be that the lower GI of brown rice is a result of other bran constituents. Some studies showed that the rice bran fraction reduced the glycaemic response in healthy (Brand-Miller et al., 1992) and diabetic individuals (Qureshi et al., 2002). Since the bran envelopes the starchy centre it may be also acting both as a physical barrier for water entry and as an impediment to swelling. Optimum heat treatment for brown and milled rice to obtain an acceptable texture according to their minimum cooking times (24 and 14 min respectively) still produced a slower digestion rate and blood glucose response for brown than milled rice (Panlasigui & Thompson, 2006). To some extent, digestive enzymes were still not able to act on the starch substrate in brown rice which suggests that the low digestibility was due to the bran

creating a physical barrier. Differences in chemical composition and physico-chemical properties may also contribute to the different effects of brown and milled rice (Panlasigui & Thompson, 2006). The RS content was shown to be higher in brown rice (Nugent, 2005; Sajilata et al., 2006). Anti-nutrition factors such as phytic acid and polyphenols, may also contribute to slow starch digestion and lower blood glucose response associated with brown rice (Yoon et al., 1983; Thompson et al., 1987). More recently soluble viscous fibre has shown to have greater effect on carbohydrate metabolism than non-soluble fibre. Natural viscous vegetables (natto, yam and okra) in combination with white rice, eaten in Japanese-style breakfasts, delayed gastric emptying (Taniguchi-Fukatsu et al., 2011). Also, viscous vegetables lowered blood glucose and insulin secretion, unlike milk products that lower blood glucose but stimulate insulin secretion. Two recent studies incorporated a novel functional viscous fibre, PGX® into commonly consumed starchy foods including rice (Jenkins et al., 2010; Brand-Miller et al., 2010). Both studies found a dose dependant reduction in the GI when PGX was added to white rice. (19% for 2.5g dose and 30% for 5g dose).

In summary it is still unclear if inherent non-soluble fibre has a direct effect on rice GI. The GI of brown rice may be lower due to a combination of factors such as reduced digestibility, higher RS content and effects of other bran constituents. More data from well designed studies are required for firm conclusions. Co-ingestion of viscous fibre appears to reduce the GI of rice.

Particle size

Food particle size influences starch digestibility both in vivo and in vitro (Bjorck et al., 1994; Holt & Brand Miller, 1994; Snow et al., 1981; Heaton et al., 1988). O’Dea et al. (1981) noted that cooking ground rice resulted in significantly higher glycaemic and lipemic responses than that seen with whole rice in both normal and diabetic subjects. The higher surface area in ground rice increases digestion rate. Kim et al. (2004) looked at the influence of three processed forms of rice (garaeduk, bagsulgi, cooked rice) on in vitro starch hydrolysis and postprandial glucose and insulin responses in diabetics. The firmer rice product (bagsulgi) was more resistant to maceration and therefore resulted in slower in vitro and in vivo digestion compared to the softer forms which underwent the same level of maceration. Ranawana et al. (2010) investigated between-individual variations in the degree of mastication when eating rice, and its effect on the in vivo glycaemic response and found that the extent of chewing significantly impacted on the glycaemic response. Those who masticated the rice to a greater extent showed a significantly larger glycaemic response compared to those who did not. As an intact grain the starch in rice resides within cells and mechanical breakdown is therefore important for its release for digestion. The findings suggest that the degree of mastication may be a significant digestion rate limiting factor for rice, and that chewing less will result in a lower GI. The method of milling also appears to impact on the glycaemic response. Tran et al. (2011) looked at the degradation of starch structure after three levels of milling of polished rice grains. They observed that the more destructive means of grinding (grinding intensity, temperature and time) disrupts the starch crystalline structure to a greater extent and that this has nutritional implications. For instance, hammer milling caused less structural damage resulting in longer branched chain starch molecules and slower digestibility. Porridge or congee is made with broken rice and is

commonly consumed in Chinese cultures. Broken rice has a greater amount of cracking within the grain which results in increased gelatinisation and a digestibility (GI-86) (Chan et al., 2001). Indeed Srikaeo and Sopade (2010) found that instant rice porridges (made from Thai Jasmine rice) have a high potential GI (68 to 97). Porridges are also cooked for longer than conventional rice and the higher resulting gelatinisation also increase digestibility and the GI.

In conclusion the particle size of ingested rice appears to have a significant impact on its GI. The particle size of rice after milling will determine gelatinisation rates during cooking whilst the degree of breakdown during chewing will impact on the digestion rate in the small intestine.

The glycaemic index of mixed meals and impact of condiments

As rice is hardly consumed on its own, but in accompaniment with other foods (vegetables, pulses, legumes, nuts, seafood and meat) this section reviews the GI of mixed meals consumed in Asia. The published GI values of mixed meals commonly consumed in Asia are assembled in Appendix A.

Sharavathy et al. (2001) studied the rate and extent of starch digestion *in vitro* of cereal-based Indian food preparations (with/without accompaniments). Their findings indicated that the amount of RDS and slowly digestible starch (SDS) can be manipulated relatively simply by varying the type of accompaniments. Rice roti consumed on its own had lower SDS as compared

to when it was consumed with coconut chutney. Rice flour roti was classified as high GI (103) when consumed without accompaniments (Widanagamage et al., 2009).

Hettiaratchi et al. (2011) reported that the GI of Sri Lankan rice mixed meals may be reduced by including naturally occurring sources of fibre with starchy staples. The effect of consuming curry with parboiled rice (*Mottaikarupan*) in Sri Lanka was studied by Pirasath et al. (2010) where the mean GI values of parboiled rice consumed either with *Amaranthus* leaf curry, soya meat gravy or a combination of both were 47, 56 and 55 respectively. The addition of legumes such as lentils lowers the GI of rice when consumed together (Chaturvedi et al., 1997; Hettiaratchi et al., 2009; Chew et al., 1988; Araya et al., 2003).

Sugiyama et al. (2003) fed different forms of soybean (roasted and grounded, fermented, paste) with rice and reported GI lowering effects. Beans are low GI and the attenuated GI of the rice meals was attributed to soluble fibre and anti-nutrients found in beans. Similarly natto and viscous vegetables (Japanese yams and okra) combined with white rice showed improved insulin sensitivity, lipid metabolism and oxidative stress in overweight subjects with impaired glucose tolerance (Taniguchi-Fukatsu et al., 2011). Soluble fibre may be reducing the GI through their effects on gastric emptying and the creation of a barrier between starch and enzymes.

Studies have investigated the effect of other macronutrients on rice GI. Aston et al. (2008) determined the GI of Basmati, easy-to-cook Basmati and easy-to-cook American rice with the addition of margarine. The GIs reported were 43, 68 and 49 respectively for each rice type.

Although the study does not provide data from controls for comparison other studies suggest that the plain forms of these rices have a relatively higher GI (Ranawana et al, 2009). The addition of fat may have reduced the glycemic response by delaying gastric emptying and enhancing gastric inhibitory polypeptide (GIP) secretion (Collier & O’Dea, 1983; Collier et al., 1984; Ercan et al., 1994). However, Japanese low amylose (17%) rice (147g) consumed with butter (10g) had a high GI of 96 (Sugiyama et al., 2010). This suggests that the GI lowering effects are due to interactions between fat and amylose, and this concurs with previous observations (Goddard et al, 1989). Chen et al. (2010) found that rice noodles stir-fried “Singapore-style” had a GI value of 54 (low GI) and fried rice noodles with sliced beef had a GI of 66 (medium GI). The study however found no relationship between the amounts of fat and protein in the foods and their GI values suggesting that the cooking method had a greater effect on GI than the macronutrient composition. Only a limited number of studies that have investigated the effects of fats on rice GI and this aspect requires further research. Steamed glutinous rice roll (89), sticky rice wrapped in lotus leaf (83) and plain steamed vermicelli roll (90) all had high GIs (Chen et al., 2010). Stir-fried vegetables and chicken when eaten with rice had a high GI (73) (Chew et al. 1988). An in vitro study reported the glycaemic potential of Malaysian foods and estimated that fried rice and nasi lemak (coconut rice with accompaniments) had a medium GI of 59 and 66 respectively (Shanita et al., 2011). The same study showed fried bee hoon made with low amylose rice had an estimated high GI (99). However, these findings need to be confirmed in vivo.

Sugiyama et al. (2003) reported the GI of low amylose (17%) high GI (80) Koshikari rice when consumed with other ingredients. Dairy products (milk, cheese, yogurt) significantly reduced the

GI of the rice when consumed together. Rice with curry and cheese (67) had a lower GI than rice consumed with curry on its own. This may be due to macronutrient effects on either or both gastric emptying and starch digestion. However, these findings are contrary to Chen et al. (2010) who showed that fat and protein had no effect on rice GI. However, dairy products have been criticized as causing excessive insulin secretion so its use in lowering the GI of rice meals has to be considered with caution.

The addition of vinegar (acetic acid) and vinegared foods to white rice reduces its GI (Brighenti et al., 1995; Liljeberg & Bjork, 1998). One study showed that rice with the addition of vinegar or vinegared foods (pickled cucumber), reduced the GI of Koshihikari low-amylose rice (Sugiyama et al., 2003). The effective dosage of vinegar was estimated to be about 0.2-1.5g/100g which is as low as that found in sushi (Sugiyama et al., 2003). More studies are needed to confirm the mechanism of vinegar in lowering the GI of rice.

Sakuma et al. (2009) studied the glycaemic effect when rice is eaten along with barley and found that postprandial glucose and insulin levels were suppressed compared with white rice on its own. Area under the curves of plasma glucose and insulin concentrations was reduced by barley intake in a dose-dependent manner. This reduction may have been due to the effects of beta-glucan a soluble fibre found in barley and which has shown to attenuate postprandial glucose and insulin responses (Braaten et al., 1991; Cavallero et al., 2002; Granfeldt et al., 1994; Hallfrisch & Behall, 2000; Makelainen et al., 2007; Panahi et al., 2007; Thondre & Henry; Chillo et al., 2010). Beta glucan has been shown to increase viscosity in the upper digestive tract (Wood et al., 1994).

It may therefore be slowing gastric emptying and digestion rate but this remains to be confirmed. Consuming rice with low GI foods will reduce overall meal GI. But the cooking method and degree of processing also play an important role in the final glycaemic response.

Emulsifiers appear to reduce the digestibility of starch. Guraya et al. (1997) studied the effect of rice starch-lipid complexes on in vitro digestibility of non waxy and waxy rice (100% amylopectin) with different amylose:amylopectin ratios. Long chain saturated emulsifiers reduced digestibility more than short chain saturated and unsaturated emulsifiers when complexed with nonwaxy and waxy rice starch. The largest decrease in digestibility was seen with Polyaldo (100% C18:0 with decaglyceryl monostearate modification) for non waxy rice. Waxy rice starch did not complex with most of the emulsifiers. Most emulsifiers that reduced digestibility by about 10% were composed of unsaturated monoglycerides including some acetylated and succinylated monoglycerides..

The addition of other starches (corn, tapioca, potato) to rice starch affected digestibility and improved textural qualities of noodles (Sato et al., 2010). The authors found that rice vermicelli made from long grain rice, tapioca and corn starch had a GI of 35. Tapioca and corn starch has GIs ranging between 46-70 and 37-62 respectively (Foster-Powell et al., 2002). Combining these two starches with rice starch to make rice vermicelli may have produced a synergistically lower GI compared to pure rice vermicelli which had a GI of 55. Instant rice vermicelli (made from long grain rice, tapioca and potato starch) had a GI of 59, kway teow (made from long grain rice and potato starch) had a GI of 60 and 'pho' (made from long grain rice and tapioca starch) had a

GI of 62 (Sato et al., 2010). Other studies showed that plain kway teow had a GI of 50 (Sato et al., 2010). The addition of potato starch may have increased the glycaemic response of the kway teow as potato has a medium to high GI. However, like rice, potato starch varies in amylose content (14-31% amylose in normal genotypes and up to 70% amylose in waxy genotypes) (Singh et al., 2010) and the use of a high amylose variety with rice could bring down the overall GI of a meal. The presence of other additives could also influence the functional and digestion properties of rice and rice products, and the GI. For example, studies on bread and lentils have shown that salt increases postprandial glucose and insulin excursions (Thorburn et al., 1986).

In summary the literature shows that the type and quantity of condiments and accompaniments eaten with rice impacts on the GI. The effect of proteins and fats on the GI of rice is still unclear and further work in this area is required. In general, the addition of acidic condiments, viscous fibre, emulsifiers, dairy products, vegetables and pulses appears to reduce the GI of rice.

CONCLUDING REMARKS

The studies reviewed in this article appeared to use the recommended GI testing method (Brouns et al., 2005, Fao/Who, 1998, Iso, 2010) and fulfilled the minimum requirements for study design, subject number, reference food and test protocol (Appendix A). The most used reference foods were glucose and white bread. The feasibility of using white rice as the reference was evaluated in a study testing the GI of Japanese mixed meals (Sugiyama et al., 2003). The authors suggested that white rice might be a more applicable reference food in Asian GI studies as it is the principle

staple food in the region. Although the authors found a strong correlation between the GI values obtained using rice and glucose standards more research is required before it can be adopted conclusively. Other studies on Asian foods have evaluated the use of wheat chapatti as a reference (Rahman et al., 2009; Dilwari et al., 1981). Using commonly consumed regional staples as the reference in GI studies is a novel and pragmatic concept which is worthy of prosecution in future research.

Rice is the main constituent in the diets of a large population segment in the world. Whilst it contributes a large GL to the diet and thereby potentially promotes impaired glucose tolerance (notably in countries and communities accustomed to rice consumption), advocacy to reduce consumption may not be a sustainable solution. Reducing the GI of rice may be the more practical approach. Whilst the review highlights the large number of variables influencing the GI of rice, it shows that rice is more often than not in the medium to high GI range. Especially white rice which is the predominant form eaten in the world and the principle type associated with poor glycaemic control. The literature suggests that certain processes can reduce its GI to a low level. However, these processes almost always also change the physicochemical and sensory properties of the rice thus reducing their acceptance. It is harder to convince individuals to accept new types of rice or change traditional dietary habits. The more practical strategy may be the development of rice-accompaniment combinations that reduce the overall GI of rice meals. Future studies need to focus on macronutrient and condiment interactions with rice and their effects on the GI, and the development of optimum rice and accompaniment combinations that will be both culturally acceptable and reduce meal GI. Such developments will not only secure better health outcomes but also enable consumers to enjoy the delights of rice.

REFERENCES

- Al-Mssallem, M. Q., S. M. Hampton, et al. (2011). "A study of Hassawi rice (*Oryza sativa* L.) in terms of its carbohydrate hydrolysis (in vitro) and glycaemic and insulinaemic indices (in vivo)." Eur J Clin Nutr **65**(5): 627-634.
- Araya, H., Contreras, P., Alvina, M., Vera, G., & Pak, N. (2002). "A comparison between an in vitro method to determine carbohydrate digestion rate and the glyceemic response in young men." Eur J Clin Nutr **56**: 735-739.
- Aston, L. M., J. M. Gambell, et al. (2007). "Determination of the glycaemic index of various staple carbohydrate-rich foods in the UK diet." Eur J Clin Nutr **62**(2): 279-285.
- Atkinson, F. S., K. Foster-Powell, et al. (2008). "International Tables of Glycemic Index and Glycemic Load Values: 2008." Diabetes Care **31**(12): 2281-2283.
- Augustin, L. S., Franceschi, S., Jenkins, D.J.A., Kendall, C.W.C., & Vecchia, C.La. (2002). "Glycaemic index in chronic disease: A review." Eur J Clin Nutr **56**: 1049-1071.
- Azudin, M. N., & Morrison, W. K. , 4, 23–31. (1986). "Non-starch lipids and starch lipids in milled rice." Journal of Cereal Science **4**: 23-31.
- Baik, M.-Y., K.-J. Kim, et al. (1997). "Recrystallization Kinetics and Glass Transition of Rice Starch Gel System." Journal of Agricultural and Food Chemistry **45**(11): 4242-4248.
- Baldwin, P. M. (2001). "Starch-granule associated proteins and polypeptides: a review." Starch/Starke **53**: 475-503.
- Bao, J., & Bergman, C.J. (2004). The functionality of rice starch. In: Starch in food: Structure, function and applications. Boca Raton, Fl, Woodhead Publishing Ltd. and CRC Press

LLC.

Bao, J., Z. Ao, et al. (2005). "Characterization of Physical Properties of Flour and Starch

Obtained from Gamma-Irradiated White Rice." Starch - Stärke **57**(10): 480-487.

Berry, C. S. (1986). "Resistant starch: Formation and measurement of starch that survives

exhaustive digestion with amylolytic enzymes during the determination of dietary fibre."

Journal of Cereal Science **4**(4): 301-314.

Bhattacharjee, P., R. S. Singhal, et al. (2002). "Basmati rice: a review." International Journal of

Food Science & Technology **37**(1): 1-12.

Bhattacharya K, R. (2004). "Parboiling of rice." Rice Chemistry and Technology.

Biliaderis, C. G. (1991). "The structure and interactions of starch with food constituents."

Canadian Journal of Physiology & Pharmacology **69**: 60-78.

Bjorck, I. and N. G. Asp (1983). "The effects of extrusion cooking on nutritional value -" A

literature review." Journal of Food Engineering **2**(4): 281-308.

Bjorck, I., Y. Granfeldt, et al. (1994). "Food properties affecting the digestion and absorption of

carbohydrates." The American Journal of Clinical Nutrition **59**(3): 699S-705S.

Blakeney, A. B. and N. K. Matheson (1984). "Some Properties of the Stem and Pollen Starches

of Rice." Starch - Stärke **36**(8): 265-269.

Blanshard, J. M. V. (1987). Starch granule structure and function: a physicochemical approach.

In: Starch: Property and Potential. Chichester, John Wiley.

Bornet, F. (1993). "Technological treatments of cereals. Repercussions on the physiological

properties of starch." Carbohydrate Polymers **21**(2-3): 195-203.

Braaten, J. T., P. J. Wood, et al. (1991). "Oat gum lowers glucose and insulin after an oral

- glucose load." The American Journal of Clinical Nutrition **53**(6): 1425-1430.
- Brand, J. C., S. Colagiuri, et al. (1991). "Low-Glycemic Index Foods Improve Long-Term Glycemic Control in NIDDM." Diabetes Care **14**(2): 95-101.
- Brand, J. C., P. L. Nicholson, et al. (1985). "Food processing and the glycemic index." The American Journal of Clinical Nutrition **42**(6): 1192-1196.
- Brand-Miller J, W. T., Foster-Powell K, & Colagiuri, S. (2007). The New Glucose Revolution, 3rd edn. London, UK., Hodder & Stoughton.
- Brand-Miller, J. C., F. S. Atkinson, et al. (2010). "Effects of added PGX[®], a novel functional fibre, on the glycaemic index of starchy foods." British Journal of Nutrition **FirstView**: 1-4.
- Brennan, C. S., D. E. Blake, et al. (1996). "Effects of Guar Galactomannan on Wheat Bread Microstructure and on the In vitro and In vivo Digestibility of Starch in Bread." Journal of Cereal Science **24**(2): 151-160.
- Brighenti, F., Castellani, G., Benini, L., Casiraghi, M., Leopardi, E., Crovetto, R., & Testolin, G. (1995). "Effect of neutralized and native vinegar on blood glucose and acetate responses to a mixed meal in healthy subjects." European Journal of Clinical Nutrition **49**: 242-247.
- Brouns, F., I. Bjorck, et al. (2005). "Glycaemic index methodology." Nutr Res Rev **18**(1): 145-71.
- Capriles, V. D., K. D. Coelho, et al. (2008). "Effects of Processing Methods on Amaranth Starch Digestibility and Predicted Glycemic Index." Journal of Food Science **73**(7): H160-H164.
- Casiraghi, M. C., Brighenti, F., Pellegrini, N., Leopardi, E., & Testolin, G. (1993). "Effect of processing on rice starch digestibility evaluated by in vivo and in vitro methods." Journal

- of Cereal Science **17**: 147–156.
- Casiraghi, M. C., F. Brighenti, et al. (1992). "Lack of effect of high temperature drying on digestibility of starch in spaghetti." Journal of Cereal Science **15**(2): 165-174.
- Cavallero, A., S. Empilli, et al. (2002). "High ((1→3) (1→4)-β-D-Glucan Barley Fractions in Bread Making and their Effects on Human Glycemic Response." Journal of Cereal Science **36**(1): 59-66.
- Champagne, E. T. (1996). "Rice starch composition and characteristics." Cereal Foods World **41**: 833-838.
- Chan, H. M. S., Brand-Miller, J.C., Holt, S.H.A., Wilson, D., Rozman, M., & P Petocz (2001). "Original Communication: The glycaemic index values of Vietnamese foods." Eur J Clin Nutr **55**: 1076-1083.
- Chan Jn, M. V. J. W. and et al. (2009). "Diabetes in Asia: Epidemiology, risk factors, and pathophysiology." JAMA: The Journal of the American Medical Association **301**(20): 2129-2140.
- Chang, T. M., Passaro, E. Jr., Shain, L.R., & Chen, W.L. (1991). "Physical properties of starch meals in vivo and in vitro and their influence on gastric emptying and oral glucose tolerance test." Nutrition Research **7**: 410-416.
- Chaturvedi, A., Sarojini, G., Nirmala, G., Nirmalamma N. & Satyanarayana D. (1997). "Glycemic index of grain amaranth, wheat and rice in NIDDM subject." J Plant Foods Hum Nutr **50**: 171-8.
- Chen, Y. J., Sun, F.H., Wong, S.H.S, & Huang, Y.J. (2010). "Glycemic index and glycemic load of selected Chinese traditional foods." World J Gastroenterol **16**: 1512-1517.

- Chew, I., J. C. Brand, et al. (1988). "Application of glycemic index to mixed meals." The American Journal of Clinical Nutrition **47**(1): 53-6.
- Chillo, S., J. A. Monro, et al. (2010). "Effect of incorporating legume flour into semolina spaghetti on its cooking quality and glycaemic impact measured in vitro." International Journal of Food Sciences and Nutrition **61**(2): 149-160.
- Chung, H. J. and Q. Liu (2009). "Effect of Gamma Irradiation on Molecular Structure and Physicochemical Properties of Corn Starch." Journal of Food Science **74**(5): C353-C361.
- Chung, H.-J., Q. Liu, et al. (2010). "Physicochemical Properties and In Vitro Starch Digestibility of Cooked Rice from Commercially Available Cultivars in Canada." Cereal Chemistry Journal **87**(4): 297-304.
- Chung, H.-J., Q. Liu, et al. (2011). "Relationship between the structure, physicochemical properties and in vitro digestibility of rice starches with different amylose contents." Food Hydrocolloids **25**(5): 968-975.
- Collier, G., A. McLean, et al. (1984). "Effect of co-ingestion of fat on the metabolic responses to slowly and rapidly absorbed carbohydrates." Diabetologia **26**(1): 50-54.
- Collier, G. and K. O'Dea (1983). "The effect of coingestion of fat on the glucose, insulin, and gastric inhibitory polypeptide responses to carbohydrate and protein." The American Journal of Clinical Nutrition **37**(6): 941-944.
- Collings, P., Williams, C., & Maodonald, I. (1981). "Effects of cooking on serum glucose and insulin responses to starch. ." British Medical Journal(282): 1032.
- Crowe, T. C., S. A. Seligman, et al. (2000). "Inhibition of Enzymic Digestion of Amylose by Free Fatty Acids In Vitro Contributes to Resistant Starch Formation." The Journal of

Nutrition **130**(8): 2006-2008.

Cuevas, R. P., & Fitzgerald, M. (2007-08). "Linking starch structure to rice cooking quality."

IREC Farmers' Newsl **177**: 16–17.

Cummings, D. E., J. Q. Purnell, et al. (2001). "A Preprandial Rise in Plasma Ghrelin Levels Suggests a Role in Meal Initiation in Humans." Diabetes **50**(8): 1714-1719.

Cummings, J. H. (1978). "Nutritional implications of dietary fiber." The American Journal of Clinical Nutrition **31**(10): S21-S29.

Dang, J. M. C. and L. Copeland (2003). "Imaging Rice Grains Using Atomic Force Microscopy." Journal of Cereal Science **37**(2): 165-170.

Daniel, H. R. (2008). "Chemical in germinated brown rice could benefit diabetics." Journal of Lipid Research:source: defeat-diabetes foundation

Daomukda, N., Moongngarm, A., Payakapol, L., & Noisuwan, A. (2011). Effect of Cooking Methods on Physicochemical Properties of Brown Rice. 2nd International Conference on Environmental Science and Technology. Singapore, IACSIT Press. **IPCBEE vol.6 (2011)**.

Denver, K., L. M. Barber, et al. (1995). "The isolation and characterization of novel low-amylose mutants of *Pisum sativum* L." Plant, Cell & Environment **18**(9): 1019-1026.

Denyer, K., Barber, L. M., Burton, R., Hedley, C. L., Hylton, C. M., S. Johnson, Jones, D. A., Marshall, J., Smith, A. M., Tatge, H., et al. (1995). "The isolation and characterisation of novel low-amylose mutants of *Pisum sativum* L. Plant, Cell and Environment." **18**: 1019-1026.

Dilawari, J. B., P. S. Kamath, et al. (1981). "Reduction of postprandial plasma glucose by Bengal

- gram dal (*Cicer arietinum*) and rajmah (*Phaseolus vulgaris*)." The American Journal of Clinical Nutrition **34**(11): 2450-3.
- Donald, A. M., Waigh, T.A., Jenkins, P.J., Gidley, M.J., & Smith., A. (1997). "Internal structure of starch granules revealed by scattering studies." Starch/Stärke **50**: 172-179.
- Dreher, M. L., C. J. Dreher, et al. (1984). "Starch digestibility of foods: A nutritional perspective." C R C Critical Reviews in Food Science and Nutrition **20**(1): 47-71.
- Eerlingen, R. C., Crombez, M., & Delcour, J.A. (1993). "Enzyme-resistant starch. I. Quantitative and qualitative influence of incubation time and temperature of autoclaved starch on resistant starch formation." Cereal Chemistry **70** . 339-344.
- Effersen, J. N. (1985). Rice quality in world markets. In: Rice grain quality and marketing. International Rice Research Institute. Los Banos, Philippines.
- Eggum, B. O., B. O. Juliano, et al. (1993). "The Resistant Starch, Undigestible Energy and Undigestible Protein Contents of Raw and Cooked Milled Rice." Journal of Cereal Science **18**(2): 159-170.
- Englyst, H. N., Kingman, S.M., & Cummings, J.H. (1992). "Classification and measurement of nutritionally important starch fractions." European Journal of Clinical Nutrition **44**: 423-431.
- Englyst, H. N., Veenstra, J., & Hudson, G. J. (1996). "Measurement of rapidly available glucose (RAG) in plant foods: A potential in vitro predictor of the glycaemic response." British Journal of Nutrition **75** 327-337.
- Englyst, H. N. and J. H. Cummings (1984). "Simplified method for the measurement of total non-starch polysaccharides by gas-liquid chromatography of constituent sugars as alditol

acetates." Analyst **109**(7): 937-942.

Englyst, K. N., Vinoy, S., Englyst, H. N., & Lang, V. (2003). "Glycaemic index of cereal products explained by their content of rapidly and slowly available glucose." British Journal of Nutrition **89**: 329-339.

Englyst, K. N., & Englyst, H.N. (2005). "Carbohydrate bioavailability." British Journal of Nutrition **94**: 1-11.

Englyst, K. N., H. N. Englyst, et al. (1999). "Rapidly available glucose in foods: an in vitro measurement that reflects the glycemic response." The American Journal of Clinical Nutrition **69**(3): 448-454.

Englyst, K. N., H. N. Englyst, et al. (1999). "Rapidly available glucose in foods: an in vitro measurement that reflects the glycemic response." The American Journal of Clinical Nutrition **69**(3): 448-454.

Englyst, K. N., G. J. Hudson, et al. (2000). Starch Analysis in Food. Encyclopedia of Analytical Chemistry, John Wiley & Sons, Ltd.

Englyst, K. N., S. Liu, et al. (2007). "Nutritional characterization and measurement of dietary carbohydrates." Eur J Clin Nutr **61**(S1): S19-S39.

Ercan, N., M. C. Gannon, et al. (1994). "Effect of Added Fat on the Plasma Glucose and Insulin Response to Ingested Potato Given in Various Combinations as Two Meals in Normal Individuals." Diabetes Care **17**(12): 1453-1459.

Faisant, N., Champ, M., Colonna, P., Buleon, A., Molis, C., Langkilde, A.M., Schweizer, T., Flourie, B., & Galmiche, J.P. (1993). "Structural features of resistant starch at the end of the human small intestine." European Journal of Clinical Nutrition **47**: 285-296.

- FAO/WHO (1998). "Carbohydrates in human nutrition. Report of a Joint FAO/WHO Expert Consultation." FAO Food Nutr Pap **66**: 1-140.
- Fardet Dr, A., C. Hoebler Dr, et al. (1999). "In vitro starch degradation from wheat-based products in the presence of lipid complex emulsions." Nutrition Research **19**(6): 881-892.
- Fernandes, G., A. Velangi, et al. (2005). "Glycemic index of potatoes commonly consumed in North America." Journal of the American Dietetic Association **105**(4): 557-562.
- Flint, A., Moller, B.K., Raben, A., Pedersen, D., Tetens, I., Holst, J.J., & Astrup, A. (2004). "The use of glycemic index tables to predict glycemic index of composite meals. ." British Journal of Nutrition **91**: 979-989.
- Foster-Powell, K., S. H. A. Holt, et al. (2002). "International table of glycemic index and glycemic load values: 2002." The American Journal of Clinical Nutrition **76**(1): 5-56.
- Foster-Powell, K. and J. B. Miller (1995). "International tables of glycemic index." The American Journal of Clinical Nutrition **62**(4): 871S-893S.
- Frei, M., P. Siddhuraju, et al. (2003). "Studies on the in vitro starch digestibility and the glycemic index of six different indigenous rice cultivars from the Philippines." Food Chemistry **83**(3): 395-402.
- Frost, G. S., A. E. Brynes, et al. (2003). "The effects of fiber enrichment of pasta and fat content on gastric emptying, GLP-1, glucose, and insulin responses to a meal." Eur J Clin Nutr **57**(2): 293-298.
- García-Alonso, A. and I. Goñi (2000). "Effect of processing on potato starch: In vitro availability and glycaemic index." Food / Nahrung **44**(1): 19-22.
- García-Alonso, A., A. Jiménez-Escrig, et al. (1999). "Assessment of some parameters involved

- in the gelatinization and retrogradation of starch." Food Chemistry **66**(2): 181-187.
- Goddard, M. S., G. Young, et al. (1984). "The effect of amylose content on insulin and glucose responses to ingested rice." The American Journal of Clinical Nutrition **39**(3): 388-392.
- Goñi, I., A. Garcia-Alonso, et al. (1997). "A starch hydrolysis procedure to estimate glycemic index." Nutrition Research **17**(3): 427-437.
- Granfeldt, Y., Bjorck, I., Drews, A., & Tovar, J. (1992). "An in vitro procedure based on chewing to predict metabolic response to starch in cereal and legume products." European Journal of Clinical Nutrition **46** 649–660.
- Granfeldt, Y., H. Liljeberg, et al. (1994). "Glucose and insulin responses to barley products: influence of food structure and amylose-amylopectin ratio." The American Journal of Clinical Nutrition **59**(5): 1075-1082.
- Grieb, P., B. Kłapcińska, et al. (2008). "Long-term consumption of a carbohydrate-restricted diet does not induce deleterious metabolic effects." Nutrition research (New York, N.Y.) **28**(12): 825-833.
- Guraya, H. S., R. S. Kadan, et al. (1997). "Effect of Rice Starch-Lipid Complexes on In Vitro Digestibility, Complexing Index, and Viscosity." Cereal Chemistry Journal **74**(5): 561-565.
- Hallfrisch, J. and K. M. Behall (2000). "Mechanisms of the Effects of Grains on Insulin and Glucose Responses." Journal of the American College of Nutrition **19**(suppl 3): 320S-325S.
- Han, J.-A. and S.-T. Lim (2009). "Effect of Presoaking on Textural, Thermal, and Digestive Properties of Cooked Brown Rice." Cereal Chemistry Journal **86**(1): 100-105.

- Hatcher, D. W., M. J. Anderson, et al. (2002). "Effects of Flour Particle Size and Starch Damage on Processing and Quality of White Salted Noodles." Cereal Chemistry Journal **79**(1): 64-71.
- Heaton, K. W., S. N. Marcus, et al. (1988). "Particle size of wheat, maize, and oat test meals: effects on plasma glucose and insulin responses and on the rate of starch digestion in vitro." The American Journal of Clinical Nutrition **47**(4): 675-682.
- Heinemann, R. J. B., J. H. Behrens, et al. (2006). "A study on the acceptability and consumer attitude towards parboiled rice." Int J Food Sci Tech **41**: 627-34.
- Hemavathy, J. and J. Prabhakar (1987). "Lipid composition of rice (*Oryza sativa* L.) bran." Journal of the American Oil Chemists' Society **64**(7): 1016-1019.
- Henry, C. J., Lightowler, H.J., Strik, C.M., Renton, H., & Hails, S. (2005). "Glycaemic index and glycaemic load values of commercially available products in the UK." British Journal of Nutrition **94**: 922-930.
- Henry, C. J. K., H. J. Lightowler, et al. (2006). "The impact of the addition of toppings//fillings on the glycaemic response to commonly consumed carbohydrate foods." Eur J Clin Nutr **60**(6): 763-769.
- Hettiaratchi, U. P. K., Ekanayake, S. & Welihinda, J. (2009). "Do Sri Lankan meals help decrease blood glucose response?" Ceylon Medical Journal **54**: 39-43.
- Hettiaratchi, U. P. K., Ekanayake, S., & Welihinda, J. (2011). "Sri Lankan rice mixed meals: Effect on glycaemic index and contribution to daily dietary fibre requirement." Malaysian Journal of Nutrition **17**: 97-104.
- Hizukuri, S. (1986). "Polymodal distribution of the chain lengths of amylopectins, and its

- significance." Carbohydrate Research **147**(2): 342-347.
- Hizukuri, S., K. Shirasaka, et al. (1983). "Phosphorus and Amylose Branching in Rice Starch Granules." Starch - Stärke **35**(10): 348-350.
- Hodge, A. M., D. R. English, et al. (2004). "Glycemic Index and Dietary Fiber and the Risk of Type 2 Diabetes." Diabetes Care **27**(11): 2701-2706.
- Holm, J., I. Björck, et al. (1983). "Digestibility of Amylose-Lipid Complexes in-vitro and in-vivo." Starch - Stärke **35**(9): 294-297.
- Holt, S., J. Brand, et al. (1992). "Relationship of satiety to postprandial glycaemic, insulin and cholecystokinin responses." Appetite **18**(2): 129-141.
- Holt, S. H. A. and J. B. Miller (1995). "Increased insulin responses to ingested foods are associated with lessened satiety." Appetite **24**(1): 43-54.
- Hoseney, R. C. (1990). Principles of Cereal Science and Technology, 2nd. Ed. Am. Assoc. Cereal Chem., St. Paul, MN.
- Hou, G. G., S. Otsubo, et al. (2010). Noodle Processing Technology. Asian Noodles, John Wiley & Sons, Inc.: 99-140.
- Hu, E. A., Oand, Pan, A., Malik, V. & Sun, Q. (2012). "White rice consumption and risk of type 2 diabetes: meta-analysis and systematic review." BMJ **344**: 1-9.
- Hu, P., H. Zhao, et al. (2004). "Starch digestibility and the estimated glycemic score of different types of rice differing in amylose contents." Journal of Cereal Science **40**(3): 231-237.
- Ingwersen, J., M. A. Defeyter, et al. (2007). "A low glycaemic index breakfast cereal preferentially prevents children's cognitive performance from declining throughout the morning." Appetite **49**(1): 240-244.

IRRI. "Rice Basics." Retrieved 21 March 2012, from <http://irri.org/about-rice/rice-facts/rice-basics>.

IRRI. (2009). "Milling/Processing." Retrieved 21 March 2012, from <http://www.knowledgebank.irri.org/rkb/rice-milling/commercial-rice-milling-systems/mist-polisher.html>.

ISO (2010). ISO 26642-2010. Food products- determination of the glycaemic index (GI) and recommendation for food classification. Switzerland:ISO.

Ito, K., Yoshida, K., Okazaki, N., & Kobayashi, S. (1988). "Effect of processing on the pore size distribution and digestibility of rice grain." *Agriculture & Biological Chemistry* **52**: 3001-3007.

Ito, Y., Mizukuchi, A., Kise, M., Aoto, H., Yamamoto, S., Yoshihara, R., & Yokoyama, J. (2005). "Postprandial blood glucose and insulin responses to pre-germinated brown rice in healthy subjects." *Journal of Medical Investigation* **52**: 159-164.

J.Wood, P., J. T. Braaten, et al. (1994). "Effect of dose and modification of viscous properties of oat gum on plasma glucose and insulin following an oral glucose load." *British Journal of Nutrition* **72**(05): 731-743.

Jaisut, D., S. Prachayawarakorn, et al. (2008). "Effects of drying temperature and tempering time on starch digestibility of brown fragrant rice." *Journal of Food Engineering* **86**(2): 251-258.

Jane, J., Kasemsuwan, T., Chen, J. F., & Juliano, B. O. (1996). "Phosphorus in rice and other starches." *Cereal Foods World* **41**: 827-838.

Jane, J., Chen, Y. Y., Lee, L. F., McPherson, A. E., Wong, K.S., Radosavljevic, M., &

- Kasemsuwan, T. (1999). "Effects of Amylopectin Branch Chain Length and Amylose Content on the Gelatinization and Pasting Properties of Starch." Cereal Chemistry **76**: 629-637.
- Jenkins, A. L., Kacinik, V., Lyon, M., & Wolever, T.M.S. (2010). "Effect of adding the novel fiber, PGX®, to commonly consumed foods on glycemic response, glycemic index and GRIP: a simple and effective strategy for reducing post prandial blood glucose levels - a randomized, controlled trial." Nutrition Journal **9**: 58.
- Jenkins, D., H. Ghafari, et al. (1982). "Relationship between rate of digestion of foods and post-prandial glycaemia." Diabetologia **22**(6): 450-455.
- Jenkins, D. A., A. Jenkins, et al. (1984). "THE GLYCAEMIC RESPONSE TO CARBOHYDRATE FOODS." The Lancet **324**(8399): 388-391.
- Jenkins, D. J., M. J. Thorne, et al. (1987a). "The effect of starch-protein interaction in wheat on the glycemic response and rate of in vitro digestion." The American Journal of Clinical Nutrition **45**(5): 946-951.
- Jenkins, D. J., T. M. Wolever, et al. (1987b). "Low-glycemic index diet in hyperlipidemia: use of traditional starchy foods." The American Journal of Clinical Nutrition **46**(1): 66-71.
- Jenkins, D. J., T. M. Wolever, et al. (1978). "Dietary fibres, fibre analogues, and glucose tolerance: importance of viscosity." BMJ **1**: 1392-1394.
- Jenkins, D. J., T. M. Wolever, et al. (1990). "Metabolic effects of reducing rate of glucose ingestion by single bolus versus continuous sipping." Diabetes **39**(7): 775-781.
- Jenkins, D. J., T. M. Wolever, et al. (1981). "Glycemic index of foods: a physiological basis for carbohydrate exchange." The American Journal of Clinical Nutrition **34**(3): 362-366.

- Jenkins, D. J., T. M. Wolever, et al. (1984). "The relationship between glycemc response, digestibility, and factors influencing the dietary habits of diabetics." The American Journal of Clinical Nutrition **40**(6): 1175-1191.
- Jenkins, D. J. A., C. W. C. Kendall, et al. (2006). "Almonds Decrease Postprandial Glycemia, Insulinemia, and Oxidative Damage in Healthy Individuals." The Journal of Nutrition **136**(12): 2987-2992.
- Jenkins, D. J. A., T. M. S. Wolever, et al. (1988). "Starchy Foods and Glycemic Index." Diabetes Care **11**(2): 149-159.
- Juliano, B. O., Perez, C.M., Komindr, S., & Banphotkasem, S. (1990). "Properties of Thai cooked rice and noodles differing in glycemc index in noninsulin-dependent diabetics." Plant Foods for Human Nutrition (Formerly Qualitas Plantarum) **40**: 231-232.
- Juliano, B. O., & Bechtel, D.B. (1985). The rice grain and its gross composition, In Rice: Chemistry and technology. St. Paul, M.N, AACC. Inc.
- Juliano, B. O. (1993). Rice in human nutrition. Food and Agriculture Organization of the United Nations, International Rice Research Institute.
- Juliano, B. O. and C. Editor-in-Chief: Benjamin (2003). RICE. Encyclopedia of Food Sciences and Nutrition (Second Edition). Oxford, Academic Press: 4995-5001.
- Juliano, B. O. and M. S. Goddard (1986). "Cause of varietal difference in insulin and glucose responses to ingested rice." Plant Foods for Human Nutrition (Formerly Qualitas Plantarum) **36**(1): 35-41.
- Juliano, B. O., C. M. Perez, et al. (1989). "Properties of Thai cooked rice and noodles differing in glycemc index in noninsulin-dependent diabetics." Plant Foods for Human Nutrition

(Formerly Qualitas Plantarum) **39**(4): 369-374.

Kalergis, M., Pacaud, D., & Yale, J-F. (1998). "Attempts to control the glycemc response to carbohydrate in diabetes mellitus: Overview and practical implications." Canadian Journal of Diabetes Care **22**: 20-20.

Kaur, L., J. Singh, et al. (2007). "Physico-chemical, rheological and structural properties of fractionated potato starches." Journal of Food Engineering **82**(3): 383-394.

Kavita, M. S., & Prema, L. (1997). "Glycaemic response to selected cereal-based South Indian meals in non-insulin dependent diabetics." J Nutr Environ Med **7**: 287-94.

Kennedy G, B. B., Nguyen VN. (2002). Nutritional contribution of rice and impact of biotechnology and biodiversity in rice-consuming countries. Proceedings of the 20th Session of the International Rice Commission, Bangkok; Thailand, FAO corporate document repository; 2003. .

Kim, J. C., Mulla, B. P., Hampson, D. J., & Pluske J. R. (2006). "Effects of amylose content, autoclaving, parboiling, extrusion, and post-cooking treatments on resistant starch content of different rice cultivars." Aust. J. Agri. Res **57**: 1291-1296.

Kim, J. C., J.-I. Kim, et al. (2004). "Influence of the Physical Form of Processed Rice Products on the Enzymatic Hydrolysis of Rice Starch in Vitro and on the Postprandial Glucose and Insulin Responses in Patients with Type 2 Diabetes Mellitus." Bioscience, Biotechnology, and Biochemistry **68**(9): 1831-1836.

Kim, K., Yun, S.H., Choi, B.Y., & Kim, M.K. (2008a). "Cross-sectional relationship between dietary carbohydrate, glycaemic index, glycaemic load and risk of the metabolic syndrome in a Korean population." British Journal of Nutrition (2008), **100** 576–584.

- Kim, Y., Hertzler, S.R., Byrne, H.K., Mattern, C.O. (2008). "Raisins are a low to moderate glycemic index food with a correspondingly low insulin index." Nutrition Research **28**: 304-308.
- Klein, S., N. F. Sheard, et al. (2004). "Weight Management Through Lifestyle Modification for the Prevention and Management of Type 2 Diabetes: Rationale and Strategies." Diabetes Care **27**(8): 2067-2073.
- Kumar, S., R. Mohanraj, et al. (2011). "Perceptions about varieties of brown rice: a qualitative study from Southern India." Journal of American Dietetic Association **111**(10): 1517-22.
- Lamberts, L., S. V. Gomand, et al. (2009). "Presence of Amylose Crystallites in Parboiled Rice." Journal of Agricultural and Food Chemistry **57**(8): 3210-3216.
- Langworthy, C. F., & H.J. Deuel. (1992). "Digestibility of raw rice, arrow-root, canna, cassava, taro, tree-fern, and potato starches." Journal of Biological Chemistry **52**: 251-261.
- Larsen, H. N., Christensen, C., Rasmussen, O.W., Tetens, I.H., Choudhury, N.H., Thilsted, S.H., & Hermansen, K. (1996). "Influence of parboiling and physico-chemical characteristics of rice on the glycaemic index in non-insulin-dependent diabetic subjects." European Journal of Clinical Nutrition **50**: 22-7.
- Larsen, H. N., Rasmussen, O.W., Rasmussen, P.H., Alstrup, K.K., Biswas, S.K., Tetens, I., Thilsted, S.H., & Hermansen, K. (2000). "Glycaemic index of parboiled rice depends on the severity of processing: study in type 2 diabetic subjects." European Journal of Clinical Nutrition **54**: 380-5.
- Larsen, T. M., S. Dalskov, et al. (2010). "The Diet, Obesity and Genes (Diogenes) Dietary Study in eight European countries – a comprehensive design for long-term intervention."

Obesity Reviews **11**(1): 76-91.

Lebet, V., E. Arrigoni, et al. (1998). "Digestion Procedure Using Mammalian Enzymes to Obtain Substrates for In Vitro Fermentation Studies." LWT - Food Science and Technology **31**(6): 509-515.

Leeds, A. R. (1982). Modification of intestinal absorption by dietary fiber and fiber components. New York, Plenum Press.

Leeman, M., E. Ostman, et al. (2005). "Vinegar dressing and cold storage of potatoes lowers postprandial glycaemic and insulinaemic responses in healthy subjects." Euro J Clin Nutr **59**(11): 1266-1271.

Lehmann, U. and F. Robin (2007). "Slowly digestible starch- its structure and health implications: a review." Trends in Food Science & Technology **18**(7): 346-355.

Liatis, S., S. Grammatikou, et al. (2010). "Vinegar reduces postprandial hyperglycaemia in patients with type II diabetes when added to a high, but not to a low, glycaemic index meal." Eur J Clin Nutr **64**(7): 727-732.

Liljeberg, H., & Björck, I. (1998). "Delayed gastric emptying rate may explain improved glycaemia in healthy subjects to a starchy meal with added vinegar." European Journal of Clinical Nutrition **52** 368-371.

Lin, M., Wu, M.C., Lu, S., & Lin, J. (2010). "Glycemic index, glycemic load and insulinemic index of Chinese starchy foods." World Journal of Gastroenterology **39**: 4973-4979.

Lin, P., Nhung, B.T., Khan, N.C., Sarukura, N., Kunii, D., Sakai, T., Kassu, A., & Yamamoto, S. (2007). "Effect of Vietnamese Common Diet on Postprandial Blood Glucose Level in Adult Females." Journal of Nutritional Science and Vitaminology **53** 247-252.

- Lin, Y. H., Yeh, C.S., & Lu, S. (2003). "Extrusion processing of rice-based breakfast cereals enhanced with tocopherol from a Chinese medical plant." Cereal Chemistry **80**: 491-494.
- Lindeboom, N., P. R. Chang, et al. (2004). "Analytical, Biochemical and Physicochemical Aspects of Starch Granule Size, with Emphasis on Small Granule Starches: A Review." Starch - Stärke **56**(3-4): 89-99.
- Liu, S. and E. L. Chou (2010). "Dietary glycemic load and type 2 diabetes: modeling the glucose-raising potential of carbohydrates for prevention." The American Journal of Clinical Nutrition **92**(4): 675-677.
- Liu, S., W. C. Willett, et al. (2000). "A prospective study of dietary glycemic load, carbohydrate intake, and risk of coronary heart disease in US women." The American Journal of Clinical Nutrition **71**(6): 1455-1461.
- Livesey, G. and H. Tagami (2009). "Interventions to lower the glycemic response to carbohydrate foods with a low-viscosity fiber (resistant maltodextrin): meta-analysis of randomized controlled trials." The American Journal of Clinical Nutrition **89**(1): 114-125.
- Lok, K. Y., Chan, R., Chan, D., Li, L., Leung, G., Woo, J., Lightowler, H.J., & C.J.K., Henry (2010). "Glycaemic index and glycaemic load values of a selection of popular foods consumed in Hong Kong. ." British Journal of Nutrition **103**: 556-560
- Lopez, H. W., M.-A. Levrat-Verny, et al. (2001). "Class 2 Resistant Starches Lower Plasma and Liver Lipids and Improve Mineral Retention in Rats." The Journal of Nutrition **131**(4): 1283-1289.
- Lu, S., Chen, L. N., & Lii, C. Y. (1997). "Correlations between the fine structure,

- physicochemical properties, and retrogradation of amylopectins from Taiwan rice varieties." Cereal Chemistry **74**: 34-39.
- Ludwig, D. S. (2002). "The Glycemic Index." JAMA: The Journal of the American Medical Association **287**(18): 2414-2423.
- Ludwig, D. S., M. A. Pereira, et al. (1999). "Dietary Fiber, Weight Gain, and Cardiovascular Disease Risk Factors in Young Adults." JAMA: The Journal of the American Medical Association **282**(16): 1539-1546.
- Ma, Y., B. Olendzki, et al. (2005). "Association between Dietary Carbohydrates and Body Weight." American Journal of Epidemiology **161**(4): 359-367.
- Madar, Z. and I. Shomer (1990). "Polysaccharide composition of a gel fraction derived from fenugreek and its effect on starch digestion and bile acid absorption in rats." Journal of Agricultural and Food Chemistry **38**(7): 1535-1539.
- Magaletta, R. L., DiCataldo, S. N., Liu, D., Li, H. L., Borwankar, R. P., & Martini, M. C (2010). "In vitro method for predicting glycemic index of foods using simulated digestion and an artificial neural network." Cereal Chemistry **87**: 363-369.
- Mahasukhonthachat, K., P. A. Sopade, et al. (2010). "Kinetics of starch digestion in sorghum as affected by particle size." Journal of Food Engineering **96**(1): 18-28.
- Makelainen, H., H. Anttila, et al. (2006). "The effect of [beta]-glucan on the glycemic and insulin index." Eur J Clin Nutr **61**(6): 779-785.
- Manners, D. J. (1979). The enzymic degradation of starches. Polysaccharides in Food. London, Butterworths.
- Meng, F., Y. Wei, et al. (2005). "Iron content and bioavailability in rice." Journal of Trace

Elements in Medicine and Biology **18**(4): 333-338.

Meyer, K. A., L. H. Kushi, et al. (2000). "Carbohydrates, dietary fiber, and incident type 2 diabetes in older women." The American Journal of Clinical Nutrition **71**(4): 921-930.

Miller, J. B., E. Pang, et al. (1992). "Rice: a high or low glycemic index food?" The American Journal of Clinical Nutrition **56**(6): 1034-1036.

Mishra, S., J. Monro, et al. (2008). "Effect of Processing on Slowly Digestible Starch and Resistant Starch in Potato." Starch - Stärke **60**(9): 500-507.

Mishra, S. and J. A. Monro (2009). "Digestibility of starch fractions in wholegrain rolled oats." Journal of Cereal Science **50**(1): 61-66.

Mohan, V., G. Radhika, et al. (2009). "Dietary carbohydrates, glycaemic load, food groups and newly detected type 2 diabetes among urban Asian Indian population in Chennai, India (Chennai Urban Rural Epidemiology Study 59)." Br J Nutr **102**(10): 1498-506.

Monro, J. A., S. Mishra, et al. (2010). "Baselines representing blood glucose clearance improve in vitro prediction of the glycaemic impact of customarily consumed food quantities." British Journal of Nutrition **103**(02): 295-305.

Monro, J. A. and M. Shaw (2008). "Glycemic impact, glycemic glucose equivalents, glycemic index, and glycemic load: definitions, distinctions, and implications." The American Journal of Clinical Nutrition **87**(1): 237S-243S.

Muir, J. G., A. Birkett, et al. (1995). "Food processing and maize variety affects amounts of starch escaping digestion in the small intestine." The American Journal of Clinical Nutrition **61**(1): 82-89.

Murakami, K., S. Sasaki, et al. (2006). "Dietary glycemic index and load in relation to metabolic

- risk factors in Japanese female farmers with traditional dietary habits." The American Journal of Clinical Nutrition **83**(5): 1161-1169.
- Nakashima, M., M. Sakurai, et al. (2010). "Dietary Glycemic Index, Glycemic Load and Blood Lipid Levels in Middle-Aged Japanese Men and Women." Journal of Atherosclerosis and Thrombosis **17**(10): 1082-1095.
- Nanri, A., T. Mizoue, et al. (2010). "Rice intake and type 2 diabetes in Japanese men and women: the Japan Public Health Center-based Prospective Study." The American Journal of Clinical Nutrition **92**(6): 1468-1477.
- Nanri, A., T. Mizoue, et al. (2008). "Dietary Patterns and A1C in Japanese Men and Women." Diabetes Care **31**(8): 1568-1573.
- Niba, L. L. (2003a). "Processing effects on susceptibility of starch to digestion in some dietary starch sources." International Journal of Food Sciences and Nutrition **54**(1): 97-109.
- Niba, L. L., Niba, S.N. (2003b). "Role of non-digestible carbohydrates in colon cancer protection." Nutrition & Food Science **33**: 28 - 33.
- Nugent, A. P. (2005). "Health properties of resistant starch." Nutrition Bulletin **30**(1): 27-54.
- Nuttall, F. Q., A. D. Mooradian, et al. (1984). "Effect of Protein Ingestion on the Glucose and Insulin Response to a Standardized Oral Glucose Load." Diabetes Care **7**(5): 465-470.
- Oates, C. G. (1997). "Towards an understanding of starch granule structure and hydrolysis." Trends in Food Science & Technology **8**(11): 375-382.
- O'Dea, K., P. Snow, et al. (1981). "Rate of starch hydrolysis in vitro as a predictor of metabolic responses to complex carbohydrate in vivo." The American Journal of Clinical Nutrition **34**(10): 1991-1993.

- O'Dea, K. N., P. J.; & Antonoff, L. (1980). "Physical factors influencing postprandial glucose and insulin responses to starch." The American Journal of Clinical Nutrition **33** 760-765
- Ong, M. H., & Blanshard, J. M. V. (1995). "Texture determinants of cooked, parboiled rice. I. Rice starch amylose and the fine structure of amylopectin." Journal of Cereal Science **21**: 251-260.
- Orford, P. D., S. G. Ring, et al. (1987). "The effect of concentration and botanical source on the gelation and retrogradation of starch." Journal of the Science of Food and Agriculture **39**(2): 169-177.
- Ostman, E. M., Liljeberg, H., Björck, I. (2001). "Inconsistency between glycemic and insulinemic responses to regular and fermented milk products." American Journal of Clinical Nutrition **74**: 96-100.
- Pal, S., Lim, S., Egger, G.. (2008). "The effect of a low glycaemic index breakfast on blood glucose, insulin, lipid profiles, blood pressure, body weight, body composition and satiety in obese and overweight individuals: a pilot study." Journal of the American College of Nutrition **27**: 387-93.
- Panahi, S., A. Ezatagha, et al. (2007). " β -Glucan from Two Sources of Oat Concentrates Affect Postprandial Glycemia in Relation to the Level of Viscosity." Journal of the American College of Nutrition **26**(6): 639-644.
- Panlasigui, L. N. and L. U. Thompson (2006). "Blood glucose lowering effects of brown rice in normal and diabetic subjects." International Journal of Food Sciences and Nutrition **57**(3-4): 151-158.
- Panlasigui, L. N., L. U. Thompson, et al. (1992). "Extruded rice noodles: Starch digestibility and

glycemic response of healthy and diabetic subjects with different habitual diets."

Nutrition Research **12**(10): 1195-1204.

Panlasigui, L. N., L. U. Thompson, et al. (1991). "Rice varieties with similar amylose content differ in starch digestibility and glycemic response in humans." The American Journal of Clinical Nutrition **54**(5): 871-877.

Parastouei, K., Shahaboddin, M.E., Motalebi, M., Mirhashemi, S.M., Faraji, A.M., & Seyyedi, F. (2011). "Glycemic index of Iranian rice." Scientific Research and Essays **6**: 5302-5307.

Park, E. Y., B.-K. Baik, et al. (2009). "Influences of temperature-cycled storage on retrogradation and in vitro digestibility of waxy maize starch gel." Journal of Cereal Science **50**(1): 43-48.

Park, J. K., Kim, S.S., & Kim, K.O. (2001). "Effect of milling ratio on sensory properties of cooked rice and on physicochemical properties of milled and cooked rice." Cereal Chemistry **78**: 151-156.

Pathiraje, P., W. M. T. Madhujith, et al. (2010). The Effect of Rice Variety and Parboiling on in vivo Glycemic Response.

Philpot, K., M. Martin, et al. (2006). "Environmental Factors that Affect the Ability of Amylose to Contribute to Retrogradation in Gels Made from Rice Flour." Journal of Agricultural and Food Chemistry **54**(14): 5182-5190.

Pirasath, S., Thayaanathan, K., Balakumar, S. & Arasaratnam, V. (2010). "Effect of dietary curries on the glycaemic index." Ceylon Medical Journal 2010; **55**: 118-122.

Politz, M. L., Timpa, J.D., & Wasserman, B.P. (1994). "Quantitative Measurement of Extrusion-Induced Starch Fragmentation Products in Maize Flour Using Nonaqueous Automated

- Gel-Permeation Chromatography." Cereal Chemistry **71**: 532-536.
- Qureshi, A. A., S. A. Sami, et al. (2002). "Effects of stabilized rice bran, its soluble and fiber fractions on blood glucose levels and serum lipid parameters in humans with diabetes mellitus Types I and II." The Journal of Nutritional Biochemistry **13**(3): 175-187.
- Rahman, M., M. A. Malik, et al. (1992). "Glycaemic index of Pakistani staple foods in mixed meals for diabetics." J Pak Med Assoc **42**(3): 60-2.
- Ranawana, D. V., C. J. K. Henry, et al. (2009). "Glycaemic index of some commercially available rice and rice products in Great Britain." International Journal of Food Sciences and Nutrition **60**(s4): 99-110.
- Ranawana, V., M. E. Clegg, et al. (2011). "Postmastication digestion factors influence glycemic variability in humans." Nutrition Research **31**(6): 452-459.
- Ranawana, V., C. J. K. Henry, et al. (2010b). "Degree of habitual mastication seems to contribute to interindividual variations in the glycemic response to rice but not to spaghetti." Nutrition Research **30**(6): 382-391.
- Ranawana, V., J. A. Monro, et al. (2010b). "Degree of particle size breakdown during mastication may be a possible cause of interindividual glycemic variability." Nutrition Research **30**(4): 246-254.
- Rani, M. R. S. and K. R. Bhattacharya (1995). "Microscopic of Rice Starch Granules During Cooking." Starch - Stärke **47**(9): 334-337.
- Ray, K. and P. Singhanian (2011). "Glycemic and insulinemic responses to carbohydrate rich whole foods." Journal of Food Science and Technology: 1-6.
- Regand, A., Z. Chowdhury, et al. (2011). "The molecular weight, solubility and viscosity of oat

- beta-glucan affect human glyceic response by modifying starch digestibility." Food Chemistry **129**(2): 297-304.
- Resurreccion, A. P., Li, X., Okita, T. W., & Juliano, B. O. (1993). "Characterisation of poorly digested protein of cooked rice protein bodies." Cereal Chemistry **70**: 101-104.
- Resurreccion, A. P., B. O. Juliano, et al. (1979). "Nutrient content and distribution in milling fractions of rice grain." Journal of the Science of Food and Agriculture **30**(5): 475-481.
- Robert, D. S., Ismail, A. Al-Safi., & Wan Nikk., W.S. (2009). Determination of the glycaemic index of selected Malaysian foods. Kelantan, Malaysia, University of Science Malaysia, Department of Dietetics, School of Health Sciences.
- Sajilata, M. G., R. S. Singhal, et al. (2006). "Resistant Starch—A Review." Comprehensive Reviews in Food Science and Food Safety **5**(1): 1-17.
- Sakuma, M., Yamanaka-Okumura, H., Naniwa, Y., Matsumoto, D., Tsunematsu, M., Yamamoto, H., Taketani, Y., & Takeda, E. (2009). "Dose-dependent effects of barley cooked with white rice on postprandial glucose and desacyl ghrelin levels." Journal of Clinical Biochemistry and Nutrition **44**: 151-159.
- Salmeron, J., A. Ascherio, et al. (1997). "Dietary Fiber, Glycemic Load, and Risk of NIDDM in Men." Diabetes Care **20**(4): 545-550.
- Salmeron, J., J. E. Manson, et al. (1997a). "Dietary Fiber, Glycemic Load, and Risk of Noninsulin-dependent Diabetes Mellitus in Women." JAMA: The Journal of the American Medical Association **277**(6): 472-477.
- Sano, Y. (1984). "Differential regulation of waxy gene expression in rice endosperm." Theoretical and Applied Genetics, **68**: 467-473.

- Sasaki, T., K. Kohyama, et al. (2009). "Physicochemical characteristics of waxy rice starch influencing the in vitro digestibility of a starch gel." Food Chemistry **116**(1): 137-142.
- Sato, S., K. Fukumura, et al. (2010). "Glycemic Index and Glucose Utilization of Rice Vermicelli in Healthy Subjects." Biological and Pharmaceutical Bulletin **33**(8): 1385-1393.
- Schulze, M. B., S. Liu, et al. (2004). "Glycemic index, glycemic load, and dietary fiber intake and incidence of type 2 diabetes in younger and middle-aged women." The American Journal of Clinical Nutrition **80**(2): 348-356.
- Schulze, M. B., M. Schulz, et al. (2007). "Fiber and Magnesium Intake and Incidence of Type 2 Diabetes: A Prospective Study and Meta-analysis." Arch Intern Med **167**(9): 956-965.
- Seki, T., R. Nagase, et al. (2005). "Insoluble Fiber Is a Major Constituent Responsible for Lowering the Post-Prandial Blood Glucose Concentration in the Pre-Germinated Brown Rice." Biological and Pharmaceutical Bulletin **28**(8): 1539-1541.
- Shanita, S. N., Hasnah, H. & Khoo, C.W. (2011). "Amylose and amylopectin in selected Malaysian foods and its relationship to glycemic index." Sains Malaysiana **40**: 865-870.
- Sharavathy, M. K., A. Urooj, et al. (2001). "Nutritionally important starch fractions in cereal based Indian food preparations." Food Chemistry **75**(2): 241-247.
- Shaw, J. E., R. A. Sicree, et al. (2010). "Global estimates of the prevalence of diabetes for 2010 and 2030." Diabetes Research and Clinical Practice **87**(1): 4-14.
- Shibanuma, K., Takeda, Y., Hizukuri, S., & Shibata, S. (1994). "Molecular structures of some wheat starches." Carbohydrate Polymers **25**: 111-116.
- Shirani, G. and R. Ganesharane (2009). "Extruded products with Fenugreek (Trigonella

- foenum-graecium) chickpea and rice: Physical properties, sensory acceptability and glycaemic index." Journal of Food Engineering **90**(1): 44-52.
- Shobana, S., N. G. Malleshi, et al. (2011). "Nutritional and sensory profile of two Indian rice varieties with different degrees of polishing." Int J Food Sci Nutr **62**(8): 800-10.
- Shu, X., L. Jia, et al. (2009). "Slow Digestion Properties of Rice Different in Resistant Starch." Journal of Agricultural and Food Chemistry **57**(16): 7552-7559.
- Sieri, S., V. Pala, et al. (2007). "Dietary glycemic index, glycemic load, and the risk of breast cancer in an Italian prospective cohort study." The American Journal of Clinical Nutrition **86**(4): 1160-1166.
- Sievert, D., & Pomeranz, Y. (1989). "Enzyme-resistant starch. I. Characterization and evaluation by enzymatic, thermoanalytical, and microscopic methods." Cereal Chemistry **66**: 342-347.
- Silvester, K. R., H. N. Englyst, et al. (1995). "Ileal recovery of starch from whole diets containing resistant starch measured in vitro and fermentation of ileal effluent [published erratum appears in Am J Clin Nutr 1996 Mar;63(3):407]." The American Journal of Clinical Nutrition **62**(2): 403-411.
- Singh, J., A. Dartois, et al. (2010). "Starch digestibility in food matrix: a review." Trends in Food Science & Technology **21**(4): 168-180.
- Singh, J., L. Kaur, et al. (2007). "Factors influencing the physico-chemical, morphological, thermal and rheological properties of some chemically modified starches for food applications—A review." Food Hydrocolloids **21**(1): 1-22.
- Singh, J., O. J. McCarthy, et al. (2008). "Low temperature post-harvest storage of New Zealand

- Taewa (Maori potato): Effects on starch physico-chemical and functional characteristics." Food Chemistry **106**(2): 583-596.
- Slavin, J. L., M. C. Martini, et al. (1999). "Plausible mechanisms for the protectiveness of whole grains." The American Journal of Clinical Nutrition **70**(3): 459S-463S.
- Snow, P. and K. O'Dea (1981). "Factors affecting the rate of hydrolysis of starch in food." The American Journal of Clinical Nutrition **34**(12): 2721-2727.
- Sopade, P. A. and M. J. Gidley (2009). "A Rapid In-vitro Digestibility Assay Based on Glucometry for Investigating Kinetics of Starch Digestion." Starch - Stärke **61**(5): 245-255.
- Srikaeo, K., J. E. Furst, et al. (2005). "Wheat grain cooking process as investigated by modulated temperature differential scanning calorimetry." Carbohydrate Polymers **61**(2): 203-210.
- Srikaeo, K. and P. A. Sopade (2010). "Functional properties and starch digestibility of instant Jasmine rice porridges." Carbohydrate Polymers **82**(3): 952-957.
- Srinivasan, K. (2008). "Reason to season: spices as functional food adjuncts with multiple health effects." Indian Food Industry **27** 36-46.
- Sugiyama, M., A. C. Tang, et al. (2003). "Glycemic index of single and mixed meal foods among common Japanese foods with white rice as a reference food." European Journal of Clinical Nutrition **57**(6): 743-752.
- Sun, Q., D. Spiegelman, et al. (2010). "White rice, brown rice, and risk of type 2 diabetes in US men and women." Arch Intern Med **170**(11): 961-9.
- Taira, H., M. Nakagahra, et al. (1988). "Fatty acid composition of Indica, Sinica, Javanica, Japonica groups of nonglutinous brown rice." Journal of Agricultural and Food

Chemistry **36**(1): 45-47.

Takeda, Y., Hizukuri, S., & Juliano, B. (1987). "Structures of rice amylopectins with low and high affinities for iodine." Carbohydrate Research **168**: 79-88.

Takeda, Y., S. Shibahara, et al. (2003). "Examination of the structure of amylopectin molecules by fluorescent labeling." Carbohydrate Research **338**(5): 471-475.

Takeda, Y., S. Tomooka, et al. (1993). "Structures of branched and linear molecules of rice amylose." Carbohydrate Research **246**(1): 267-272.

Taki, Y., H. Hashizume, et al. (2010). "Breakfast Staple Types Affect Brain Gray Matter Volume and Cognitive Function in Healthy Children." PLoS ONE **5**(12): e15213.

Taniguchi-Fukatsu, A., Yamanaka-Okumura, H., Naniwa-Kuroki, Y., Nishida, Y., Yamamoto H., Taketani, Y., & Takeda, E. (2011). "Natto and viscous vegetables in a Japanese-style breakfast improved insulin sensitivity, lipid metabolism and oxidative stress in overweight subjects with impaired glucose tolerance." British Journal of Nutrition: 1-8.

Tester, R. F., X. Qi, et al. (2006). "Hydrolysis of native starches with amylases." Animal Feed Science and Technology **130**(1-2): 39-54.

Theander, O. and E. Westerlund (1987). "Studies on Chemical Modifications in Heat-processed Starch and Wheat Flour." Starch - Stärke **39**(3): 88-93.

Thompson, L. U., C. L. Button, et al. (1987). "Phytic acid and calcium affect the in vitro rate of navy bean starch digestion and blood glucose response in humans." The American Journal of Clinical Nutrition **46**(3): 467-73.

Thondre, P. S., J. A. Monro, et al. (2010). "High molecular weight barley β -glucan decreases particle breakdown in chapattis (Indian flat breads) during in vitro digestion." Food

- Research International **43**(5): 1476-1481.
- Thorburn, A. W., J. C. Brand, et al. (1986). "Salt and the glycaemic response." Br Med J (Clin Res Ed) **292**(6537): 1697-9.
- Tran, T. T. B., K. J. Shelat, et al. (2011). "Milling of Rice Grains. The Degradation on Three Structural Levels of Starch in Rice Flour Can Be Independently Controlled during Grinding." Journal of Agricultural and Food Chemistry **59**(8): 3964-3973.
- Trinidad, T. P., A. C. Mallillin, et al. (2010). "Glycemic index of commonly consumed carbohydrate foods in the Philippines." Journal of Functional Foods **2**(4): 271-274.
- Tuntipopipat, S., C. Zeder, et al. (2009). "Inhibitory effects of spices and herbs on iron availability." International Journal of Food Sciences and Nutrition **60**(s1): 43-55.
- Urooj, A. and S. Puttraj (1999). "Digestibility index and factors affecting rate of starch digestion in vitro in conventional food preparation." Food / Nahrung **43**(1): 42-47.
- Vandeputte, G. E., Vermeulen, R., Geeroms, J., & Delcour, J. A. (2003a). "Rice starches. I. Structural aspects provide insight into crystallinity characteristics and gelatinisation behaviour of granular starch." Journal of Cereal Science **38**: 43-52.
- Vandeputte, G. E., Derycke, V., Geeroms, J., & Delcour, J. A. (2003b). "Rice starches. II. Structural aspects provide insight into swelling and pasting properties." Journal of Cereal Science **38**: 43-52.
- Venn, B. J. and T. J. Green (2007). "Glycemic index and glycemic load: measurement issues and their effect on diet-disease relationships." Eur J Clin Nutr **61**(S1): S122-S131.
- Villegas, R., S. Liu, et al. (2007). "Prospective Study of Dietary Carbohydrates, Glycemic Index, Glycemic Load, and Incidence of Type 2 Diabetes Mellitus in Middle-aged Chinese

- Women." Arch Intern Med **167**(21): 2310-2316.
- Vonk, R. J., R. E. Hagedoorn, et al. (2000). "Digestion of so-called resistant starch sources in the human small intestine." The American Journal of Clinical Nutrition **72**(2): 432-438.
- Wachters-Hagedoorn, R. E., M. G. Priebe, et al. (2006). "The Rate of Intestinal Glucose Absorption Is Correlated with Plasma Glucose-Dependent Insulinotropic Polypeptide Concentrations in Healthy Men." The Journal of Nutrition **136**(6): 1511-1516.
- Wang, L., B. Xie, et al. (2011). "Study on the granular characteristics of starches separated from Chinese rice cultivars." Carbohydrate Polymers **87**(2): 1038-1044.
- Welch, I. M. L., Bruce, C., Hill, S.E., & Read, N.W. (1987). "Duodenal and ileal lipid suppresses postprandial blood glucose and insulin responses in man: possible implications for the dietary management of diabetes mellitus." Am J Clin Nutr **43**: 167-172.
- Weurding, R. E., A. Veldman, et al. (2001). "In Vitro Starch Digestion Correlates Well with Rate and Extent of Starch Digestion in Broiler Chickens." The Journal of Nutrition **131**(9): 2336-2342.
- Widanagamage, R. D., S. Ekanayake, et al. (2009). "Carbohydrate-rich foods: glycaemic indices and the effect of constituent macronutrients." International Journal of Food Sciences and Nutrition **60**(s4): 215-223.
- Wild, S. H., G. Roglic, et al. (2004). "Global Prevalence of Diabetes: Estimates for the Year 2000 and Projections for 2030." Diabetes Care **27**(10): 2569.
- Wolever, T. M., & Bolognesi, C. (1996). "Prediction of glucose and insulin responses of normal subjects after consuming mixed meals varying in energy, protein, fat, carbohydrate and glycemic index." Journal of Nutrition **126**: 2807-2812.

- Wolever, T. M., D. J. Jenkins, et al. (1991). "The glycemic index: methodology and clinical implications." The American Journal of Clinical Nutrition **54**(5): 846-854.
- Wolever, T. M., P. M. Nguyen, et al. (1994). "Determinants of diet glycemic index calculated retrospectively from diet records of 342 individuals with non-insulin-dependent diabetes mellitus." The American Journal of Clinical Nutrition **59**(6): 1265-1269.
- Wolever, T. M. S., & Jenkins, D.J.A. (1986). Effect of fiber and foods on carbohydrate metabolism. In: Spiller G, ed. Handbook of dietary fiber in human nutrition. Boca Raton, CRC Press Inc.
- Wolever, T. M. S. (2006). The glycaemic index A physiological classification of dietary carbohydrate. MA: CABI, Cambridge.
- Wolever, T. M. S., J. C. Brand-Miller, et al. (2008). "Measuring the glycemic index of foods: interlaboratory study." The American Journal of Clinical Nutrition **87**(1): 247S-257S.
- Wolever, T. M. S., D. J. A. Jenkins, et al. (1986). "Comparison of regular and parboiled rices: explanation of discrepancies between reported glycemic responses to rice." Nutrition Research **6**(4): 349-357.
- Wolever, T. M. S., D. J. A. Jenkins, et al. (1990). "Glycemic Index of Foods in Individual Subjects." Diabetes Care **13**(2): 126-132.
- Wolever, T. M. S., H. H. Vorster, et al. (2003). "Determination of the glycaemic index of foods: interlaboratory study." Eur J Clin Nutr **57**(3): 475-482.
- Wong, J. M. W. and D. J. A. Jenkins (2007). "Carbohydrate Digestibility and Metabolic Effects." The Journal of Nutrition **137**(11): 2539S-2546S.
- Wood, P. J., Braaten, J.T., Scott, F.W., Riedel, K.D., Wolynetz, M.S., & Collins, M.W. (1994).

- " Effect of dose and modification of viscous properties of oat gum on plasma glucose and insulin following an oral glucose load." British Journal of Nutrition **72**: 731-743.
- Wood, P. J., M. U. Beer, et al. (2000). "Evaluation of role of concentration and molecular weight of oat β -glucan in determining effect of viscosity on plasma glucose and insulin following an oral glucose load." British Journal of Nutrition **84**(01): 19-23.
- Woolnough, J. W., J. A. Monro, et al. (2008). "Simulating human carbohydrate digestion in vitro: a review of methods and the need for standardisation." International Journal of Food Science & Technology **43**(12): 2245-2256.
- Yong, L. Z., Chan, C. H. & Sopade, P. A. (2008). Changes in digestibility and bioactivity of a model high-protein-low-carbohydrate (HPLC) food as a result of extrusion. In: 41st Australian Institute of Food Science and Technology. Sydney, Australia.
- Yoon, J. H., L. U. Thompson, et al. (1983). "The effect of phytic acid on in vitro rate of starch digestibility and blood glucose response." The American Journal of Clinical Nutrition **38**(6): 835-42.
- Zai, H., M. Kusano, et al. (2009). "Monosodium l-glutamate added to a high-energy, high-protein liquid diet promotes gastric emptying." The American Journal of Clinical Nutrition **89**(1): 431-435.
- Zhang, G., Malik, V.S., Pan, A., Kumar, S., Holmes, M.D., Spiegelman, D., Lin, Xu., & Hu, F.B. (2010). "Substituting brown rice for white rice to lower diabetes risk: A focus-group study in Chinese adults." American Dietetic Association **110**: 1216-1221.
- Zhang, G., Z. Ao, et al. (2008). "Nutritional Property of Endosperm Starches from Maize Mutants: A Parabolic Relationship between Slowly Digestible Starch and Amylopectin

Fine Structure." Journal of Agricultural and Food Chemistry **56**(12): 4686-4694.

Zhang, G. and B. R. Hamaker (2009). "Slowly Digestible Starch: Concept, Mechanism, and Proposed Extended Glycemic Index." Critical Reviews in Food Science and Nutrition **49**(10): 852-867.

Zhou, Z., K. Robards, et al. (2002). "Composition and functional properties of rice." International Journal of Food Science & Technology **37**(8): 849-868.

Zhu, L.-J., Q.-Q. Liu, et al. (2011). "Digestibility and physicochemical properties of rice (*Oryza sativa* L.) flours and starches differing in amylose content." Carbohydrate Polymers **86**(4): 1751-1759.

Appendix A

GI of rice and rice products published (1981 to 2012)

Food	GI	Subjects (Type* and Number)	Reference food and time duration	Reference
Rice, white				
White (<i>Oryza sativa</i>), boiled (India)	69	Type 2, 6	Glucose, 3h	<i>Chaturvedi et al., 1997</i>
Rice, boiled white, type NS (India)	68	Healthy, 6	Wheat chappati, 2h	<i>Dilwari et al., 1981</i>
Rice, boiled white, type NS (Canada)	80	Type 2, 6	Bread, 3h	<i>Jenkins et al., 1983</i>
Rice, boiled white, type NS (Pakistan)	98	Type 2, 22	Wheat chappati, 3h	<i>Rahman et al., 1992</i>
Rice, boiled white, type NS, boiled 13 min (Italy)	102	Healthy, 14	Glucose, 2h	<i>Gatti et al., 1987</i>
Long grain, boiled 5 min (Canada)	58	Type 1, 5; Type 2, 13	Bread, 3h	<i>Wolever et al., 1986</i>
Long grain, white, unconverted, boiled, 15 min (Mahatma Brand; Australia)	50	Healthy, 6	Glucose, 2h	<i>Brand et al., 1985</i>
Gem long grain (Dainty Food Inc, Toronto, Canada)	79	Type 2, 10	Bread, 3h	<i>Wolever et al., 1985</i>
Gem long grain (Dainty Food Inc, Toronto, Canada)	82	Type 1, 6	Bread, 3h	<i>Wolever et al., 1985</i>
Long grain, white (Uncle Bens, Auckland, New Zealand)	56	Healthy, 14	Glucose, 2h	<i>Giacco et al., 2000</i>
Long grain, boiled 15 min	83	Type 1, 5; Type 2, 13	Bread, 3h	<i>Wolever et al., 1986</i>
Long grain, boiled 25 min (Surinam)	56	Type 2,3	Glucose, 3h	<i>Liu et al., 2000</i>
Gem long grain (Dainty Food Inc, Toronto, Canada)	86	Type 1, 6	Bread, 3h	<i>Gavin, 2001</i>
Long grain rice (Purchased in Oxford,	47	Healthy, 14	Glucose, 2h	<i>Ranawana et al., 2009</i>

UK)

BR 16, 28% high-amylose (Bangladesh)	53	Type 2, 12	Bread, 3h	<i>Larsen et al., 1996</i>
BR 16, white long grain, 27% high-amylose, boiled 17.5min (Bangladesh)	55	Type 2, 9	Bread, 3h	<i>Larsen et al., 2000</i>
Milagrosa, high-amylose, 26.9% amylose, (Philippines, Aklan Province)	68		<i>In vitro</i>	<i>Frei et al., 2003</i>
Manumbaeay, high-amylose, 29.9% amylose, (Philippines, Aklan Province)	87		<i>In vitro</i>	<i>Frei et al., 2003</i>
Kutisyam, intermediate-amylose, 18.7% amylose, (Philippines, Aklan Province)	69		<i>In vitro</i>	<i>Frei et al., 2003</i>
Kinaures, low-amylose, 9.8% amylose (Philippines, Aklan Province)	97		<i>In vitro</i>	<i>Frei et al., 2003</i>
Bagoean, waxy cultivar, 0% amylose, (Philippines, Aklan Province)	92		<i>In vitro</i>	<i>Frei et al., 2003</i>
Karaya, waxy cultivar, 0% amylose, (Philippines, Aklan Province)	109		<i>In vitro</i>	<i>Frei et al., 2003</i>
ZF201, indica milled rice, 26.8% high-amylose, (China)	63		<i>In vitro</i>	<i>Hu et al., 2004</i>
Jiayu293, indica milled rice, 21.3% intermediate-amylose, (China)	79		<i>In vitro</i>	<i>Hu et al., 2004</i>
Zhefu504, indica milled rice, 13.1% low-	99		<i>In vitro</i>	<i>Hu et al., 2004</i>

amylose (China) Yunuo No. 1, indica milled rice, 1.1% waxy, (China)	106			<i>In vitro</i>	<i>Hu et al., 2004</i>
JIN3, japonica milled rice, 25.8% high- amylose (China)	78			<i>In vitro</i>	<i>Hu et al., 2004</i>
Xiushui 11, japonica milled rice, 20.1% intermediate-amylose (China)	69			<i>In vitro</i>	<i>Hu et al., 2004</i>
JIN1, japonica milled rice, low-amylose, 13.8% (China)	89			<i>In vitro</i>	<i>Hu et al., 2004</i>
Shaonuo, indica milled rice, waxy, 0%, (China)	102			<i>In vitro</i>	<i>Hu et al., 2004</i>
Ilyou3027, hybrid milled rice, high- amylose, 26.5% (China)	78			<i>In vitro</i>	<i>Hu et al., 2004</i>
Xieyou46, hybrid milled rice, 21.6% intermediate-amylose (China)	63			<i>In vitro</i>	<i>Hu et al., 2004</i>
Fenyouxiangzan, hybrid milled rice, 14.3% low- amylose (China)	92			<i>In vitro</i>	<i>Hu et al., 2004</i>
Zanuo, hybrid milled rice, 0.7%, waxy, (China)	100			<i>In vitro</i>	<i>Hu et al., 2004</i>
Red rice (Thailand)	76	Healthy, 14	Glucose, 2h		<i>Ranawana et al., 2009</i>
Long grain rice, Indica rice (Thailand)	60	Healthy, 15	Glucose, 2h		<i>Sato et al., 2010</i>
White rice, (Golestan company),boiled 10 min (Iran)	55	Healthy, 10	Glucose, 2h		<i>Parastouei et al., 2011</i>
White rice, (Golestan company), boiled 5- 8min, then simmered 30 min (Iran)	66	Healthy, 10	Glucose, 2h		<i>Parastouei et al., 2011</i>
Sona Masuri rice (India)	72	Healthy, N.A.	Glucose, 2h		<i>Shobana et al., 2012</i>

Ponni rice (India)	70	Healthy, N.A.	Glucose, 2h	<i>Shobana et al., 2012</i>
Surti Kolam (India)	77	Healthy, N.A.	Glucose, 2h	<i>Shobana et al., 2012</i>
White rice, Japonica, normal grain (Koshihikari, Japan)	89	Healthy, 15	Glucose, 2h	<i>Sato et al., 2010</i>
Koshihikari, low-amylose, 17% (Japan)	80	Healthy, subgroups of 9-11 (n=58)	Glucose, 2h	<i>Sugiyama et al.2003</i>
Bg 300 unparboiled rice, high amylose 27.1% (Sri Lanka)	61	Healthy, 10	Glucose, 2h	<i>Pathiraje et al., 2010</i>
Bg 352 unparboiled rice, high amylose 27.0% (Sri Lanka)	67	Healthy, 10	Glucose, 2h	<i>Pathiraje et al., 2010</i>
Bg 358 unparboiled rice, high amylose 29.0% (Sri Lanka)	67	Healthy, 10	Glucose, 2h	<i>Pathiraje et al., 2010</i>
Bg 406 unparboiled rice, high amylose 24.5% (Sri Lanka)	7	Healthy, 10	Glucose, 2h	<i>Pathiraje et al., 2010</i>
LD 356, unparboiled rice, intermediate-amylose 21.5% (Sri Lanka)	70	Healthy, 10	Glucose, 2h	<i>Pathiraje et al., 2010</i>
Rathkaral unparboiled, high amylose 29.5% (Sri Lanka)	60	Healthy, 10	Glucose, 2h	<i>Pathiraje et al., 2010</i>
Wedaheenati unparboiled, high amylose 29.0% (Sri Lanka)	57	Healthy, 10	Glucose, 2h	<i>Pathiraje et al., 2010</i>
Heendikwel unparboiled, high amylose 27.7% (Sri Lanka)	62	Healthy, 10	Glucose, 2h	<i>Pathiraje et al., 2010</i>
Red rice (Sri Lanka)	99	<i>In vitro</i>		<i>Hettiaratchi et al., 2009</i>
Commercial rice varieties (Sri Lanka)	55-73 (majority medium)	Healthy, 12	White bread, 2h	<i>Hettiaratchi et al., 2010</i>

	GI			
Rice, brown				
Brown rice (Canada)	66	Healthy, 7	Glucose, 2h	<i>NHMRC, 1999</i>
Brown, steamed (USA)	50	Healthy, 8	Glucose, 3h (AUC calculated over 3h for 5 time points only)	<i>Potter et al., 1981</i>
Brown (<i>Oryza sativa</i>), boiled (South India)	50	Healthy, 12-15	Glucose, 3h (AUC calculated over 3h for 5 time points only)	<i>Kurup et al., 1992</i>
Calrose brown (Rice Growers Co-op, Australia)	124	Healthy, 8	Bread, 2h	<i>Miller et al., 1992</i>
Doongara brown, high- amylose (Rice Growers Co-op, Australia)	94	Healthy, 8	Bread, 2h	<i>Miller et al., 1992</i>
Pelde brown (Rice Growers Co-op, Australia)	109	Healthy, 8	Bread, 2h	<i>Miller et al., 1992</i>
Sunbrown Quick (Rice Growers Co-op, Australia)	114	Healthy, 8	Bread, 2h	<i>Miller et al., 1992</i>
Tai Ken, brown rice, (Union Rice Company; Taipei, Taiwan)	82	Healthy, 10	Bread, 2h	<i>Lin et al., 2010</i>
Rice, basmati				
Basmati, white, boiled (Mahatma brand, Sydney, Australia)	83	Healthy, 9	Bread, 2h	<i>Holt & Brand Miller, 1995</i>
Brown basmati rice	75	Healthy, 14	Glucose, 2h	<i>Ranawana et al., 2009</i>
White and brown basmati rice (mixture 60% white basmati, 40% brown basmati)	59	Healthy, 14	Glucose, 2h	<i>Ranawana et al., 2009</i>
Basmati with wild rice (83% easy-cook)	63	Healthy, 14	Glucose, 2h	<i>Ranawana et al., 2009</i>

basmati) and 17%
North American Wild
rice

Rice, parboiled

Long grain, boiled 10 min (USA)	61	Type 2, 8	Glucose, 3h (GI calculated from the AUC of glucose)	<i>Miller et al., 2001</i>
Long grain, boiled 12 min (Denmark)	60	Type 2, 7	Bread, 2h	<i>Rasmussen et al, 1992</i>
Long grain, boiled 5 min (Canada)	54	Type 1, 5 Type 2, 13	Bread, 3h	<i>Wolever et al., 1986</i>
Long grain, boiled 15 min (Canada)	67	Type 1, 5 Type 2, 13	Bread, 3h	<i>Wolever et al., 1986</i>
Long-grain white, parboiled (Uncle Ben's; Masterfoods, Brucargo-Zaventem, Belgium)	54	Healthy, 13	Glucose, 2h	<i>Al-Mssallem et al, 2011</i>
Hassawi rice (Al-Hassa, Saudi Arabia)	59	Healthy, 13	Glucose, 2h	<i>Al-Mssallem et al, 2011</i>
Bg 352 parboiled rice, amylose content not determined (Sri Lanka)	60	Healthy, 10	Glucose, 2h	<i>Pathiraje et al., 2010</i>
Bg 358 parboiled rice, amylose content not determined (Sri Lanka)	62	Healthy, 10	Glucose, 2h	<i>Pathiraje et al., 2010</i>
Bg 356 parboiled rice, amylose content not determined (Sri Lanka)	64	Healthy, 10	Glucose, 2h	<i>Pathiraje et al., 2010</i>
Bg 406 parboiled rice, amylose content not determined (Sri Lanka)	71	Healthy, 10	Glucose, 2h	<i>Pathiraje et al., 2010</i>
Rice, glutinous				
Glutinous rice (Thailand)	94	Healthy, 12	Glucose, 2h	<i>Chan et al., 2001</i>
Glutinous rice (Thailand)	92	Healthy, 14	Glucose, 2h	<i>Ranawana et al., 2009</i>
Glutinous rice (Japan)	105	Healthy, subgroups of 9-11 (n=58)	White rice, 2h	<i>Sugiyama et al., 2003</i>

Rice noodles

Rice vermicelli, Kongmoon (China)	83	Type 1 and 2, 9	Bread ,3h	<i>Jenkins et al., 1981</i>
Jianxi rice vermicelli (rice, water), boiled 8 min (Hong Kong)	55	Healthy, 10	Glucose,2h	<i>Lok et al., 2010</i>
Taiwan vermicelli (rice, water, maize starch), boiled 2 min (Hong Kong)	68	Healthy, 10	Glucose,2h	<i>Lok et al., 2010</i>
Vermicelli, Produced from 100% long grain rice	55	Healthy, 15	Glucose, 2h	<i>Sato et al., 2010</i>
Vermicelli, Produced from 99% long grain rice+1% calcium	50	Healthy, 15	Glucose, 2h	<i>Sato et al., 2010</i>
Bihon (Philippines)	49	Healthy, 10	Glucose,2h	<i>Trinidad et al., 2010</i>
Guilin rice vermicelli (purchased from Oxford, UK), cooked 8 min	37	Healthy, 14	Glucose, 2h	<i>Ranawana et al., 2009</i>
Jiangxi rice vermicelli (purchased from Oxford, UK), cooked 8 min	40	Healthy, 14	Glucose, 2h	<i>Ranawana et al., 2009</i>
Rice noodles, dried, boiled (Thai World, Bangkok, Thailand)	61	Healthy, 12	Glucose,2h	<i>Chan et al., 2001</i>
Rice noodles, freshly made, boiled (Australia)	40	Healthy, 12	Glucose,2h	<i>Chan et al., 2001</i>
Instant rice vermicelli (made from Thai long grain rice, tapioca starch, potato starch)	59	Healthy, 15	Glucose, 2h	<i>Sato et al., 2010</i>
Kway teow (made from Thai long grain rice and potato starch)	60	Healthy, 15	Glucose, 2h	<i>Sato et al., 2010</i>
Pho (made from Thai long grain rice and tapioca starch)	62	Healthy, 15	Glucose, 2h	<i>Sato et al., 2010</i>

Rice, speciality rices

Cajun Style (Uncle Ben's; Effem Foods Ltd, Canada)	51	Type 1 and 2, 9	Bread ,3h	<i>Jenkins et al., 1981</i>
Garden Style (Uncle Ben's; Effem Foods Ltd, Canada)	55	Type 1 and 2, 9	Bread ,3h	<i>Jenkins et al., 1981</i>
Long grain and Wild (Uncle Ben's; Effem Foods Ltd, Canada)	54	Type 1 and 2, 9	Bread ,3h	<i>Jenkins et al., 1981</i>
Mexican Fast and Fancy (Uncle Ben's; Effem Foods Ltd, Canada)	58	Type 1 and 2, 9	Bread ,3h	<i>Jenkins et al., 1981</i>
Saskatchewan wild rice (Canada)	57	Type 1 and 2, 9	Bread ,3h	<i>Jenkins et al., 1981</i>
Easy-cook Basmati rice	80	Healthy, 14	Glucose, 2h	<i>Ranawana et al., 2009</i>
Easy-cook long grain rice	47	Healthy, 14	Glucose, 2h	<i>Ranawana et al., 2009</i>
Broken rice	86	Healthy, 12	Glucose,2h	<i>Chan et al., 2001</i>
Instant rice, white, boiled 1 min (Canada)	65	Type 1, 5 Type 2, 13	Bread, 3h	<i>Wolever et al., 1986</i>
Instant rice, white, cooked 6 min (Trice Brand, Australia)	87	Healthy, 6	Glucose, 2h	<i>Brand et al., 1985</i>
Instant Doongara, white, cooked 5 min (Rice Growers Co-op, Australia)	132	Healthy, 9	Bread, 2h	<i>Holt & Brand-Miller, 1995</i>
Other rice products				
Rice Bubbles (Puffed rice) (Kellogg's Australia)	116	Healthy, 7	Bread, 3h	<i>Holt et al., 1992</i>
Rice Bubbles (Puffed rice) (Kellogg's Australia)	95	Healthy, 6	Glucose, 2h	<i>Brand et al., 1985</i>
Puffed rice cakes, white (Rice Growers Co-op, Australia)	117	Healthy, 6	Bread, 2h	<i>Miller et al., 1992</i>
Puffed rice cakes, Calrose rice, low amylose (Rice Growers Co-op, Australia)	128	Healthy, 9	Bread, 2h	<i>Holt & Brand-Miller, 1995</i>
Puffed rice cakes,	85	Healthy, 9	Bread, 2h	<i>Holt & Brand-Miller,</i>

Doongara rice, high
amylose (Rice Growers
Co-op, Australia)

1995

*GI= Glycaemic Index ; *Type 1= Insulin dependent diabetes mellitus (IDDM) ; Type 2= Non-insulin dependent diabetes mellitus (NIDDM) ; N.A.=Not available*

The glycaemic index of rice mixed meals consumed in Asia

Food	GI	Subjects (Type and Number)*	Reference food and time duration	Classification
<u>Chinese</u>				
Fried rice vermicelli in Singapore-style	54	Healthy, 15	Glucose, 2h	Low GI
Fried rice noodles with sliced beef	66	Healthy, 15	Glucose, 2h	Medium GI
Glutinous rice ball	61	Healthy, 15	Glucose, 2h	Medium GI
Salted meat rice dumpling	69	Healthy, 15	Glucose, 2h	Medium GI
Fried rice in Yangzhou-style	80	Healthy, 15	Glucose, 2h	High GI
Sticky rice wrapped in lotus leaf	83	Healthy, 15	Glucose, 2h	High GI
Steamed glutinous rice roll	89	Healthy, 15	Glucose, 2h	High GI
Plain steamed vermicelli roll	90	Healthy, 15	Glucose, 2h	High GI
Rice with stir-fried vegetables and chicken	73	Healthy, 8	Glucose, 3h	High GI
<u>Japanese</u>				
Sushi, roasted sea algae, vinegar – low amylose content rice (17%)	55	Healthy, subgroups of 9-11 (n=58)	White rice, 2h	Low GI
White rice with milk 100ml (taken together) low-amylose content rice (17%)	59	Healthy, subgroups of 9-11 (n=58)	White rice, 2h	Medium GI
Butter rice low-amylose content (17%)	96	Healthy, subgroups of 9-11 (n=58)	White rice, 2h	High GI
Curry rice low-amylose content (17%)	82	Healthy, subgroups of 9-11	White rice, 2h	High GI

White rice low-amylose content (17%) with curry and cheese	67	(n=58) Healthy, subgroups of 9-11	White rice, 2h	Medium GI
White rice low-amylose content (17%) and salted plum (umeboshi)	98	(n=58) Healthy, subgroups of 9-11	White rice, 2h	High GI
White rice low-amylose content (17%) and dried fish strip	115	(n=58) Healthy, subgroups of 9-11	White rice, 2h	High GI
White rice low-amylose content (17%) rolled in toasted algae	94	(n=58) Healthy, subgroups of 9-11	White rice, 2h	High GI
White rice low-amylose content (17%) with raw egg, with soy sauce	114	(n=58) Healthy, subgroups of 9-11	White rice, 2h	High GI
White rice low-amylose content (17%) and pickled food (sliced cucumber mixed with vinegar)	75	(n=58) Healthy, subgroups of 9-11	White rice, 2h	High GI
White rice and roasted, ground soybean (beihan, kinako) low-amylose content (17%)	68	(n=58) Healthy, subgroups of 9-11	White rice, 2h	Medium GI
White rice and fermented soybean (natto) low-amylose content (17%)	68	(n=58) Healthy, subgroups of 9-11	White rice, 2h	Medium GI
Rice with soybean paste soup (miso shiru) low-amylose content (17%)	74	(n=58) Healthy, subgroups of 9-11	White rice, 2h	High GI
Glutinous rice cake (mochi) low-amylose content (17%)	101	(n=58) Healthy, subgroups of 9-11	White rice, 2h	High GI

Indian, Sri Lankan

Rice with lentil and cauliflower curry	60	Healthy, 8	Glucose,3h	Medium GI
Rice, boiled served with <i>Lagenaria vulgaris</i> (bottle gourd) and (<i>Lycopersicon esculentum</i>) tomato curry	69	Type 2, 6	Glucose, 3h	Medium GI
Parboiled rice with green leaf curry (<i>Amaranthus</i>)	48	Healthy, 20	Glucose, 2h	Low GI
Parboiled rice with gravy (soya meat)	56	Healthy, 20	Glucose, 2h	Medium GI
Parboiled rice with green leaf curry and gravy	55	Healthy, 20	Glucose, 2h	Low GI
Red rice with lentil curry, boiled egg + <i>Centella asiatica</i> (<i>Gotukola</i>) salad and coconut gravy (Kiri hodi)	63	Healthy, 12	White bread,2h	Medium GI
Red rice with lentil curry, boiled egg + <i>Centella asiatica</i> (<i>Gotukola</i>) salad and coconut gravy (Kiri hodi) and <i>Lasia spinosa</i> (<i>kohila</i>) salad	57	Healthy, 12	White bread,2h	Medium GI
Red rice with lentil curry, boiled egg + <i>Centella asiatica</i> (<i>Gotukola</i>) salad, coconut gravy and <i>Trichosanthes cucumerina</i> (<i>snake gourd</i>) salad	61	Healthy, 12	White bread,2h	Medium GI
Red rice with lentil curry- Red rice (82% starch), Lentils curry (18% starch).	60		<i>In vitro</i>	Medium GI

Malaysian

Fried rice	59		<i>In vitro</i>	Medium GI
Nasi lemak	66		<i>In vitro</i>	Medium GI
Fried beehoon	99		<i>In vitro</i>	High GI

Other studies

Lentil-rice meal (70:30) (Chile)	49		<i>In vitro</i>	Low GI
----------------------------------	----	--	-----------------	--------

Sources: Lok et al.; 2009; Pirasath et al.,2010; Chaturvedi et al., 1997; Chew et al., 1988;

Sugiyama et al., 2003; Araya et al., 2003; Hettiaratchi et al., 2009; Shanita et al., 2011; Chen et

al., 2010 ; GI= Glycaemic Index ; *Type 1= Insulin dependent diabetes mellitus (IDDM) ; Type

2= Non-insulin dependent diabetes mellitus (NIDDM)

Table 1: Apparent amylose content type and other properties preferred for various processed rice products

Rice product	Apparent amylose content type and other properties
Parboiled rice	High and intermediate-AC
Precooked/quick cooking rice	Waxy, Low, Intermediate and High-AC
Extruded and flat rice noodles	Mainly aged, high-AC low-GT rice
Rice made into puddings, breads, cakes, beer adjuncts	Low starch GT
Rice wines and rice based sauces, desserts, snacks and sweets	Waxy, low-AC
Rice cakes	Steamed rice cakes (All AC types except High) Bake rice cakes (All AC types)
Rice crackers	Waxy and non waxy
Idli (Indian steamed rice), dosai (Indian rice pancake)	High and intermediate AC (Parboiled rice preferred over raw rice)
Rice based infant food	Intermediate or high

(Source: Juliano, 2003)

Table 2: Yield and composition of defatted and protease-amylase treated cell wall preparations obtained from different histological fractions of milling of brown rice

Rice fraction	Yield (% defatted tissue)	Composition (% of total)				Uronic acid in pectin (%)	Arabinose:xylose ratio	
		Pectic substances	Hemi-cellulose	α -cellulose	Lignin		Pectic substances	Hemi-cellulose
Caryopsis coat	29	7	38	27	32	32	1.63	0.82
Aleurone tissue	20	11	42	16	25	25	1.78	0.84
Germ	12	23	47	9	16	16	2.29	0.96
Endosperm	0.3	27	49	1	34	34	1.09	0.64

(Source: Shibuya, 1989)

Table 3: Cooking time and effect on GI

Test meal	Cooking time (min)	GI
Regular rice	5	58 ± 4
Regular rice	15	83 ± 4
Parboiled rice	5	54 ± 5
Parboiled rice	15	67 ± 5
Parboiled rice	25	66 ± 4

(Source: Wolever et al., 1986)

Figure Legends

Figure 1: World production and consumption of domestic milled rice (1980-1982) (Source: IRRI, http://ricestat.irri.org/vis/wrs_quickCharts.php)

